

# UC Davis

## UC Davis Previously Published Works

### Title

Stratification requirements for seed dormancy alleviation in a wetland weed.

### Permalink

<https://escholarship.org/uc/item/1f2026d9>

### Journal

PloS one, 8(9)

### ISSN

1932-6203

### Authors

Boddy, Louis G

Bradford, Kent J

Fischer, Albert J

### Publication Date

2013

### DOI

10.1371/journal.pone.0071457

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

# Stratification Requirements for Seed Dormancy Alleviation in a Wetland Weed

Louis G. Boddy<sup>1\*</sup>, Kent J. Bradford<sup>2</sup>, Albert J. Fischer<sup>2</sup>

**1** Plant Science Department, Marrone Bio Innovations, Davis, California, United States of America, **2** Department of Plant Sciences, University of California Davis, Davis, California, United States of America

## Abstract

*Echinochloa oryzicola* (syn. *E. phyllopogon*) is an exotic weed of California rice paddies that has evolved resistance to multiple herbicides. Elimination of seedling through certain weed control methods can limit the spread of this weed, but is contingent on accurate predictions of germination and emergence timing, which are influenced by seed dormancy levels. In summer annuals, dormancy can often be relieved through stratification, a period of prolonged exposure to cold and moist conditions. We used population-based threshold models to quantify the effects of stratification on seed germination of four *E. oryzicola* populations at a range of water potential ( $\Psi$ ) and oxygen levels. We also determined how stratification temperatures, moisture levels and durations contributed to dormancy release. Stratification released dormancy by decreasing base  $\Psi$  and hydrotimerequired for germination and by eliminating any germination sensitivity to oxygen. Stratification also increased average germination rates (GR), which were used as a proxy for relative dormancy levels. Alternating temperatures nearly doubled GR in all populations, indicating that seeds could be partially dormant despite achieving high final germination percentages. Stratification at  $\Psi=0$  MPa increased GR compared to stratification at lower water potentials, demonstrating that  $\Psi$  contributed to regulating dormancy release. Maximum GR occurred after 2-4 weeks of stratification at 0 MPa; GR were often more rapid for herbicide-resistant than for herbicide-susceptible seeds, implying greater dormancy in the latter. Manipulation of field conditions to promote dormancy alleviation of *E. oryzicola* seeds might improve the rate and uniformity of germination for seed bank depletion through seedling weed control. Our results suggest field soil saturation in winter would contribute towards *E. oryzicola* dormancy release and decrease the time to seedling emergence.

**Citation:** Boddy LG, Bradford KJ, Fischer AJ (2013) Stratification Requirements for Seed Dormancy Alleviation in a Wetland Weed. PLoS ONE 8(9): e71457. doi:10.1371/journal.pone.0071457

**Editor:** Dorian Q. Fuller, University College London, United Kingdom

**Received:** January 6, 2013; **Accepted:** July 2, 2013; **Published:** September 5, 2013

**Copyright:** © 2013 Boddy et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Funding:** This study was partially funded by the California Rice Research Board. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript. No additional external funding was received for this study.


**Competing Interests:** Some funding for this research was provided by the California Rice Research Board. The affiliation of the first author with Marrone Bio Innovations occurred after all the research in this article was complete; this does not alter the authors' adherence to all the PLOS ONE policies on sharing data and materials.

\* E-mail: lboddy@marronebio.com

## Introduction

The temperate region summer annual weed *Echinochloa oryzicola* Vasing. (syn. *E. phyllopogon* Stapf ex Kossenko) is a morphological mimic of rice (*Oryza sativa*) that can germinate and initiate shoot growth under hypoxia in flooded paddies [1] and causes up to 50% rice yield losses in California if not controlled [2,3]. Decades of heavy reliance on herbicides for *E. oryzicola* control [3] have resulted in the widespread occurrence of populations with simultaneous resistance to most available grass herbicides for selective use in rice [4–7]. Successful control of herbicide-resistant *E. oryzicola* now hinges on maximizing weed seedling recruitment in order to eliminate such seedlings prior to planting the crop [8,9]. The stale seedbed approach entails recruiting and treating weeds prior to planting rice with a mechanical method or a broad-spectrum herbicide for which resistance does not exist in these weeds [9,10]. The effectiveness of this approach would be optimized if the timing of weed seedling emergence under varying temperatures and irrigation regimes could be accurately predicted and if the conditions for maximizing emergence rate and synchrony could be identified [11].

Population-based threshold models (PBTM) have been developed to describe germination responses to temperature, water potential [12] and oxygen [13], and have been used to predict crop seedling emergence [14,15]. For non-dormant *E. oryzicola* seed, the PBTM approach predicted with useful accuracy the germination responses of seeds to shifting temperature and water availability and their subsequent emergence from field soils [16]. However, *Poaceae* seeds typically possess non-deep physiological dormancy (NDPD), which indicates that seed dormancy release and increases in germination rates (speed) vary along a continuum of time and environmental conditions [17,18]. NDPD may be released by stratification, after-ripening, scarification, excision of the embryo or addition of gibberellin [17] and by various environmental signals including light, fluctuating temperatures and soil nitrate [19]. In addition, the environmental requirements for dormancy alleviation are often population- rather than species-specific [20–22], thus requiring analysis at the population level. While non-dormant seeds of selected herbicide-resistant (R) and herbicide-susceptible (S) populations of *E. oryzicola* germinated similarly [16], information on differences in seed dormancy between R and S populations is lacking. Herbicide-resistant *E.*



**Figure 1. Final germination of stratified (solid lines and symbols) and nonstratified (dashed lines, open symbols) seeds of herbicide-resistant (R) populations (KS and SW) and -susceptible (S) *E. oryzicola* populations (CR and HR) across a range of water potentials.** Seeds were germinated for 14 days at 25°C and 21% oxygen following three months of chilling at 3°C. Symbols represent averages of 6 observations  $\pm$ SE; the  $LSD_{0.05}$  for the interaction between population and stratification treatment was 7% with 192 d.f.; lines are polynomial regressions.

doi:10.1371/journal.pone.0071457.g001

*oryzicola* populations trace their origin to a single introduced biotype dispersed throughout California rice fields [23] suggesting that R populations may respond similarly to environmental variables affecting germination and dormancy.

As in many summer annual species with NPD [24], innate dormancy of *E. oryzicola* seed populations that emerge in spring [1] is alleviated by cold stratification when exposed to a period of moisture at wintertime temperatures in California. Thus, hydration and dark storage at 3°C alleviated dormancy of most seeds in this species [16]. In California, yearly wintertime variation in field temperatures may be less than year-to-year variation in moisture levels, which may range from sporadic rain to prolonged periods of flooding [25]. Adaptation to these conditions would suggest that stratification moisture levels may influence the magnitude of *E. oryzicola* seed dormancy release and that dormancy levels could perhaps be manipulated using wintertime irrigation to increase the rate of springtime germination and weed seedling recruitment [26]. The median base water potential estimated using hydrotime germination models is often a measure of the relative dormancy status of a seed population [12], and because dormancy removal enables *E. oryzicola* seeds to transition from aerobic respiration to anaerobic alcoholic fermentation [1], oxygen-time germination models [13] might also provide a means of assessing dormancy levels in seeds of this species.


To understand the environmental requirements for *E. oryzicola* seed dormancy alleviation, we sought here to: 1) quantify stratification effects upon germination of seeds of R and S populations of *E. oryzicola* across a range of moisture and oxygen levels; and 2) ascertain the relative contributions of alternating temperatures and of stratification temperature, water potential and duration towards dormancy release in R and S *E. oryzicola* populations. This knowledge will contribute towards the accuracy

of germination-based predictions of seedling emergence as affected by the dormancy status of the seed and thus improve the timing and efficacy of weed control programs.

## Materials and Methods

### Plant material and general experimental conditions

*E. oryzicola* seeds (spikelets) of four populations (CR, HR, KS and SW) representing the range of phenotypic variability previously reported in California [23] were mass collected from Sacramento Valley, California, rice fields (with consent of field owners) between 1997 and 2002 [16] and used in all experiments of this study. Populations CR and HR were subsequently classified as herbicide-susceptible (S) and populations KS and SW as herbicide-resistant (R) [16,27]. In the summers of 2007 and 2009, 38 plants from each population were placed in separate greenhouses for seed multiplication at the University of California, Davis. Plants were grown in 2-L pots filled with soil placed in flooded basins under conditions set to approximate mid-springtime field conditions in the Sacramento Valley [25]: 28/14°C day/night temperatures, 50% relative humidity; natural light was supplemented by 900  $\mu\text{mol m}^{-2} \text{s}^{-1}$  of photosynthetic photon flux density (PPFD) from metal halide and high pressure sodium lamps to maintain a 16-h day length; soluble fertilizer (Grow More, Inc., Gardena, CA) was applied through irrigation as needed. Seeds were harvested from panicles at the time of seed shattering in early fall, stored at 20°C for 3 weeks to approximate typical early autumn temperatures and thereafter stored at 3°C, approximating mid-winter temperatures. Water content of seeds kept in dry storage was 7 to 9% (dry weight basis).



**Figure 2. Average observed (symbols) and predicted (lines) germination among herbicide-resistant (R) and -susceptible (S) *E. oryzicola* populations across a gradient of  $\Psi$  for stratified (solid) and nonstratified (open/dashed) seeds.** Seeds were germinated following three months of chilling at 3°C in water (stratified) or under dry conditions (nonstratified). Hydrotime germination models were fit by replicate with the equation  $probit(g) = [\Psi - \theta_H/t_g - \Psi_b(50)]/\sigma_{\Psi b}$ , where  $\Psi$  is experimental water potential,  $\theta_H$  is the hydrotime constant to germination,  $t_g$  is time to germination of fraction  $g$  of the seed population,  $\Psi_b(50)$  is median base water potential and  $\sigma_{\Psi b}$  is the standard deviation in  $\Psi_b$  among seeds in a population. Average stratified and nonstratified root mean squared errors (RMSE)  $\pm$  SE were: 0.079  $\pm$  0.009 and 0.082  $\pm$  0.009 for CR, 0.065  $\pm$  0.006 and 0.107  $\pm$  0.003 for HR, 0.082  $\pm$  0.005 and 0.116  $\pm$  0.014 for KS, and 0.077  $\pm$  0.006 and 0.126  $\pm$  0.007 for SW, respectively. Symbols represent averages of observations and bars represent SE based on six replicate sets of 35 seeds. doi:10.1371/journal.pone.0071457.g002

### Effects of Dormancy on Germination

The role of dormancy in determining the range of environmental conditions that allow germination was assessed in spring 2008 in two concurrent experiments by comparing germination responses to gradients of water potential ( $\Psi$ ) or oxygen concentration ( $O_x$ ) in seeds that had been stratified (wet-chilled) for dormancy removal or not stratified (dry-chilled). Seeds of each population were subjected during 90 days to either stratification by 10 cm immersion in 500 ml water and storage at 3°C to simulate wintertime stratification [24] or to dry chilling at 3°C. Seeds were briefly surface sterilized by 0.5% NaOCl to reduce microbial growth during germination tests, but this did not affect dormancy without stratification. The  $\Psi$  and  $O_x$  experiments that followed were conducted in a growth chamber at a constant 25°C, as this falls within the range of optimal temperatures for germination in these populations [unpublished data], and 12 h day length under 200  $\mu\text{mol m}^{-2} \text{s}^{-1}$  PPF (halogen lights) to satisfy any light requirements for germination. Germinated seeds were counted and removed daily for the first 5 days of the experiment and every 2–3 days thereafter until day 24. Seeds were counted as germinated whenever coleoptile growth reached  $\geq 1$  mm. At the end of each experiment non-germinating seeds were tested for viability using tetrazolium [28].

**Water potential experiment.** Because of this species' association with flooded environments [1,2], a range of fairly moist conditions was simulated using polyethylene glycol 8000 (PEG, Fisher Scientific, Pittsburgh, PA) solutions prepared according to Michel [29] to create  $\Psi$  levels of 0 (pure DI water),  $-0.2$ ,  $-0.4$  and  $-0.7$  MPa. These  $\Psi$  levels were verified with a

dewpoint water potential meter (WP4 DewpointPotentialMeter, DecagonDevices, Pullman, WA) at the beginning and end of the experiment. Experimental units were designed to prevent PEG or moisture loss through evaporation, and there was no change in  $\Psi$  levels over the period of germination. The experimental unit was a 14  $\times$  14  $\times$  5 cm transparent plastic container placed inside a 3.8 L airtight re-sealable clear plastic bag [16]. In each container, a set of 35 seeds for each combination of population and chilling treatment was attached to a 2.5  $\times$  2.5 cm section of Velcro acrylic-based adhesive industrial strip (Uline, Pleasant Prairie, WI) that was fixed to the bottom surface, thus allowing seeds to remain in place 2 cm deep under 300 mL of either DI water or PEG solution per container. Pressurized 21% oxygen was humidified and equally distributed via a flow board through latex tubes (4.8 mm inner diameter, 9.5 mm outer diameter; Kent Elastomer Products, Inc., Kent, OH) at a rate of 600 mL  $\text{min}^{-1}$  and bubbled into containers [16]. Treatments were arranged in a split-plot factorial and replicated six times;  $\Psi$  levels were main plots and populations  $\times$  chilling treatment were randomized sub-plots.

**Oxygen concentration experiment.** Using the same experimental arrangement described in the preceding section and following Al-Ani *et al.* [30], dormancy modification of germination responses to oxygen was evaluated using oxygen concentrations ( $O_x$ ) of 21, 10, 1, 0.01 and 0.001% created using pressurized 21%  $O_2$  and premixed  $O_2/N_2$  ratios (Airgas NCN, Sacramento, CA), and supplied to containers holding 300 mL of DI water. Upon being established in each container through an initial 5-minute flush of gas at a rate of 3000 mL  $\text{min}^{-1}$ ,  $O_x$  treatments were continuously distributed into containers as described above.

**Table 1.** Parameters of the hydrotime model (Equation 2) characterizing the responses of herbicide-resistant (R) and – susceptible (S) *E. oryzipol* populations germinated at 25°C, 21% oxygen and  $\Psi$  of 0, –0.2, –0.4 or –0.7 MPa, after being stratified [16] or not for three months at 3°C;  $\theta_H$  is the hydrotime constant;  $\Psi_b(50)$  is median base water potential, and  $\sigma_{\Psi_b}$  is the standard deviation in  $\Psi_b$  among seeds around  $\Psi_b(50)$ .

<i>E. oryzipol</i>	$\theta_H$	$\Psi_b(50)$	$\sigma_{\Psi_b}$
Population	(MPa d <sup>-1</sup> ±SE)	(MPa ±SE)	(MPa ±SE)
<i>Stratified</i>			
CR (S)	2.91±0.13	–1.12±0.03	0.27±0.02
HR (S)	1.97±0.21	–1.13±0.03	0.17±0.07
KS (R)	1.81±0.10	–1.06±0.04	0.15±0.04
SW (R)	1.75±0.04	–1.07±0.01	0.11±0.02
Average	2.11	–1.10	0.18
<i>Nonstratified</i>			
CR (S)	4.74±0.90	0.27±0.19	0.84±0.22
HR (S)	2.94±0.14	–0.63±0.02	0.15±0.01
KS (R)	3.22±0.12	–0.73±0.02	0.11±0.01
SW (R)	2.62±0.15	–0.66±0.02	0.10±0.00
Average	3.38	–0.37	0.30
<i>LSD<sub>0.05</sub> (error d.f. = 48)</i>			
Population (A)	0.23		
Stratification (B)	0.32		
A × B	NS	0.15	0.09

Models were fit for each of six replicates; values are parameter averages ±SE. doi:10.1371/journal.pone.0071457.t001

Immediately before inflow into the containers, O<sub>2</sub> concentrations in gas mixtures were verified with a headspace trace oxygen analyzer (Pac Check 650, Mocon Inc., Minneapolis, MN). Gas entered the containers via an airtight grommet-lined opening and emerged through an aquarium bubbler fixed to the bottom center of the container. Inflow of gas created a positive pressure inside the sealed bag that permanently pushed air outwards through a syringe needle [16]. Treatments were arranged in a split plot factorial and replicated six times; O<sub>x</sub> levels were main plots and populations × chilling treatment were subplots randomized within.

**Germination data analysis.** Final germination was expressed as a percentage of the total number of seeds in a treatment.

The hydrotime population-based threshold model proposed by Gummerson [31] and described by Bradford [32] was used to quantify stratification effects upon germination responses to moisture stress. The model assumes that all seeds in a population require a constant hydrotime ( $\theta_H$ ) for germination that is defined as:

$$\theta_H = [\psi - \psi_b(g)]t_g \tag{1}$$

where  $\Psi$  is experimental water potential,  $\Psi_b(g)$  is the base water potential below which germination is prevented for a given fraction  $g$  of the seed population, and  $t_g$  is time to germination of  $g$ . The model also assumes  $\Psi_b$  follows a normal distribution among seeds and is thus responsible for within-population variation in

germination timing. Probit analysis was used to estimate the parameters of equation 1 [33]:

$$probit(g) = [\Psi - \theta_H/t_g - \Psi_b(50)]/\sigma_{\Psi_b} \tag{2}$$

where  $\Psi_b(50)$  is median  $\Psi_b$  and  $\sigma_{\Psi_b}$  is the standard deviation in  $\Psi_b$  among seeds in a population. We combined germination data from the range of experimental  $\Psi$  treatments and used equation 2 to predict and plot germination [34]. Following Huarte and Benech-Arnold [35], we used the Solver tool of Microsoft Excel (2003–2010) to derive parameters for this function by minimizing the root-mean-square error (RMSE) between simulated and observed data. ANOVA was conducted for each parameter and protected LSD ( $P < 0.05$ ) values obtained using JMP 8.0 software (SAS Institute Inc. Cary, NC); orthogonal contrasts of means were subsequently used to compare between R and S groups of populations.

Stratification effects on the germination response to decreasing oxygen were analyzed using an oxygen-time threshold model analogous to the hydrotime model [13]. This model assumes a normal distribution of base oxygen threshold levels ( $O_{x_b}$ ) across the seed population and a constant oxygen time ( $\theta_{O_x}$ ) to germination for all seeds. Germination responses are described as a function of the logarithm of the oxygen concentration:

$$\theta_{O_x} = [\log O_x - \log O_{x_b}(g)]t_g \tag{3}$$

where  $O_x$  is daily average oxygen percentage in the micro-environment surrounding the seed,  $O_{x_b}(g)$  is the base or minimum level of oxygen just allowing germination of a given fraction  $g$  of seeds, and  $t_g$  is time to germination for that same fraction. Parameter values for equation 3 were also derived from probit analysis:


$$probit(g) = [\log O_x - \theta_{O_x}/t_g - \log O_{x_b}(50)]/\sigma_{O_{x_b}} \tag{4}$$

where  $O_{x_b}(50)$  is the median base oxygen level, and  $\sigma_{O_{x_b}}$  is the standard deviation of  $\log O_{x_b}$  distribution among individual seeds in the population. Further data analysis was as described earlier for the hydrotime model.

### Dormancy Release Experiments

Three experiments evaluated the roles of temperature fluctuations and stratification conditions (temperature, moisture level, and duration) in releasing dormancy. Populations AM (S) and RD (R) [16,27] were added to the previous set of four populations to better distinguish between R and S seeds. After harvest in fall 2009, seeds were stored dry at 20°C until March 2010, when they were surface-sterilized and exposed to dormancy-release treatments as described in the following sections. Upon completion of those treatments, sets of 50 seeds were removed from each dish, washed with 0.2% Captan fungicide for 1 minute, sprayed with 70% ethanol, rinsed with DI water for 1 minute, transferred to 3-cm Petri dishes containing 2 mL DI water, sealed with Parafilm, and placed in a growth chamber for germination. Germination conditions, set to approximate springtime conditions in the mid Sacramento Valley [25], were 14/26°C night/day, 14-h day length under 390  $\mu\text{mol m}^{-2} \text{s}^{-1}$  PPFD, and 80% RH. Germinated seeds were counted and removed daily over two weeks.

**Alternating temperatures.** Sets of dry-stored non-chilled seeds, not exposed to any dormancy-releasing treatment, were placed in Petri dishes and simultaneously germinated at either constant 20°C or at alternating 14/26°C night/day using growth



**Figure 3. Final germination of stratified (solid lines and symbols) and nonstratified (dashed, open symbols) seeds of herbicide-resistant (R) and -susceptible (S) *E. oryzicola* populations across a range of oxygen levels.** Seeds were germinated at 25°C and 0 MPa, following three months of chilling at 3°C in water (stratified) or under dry conditions (nonstratified). Symbols represent averages of observations  $\pm$ SE based on six replicate sets of 35 seeds; lines are linear regressions. The  $LSD_{0.05}$  for the interaction between population and stratification treatment was = 7% with 192 d.f.

doi:10.1371/journal.pone.0071457.g003

chambers set as described above. Four replicate sets of seeds per population were randomly distributed within each temperature regime.


**Stratification temperature.** Seeds of each population were placed in Parafilm-sealed Petri dishes containing 50 ml of DI water, wrapped in aluminum foil to control for light effects on dormancy release, and randomly arranged in covered 6.1 L plastic storage boxes in dark rooms set for constant 2.5, 5, and 7.5°C temperature regimes. These temperatures were selected because 5°C is the optimum chilling temperature for cold stratification in many species [24] and also because they all fall below the base temperature of approximately 9°C for germination in these populations [16], thus avoiding confounding effects of dormancy release and germination. After 24 days of stratification, seeds were transferred to a growth chamber for germination, as described above. Three replicates per population were arranged in a completely randomized design at each temperature.

**Stratification duration and moisture stress.** Seeds of each population were placed in Parafilm-sealed Petri dishes containing 50 mL PEG solutions of either 0 (DI water), -0.4, -0.8 or -1.6 MPa and kept under a constant 5°C for either 0, 3, 4, 7, 10, 14, 17, 28, 35, 57 or 92 days. The  $\Psi$  spanned the range from no water stress to stress below -1.5 MPa, which is the permanent wilting point of many herbaceous species [36]. The  $\Psi$  levels were established and verified as described earlier, and re-verified every three weeks. Upon completion of stratification treatments, seeds were germinated as described earlier. Treatments were arranged in a split-plot factorial where population  $\times$   $\Psi$  combinations were in main plots and stratification durations were in sub-plots; there were two replicate sets of seeds in a completely randomized design.

**Dormancy release data analysis.** Dormancy-releasing effects of alternating temperatures and of stratification temperature, moisture stress and stratification duration were assessed by comparing germination rates (GR). The GR for a given treatment was calculated as the inverse time to median germination (i.e. 50% of the total seed population), which was estimated for each blocked replicate by fitting cumulative germination data with a three-parameter log-logistic equation using the R statistical program with the *drc* add-on package [37]:

$$G = a / (1 + (x/c)^b) \quad (5)$$

where  $G$  is cumulative germination,  $a$  is the upper asymptote (the lower asymptote is assumed to be 0),  $x$  is days,  $c$  is time to 50% germination and  $b$  is the slope. The inverse of the thus calculated  $c$  parameter equals the GR [32]. Final germination (% of total seeds germinated) and GR were subjected to ANOVA of Box-Cox transformed data (when required to meet assumptions of homogeneity of variance). Orthogonal contrasts between means were subsequently used to compare between R and S groups of populations. All treatments achieved complete final germination, thus equation 5 yielded an accurate estimate of the median germination time for the population. Germination rates (GR) were used for comparisons among treatments as they vary linearly with temperature and  $\Psi$ , whereas median times to germination vary non-linearly with respect to these factors; while germination tests can be done at warmer temperatures, germination time course at all suboptimal temperatures can be predicted based on the GR and base temperatures [19].



**Figure 4. Average observed (symbols) and predicted (lines) germination among herbicide-resistant (R) and -susceptible (S) *E. oryzicola* populations across a gradient of oxygen levels ( $Ox$ ) in stratified (solid) and nonstratified (open/dashed) seeds.** Modeling of oxygen time was only applied when an oxygen dose response was observed (nonstratified seeds in the 21-1%  $Ox$  range) by fitting to each replicate the equation  $probit(g) = [\log Ox - \theta_{Ox}/t_g - \log Ox_b(50)]/\sigma_{Ox_b}$  where  $Ox$  is daily average oxygen percentage in the micro-environment surrounding the seed,  $\theta_{Ox}$  is the oxygen time constant,  $t_g$  is time to germination for fraction  $g$  of the seed population,  $Ox_b(50)$  is the median base oxygen level, and  $\sigma_{Ox_b}$  is the standard deviation of the  $\log Ox_b$  distribution among individual seeds in the population. Average RMSE  $\pm$  SE for nonstratified seeds were  $0.084 \pm 0.003$ ,  $0.075 \pm 0.010$ ; and  $0.077 \pm 0.011$  for HR, KS and SW, respectively. A hydrotime model for 0 MPa (Equation 2) was fit to data from germination time-courses of stratified seeds; when compared to observed germination across  $Ox$ , RMSE for these models were  $0.096 \pm 0.005$ ,  $0.093 \pm 0.010$ ,  $0.087 \pm 0.012$  and  $0.128 \pm 0.016$  for CR, HR, KS and SW, respectively. Symbols represent averages of observations and bars represent SE based on six replicate sets of 35 seeds.  
doi:10.1371/journal.pone.0071457.g004

**Table 2. Effects of alternating temperatures on final germination (G) and germination rates (GR, calculated by replicate as the inverse of median time to germination ( $1/t_{50}$ ) using Equation 5) of herbicide-resistant (R) and -susceptible (S) *E. oryzicola*.**

<i>E. oryzicola</i> Population	Constant Temperature		Alternating Temperature	
	Final G(% $\pm$ SE)	GR ( $1/t_{50} \pm$ SE)	Final G(% $\pm$ SE)	GR( $1/t_{50} \pm$ SE)
AM (S)	98 $\pm$ 3	0.12 $\pm$ 0.00	100 $\pm$ 0	0.19 $\pm$ 0.01
CR (S)	94 $\pm$ 3	0.11 $\pm$ 0.02	100 $\pm$ 0	0.19 $\pm$ 0.01
HR (S)	98 $\pm$ 2	0.14 $\pm$ 0.00	100 $\pm$ 0	0.21 $\pm$ 0.01
KS (R)	100 $\pm$ 0	0.15 $\pm$ 0.00	100 $\pm$ 0	0.23 $\pm$ 0.00
RD (R)	100 $\pm$ 0	0.16 $\pm$ 0.00	100 $\pm$ 0	0.25 $\pm$ 0.01
SW (R)	100 $\pm$ 0	0.15 $\pm$ 0.00	100 $\pm$ 0	0.25 $\pm$ 0.01
Average	98	0.13	100	0.22
<i>LSD</i> <sub>0.05</sub> (error d.f. = 24)				
Final G			2.2	
GR Population x Temperature regime			0.01	

Nonstratified seeds were germinated at 0 MPa and 21% oxygen, with temperatures either held constant at 20°C or set to 14/26°C day/night regime. Values are averages  $\pm$  SE of four replicate sets of 50 seeds.  
doi:10.1371/journal.pone.0071457.t002



**Table 3.** Germination rates (GR, calculated by replicate as the inverse of median time to germination ( $1/t_{50}$ ) using Equation 5) following 24 days of stratification at three temperatures  $<T_b$  for germination.

<i>E. oryzicola</i>	Temperature regime		
	2.5°C	5°C	7.5°C
<b>Population</b>			
	<i>GR (1/t<sub>50</sub>)</i>		
AM (S)	0.67±0.04	0.67±0.02	0.78±0.04
CR (S)	0.56±0.02	0.62±0.02	0.74±0.06
HR (S)	0.60±0.06	0.65±0.04	0.85±0.09
KS (R)	0.66±0.05	0.67±0.04	0.81±0.01
RD (R)	0.64±0.03	0.61±0.02	0.74±0.09
SW (R)	0.66±0.03	0.60±0.01	0.79±0.05
Average	0.63	0.64	0.79
LSD <sub>0.05</sub> (d.f. = 18)	NS	NS	NS

Herbicide-resistant (R) and -susceptible (S) *E. oryzicola* seeds were germinated at 14/26°C night/day, 0 MPa and 21% oxygen. Values are averages ±SE of 3 replicate sets of 50 seeds.

doi:10.1371/journal.pone.0071457.t003

## Results

### Effects of dormancy on germination

All populations germinated to high percentages (85–90%) across  $\Psi$  when stratified (Figure 1). However, final germination of nonstratified seeds approached 0% as  $\Psi$  decreased to  $-0.7$  MPa, and was notably low for the S population CR. Tetrazolium testing of nonstratified seeds that failed to germinate at  $-0.7$  MPa indicated generally high seed viability across populations:  $91 \pm 3$ ,  $97 \pm 2$ ,  $65 \pm 10$  and  $92 \pm 3\%$  for CR, HR, KS and SW, respectively. Dormancy release by stratification enabled faster germination and under drier conditions (Figure 2) by reducing average  $\theta_H$  by 38% and decreasing  $\Psi_b(50)$  compared to nonstratified seeds (Table 1). Stratification only improved germination synchrony ( $\sigma_{\Psi_b}$ ) for the CR population. The hydrotime model fitted better the germination of stratified seeds, as indicated by the lower RMSE values (Figure 2). Model parameters suggested no consistent differences between the R and S seeds in their germination time course responses to moisture availability (Table 1). The two S populations differed from each other ( $p < 0.001$ ) in all three hydrotime model parameters, while differences between the two R populations were not significant.

Stratified seeds were insensitive to changes in oxygen availability (Figures 3–4) even at levels as low as those of flooded paddy fields [38]. Stratification enhanced final germination of S seeds only (Figure 3). Thus germination of nonstratified HR seeds decreased by 15% when  $O_x$  dropped to 1%, and final germination of nonstratified CR seeds was notably low and insensitive to oxygen availability (Figure 3). Tetrazolium testing confirmed viability of non-germinating HR and CR seeds (on average  $72 \pm 6\%$  and  $72 \pm 12\%$  across  $O_x$  for HR and CR, respectively). Germination time courses of stratified *E. oryzicola* seeds in all populations were almost insensitive to hypoxia and sensitivity was even limited in nonstratified seeds (Figure 4), which had an average  $O_{x_b}(50)$  of 0.04 ppm (not shown). Given this general lack of responses, an oxygen threshold model could only be fit to nonstratified seeds of three of the populations germinating under  $O_x$  of 1 to 21%, but model parameters still reflected only minor sensitivity to  $O_x$  and were not statistically different across populations (not shown). This minor

sensitivity disappeared under stratification and germination rates were faster than for non-stratified seeds under all  $O_x$  levels (Figure 4).

### Dormancy release

**Effect of alternating temperatures.** Full final germination was achieved by all populations under both constant and alternating temperatures, but alternating temperatures nearly doubled germination rates (GR) in all populations (Table 2). Average GR for R populations (KS, RD, SW) were greater ( $p < 0.001$ ) than for S populations under both constant ( $0.15 \pm 0.00$  vs.  $0.13 \pm 0.00$ , respectively) and alternating ( $0.24 \pm 0.00$  vs.  $0.20 \pm 0.01$ , respectively) temperature regimes.

**Decreasing stratification temperature.** Germination rates were similar at stratification temperatures of 2.5 and 5°C, but increased when temperatures rose from 5 to 7.5°C (Table 3). There were no differences ( $p = 0.890$ ) in GR among R and S *E. oryzicola* populations within the range of stratification temperatures studied (Table 3).

**Stratification duration and moisture stress.** Final germination percentages were consistently  $\geq 95\%$  (not shown) and the enhancement of GR due to stratification varied ( $p < 0.05$ ) with the duration of wet-chilling and moisture stress during that period (Figure 5). Thus GR peaked after 17–30 days of immersion in water ( $\Psi = 0$  MPa), and maximum GR values declined with  $\Psi < 0$  MPa (Table 4). Lengthening stratification duration did not mitigate decreases in GR as  $\Psi$  became more negative (Figure 5). Average GR was higher for R populations than for S populations ( $0.46 \pm 0.01$  and  $0.39 \pm 0.01$ , respectively;  $p < 0.001$ ) across stratification durations and  $\Psi$  levels. Stratification in water beyond the time to maximum GR ultimately led to greater reductions in GR for the R than for the S populations (Figure 5).


## Discussion

### Effects of stratification on germination responses to moisture and oxygen

Reductions in both  $\theta_H$  and  $\Psi_b(50)$  indicated dormancy removal and broadened the range of water potentials at which germination could occur following stratification [39,40]. Thus dormancy was partly associated with reduced ability to germinate under drier conditions, which is a logical adaptive trait for an aquatic species like *E. oryzicola* [1]. This was further reflected in the dramatic drop in germination of nonstratified seeds when  $\Psi$  fell to  $-0.7$  MPa (Figure 1), and in the positive  $\Psi_b(50)$  that reflected the inability of nonstratified CR seeds to germinate (Table 1, Figure 1). The overall reduction in  $\Psi_b(50)$  due to stratification is consistent with similar reductions from after-ripening in red rice [41] and stratification in *Polygonum* [40]. Poorer hydrotime model predictions of germination time courses for nonstratified seeds (Figure 2) suggest sensitivity to other dormancy-related variables, such as light, nitrate or hormone levels, which may complicate the application of this model when dormancy is present in a population [42].

A transition from aerobic respiration to anaerobic metabolism enables adaptation of this weed to the flooding of paddy fields, and previous research has linked dormancy removal in *E. oryzicola* to increased ability to germinate under hypoxia through anaerobic fermentation [1]. In our study, stratification removed a trend towards slower germination at the lower  $O_x$  levels in three nonstratified populations (HR, KS and SW) (Figure 4). Nevertheless, the range of germination responses to  $O_x$  levels for the nonstratified seeds of these three populations was narrow and the estimated  $O_{x_b}(50)$  values ( $\sim 0.04$  ppm) fell well below those of most





**Figure 5. Herbicide-resistant (R) and -susceptible (S) *E. oryzae* germination in response to stratification duration and water potential.** Seeds were immersed in PEG solutions of 0, -0.4, -0.8 and -1.6 MPa at a constant 5°C for 3, 4, 7, 10, 14, 17, 23, 30, 57 and 92 days prior to germination at 21°C and 0 MPa. Final germination for all treatments was  $\geq 95\%$ . Germination rates were calculated by replicate as the inverse of median time to germination ( $1/t_{50}$ ), which was determined from Equation 5. Seeds were germinated at 17/24°C night/day temperatures, 0 MPa and 21% oxygen. Symbols are averages of two replicates of 50 seeds per treatment. Peak GR was attained at 30, 23, 30, 23, 23 and 17 days of stratification in 0 MPa for AL, CR, HR, KS, RD and SW, respectively. The  $LSD_{0.05}$  for the interaction between population, stratification duration and stratification  $\Psi$  was 0.06; d.f. = 478.

doi:10.1371/journal.pone.0071457.g005

**Table 4.** Maximum germination rates (GR, calculated by replicate as the inverse of median time to germination ( $1/t_{50}$ ) using Equation 5) after stratification at 5°C under various water potentials.

<i>E. oryzicola</i> Population	Water Potential				Average
	0 MPa	-0.4 MPa	-0.8 MPa	-1.6 MPa	
	<i>GR (1/t<sub>50</sub>) ± SE</i>				
AM (S)	0.67±0.04	0.56±0.01	0.55±0.02	0.48±0.01	0.57±0.03
CR (S)	0.59±0.02	0.51±0.01	0.50±0.01	0.46±0.03	0.52±0.02
HR (S)	0.71±0.00	0.59±0.01	0.54±0.01	0.52±0.01	0.59±0.03
KS (R)	0.72±0.01	0.66±0.05	0.58±0.03	0.55±0.01	0.63±0.03
RD (R)	0.77±0.07	0.61±0.01	0.57±0.01	0.56±0.03	0.64±0.03
SW (R)	0.78±0.08	0.66±0.05	0.58±0.02	0.57±0.02	0.64±0.04
Average	0.71±0.02	0.60±0.02	0.55±0.01	0.52±0.01	
<i>LSD<sub>0.05</sub></i> (error d.f. = 48) <sup>a</sup>					
Population (A)		0.03			
Water Potential (B)		0.03			

GR was assessed for each herbicide-resistant (R) and -susceptible (S) *E. oryzicola* population after 0, 3, 4, 7, 10, 14, 17, 28, 35, 57 and 92 days of stratification and values are averages ±SE of the maximum GR for each replicate.

<sup>a</sup>To meet ANOVA assumptions, a Box-Cox transformation ( $\lambda = -1.8$ ) was applied.

doi:10.1371/journal.pone.0071457.t004

other species [13]. Thus *E. oryzicola* exhibited limited sensitivity to hypoxia consistent with the known ability of this weed to germinate under flooded conditions [1]. The somewhat lower final germination of nonstratified HR (S) seeds at reduced  $O_x$  levels (Figure 3) could suggest that a fraction of that seed population had comparatively greater requirements of aerobic respiration to germinate [13]. The timing of the *E. oryzicola* transition from aerobic to fermentative metabolism and its relationship to stratification and dormancy removal requires further study. The lower final germination of nonstratified CR (S) seeds across  $O_x$  levels (Figure 3) could be explained by their positive  $\Psi_b(50)$  value (Table 1) rather than as an oxygen response.

Stronger similarities between the R populations KS and SW than between the S populations CR and HR in hydrotime model parameters (Table 1) and in final germination across  $\Psi$  and  $O_x$  levels (Figures 1 and 3) corroborate previous reports of reduced phenotypic variation among R populations compared to S [23,27].

### Dormancy release by alternating temperatures

Dormancy release by alternating temperatures typical of springtime at the time of rice seeding in the Sacramento Valley of California [25] was expressed as an increase in GR for seeds of all populations used in these experiments, rather than by a change in final germination percentage (Table 2). These observations are consistent with previous reports that fluctuating temperatures aid in alleviating dormancy [43–45], and indicate that dormancy was present in our seeds [46]. Thus our results corroborate previous reports on the presence of seed dormancy in *E. oryzicola* [1]. However, as seeds were stored at room temperature for a few months prior to experimentation and NDPD release can already begin during dry after-ripening [47–49], some loss of dormancy could already have occurred.

### Dormancy release by stratification at different temperatures and $\Psi$

Dormancy release by stratification was expressed primarily through increases in germination rates. Studies with other species have documented enhancement of stratification effects as

stratification temperatures were reduced below  $T_b$ ; such a relationship has been the basis for the application of population-based stratification-time models [50,51]. However, GR in our seeds was not enhanced by lowering stratification temperatures (Table 3) but, instead, by raising them to 7.5, which is close to the  $T_b(50)$  for *E. oryzicola* [16]. Further experimentation would be required to define whether this increase in GR limited to a narrow temperature range signifies a the beginning of a trend toward enhanced dormancy release with rising temperatures, or the onset of germination, or both.

In California rice growing areas, winter temperatures will remain within 5°C of  $T_b$  for *E. oryzicola* germination [25], but the degree of wintertime chilling is beyond the control of land managers. However, growers may modify the hydric status of their fields through irrigation; thus our research also focused on the effects of moisture on dormancy release. We found lower maximum GR values following stratification under decreasing  $\Psi$  (Table 4), which implies that dormancy release may be in part controlled by winter moisture levels in this wetland species. Stratification duration to full dormancy removal, as represented by maximum GR in these populations (Figure 5), was less than that reported for *Polygonum aviculare* [40], *Bromus tectorum* [47] and for weedy rice [41], suggesting dormancy is either lower or more readily removed in our *E. oryzicola* populations.

Since seeds used for the germination studies were stratified for 90 days, some measure of secondary dormancy may have been induced [51–53]. However, the large differences in observed germination responses between stratified and nonstratified seeds are evidence that significant levels of dormancy had been removed by stratification. Secondary dormancy can lead *E. oryzicola* seeds to persist in seedbanks for as many as ten years [1]; thus although weed control techniques for this weed can be successful [8,9], the complete eradication of the weed from a rice field is difficult [1].

Applications of this research to field conditions must take into account that dormancy levels may have been influenced to varying degrees by each of the general experimental conditions to which seeds were subjected [24]. These include environmental factors, such as fixed temperature regimes and photoperiods, experienced

by seeds during their development and maturation [21,24,54], as well as seed cleaning procedures and storage conditions after harvest but before experiments began [20,24,55,56]. Post-harvest seed storage temperatures may have influenced dormancy levels, as could the length of storage prior to experimentation. Indeed, there is evidence that *Arabidopsis* germination may be more affected by the postdispersal environment than by certain conditions affecting seed maturation [57]; however, such results were not explicitly tied to dormancy in grasses. Therefore, any inferences made to field conditions should be adequately tested through field experiments.

Winter flooding is often practiced by California rice growers to facilitate stubble decomposition [26], a practice that may help *E. oryzicola* control by hastening its springtime GR, provided winter-time soil saturation can be maintained for an extended period of time (Figure 5). Herbicide-resistant populations generally exhibited higher germination rates under either constant temperatures (Table 2) and when stratified at sub-optimal  $\Psi$  (<0 MPa), suggesting lower dormancy levels. Greater basal ethylene levels, which can play a causal role in the metabolic response to submergence [58], have been detected in R plants compared to S plants [59]. Since increased ethylene has been shown to correlate to increased dormancy release and germination in some species [60], further research might explore whether higher basal ethylene levels in R seeds might contribute to their higher GR.

In summary, *E. oryzicola* seed dormancy was manifested primarily by reduced GR rather than by lower final germination percentages, and was released by alternating temperatures and by

stratification. Stratification led to dormancy release characterized by hastened germination rates of unstressed seeds, enhanced ability to accrue hydrotime and germinate under drier conditions, and a minor increment of the ability to germinate under hypoxia. There was, otherwise, little germination response to oxygen availability in the *E. oryzicola* seeds tested, consistent with the wetland habit of this weed. The effects of stratification on GR were generally more sensitive to changes in  $\Psi$  than to temperature. Dormancy levels may have been lower among R populations compared to S populations in these experiments, but R seeds may also be more prone to secondary dormancy induction as stratification duration increases. A weed seed bank depletion program based on favoring weed emergence for subsequent control, such as with the stale seedbed technique, would benefit from optimizing environmental conditions for weed seed germination. Based on our results, and depending on the degree of dormancy of the population, *E. oryzicola* dormancy release to enhance weed seed germination would benefit from field soil saturation in winter. This will decrease time to seedling emergence, allowing for early-season weed control and a shortened crop planting delay.

## Author Contributions

Conceived and designed the experiments: LGB KJB AJF. Performed the experiments: LGB. Analyzed the data: LGB KJB AJF. Contributed reagents/materials/analysis tools: LGB AJF. Wrote the paper: LGB KJB AJF.

## References

- Yamasue Y (2001) Strategy of *Echinochloa oryzicola* Vasing. for survival in flooded rice. *Weed Biol Manage* 1: 28–36.
- Barrett SCH (1983) Crop mimicry in weeds. *Econ Bot* 37: 255–282.
- Hill JE, Le Strange ML, Bayer DE, Williams JF (1985) Integrated weed management in California. *Proc Western Weed Sci Soc* 38: 100–104.
- Fischer AJ, Ateh CM, Bayer DE, Hill JE (2000) Herbicide-resistant *Echinochloa oryzoides* and *E. phylloipogon* in California *Oryza sativa* fields. *Weed Sci* 48: 225–230.
- Yun MS, Yogo Y, Miura R, Yamasue Y, Fischer AJ (2005) Cytochrome P-450 monooxygenase activity in herbicide-resistant and -susceptible late watergrass (*Echinochloa phylloipogon*). *Pest Biochem Physiol* 83: 107–114.
- Ruiz-Santaella JP, De Prado R, Wagner J, Fischer AJ, Gerhards R (2006) Resistance mechanisms to cyhalofop-butyl in a biotype of *Echinochloa phylloipogon* (Stapf) Koss. from California. *J Plant Diseases Prot* 20: 95–100.
- Yasuor H, TenBrook PL, Tjeerdema RS, Fischer AJ (2008) Responses to clomazone and 5-ketoclomazone by *Echinochloa phylloipogon* resistant to multiple herbicides in Californian rice fields. *Pest Manage Sci* 64: 1031–1039.
- Linquist BA, Fischer A, Godfrey L, Greer C, Hill J, et al. (2008) Minimum tillage could benefit California rice farmers. *Calif Ag* 62: 24–29.
- Pittelkow CM, Fischer AJ, Moechnig MJ, Hill JE, Koffler KB, et al. (2012) Agronomic productivity and nitrogen requirements of alternative tillage and crop establishment systems for improved weed control in direct-seeded rice. *Field Crops Res* 130: 128–137.
- Johnson WC, Mullinix BG (1995) Weed management in peanut using stale seedbed techniques. *Weed Sci* 43: 293–297.
- Fischer AJ, Linquist B, Moechnig M, Mutters R, Hill JE, et al. (2009) Alternative rice stand establishment systems to manage herbicide resistant weeds. *Proc Ann Mtg Weed Sci Soc Am No. 538*.
- Bradford KJ (2002) Applications of hydrothermal time to quantifying and modeling seed germination and dormancy. *Weed Sci* 50: 248–260.
- Bradford KJ, Come D, Corbineau F (2007) Quantifying the oxygen sensitivity of seed germination using a population-based threshold model. *Seed Sci Res* 17: 33–43.
- Finch-Savage WE, Phelps K (1993) Onion (*Allium cepa* L.) seedling emergence patterns can be explained by the influence of soil temperature and water potential on seed germination. *J Exp Bot* 44: 407–414.
- Finch-Savage WE, Rowse HR, Dent KC (2005) Development of combined imbibition and hydrothermal threshold models to simulate maize (*Zea mays*) and chickpea (*Cicer arietinum*) seed germination in variable environments. *New Phytologist* 165: 825–838.
- Boddy LG, Bradford KJ, Fischer AJ (2012) Population-based threshold models describe weed germination and emergence patterns across varying temperature, moisture and oxygen conditions. *J Appl Ecol* 49: 1225–1236.
- Baskin JM, Baskin CC (2004) A classification system for seed dormancy. *Seed Sci Res* 14: 1–16.
- Gordon AG (1973) The rate of germination. In: Heydecker W, editor. *Seed Ecology*. London: Butterworths. 391–409.
- Bewley JD, Bradford KJ, Hilhorst HWM, Nonogaki H (2013) *Seeds: Physiology of Development, Germination and Dormancy* 3<sup>rd</sup> edition. New York: Springer.
- Finch-Savage WE, Leubner-Metzger G (2006) Seed dormancy and the control of germination. *New Phytologist* 171: 501–523.
- Donahue K (2005) Seeds and seasons: interpreting germination timing in the field. *Seed Sci Res* 15: 175–187.
- Grundy AC (2003) Predicting weed emergence: a review of approaches and future challenges. *Weed Res* 43: 1–11.
- Tsuji R, Fischer AJ, Yoshino M, Roel A, Hill JE, et al. (2003) Herbicide-resistant late watergrass (*Echinochloa phylloipogon*): similarity in morphological and amplified fragment length polymorphism traits. *Weed Sci* 51: 740–747.
- Baskin CC, Baskin JM (2001) *Seeds: ecology, biogeography, and evolution of dormancy and germination*. San Diego: Academic Press. 16–17, 57–70, 187.
- University of California IPM Online. Available: <http://www.ipm.ucdavis.edu/calludt.cgi/WXSTATIONAVG?MAP=butte.html&STN=DURHAM>. Accessed March, 2012.
- Fitzgerald GJ, Scow KM, Hill JE (2000) Fallow season straw and water management effects on methane emissions in California rice. *Glob Biogeochem Cycles* 18: 767–776.
- Boddy LG, Streibig JC, Yamasue Y, Fischer AJ (2012) Biomass, fecundity and interference ability of multiple herbicide-resistant and -susceptible late watergrass (*Echinochloa phylloipogon*). *Weed Sci* 60: 401–410.
- Peters J (2000) *Tetrazolium testing handbook*. Contribution No. 29: The handbook on seed testing. Association of Official Seed Analysts, Lincoln, NE, USA.
- Michel BE (1983) Evaluation of the water potentials of solutions of polyethylene glycol 8000 both in the absence and presence of other solutes. *Plant Physiol* 72: 66–70.
- Al-Ani A, Bruzau F, Raymond P, Saintes V, Leblanc JM, Pradet A (1985) Germination, respiration, and adenylate energy charge of seeds at various oxygen partial pressures. *Plant Physiol* 79: 885–890.
- Gummerson RJ (1986) The effect of constant temperatures and osmotic potentials on the germination of sugar beet. *J Exp Bot* 37: 729–741.
- Bradford KJ (1990) A water relations analysis of seed germination rates. *Plant Physiol* 94: 840–849.
- Dahal P, Bradford KJ (1990) Effects of priming and endosperm integrity on seed germination rates of tomato genotypes II: germination at reduced water potential. *J Exp Bot* 41: 1431–1439.

34. Chantre GR, Sabbatini MR, Orioli GA (2010). An after-ripening thermal-time model for *Lithospermum arvense* seeds based on changes in population hydrotime parameters. *Weed Res* 50: 218–227.
35. Huarte HR, Benech-Arnold RA (2010) Hormonal nature of seed responses to fluctuating temperatures in *Cynaracardunculus* (L.). *Seed Sci Res* 20: 39–45.
36. Lambers H, Chapin FS, Pons TL (2008) *Plant physiological ecology*. New York: Springer, Berlin, Heidelberg. 169.
37. Ritz C, Streibig JC (2005) Bioassay analysis using R. *J Stat Software* 12: 1–22.
38. Patrick WH, Mikkelsen DS (1971) Plant nutrient behavior in flooded soil. In: Patrick WH, Mikkelsen DS, Wells BR, editors. *Fertilizer technology and use* 2<sup>nd</sup> ed. Soil Science Society of America. 187–215.
39. Alvarado V, Bradford KJ (2005) Hydrothermal time analysis of seed dormancy in true (botanical) potato seeds. *Seed Sci Res* 15: 77–88.
40. Batlla D, Benech-Arnold RL (2004) A predictive model for dormancy loss in *Polygonum aviculare* L. seeds based on changes in population hydrotime parameters. *Seed Sci Res* 14: 277–286.
41. Gianinetti A, Cohn MC (2007) Seed dormancy in red rice. XII: Population based analysis of dry after-ripening with a hydrotime model. *Seed Sci Res* 17: 253–271.
42. Allen PS, Benech-Arnold RL, Batlla D, Bradford KJ (2007) Modeling of seed dormancy. In: K.J. Bradford and H. Nonogaki, *Seed Development, Dormancy and Germination*. *Annu Plant Rev* 27: 72–112.
43. Chao WS, Foley ME, Dogramaci M, Anderson JV, Horvath DP (2011) Alternating temperature breaks dormancy in leafy spurge seeds and impacts signaling networks associated with HY5. *FunctIntegGenom* 11: 637–649.
44. Probert RJ (2000) The role of temperature in the regulation of seed dormancy and germination. In: Fenner M, editor. *Seeds: the ecology of regeneration in plant communities* 2<sup>nd</sup> edition. Wallingford: CABI Publishing. 261–292.
45. Totterdell S, Roberts EH (1980) Characteristics of alternating temperatures which stimulate loss of dormancy in seeds of *Rumex obtusifolius* L. and *Rumex crispus* L. *Plant Cell Env* 3: 3–12.
46. Hilhorst HWM (2010) Standardizing seed dormancy research. In: Kermod AR, editor. *Seed dormancy: methods and protocols*. *Methods in Molecular Biology*, vol. 773. Springer Science. 43–52.
47. Bair NB, Meyer SE, Allen PS (2006) A hydrothermal after-ripening time model for seed dormancy loss in *Bromustectorum* L. *Seed Sci Res* 16: 17–28.
48. Chantre GR, Batlla D, Sabbatini MR, Orioli G (2009) Germination parameterization and development of an after-ripening thermal-time model for primary dormancy release of *Lithospermum arvense* seeds. *Ann Bot* 103: 1291–1301.
49. Steadman KJ, Crawford AD, Gallagher RS (2003) Dormancy release in *Lolium rigidum* seeds is a function of thermal after-ripening time and seed water content. *Funct Plant Bio* 30: 345–352.
50. Batlla D, Benech-Arnold RL (2003) A quantitative analysis of dormancy loss dynamics in *Polygonum aviculare* L. seeds: Development of a thermal time model based on changes in seed population thermal parameters. *Seed Sci Res* 13: 55–68.
51. Wang WQ, Song SQ, Li SH, Gan YY, Wu JH, et al. (2009) Quantitative description of the effect of stratification on dormancy release of grape seeds in response to various temperatures and water contents. *J Exp Bot* 60: 3397–3406.
52. Totterdell S, Roberts EH (1979) Effects of low temperatures on the loss of innate dormancy and the development of induced dormancy in seeds of *Rumex obtusifolius* L. and *Rumex crispus* L. *Plant Cell Env* 2: 131–137.
53. Willemsen RW (1975) Effect of stratification temperature and germination temperature on germination and the induction of secondary dormancy in common ragweed seeds. *Am J Bot* 62: 1–5.
54. Donohue K (2009) Completing the cycle: maternal effects as the missing link in plant life histories. *Phil Trans R Soc B* 364: 1059–1074.
55. Allen PS, Meyer SE, Beckstead J (1995) Patterns of seed after-ripening in *Bromustectorum* L. *J Exp Bot* 46: 1737–1744.
56. Benech-Arnold RL, Sanchez RA, Forcella F, Kruk BC, Ghersa CM (2000) Environmental control of dormancy in weed seed banks in soil. *Field Crops Res* 67: 105–122.
57. Donohue K, Dorn L, Griffith C, Kim E, Aguilera A, et al. (2005) Environmental and genetic influences on the germination of *Arabidopsis thaliana* in the field. *Evol* 59: 740–757.
58. Bailey-Serres J, Fukao T, Gibbs DJ, Holdsworth MJ, Lee SC, et al. (2012) Making sense of low oxygen sensing. *Trends Plant Sci* 17: 129–138.
59. Yasuor H, Milan M, Eckert JW, Fischer AJ (2012) Quinlorac resistance: a concerted hormonal and enzymatic effort in *Echinochloa phyllipogon*. *Pest Manage Sci* 68: 108–115.
60. Matilla AJ (2000) Ethylene in seed formation and germination. *Seed Sci Res* 10: 111–126.