

UC Office of the President

Recent Work

Title

How is exposure to tobacco outlets within activity spaces associated with daily tobacco use among youth? A mediation analysis

Permalink

<https://escholarship.org/uc/item/1hh300nb>

Authors

Kowitt, Sarah D
Lipperman-Kreda, Sharon

Publication Date

2019-05-23

Peer reviewed

How is exposure to tobacco outlets within activity spaces associated with daily tobacco use among youth? A mediation analysis

Sarah D. Kowitt, PhD, MPH ¹

Sharon Lipperman-Kreda, PhD ²

1. Department of Family Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC, USA
2. Prevention Research Center, Pacific Institute for Research and Evaluation, Berkeley, CA, USA

Correspondence: Sarah D. Kowitt, PhD MPH, University of North Carolina, Department of Family Medicine, School of Medicine, University of North Carolina at Chapel Hill, Chapel Hill, NC, 27599, USA. Telephone: 704-443-1811. Email: kowitt@email.unc.edu.

Main Text Word Count: 4,022

Abstract Word Count: 245

References: 48

Tables: 2

Figures: 2

Supplemental tables: 1

Supplemental figure: 1

Abstract:

Introduction: We investigated associations between exposure to tobacco outlets within activity spaces and daily tobacco use, and whether exposure to adults or peers using tobacco mediate these relationships.

Methods: We used Geographic Ecological Momentary Assessment data over 14 days from 85 youth aged 16-20 years in 8 mid-sized California city areas. Tobacco outlet addresses and GPS locations were geocoded and activity spaces were constructed by joining sequential points. We assessed daily number of tobacco outlets within 50 or 100 meters of activity space polylines and number of minutes participants were within 50 or 100 meters of tobacco outlets each day; daily use of tobacco; and whether participants saw 1) adults and 2) people their age (peers) using tobacco each day.

Results: Controlling for demographics, results of multilevel structural equation models showed no association between number of tobacco outlets within 50 meters of polylines and tobacco use (probit regression coefficient: 0.01, $p=0.82$). However, we found evidence of an indirect effect ($p=0.001$) through daily exposure to peers using tobacco. Specifically, greater number of tobacco outlets within 50 meters of polylines was positively associated with seeing peers use tobacco (probit regression coefficient: 0.10, $p<0.001$). In turn, seeing peers use tobacco was positively associated with tobacco use on that day (probit regression coefficient: 2.23, $p<0.001$). Similar results were found for number of tobacco outlets within 100 meters of polylines.

Conclusions: Exposure to tobacco outlets within activity spaces affects youth tobacco use through daily exposure to peers who use tobacco.

Keywords: youth tobacco use; retail outlets; norms; geographic ecological momentary assessment; longitudinal research; mediation

Implications

Using real time-ordered data, this paper examines whether exposure to adults and peers using tobacco mediate associations between exposure to tobacco outlets within activity spaces and daily tobacco use among youth. Results suggest that exposure to tobacco outlets within activity spaces affects daily tobacco use through exposure to peers who use tobacco. These findings provide additional significant support for policymakers who are considering regulating the number and density of tobacco retailers and point to the importance of interventions focused on peer tobacco use and youths' daily environments to reduce tobacco use.

INTRODUCTION

Tobacco use is the leading cause of preventable death in the United States (US).¹ Since most tobacco use starts during adolescence,^{1,2} prevention efforts need to focus on youth. One important determinant of youth tobacco use is exposure to tobacco retail outlets. A growing number of studies have found associations between density of tobacco outlets near homes and schools and cigarette smoking.³⁻⁹

Recent research has also begun to explore how to best measure exposure to tobacco outlets and its associations with youth tobacco use. For instance, results of an exploratory study in California suggest that focusing only on exposure to tobacco retail outlets around homes and schools may underestimate the extent to which youth are exposed to tobacco outlets in their daily lives and that exposure to tobacco outlets was more accurately measured by considering activity spaces.¹⁰ Activity spaces refer to locations that people commonly frequent as part of their daily routines. Activity spaces can be defined *a priori* as places in the community (e.g., parks, malls, urban centers, schools) or based on routes an individual frequents based on his/her travel patterns. Research shows that individuals engage in many daily activities outside of their home or residential environment.^{11,12} Few other studies have examined exposure to tobacco outlets considering activity spaces,¹³⁻¹⁵ however, research on its use and applications are growing.¹⁶

Also, questions remain about *how* exposure to tobacco outlets affects youth tobacco use. Exposure to tobacco outlets may increase tobacco use through perceptions of others' tobacco use. That is, daily exposure to tobacco outlets may increase exposure to adults or peers who use tobacco, who are likely to be seen around tobacco outlets, which in turn may influence tobacco use. Though the Social Cognitive theory^{17,18} and previous research^{2,19,20} also suggest that perceptions of others' tobacco use are strong determinants of youth tobacco use, no research, to our knowledge, has examined if perceptions of tobacco use mediate associations between exposure to tobacco outlets and tobacco use.

To address these research gaps, this study examined whether daily exposure to tobacco outlets within activity spaces is associated with daily tobacco use among youth and if perceptions of daily tobacco use by adults or peers mediate this relationship. We hypothesized that youth who were exposed to

more tobacco outlets and spent more time around tobacco outlets in their daily activity spaces would report seeing more peers and adults use tobacco and in turn, seeing more peers and adults use tobacco would be associated with daily tobacco use.

METHODS

Study cities and participants

We collected Geographic Ecological Momentary Assessment (GEMA) data from youth aged 16-20 years (n=101 participants) in 8 mid-sized California city areas. GEMA data combine ecological momentary assessment data (EMA) with global positioning systems (GPS) and geographic information systems (GIS) and allow researchers to collect real-time data on participants' behaviors and environments.²¹ Cities were selected from an existing geographically diverse sample of 50 noncontiguous California cities (population range: 50,000 to 500,000).^{22,23} To select the 8 cities, we considered cities within a 50-mile radius of the city of Oakland, where our research center was located at the time of the study. Of the 50 cities, 11 cities met this criterion. To maximize variation in youth exposure to tobacco outlets in their living environments, we first stratified these cities based on measures of socioeconomic status (SES) (i.e., a measure derived from: median household income, percentage of population with a college education, and percentage of population unemployed) and tobacco outlet density (i.e., number of licensed tobacco outlets per 10,000 persons) and then randomly selected 8 cities representing low versus high SES and low versus high tobacco outlet density. We recruited participants who lived in these 8 cities or in cities that were within a 10-mile buffer of the 8 cities.

At the time of data collection (February 2017 – May 2018), California had raised the minimum tobacco sales age to 21, which applied to all cities in the study. Also, in all 8 cities, there were no local policies restricting point of sale tobacco promotions at the time of the study. Self-service displays of tobacco products were prohibited by state law though there are some exceptions (e.g., pipe tobacco, snuff).²⁴ Of the study cities, four implemented one or more local tobacco sales policies including stronger tobacco retailer licensing, tobacco retailer location restrictions, licensing ordinances that include

emerging products (e.g., e-cigarettes), prohibitions on sales of flavored tobacco products, or prohibitions on sales of single cigars.²⁵

We used a multi-tiered approach to recruit youth participants (50% past month tobacco users). We recruited participants through internet and social media advertisements including Craigslist, Facebook, Twitter and Myspace. Also, participants were recruited through flyers distributed to youth organizations in the study cities and by referral. Potential participants were screened for eligibility (i.e., age, city of residence, speak English, and tobacco use). Researchers obtained parental consent for participation for those younger than 18 years old. All participants provided signed consent or assent to participate in the research. The Pacific Institute for Research and Evaluation (PIRE) institutional review board (Federal-wide Assurance #FWA00003078) approved the study prior to implementation.

Procedures

At the beginning of the study, participants completed an online initial survey (30 min) which included questions about demographic characteristics, personal risk factors, and tobacco use and beliefs. Using GPS-enabled smartphones with a survey application, participants then responded to brief daily surveys and location coordinates (latitude and longitude) were obtained at one-minute intervals for 14 days. Research team members provided GPS-enabled phones to participants and briefed them about study procedures. Participants completed the daily surveys using a phone survey application programed to send reminders to complete the survey each evening at 8pm. Youth had a 3-hour window to respond to the survey each day.

As compensation for their participation, participants received \$10 for completing the initial survey, \$5 for each daily survey, and a \$20 bonus if they completed all surveys. Additionally, they received \$40 for return of the phone at the end of the study and \$10 for return of the charger. Participants could use the phones with unlimited voice and text during the study. All participants received a resource card upon completion of the study.

Analytical sample

We excluded geocoded data for days in which participants were tracked for less than 360 min (n=76). Overall, of the 1483 days, 363 days were missing on observed exogenous variables (i.e., control variables and daily neighborhood exposures) and 170 days missing on the endogenous variable (i.e., daily tobacco use) and excluded. The final analytic sample therefore included 950 days, which were clustered within 85 participants. After excluding days with missing data, each participant in the study had, on average, 11.12 days of data (SD: 2.52).

Measures

Daily tobacco use

Our main outcome of interest was tobacco use, which was measured daily. Participants were asked, “Since this time yesterday, did you...”

- 1) smoke at least one cigarette?
- 2) smoke any cigar, cigarillo, or little cigar?
- 3) use chewing tobacco snuff or dip, such as Redman, Levi Garrett, Beechnut, Skoal, Skoal Bandits, or Copenhagen?
- 4) use an e-cigarette or vape device, including e-pen, vape pen, cigalikes, e-hookah, personal vaporizers, or mods to get nicotine?
- 5) smoke part or all of a cigar, cigarillo or little cigar with marijuana in it (a blunt)?

We included blunt use in our measure since 1) blunts contain nicotine even if all of the tobacco filler is removed from the cigar²⁶ and 2) cigars, cigarillos, or little cigars used to make blunts are sold in tobacco outlets. For each question, participants could respond yes (coded as 1) or no (coded as 0). If a participant reported use of any of those products on a specific day, they were coded as having used tobacco on that day.

Daily exposure to peers and adults who use tobacco

To determine whether participants were exposed to *peers* who used tobacco each day, we asked, “Since this time yesterday, did you see people your age who were smoking cigarettes, vaping e-cigarettes, or using any other tobacco or nicotine product?” To determine whether participants were exposed to

adults who used tobacco, we asked “Since this time yesterday, did you see adults (other than your parents/guardians) smoking cigarettes, vaping e-cigarettes, or using any other tobacco or nicotine products?” For both questions, participants could respond yes (coded as 1) or no (coded as 0).

Exposure to tobacco outlets

Using the Dun & Bradstreet, INC (D&B) commercial list, the North American Industry Classification System (NAICS) codes were used to identify probable tobacco outlets in the 8 cities and within 10-mile buffer of city boundaries. Specifically, probable tobacco outlets were searched using the NAICS codes of the top ten retail industries that sell tobacco products including supermarkets and other grocery (except convenience) stores (445110), convenience stores (445120), tobacco stores (453991), gasoline station with convenience stores (447110), warehouse clubs and supercenters (452910), news dealers and newsstands (451212), beer, wine, and liquor stores (445310), pharmacies and drug stores (446110), discount department stores (452112) and other gasoline stations (447190). These codes are industries that represent approximately 98% of all tobacco sales and were used in a study that validated the use of commercial lists to identify tobacco outlets in states that do not have a comprehensive list of tobacco outlet addresses.²⁷ Given that none of the selected cities prohibited sales of tobacco products in pharmacies, the NAICS code for pharmacies and drug stores were included for all cities. Next, chains with policies restricting the sale of tobacco (e.g., Target and CVS) were excluded from the list. To ensure our study included places that sell alternative nicotine delivery systems (ANDS), but do not sell other tobacco products (e.g., hookah bars), we conducted an online search of places that sell hookahs and e-cigarettes in each city. All identified tobacco outlets were contacted by phone to verify business status, sales of tobacco products, address, and hours of operation. Next, these outlets were visited by observers to record outlet GPS point locations (i.e. latitude and longitude) and obtain data about tobacco products and marketing (not reported in the current study).

Tobacco outlet addresses and participants’ GPS locations were geocoded and activity spaces were constructed by joining sequential GPS points into a polyline, which was then buffered and overlaid with tobacco outlet locations. An example of a participant’s activity space can be seen in Supplementary

Figure A. Exposure measures included the number of tobacco outlets within 1) 50m or 2) 100m of these polylines each day, as well as the number of minutes participants were within 3) 50m or 4) 100m of tobacco outlet each day. The downloaded GPS data had a field of accuracy of each point. The average accuracy was 20m. To minimize potential errors, we used 50- and 100-meter buffers. All four exposure measures were weighted by the time participants were within the study area. We ran separate models for each of these exposures since number of tobacco outlets within 50m or 100m of polylines were highly correlated with one another ($r= 0.96, p<0.001$), as were time spent within 50m or 100m of tobacco outlets ($r= 0.73, p<0.001$).

Control variables

Control variables included sex assigned at birth (male, female, or intersex), race (White or non-White), ethnicity (Latino or non-Latino), age group (less than 18 or 18+), and perceived SES with the item, “Compared with other people in America, how rich or poor do you consider yourself?” Respondents could answer on a likert scale (1=rich to 7=poor). Researchers have found that perceived SES is associated with health behaviors and health outcomes.²⁸ All control variables were assessed at baseline.

Data analysis

We first examined means, standard deviations, or frequencies of all variables using SAS version 9.4 (SAS Inc., Cary, NC, USA). We then used multilevel structural equation modeling (MSEM) with MPlus Version 7 for the mediation analysis. As the data was cross-sectional and not able to provide temporal relationships for cause and effect we have used the term "potentially mediate" throughout the article. We used MSEM for three reasons: 1) we had longitudinal data in which days over time were clustered within individuals and MSEM models may decrease bias of effects compared to traditional multilevel mediation approaches;²⁹ 2) our potential mediators and outcomes were binary, which many traditional multilevel mediation approaches cannot easily accommodate;²⁹ and 3) we were interested in examining two potential mediators simultaneously, which is relatively easy with MSEM.³⁰

Since our outcome and potential mediators were binary, we analyzed data using the probit link, which is used to transform outcome probabilities to the standard normal variable.³¹ The probit regression

coefficients give the change in the z-score or probit index for a one unit change in the predictor.³² Since the MSEM included binary outcomes, no model fit data were reported by Mplus. For all analyses, we set critical $\alpha = .05$ and used 2-tailed statistical tests.

RESULTS

Sample characteristics

Of the 85 participants, more than half identified as White ($n=48$, 56.47%) and one-fifth identified as Hispanic or Latino ($n=19$, 22.35%). A slight majority of the sample was female ($n=53$, 62.35%) and younger than 18 ($n=52$, 61.18%) (Table 1).

Of the 950 study days, participants reported using a tobacco product on 24.63% of days ($n=234$ days), seeing a peer use tobacco on 35.30% of days ($n=335$ days), and seeing an adult use tobacco on 45.09% of days ($n=427$ days). In addition, on average participants were exposed to 2.95 tobacco outlets within 50m of polylines (SD: 3.65) and 4.40 tobacco outlets within 100m of polylines (SD: 5.03) per day. Finally, participants spent, on average, 7.55 minutes within 50m of tobacco outlets (SD: 19.95) and 17.41 minutes within 100m of tobacco outlets (SD: 48.61) per day (Table 1).

Descriptive associations

Of the 234 days in which participants reported using a tobacco product, most days (72.65%) participants also reported seeing a peer use tobacco (Table 2). Participants were therefore more likely to report using tobacco on days in which they saw a peer tobacco user ($p<0.001$) than on days in which they did not see a peer tobacco user. Similar results were found for exposure to adult tobacco users. In addition, we found that number of tobacco outlets within 50m and 100m of activity space polylines was significantly associated with exposure to peer and adult tobacco users (see Supplementary Table A for these results).

Mediation

As seen in Figure 1a and 1b, our SEM contains two main pathways: 1) the pathway from tobacco retail outlet exposure to the potential mediators and 2) the pathway from the potential mediators to tobacco use.

Exposure to number of tobacco outlets within 50m and 100m of activity space polylines

Controlling for demographics, we found no association between daily tobacco use and number of tobacco outlets within 50m (probit regression coefficient: 0.01, $p=0.82$, Figure 1a) and 100m (probit regression coefficient: 0.01, $p=0.62$, Figure 1b) of activity space polylines. However, we found evidence of an indirect effect ($p=0.001$ for both the 50m and 100m models) through daily exposure to peers who use tobacco. Specifically, greater daily exposure to tobacco outlets within 50m and 100m of activity polylines was positively associated with seeing peers use tobacco on a given day (probit regression coefficient: 0.10, $p<0.001$; probit regression coefficient: 0.08, $p<0.001$, respectively). In turn, seeing peers use tobacco on a given day was positively associated with tobacco use on that day in both the 50m and 100m models (probit regression coefficient: 2.23, $p<0.001$; probit regression coefficient: 2.24, $p<0.001$, respectively).

While increased daily exposure to tobacco outlets within 50m and 100m of activity space polylines was also positively associated with daily exposure to adults who use tobacco (probit regression coefficient: 0.09, $p=0.002$; probit regression coefficient: 0.07, $p=0.001$, respectively), there was no association between daily exposure to adults who use tobacco and tobacco use on that day in the 50m and 100m models (probit regression coefficient: -0.13, $p=0.78$; probit regression coefficient: -0.15, $p=0.73$). There was, therefore, no indirect effect through daily exposure to adults who use tobacco ($p=0.78$ and $p=0.73$ for the 50m and 100m models, respectively).

Time spent within 50m and 100m of tobacco outlets

Controlling for demographics, we found no association between daily tobacco use and daily number of minutes participants were within 50m of tobacco outlets (probit regression coefficient: 0.01, $p=0.34$, Figure 2a) and daily number of minutes participants were within 100m of tobacco outlets (probit regression coefficient: 0.002, $p=0.65$, Figure 2b), nor evidence of an indirect effect ($p=0.66$; $p=0.41$, respectively) through daily exposures to peers who use tobacco. Specifically, increased number of minutes participants were within 50m and 100m of tobacco outlets was not associated with seeing peers use tobacco on a given day (probit regression coefficient: -0.003, $p=0.65$; probit regression coefficient: -

0.002, $p=0.40$). However, seeing peers use tobacco on a given day was positively associated with tobacco use on that day in both the 50m and 100m models (probit regression coefficient: 2.24, $p<0.001$; probit regression coefficient: 2.24, $p<0.001$).

Similarly, greater time participants spent within 50m and 100m of tobacco outlet was not associated with daily exposure to adults who use tobacco (probit regression coefficient: 0.01, $p=0.09$; probit regression coefficient: 0.00, $p=0.89$, respectively) and there was no association between daily exposure to adults who use tobacco and tobacco use on that day (probit regression coefficient: -0.15, $p=0.73$; probit regression coefficient: -0.17, $p=0.71$, respectively). There was, therefore, no indirect effect through daily exposure to adults who use tobacco ($p=0.74$; $p=0.90$ for the 50m and 100m models, respectively).

DISCUSSION

Extending previous findings,³⁻⁸ the current study is one of the first to examine what factors may mediate association between exposure to tobacco outlets and tobacco use and also one of the first to use time-ordered data. We found that daily exposure to tobacco outlets within activity spaces affects youth tobacco use through exposure to peers who use tobacco, but not adults. Implications for research, practice, and policy are discussed below.

In contrast to at least one previous exploratory study that used a similar population and activity space measures,¹⁰ we found no *direct* effect of exposure to tobacco outlets on daily tobacco use. It is possible that exposure to tobacco outlets promotes youth tobacco use in a more long-term cumulative manner, such as by increasing receptivity to tobacco products via advertising and promotion and increasing perceptions of the prevalence and acceptability of tobacco use in the general population. It is also likely that the mechanism through which exposure to tobacco outlets influences tobacco use depends on the population (e.g., smokers or non-smokers) and behaviors (e.g., tobacco use initiation, cessation) being examined. For instance, in studies of smokers and attempting quitters, researchers found that exposure to tobacco outlets can trigger lapses in quitting, increase impulse tobacco purchases, and reduce self-efficacy to quit smoking.³³⁻³⁵ Our study included both current and never tobacco users. It is therefore

possible that exposure to tobacco outlets would therefore not have had an immediate or direct effect on *daily* tobacco use.

Interestingly, we found that exposure to peers who use tobacco, but not adults, was associated with daily tobacco use. Social Cognitive Theory suggests that adolescents learn about tobacco use by observing peers and are reinforced to use tobacco if they gain acceptance from peers or establish a particular social identity.^{2,17,18,36} A large body of research has established that peers' tobacco use is an important determinant of youth and young adult tobacco use.^{1,20,37-39} Research also suggests that adolescents often obtain tobacco products from peers⁴⁰—therefore affording another mechanism through which peer tobacco use influences youth tobacco use. Finally, during adolescence and young adulthood, there is an increasing desire to fit in with peers, which can increase the influence of peers' substance use.^{38,41} It is important to note that our measure of 'exposure to peers who use tobacco' could have been a proxy for tobacco use norms.⁴² It is also possible that our measure of exposure to peers who use tobacco could have acted as a cue to use tobacco or that youth were going to tobacco outlets to acquire tobacco with their friends, since many youth use tobacco socially.⁴³ Regardless, it appears that exposure to peers who use tobacco is an important determinant of youth tobacco use and future research could be conducted to disentangle what this measure is capturing.

Contrary to previous research suggesting the importance of adult tobacco use as a determinant of youth tobacco use,^{19,44} we found no associations between exposure to adult tobacco users and daily tobacco use. It is possible that youth exposure to peer tobacco users increases opportunities to obtain and use tobacco with peers, whereas exposure to adult tobacco users in the community has more overall but not daily effects on youth tobacco use. Peer tobacco use would therefore be related to participants' momentary tobacco use, as we examined in the current study, whereas adult tobacco use would be more distally related to use.

We also found that exposure to number of tobacco outlets was associated with daily exposure to both peer and adult tobacco users. From previous research we know that the prevalence of cigarette smoking is higher in neighborhoods with tobacco outlets than in neighborhoods without any tobacco

outlets.³ Moreover, another study found that teens perceive more adults to be smoking if they live in a neighborhood with more tobacco outlets.⁴⁵ This is the first study, to our knowledge, that directly examines the link between exposure to tobacco outlets and exposure to tobacco users, particularly by using GEMA data. Interestingly, we found no association between *time spent* near tobacco outlets and exposure to peer and adult tobacco users suggesting that exposure effects are due to increased opportunities to see or obtain tobacco from peers rather than the time of exposure.

Results suggest that policies to reduce exposure to tobacco outlets and interventions to address social influences of tobacco use could be used to reduce youth tobacco use. Importantly, our analysis allowed us to investigate exposure to tobacco outlets, exposure to peer and adult users, and tobacco use in a time ordered design that suggests these may be causal effects. Communities in the US have already begun to regulate density and number of tobacco retailers using various strategies (e.g., prohibiting sales in certain venues like pharmacies, prohibiting sales near youth populated areas, such as schools).⁴⁶ Our findings provide additional significant support for policymakers considering these options. In addition, interventions focusing on peer tobacco use as a determinant of youth tobacco use will continue to be important and should consider how youths' daily environment increases opportunities for such social influences. For example, a phone application could send a comic strip with an anti-tobacco message from peers when youth are near tobacco outlets. To this point, some researchers have begun using “just-in-time adaptive interventions”⁴⁷—to reduce tobacco use and smoking cravings.⁴⁸

Limitations

We acknowledge a number of limitations. First, we relied on self-reported measures of tobacco use and exposure to peers and adults who use tobacco. Second, we cannot establish that exposure to tobacco outlets came before exposure to people who use tobacco or that exposure to people who use tobacco came before daily tobacco use; although using real time-ordered data helps lessen this concern. Third, we only examined two possible mediators—exposure to peers and adults using tobacco. There are other plausible mechanisms through which exposure to tobacco outlets influences youth tobacco use. Fourth, we did not control for other factors that may have influenced youth tobacco use, e.g., family

history of tobacco use, stress. Fifth, it is possible that local policies in some of the cities influenced daily tobacco use. However, the intraclass correlation for daily tobacco use by city was 0.0436 and the between-city variance in daily tobacco use was not significant, which suggests that participants' cities had small effects on daily tobacco use. Finally, data came from a convenience sample of youth in California and results may not generalize to other populations or locations.

Conclusions

Results of the current suggest that exposure to tobacco outlets within activity spaces affects youth tobacco use through exposure to peers who use tobacco. Building on these findings, future research could examine other mechanisms through which tobacco retail outlets influence tobacco use. Policies and interventions to regulate both tobacco retail outlet density and social influences of tobacco use may be helpful in preventing and reducing youth tobacco use.

Acknowledgements

Funding: This research and preparation of this manuscript were supported by grant 25IR-0029 from the California Tobacco-Related Disease Research Program (TRDRP) and grant P60-AA006282 from the National Institute on Alcohol Abuse and Alcoholism (NIAAA) of the National Institutes of Health (NIH). The content is solely the responsibility of the authors and does not necessarily represent the official views of the TRDRP, NIAAA or NIH.

Conflicts of Interest: None declared.

References

1. U.S. Department of Health and Human Services. The Health Consequences of Smoking--50 Years of Progress: A Report of the Surgeon General. Rockville, MD. 2014. Available at: <https://www.surgeongeneral.gov/library/reports/50-years-of-progress/>. Accessed January 4, 2016.
2. U.S. Department of Health and Human Services. Preventing tobacco use among youth and young adults: a report of the surgeon general. Rockville, MD. 2012. Available at: <https://www.surgeongeneral.gov/library/reports/preventing-youth-tobacco-use/index.html>. Accessed December 13, 2018.
3. Henriksen L, Feighery EC, Schleicher NC, Cowling DW, Kline RS, Fortmann SP. Is adolescent smoking related to the density and proximity of tobacco outlets and retail cigarette advertising near schools? *Prev Med*. Aug 2008;47(2):210-214.
4. Henriksen L, Feighery EC, Wang Y, Fortmann SP. Association of retail tobacco marketing with adolescent smoking. *Am J Public Health*. Dec 2004;94(12):2081-2083.
5. Henriksen L, Schleicher NC, Feighery EC, Fortmann SP. A longitudinal study of exposure to retail cigarette advertising and smoking initiation. *Pediatrics*. Aug 2010;126(2):232-238.
6. Loomis BR, Kim AE, Busey AH, Farrelly MC, Willett JG, Juster HR. The density of tobacco retailers and its association with attitudes toward smoking, exposure to point-of-sale tobacco advertising, cigarette purchasing, and smoking among New York youth. *Prev Med*. Nov 2012;55(5):468-474.
7. Shortt NK, Tisch C, Pearce J, Richardson EA, Mitchell R. The density of tobacco retailers in home and school environments and relationship with adolescent smoking behaviours in Scotland. *Tob Control*. Jan 2016;25(1):75-82.
8. Paynter J, Edwards R, Schluter PJ, McDuff I. Point of sale tobacco displays and smoking among 14-15 year olds in New Zealand: a cross-sectional study. *Tob Control*. Aug 2009;18(4):268-274.
9. Finan LJ, Lipperman-Kreda S, Abadi M, et al. Tobacco outlet density and adolescents' cigarette smoking: a meta-analysis. *Tob Control*. Mar 8 2018.
10. Lipperman-Kreda S, Morrison C, Grube JW, Gaidus A. Youth activity spaces and daily exposure to tobacco outlets. *Health Place*. Jul 2015;34:30-33.
11. Zenk SN, Schulz AJ, Matthews SA, et al. Activity space environment and dietary and physical activity behaviors: a pilot study. *Health Place*. Sep 2011;17(5):1150-1161.
12. Rainham DG, Bates CJ, Blanchard CM, Dummer TJ, Kirk SF, Shearer CL. Spatial classification of youth physical activity patterns. *Am J Prev Med*. 2012;42(5):e87-e96.
13. Shareck M, Kestens Y, Vallée J, Datta G, Frohlich KL. The added value of accounting for activity space when examining the association between tobacco retailer availability and smoking among young adults. *Tob Control*. 2015;25(4):406-412.
14. Mason MJ, Mennis J, Zaharakis NM, Way T. The Dynamic Role of Urban Neighborhood Effects in a Text-Messaging Adolescent Smoking Intervention. *Nicotine Tob Res*. May 2016;18(5):1039-1045.
15. Mason M, Mennis J, Way T, et al. Young adolescents' perceived activity space risk, peer networks, and substance use. *Health Place*. Jul 2015;34:143-149.
16. Duncan DT, Kawachi I. Chapter 1: Neighborhoods and health. In: Duncan DT, Kawachi I, eds. *Neighborhoods and Health*. New York: Oxford University Press; 2018.
17. Bandura A. *Social foundations of thought and action: A social cognitive theory*. Englewood Cliffs, NJ: Prentice-Hall; 1986.
18. Fishbein M, Ajzen I. *Predicting and changing behavior: The reasoned action approach*. New York, NY: Psychology Press, Taylor & Francis; 2010.
19. Thrul J, Lipperman-Kreda S, Grube JW, Friend KB. Community-level adult daily smoking prevalence moderates the association between adolescents' cigarette smoking and perceived smoking by friends. *J Youth Adolesc*. Sep 2014;43(9):1527-1535.
20. Villanti A, Boulay M, Juon H-S. Peer, parent and media influences on adolescent smoking by developmental stage. *Addict Behav*. 2011;36(1-2):133-136.

21. Mennis J, Mason M, Ambrus A, Way T, Henry K. The spatial accuracy of geographic ecological momentary assessment (GEMA): Error and bias due to subject and environmental characteristics. *Drug Alcohol Depend.* Sep 01 2017;178:188-193.
22. Lipperman-Kreda S, Mair C, Grube JW, Friend KB, Jackson P, Watson D. Density and proximity of tobacco outlets to homes and schools: relations with youth cigarette smoking. *Prev Sci.* Oct 2014;15(5):738-744.
23. Lipperman-Kreda S, Grube JW, Friend KB, Mair C. Tobacco outlet density, retailer cigarette sales without ID checks and enforcement of underage tobacco laws: associations with youths' cigarette smoking and beliefs. *Addiction.* Mar 2016;111(3):525-532.
24. ChangeLab Solutions. Tobacco Laws Affecting California. 2018. Available at: https://www.changelabsolutions.org/sites/default/files/2018_CA_Law_Booklet_FINAL_20180627.pdf. Accessed March 13, 2019.
25. American Lung Association in California. State of Tobacco Control 2018: California Local Grades. 2018. Available at: <https://www.lung.org/local-content/california/documents/state-of-tobacco-control/2018/2018-sotc-california-full.pdf>. Accessed March 13, 2019.
26. Peters EN, Schauer GL, Rosenberry ZR, Pickworth WB. Does marijuana “blunt” smoking contribute to nicotine exposure?: Preliminary product testing of nicotine content in wrappers of cigars commonly used for blunt smoking. *Drug Alcohol Depend.* 2016;168:119-122.
27. D'Angelo H, Fleischhacker S, Rose SW, Ribisl KM. Field validation of secondary data sources for enumerating retail tobacco outlets in a state without tobacco outlet licensing. *Health Place.* Jul 2014;28:38-44.
28. Goodman E, Adler NE, Kawachi I, Frazier AL, Huang B, Colditz GA. Adolescents' Perceptions of Social Status: Development and Evaluation of a New Indicator. *Pediatrics.* 2001;108(2):e31-e31.
29. Preacher KJ, Zhang Z, Zyphur MJ. Alternative methods for assessing mediation in multilevel data: The advantages of multilevel SEM. *Structural Equation Modeling.* 2011;18(2):161-182.
30. Tofighi D, Thoemmes F. Single-level and multilevel mediation analysis. *The Journal of Early Adolescence.* 2014;34(1):93-119.
31. Gunzler D, Chen T, Wu P, Zhang H. Introduction to mediation analysis with structural equation modeling. *Shanghai archives of psychiatry.* 2013;25(6):390.
32. UCLA Institute for Digital Research and Education. Probit Regression—MPlus Data Analysis. Available at: <https://stats.idre.ucla.edu/mplus/dae/probit-regression/>. Accessed October 5, 2018.
33. Burton S, Spanjaard D, Hoek J. An investigation of tobacco retail outlets as a cue for smoking. *Australasian Marketing Journal (AMJ).* 2013;21(4):234-239.
34. Kirchner TR, Cantrell J, Anesetti-Rothermel A, Ganz O, Vallone DM, Abrams DB. Geospatial exposure to point-of-sale tobacco: real-time craving and smoking-cessation outcomes. *Am J Prev Med.* Oct 2013;45(4):379-385.
35. Mennis J, Mason M. Tobacco outlet density and attitudes towards smoking among urban adolescent smokers. *Subst Abus.* 2016;37(4):521-525.
36. Bandura A. *Social Learning Theory.* Englewood Cliffs, NJ:: Prentice Hall; 1977.
37. Liu J, Zhao S, Chen X, Falk E, Albarracín D. The influence of peer behavior as a function of social and cultural closeness: A meta-analysis of normative influence on adolescent smoking initiation and continuation. *Psychol Bull.* 2017;143(10):1082.
38. Van Ryzin MJ, Fosco GM, Dishion TJ. Family and peer predictors of substance use from early adolescence to early adulthood: An 11-year prospective analysis. *Addict Behav.* 2012;37(12):1314-1324.
39. Powell LM, Tauras JA, Ross H. The importance of peer effects, cigarette prices and tobacco control policies for youth smoking behavior. *J Health Econ.* 2005;24(5):950-968.
40. Meyers MJ, Delucchi K, Halpern-Felsher B. Access to tobacco among California high school students: the role of family members, peers, and retail venues. *J Adolesc Health.* 2017;61(3):385-388.

41. Spear LP. The adolescent brain and age-related behavioral manifestations. *Neurosci Biobehav Rev.* Jun 2000;24(4):417-463.
42. Eisenberg ME, Forster JL. Adolescent smoking behavior: measures of social norms. *Am J Prev Med.* 2003;25(2):122-128.
43. Pepper JK, Coats EM, Nonnemaker JM, Loomis BR. How Do Adolescents Get Their E-Cigarettes and Other Electronic Vaping Devices? *Am J Health Promot.* Aug 1 2018:890117118790366.
44. Richardson A, Williams V, Rath J, Villanti AC, Vallone D. The next generation of users: prevalence and longitudinal patterns of tobacco use among US young adults. *Am J Public Health.* 2014;104(8):1429-1436.
45. Schleicher NC, Johnson TO, Fortmann SP, Henriksen L. Tobacco outlet density near home and school: Associations with smoking and norms among US teens. *Prev Med.* Oct 2016;91:287-293.
46. Ackerman A, Etow A, Bartel S, Ribisl KM. Reducing the density and number of tobacco retailers: policy solutions and legal issues. *Nicotine Tob Res.* 2017;19(2):133-140.
47. Nahum-Shani I, Smith SN, Spring BJ, et al. Just-in-time adaptive interventions (JITAIs) in mobile health: key components and design principles for ongoing health behavior support. *Ann Behav Med.* 2017;52(6):446-462.
48. Cerrada CJ, Dzibur E, Blackman KCA, Mays V, Shoptaw S, Huh J. Development of a just-in-time adaptive intervention for smoking cessation among Korean American emerging adults. *Int J Behav Med.* 2017;24(5):665-672.

Table 1. Participant Characteristics, n=85 participants and 950 days

	N	%	Mean	SD	Range
Individual-level (n=85)					
Race					
White	48	56.47%	--	--	--
Non-White	37	43.53%	--	--	--
Ethnicity					
Hispanic or Latino	19	22.35%	--	--	--
Not Hispanic or Latino	66	77.65%	--	--	--
Sex at birth					
Male	32	34.43%	--	--	--
Female	53	65.57%	--	--	--
Age					
Younger than 18	52	61.18%	--	--	--
18 or older	33	38.82%	--	--	--
Perceived SES	85	--	4.22	1.46	1-7 (higher values = more poor)
Mean number of days participants used tobacco over the study period	85	--	2.99	4.26	0-13
Mean number of days participants were exposed to peers who use tobacco over the study period	85	--	4.21	4.16	0-13
Mean number of days participants were exposed to adults who use tobacco over the study period	85	--	5.45	4.05	0-14
Mean tobacco outlets within 50m of activity space polylines per day	85	--	2.80	1.99	0-9
Mean number of tobacco outlet within 100m of activity space polylines per day	85	--	4.22	2.86	0-14
Mean amount of time spent within 50m of tobacco outlets per day (in minutes)	85	--	7.12	10.27	0-74
Mean amount of time spent within 100m of tobacco outlets per day (in minutes)	85	--	16.48	31.38	0-273
Day-level (n=950)					
Daily tobacco use					
Days in which participants did not use tobacco	716 days	75.37%	--	--	--
Days in which participants did use tobacco	234 days	24.63%	--	--	--
Daily exposure to peers who use tobacco					
Days in which participants were not exposed to peers who use tobacco	614 days	64.70%	--	--	--
Days in which participants were exposed to peers who use tobacco	335 days	35.30%	--	--	--

Daily exposure to adults who use tobacco					
Days in which participants were not exposed to adults who use tobacco	520 days	54.91%			
Days in which participants were exposed to adults who use tobacco	427 days	45.09%	--	--	--
Mean number of tobacco outlets within 50m of activity space polylines per day	950 days	--	2.95	3.65	0-21 (higher values = increasing exposure)
Mean number of tobacco outlets within 100m of activity space polylines per day	950 days	--	4.40	5.03	0-27 (higher values = increasing exposure)
Mean amount of time spent within 50m of tobacco outlets per day (in minutes)	950 days	--	7.55	19.95	0-339 (higher values = increasing exposure)
Mean amount of time spent within 100m of tobacco outlets per day (in minutes)	950 days	--	17.41	48.61	0-573 (higher values = increasing exposure)

Table 2. Descriptive associations among daily exposure to peer tobacco product users, daily exposure to adult tobacco product users, and daily tobacco use, n=950 days

Variable	Days in which participants did not use tobacco n / column n (%)	Days in which participants used tobacco n / column n (%)	P-value ^b
Daily exposure to peer tobacco users ^a			
Days in which participants were not exposed to peers who use tobacco	550 / 715 days (76.92%)	64 / 234 days (27.35%)	p<0.001
Days in which participants were exposed to peers who use tobacco	165 / 715 days (23.08%)	170 / 234 days (72.65%)	
Daily exposure to adult tobacco users ^a			
Days in which participants were not exposed to adults who use tobacco	435 / 714 days (60.92%)	85 / 233 days (36.48%)	p<0.001
Days in which participants were exposed to adults who use tobacco	279 / 714 days (39.08%)	148 / 233 days (63.52%)	
^a Column percentages are provided in parentheses			
^b P-values were calculated using chi-square tests			

Figure 1a and 1b. Mediation results from models in which daily exposure to number tobacco outlets within 50m of polylines was modeled as the exposure (Figure 1a) and daily exposure to number of tobacco outlets within 100m of polylines was modeled as the exposure (Figure 1b)

Figure 1a






Figure 1b



Notes: In Figure 1a, the Akaike information criterion value was 2358.47 and the Bayesian information criterion value was 2484.73. The indirect effect through exposure to peers who use tobacco was significant at $p = 0.001$. The indirect effect through exposure to adults who use tobacco was not significant at $p = 0.78$. In Figure 1b, the Akaike information criterion value was 2356.07 and the Bayesian information criterion value was 2482.34. Indirect effect through exposure to peers who use tobacco was significant at $p = 0.001$. The indirect effect through exposure to adults who use tobacco was not significant at $p = 0.73$.

Figure 2a and 2b. Mediation results from models in which daily time spent within 50 meters of tobacco outlet was modeled as the exposure (Figure 2a) and daily time spent within 100 meters of tobacco outlet polylines was modeled as the exposure (Figure 2b)



Notes: In Figure 2a, the Akaike information criterion value was 2393.64 and the Bayesian information criterion value was 2520.24. The indirect effect through exposure to peers who use tobacco was not significant at $p = 0.66$. The indirect effect through exposure to adults who use tobacco was not significant at $p = 0.74$. In Figure 2b, the Akaike information criterion value was 2382.69 and the Bayesian information criterion value was 2508.96. The indirect effect through exposure to peers who use tobacco was not significant at $p = 0.41$. The indirect effect through exposure to adults who use tobacco was not significant at $p = 0.90$.