

# UC Berkeley

## UC Berkeley Previously Published Works

### Title

Epigenetic control of embryo-uterine crosstalk at peri-implantation.

### Permalink

<https://escholarship.org/uc/item/1j69m4mv>

### Journal

Cellular and Molecular Life Sciences, 76(24)

### Authors

Kong, Shuangbo

Zhou, Chan

Bao, Haili

et al.

### Publication Date

2019-12-01

### DOI

10.1007/s00018-019-03245-8

Peer reviewed



# Epigenetic control of embryo–uterine crosstalk at peri-implantation

Shuangbo Kong<sup>1,2</sup> · Chan Zhou<sup>2</sup> · Haili Bao<sup>2</sup> · Zhangli Ni<sup>2</sup> · Mengying Liu<sup>2</sup> · Bo He<sup>2</sup> · Lin Huang<sup>1,2</sup> · Yang Sun<sup>1,2</sup> · Haibin Wang<sup>1,2</sup> · Jinhua Lu<sup>1,2</sup>

Received: 21 November 2018 / Revised: 28 June 2019 / Accepted: 23 July 2019 / Published online: 27 July 2019  
© Springer Nature Switzerland AG 2019

## Abstract

Embryo implantation is one of the pivotal steps during mammalian pregnancy, since the quality of embryo implantation determines the outcome of ongoing pregnancy and fetal development. A large number of factors, including transcription factors, signalling transduction components, and lipids, have been shown to be indispensable for embryo implantation. Increasing evidence also suggests the important roles of epigenetic factors in this critical event. This review focuses on recent findings about the involvement of epigenetic regulators during embryo implantation.

**Keywords** Implantation · Epigenetic regulation · Uterine receptivity · Blastocyst activation decidualization

## Introduction

Embryo implantation, the process during which blastocyst embed into the maternal uterine wall, is a critical step during mammalian pregnancy, and any disturbance to implantation will lead to pregnancy-related complications, such as infertility and recurrent pregnancy loss. Implantation only occurs within a limited time duration, which is defined as the window of implantation. During this window, the uterus is in the receptive state, and the blastocyst, must acquire implantation competency, a state of embryo differentiation that can initiate crosstalk with uterine cells to implanting into the uterus for further in vivo development of the embryo. This concept of uterine receptivity was first raised and established by studies employing the embryo transfer technique in the 1960s [1]. Based on the previous findings, uterine sensitivity to implantation-competent blastocysts is classically divided into three stages: pre-receptive, receptive,

and refractory. During the pre-receptive stage, the uterus is favourable for embryo development, but not suitable for implantation. The uterus at the receptive stage can initiate implantation when there are competent blastocysts. During the refractory stage, however, implantation-competent blastocysts cannot implant in the uterus and the uterus is even adverse to the survival and development of blastocysts [2]. The uterus is mainly composed of three different cell types, the epithelium, stroma, and myometrium. The coordination of these different endometrial cell types makes the uterus to be receptive mainly underling the regulation of progesterone (P4) and estrogen (E2) [3]. Before implanting into the uterus, the embryo must develop to form a blastocyst, which comprises of the outer trophoctoderm (TE), and the inner cell mass (ICM), which further differentiate into the primitive endoderm and epiblast (Fig. 1). The reciprocal embryo transfer experiments using the delayed implantation model also indicated that blastocyst state of activation is also a critical determinant for implantation in the receptive uterus [4].

During peri-implantation, different uterine cell types respond to changing E2 and P4 levels specifically. On day 1 (day 1 = see vaginal plug) of pregnancy in mice, the uterine epithelium undergoes extensive proliferation under the influence of a preovulatory E2 surge. From day 3 onward, P4 secreted by the newly formed corpus luteum inhibits epithelium proliferation, and initiates stroma cell proliferation. On day 4, a pre-implantation ovarian E2 surge superimposes on P4 signalling to induce the establishment of uterine receptivity and the acquisition of implantation competency


✉ Haibin Wang  
haibin.wang@vip.163.com

✉ Jinhua Lu  
jinhua888@126.com

<sup>1</sup> Reproductive Medical Center, The First Affiliated Hospital of Xiamen University, Xiamen 361003, Fujian, People's Republic of China

<sup>2</sup> Fujian Provincial Key Laboratory of Reproductive Health Research, School of Medicine, Xiamen University, Xiamen 361102, Fujian, People's Republic of China

**Fig. 1** Diagram of embryo implantation in mice. For successful implantation, the embryo at blastocyst stage must acquire implantation competency through a process defined as blastocyst activation, and the uterus, mainly involved the epithelium and stroma, must differentiate into a unique state—uterine receptivity to be conducive for embryo implantation. The synchronization between the blastocyst activation and uterine receptivity determined the window of implantation. TE trophoblast, EPI epiblast, PE primitive endoderm, LE luminal epithelium, GE gland epithelium, S stroma



of blastocysts. Ovariectomy on day 4 morning, before this pre-implantation E2 surge, induces delayed implantation and embryonic diapause, a state that can be maintained by daily P4 injection from day 5 and terminated by the re-supplement of E2 [3, 5]. Similar fluctuation of P4 and E2 hormones is also observed in humans, with high levels of E2 in the proliferation stage and increased P4 levels in the secretory phase, of which the window in human is usually regarded as 5–7 days post P4 stimulation [6]. It is still unclear whether embryonic diapause also occurs in humans.

The process of implantation is divided into three stages: apposition, attachment (adhesion), and penetration. During the apposition stage, blastocysts are distributed evenly along the whole uterine horn in rodents, and the trophoblast layer of the embryo becomes closely apposed to the uterine luminal epithelium. At the attachment stage, the association of the trophoblast and luminal epithelium is strong enough to resist dislodging of the blastocyst when the uterine lumen is flushed. The first sign of the attachment reaction, which occurs at day 4 night in mice, coincides with a localized increase in stromal vascular permeability at the site of blastocyst attachment and can be visualized by tail vein injection of blue dye before sacrifice. Penetration is the process by which the embryo invades into the uterine stroma through the luminal epithelium and basal lamina to establish a vascular relationship with the uterus. After embryo implantation, the uterus undergoes significant changes to support the further development of the embryo until the establishment of a functional placenta; specifically, the uterine stromal cells proliferate and differentiate to form the decidual cells, a process termed stromal–decidual transformation (also called decidualization) [5, 7]. While embryo implantation

is a dynamic process controlled strictly by molecules such as transcription factors and signalling pathway components, it is also regulated by epigenetic factors.

## Aspects of epigenetic regulation

Unlike genetic regulation which functions by altering the DNA sequence, epigenetic regulation is defined as stably heritable changes in a chromosome without changing the DNA sequence [8]. Epigenetic processes mainly include DNA methylation, histone modification, and non-coding RNAs, which can influence the expression of target genes [9].

### DNA methylation

DNA methylation, the best-characterized epigenetic modification, is achieved by adding a methyl group (–CH<sub>3</sub>) to the fifth carbon of the cytosine ring to form methyl cytosine, during which S-adenosylmethionine acts as the methyl donor [10]. DNA methylation is catalysed by the enzymes DNA methyltransferases (DNMTs): DNMT1, DNMT3A, DNMT3B, and DNMT3L. DNMT1 is a methylation-maintaining enzyme responsible for restoring the hemi-methylated sites to full methylation after DNA replication, whereas DNMT3A and DNMT3B are mainly involved in methylating new sites, called de novo methylation [11]. DNMT3L, without DNA methyltransferase activity in itself, is reported to regulate DNA methylation [12, 13]. In humans and other mammals, DNA methylation occurs predominantly on the cytosine base of a cytosine guanine (CG) dinucleotide in the

DNA sequence [14]. These CG dinucleotides are often clustered in small stretches of DNA, called CpG islands, which are often associated with promoter regions and at the 5' end of the gene [15]. In general, CpG islands in gene promoters are unmethylated, which allows active gene transcription, and methylated CpG islands associated with gene promoter regions would silence gene expression. However, most CpG sites outside of CpG islands are methylated, suggesting its role during the global maintenance of the genome [14, 16]. All of epigenetic modifications should be reversed biochemically, and many years after the finding of DNMT, the TET family proteins have been shown to be responsible for DNA demethylation [17]. TET proteins are capable of catalysing the conversion of 5mC into 5hmC, and then 5hmC serves as an intermediate product for DNA demethylation in an indirect demethylation manner through passive or active pathways [17].

### Histone modification

Chromatin is mainly composed of genomic DNA and the nucleosome. The nucleosome is a histone octamer consisting of two copies of each of the four core histones (H2A, H2B, H3, and H4), around which approximately 146 bp of the DNA is wrapped. These nucleosomes are linked by loops of DNA and the linker histone H1 to form chromatin. The N terminal tails of the histones protrude from the chromatin to be subjected to post-translational modification. It has been shown that nucleosomal histones are subject to numerous covalent modifications, including methylation, acetylation, phosphorylation, sumoylation, glycosylation, and ubiquitination, at specific tails of selected amino acids. A variety of enzymes are involved in the modification of histone tails, including histone methyltransferases (HMTs), acetyltransferases (HATs), kinases, and ubiquitin ligase functioning as the “writer”, as well as the eraser such as histone demethylases, deacetylases (HDACs), and phosphatases, which are able to remove the modification marks from the histone tails [18]. These modifications impose either transcriptionally repressive or transcriptionally permissive chromatin structures [19, 20]. For instance, histone acetylation usually renders active genes as does the di- or trimethylation of lysine residue four in histone H3 (H3K4me2, H3K4me3), while H3K9me2/3 and H3K27me3 modifications repress gene expression [21, 22]. Moreover, proteins, known as epigenetic “readers”, can directly bind to these specific modifications and exert repressive or stimulating effects on gene activity [23]. In general, different from DNA methylation which is believed to be a more stable and long-term silencing mechanism, various histone modifications seem to exert short-term, flexible regulation that is important for the plasticity of development [24–26].

### Non-coding RNAs

Non-coding RNAs are transcribed from the genome, but generally are not translated into proteins. They can be classified into two subgroups according to their length: long non-coding RNAs (lncRNAs) containing more than 200 nucleotides and short ncRNAs (< 200 nucleotides), which include microRNAs (miRNAs), short interfering RNAs (siRNAs), piwi-interacting RNAs (piRNAs), and so on [27–29]. Alternatively, non-coding RNAs could also be classified based on their function: housekeeping ncRNAs, including ribosomal RNA (rRNA), transfer RNA (tRNA), small nuclear RNA (snRNA), and small nucleolar RNA (snoRNA), which are expressed in all cell types and carry out essential functions in the cells and regulatory ncRNAs, including several classes of small and long molecules, are involved in the regulation of gene expression [30]. When ncRNAs act in *cis*, they are able to regulate the expression of one or more genes nearby in the same chromosome. However, when ncRNAs act in *trans*, they are able to regulate the expression of one or more genes in different chromosomes or regulate mature RNAs in the cytoplasm [31]. Examples of the well-studied mammalian lncRNAs are Xist (inactive X-specific transcript) and Tsix (X-specific transcript), which are involved in X chromosome inactivation in female mammals [9, 32]. Recent findings demonstrate that non-coding RNAs (ncRNAs) are often accompanied by epigenetic events that are important for the regulation of gene transcription, protein translation, genomic stability, and chromatin modifications [33–37].

miRNAs, which are approximately 22 nucleotides in length, are a class of small non-coding RNAs that function as post-transcriptional regulators of gene expression. They usually bind to the 3' untranslated regions (UTRs) of their target messenger RNAs (mRNAs) through complementary base pairing, and regulate target gene expression by either degrading or translationally repressing target mRNAs [38, 39]. It has been shown that a single miRNA is potentially responsible for repressing hundreds of mRNA targets, since the recognition of a target mRNA mainly depends on a small seed region within the mature miRNA. However, the targets of miRNAs responsible for different phenotypes have been proposed to be tissue or cell specific [40]. In recent years, it has been shown that small non-coding miRNAs are important components of epigenetic post-transcriptional regulators [41–45]. lncRNAs, transcripts longer than 200 nucleotides, are the most heterogeneous class of non-coding RNAs. They can regulate gene expression at different levels, including chromatin modification, alternative splicing, protein localization and activity, and increasing mRNA stability by protecting the 3' UTR of mRNAs from miRNA binding [46]. Increasing evidence has shown that lncRNAs are involved in many different biological processes, such as proliferation, differentiation, pluripotency, and cell death [47, 48].

circRNAs, covalently closed continuous loop RNA lacking both a 5' cap and a 3' tail, are a special class of endogenous non-coding RNAs (ncRNAs) with higher stability and are expressed in a tissue-specific manner [49]. It has been suggested that circRNAs function in five putative manners: (1) promoting transcription of their parental genes; (2) competitively influencing the biogenesis and processing of the mRNA transcribed from their parental genes; (3) acting as miRNA sponges to competitively bind with miRNAs by their conserved binding sites for these miRNAs to block the binding between miRNAs and their target mRNAs; (4) functioning as RNA binding protein (RBP) sponges to regulate the function of RBPs; and (5) encoding proteins due to some circRNAs containing internal ribosome entry sites that allow the binding of ribosomes for protein translation [50].

## Epigenetic regulation of the embryo–uterine crosstalk during peri-implantation

### Epigenetic control of the embryo preparation for implantation

The embryo is at the blastocyst stage when it can implant into the uterus, and the blastocyst needs to acquire implantation competency. During the pre-implantation embryo development process, the terminally differentiated gametes, the egg and sperm, fuse to form a zygote, which then undergoes several rounds of mitosis. During this process, the embryo acquires totipotency, as the single embryonic cell can contribute to both the embryonic and extra-embryonic lineage. Increasing evidence has demonstrated that many different types of epigenetic regulation, including histone modifications, DNA methylation, chromatin accessibility, and 3D chromatin organization, are all involved in this short and highly effective epigenome transforming event.

For the DNA methylation dynamics in the early embryo, the highly methylated paternal genome is actively demethylated mainly through the TET-mediated hydroxylation [51]. Passive demethylation is also thought to play a role due to the inactivation of DNA methyltransferase and rapid cell proliferation in early embryos. Both the paternal and maternal genomes lose their DNA methylation until the blastocyst stage. This highly DNA demethylation process contributes to the activation of the early embryo transcriptional network and pre-implantation embryo development. For the histone modification, the immunofluorescence staining of different histone modifications in different species has suggested dynamic changes of histone modifications in the parental pronuclei after fertilization and during pre-implantation development [52, 53]. Recently, the highly developed low-input genome-wide analysis technologies have drastically stimulated the development of this field in both mice and

humans. For example, it was found that both the distribution pattern and regulation for binding genes of H3K4me3 are not similar to the other cells. In the full-grown oocytes and metaphase II (MII) oocytes with the genome transitions to a silenced state, it was defined as a noncanonical pattern of H3K4me3 (ncH3K4me3), since it was wide spread [54]. The oocyte-specific pattern of the histone epigenome will undergo rapid erasure and re-establishment during fertilization and pre-implantation development. Other dynamic changes for epigenetic regulation, such as H3K27me3 histone modification and chromatin accessibility, have also been extensively explored during pre-implantation embryos [55, 56]. For more detailed information about the epigenome in early embryo development, please refer to other special reviews which address these progresses thoroughly [57, 58]. Based on these valuable data, the following challenges will be the functional dissection of different epigenetic regulations during totipotent acquirement and pre-implantation embryo development to form the blastocyst.

In normal pregnancy, when the embryo develops into the blastocyst stage and enters into the uterine cavity, it initiates the crosstalk with the endometrium, and the attachment reaction can usually be observed in mice on day 4 night. However, this sequence is interrupted in delayed implantation, which can be induced experimentally in mice and rats by ovariectomy before the pre-implantation ovarian E2 surge and maintained by injection of P4. In the delayed implantation, the blastocyst is metabolically dormant and incompetent to initiate attachment in the uterus. This phenomenon can also naturally occur in some species and in lactating rodents [59]. In the dormant blastocyst, beyond metabolism, the transcriptional activity generally shut down, which is consistent with the observation that the chromatin structure was well compacted with more heterochromatin in the dormant blastocyst [60]. The latest reports of manipulating the c-Myc and mTOR pathway can mimic the dormant state in delayed implantation [61, 62], further confirming the low metabolism and transcription activity for cells entering the dormant state. During the blastocyst acquiring the implantation competence, the microRNA Let-7a was downregulated, associated with upregulated miRNA processing enzyme Dicer. In addition, both forced expression of Let-7a and Dicer knockdown can compromise implantation competency of the blastocyst [63, 64]. In humans, the embryo can secrete miRNAs into the culture medium in IVF practice, and these miRNAs can be explored for human embryo reproductive competence assessment [65]. The acquisition of implantation competency for blastocysts involved the expression of molecules related to embryo–uterus crosstalk, such as the inflammatory molecule, to induce the attachment reaction. A recent report has demonstrated that the expression of inflammatory response genes in the human blastocyst stage is associated with open chromatin accessibility in their regulatory



regions [66]. How epigenetic regulation, such as chromatin accessibility and 3D chromatin organization, functions precisely during blastocyst dormancy and reactivation for implantation needs further exploration.

### Epigenetic regulation of uterine preparation for uterine receptivity and attachment reaction

One of the most fascinating tissues in mammals is the uterus, whose major function is to accept implantation-competent blastocysts when the uteri were in a receptive state. In mice, the uterus on day 4 of pregnancy is in a state of receptivity, while 7–10 days after ovulation (the mid-secretory phase of the menstrual cycle) is defined as the uterine receptive stage in humans [2]. Increasing evidence has demonstrated that the acquisition of uterine receptivity is under the control of both transcription factors and epigenetic regulators, which might be responsible for the spatial–temporal dynamic changes of tissue morphology and gene expression in different endometrial cell types during the establishment of uterine receptivity [67].


### DNA methylation and uterine receptivity

P4 and E2 are primary regulators of the cyclic changes of the uterus during the estrus cycles of animals and menstrual cycles of humans. Pregnancy is also a hormone-dominant process. The hormones during these physiological events are mainly ovarian E2 and P4, which function via their nuclear receptors E2 receptors (ER) and P4 receptors (PR), respectively. The expression of both ER and PR is accurately regulated in response to the fluctuation of their ligands. Aberrant DNA methylation of the CpG island in the promoter region

in ER [68, 69] and PRB [70] has been reported in endometrial carcinoma, suggesting that the expression of ER and PR could be regulated by DNA methylation. It has also been demonstrated that DNA methylation is correlated with PR expression in normal endometrium [71], and in mouse uterus with constitutive Notch1 activation, PR expression was shut down as DNA methylation was significantly induced at the promoter region of *Pgr* (Fig. 2a) [72]. In addition, endocrine disruptor endosulfan or folate deficiency affected the methylation states of promoters or regulatory regions in ER $\alpha$  [73, 74], leading to altered expression of uterine ER $\alpha$ .

HOXA10, another critical transcription factor for implantation, is spatio-temporally expressed in the endometrium with expression peaks during the window of implantation both in mouse and human [75, 76]. Female mice with a targeted disruption of *Hoxa10* are unable to support embryo implantation [77, 78], suggesting that HOXA10 is critical for uterine receptivity. It has been reported that HOXA10 expression is controlled by DNA methylation [79] and aberrant DNA methylation led to decreased expression of HOXA10 in the eutopic endometrium of patients with endometriosis, which caused infertility or subfertility in women (Fig. 2a) [80].

It has been reported that the loss of ER signalling triggered silencing of downstream targets by DNA methylation [81] and E2 can induce frequent methylation/demethylation on promoter CpG sites to regulate cyclic transcriptional activation of target genes [82] in breast cancer cells. This mechanism was also demonstrated to exist in endometrial cells, which are under the control of steroid hormones [83]. MUC1 is expressed in the endometrial epithelium and its important functions in endometrial receptivity and embryo attachment have been extensively studied [84, 85]. The CpG



**Fig. 2** Epigenetic regulation of critical molecule expression during implantation. **a** DNA methylation 5mC in the gene promoter or other regulatory region can negatively regulate the transcription target genes, such as *Pgr*, *Hoxa10*, and *Hand2*. **b** Nonclassical post-translational modification of critical molecules, such as HoxA10 and PR by epigenetic regulator can modulate the function or activ-

ity of these transcription factors, K Lysine residues, Ac acetylation, Ub ubiquitylation. **c** At post-transcriptional level, mRNAs of *Pgr* or *Ptgs2* (which encode the Cox2) is targeted by the miRNAs. **d** During decidualization, active chromatin modifications (such as H3K4me3 and H3K27Ac) in the gene regulatory region contribute to the highly expression of decidual marker gene *IGFBP1* and *PRL*

island of MUC1 was reported to be less methylated in the mid-secretory phase compared to the proliferative phase, which is strongly correlated with its increased gene expression in the mid-secretory phase [83].

Besides the regulation of specific gene by DNA methylation, DNA global methylation significantly decreases from the proliferative phase towards the secretory phase in glandular epithelial cells, but increases in the stromal cells of the endometrium by immunostaining [71]. Consistently, inhibition of DNA methylation by 5'-aza-2'-deoxycytidine (AZA) increased E-cadherin expression in uterine epithelial cells, in favour of establishing uterine receptivity *in vitro* [86]. Consistently, combined treatment with E2 and P4 significantly decreased the mRNA expression of DNMT3a and DNMT3b [87, 88]. However, DNMT1 mRNA expression was significantly higher in secretory-phase endometrium compared with proliferative endometrium and menstrual endometrium [89]. It was also reported that DNMT1 mRNA did not change over the menstrual cycle, while the transcription of DNMT3a and DNMT3b was significantly decreased in the secretory-phase endometrium [87]. The discrepancies may be due to the different methods of sample collection and the detection timepoints for *in vitro* experiments.

For the genome-wide change in DNA methylation during uterine receptivity establishment, the evidence is mainly from the research in human endometrium. Two research groups have reported that changes in DNA methylation were correlated with gene expression during the transition from the pre-receptive to the receptive phase in the endometrium [83, 90]. However, a recent report using the human endometrial stroma *in vitro* decidualization model, which mimics stroma cell differentiation during the establishment of uterine receptivity, has demonstrated stable DNA methylation status during decidualization, without a clear association between the differentially expressed genes and DNA methylation changes [91]. This discrepancy may be due to the difference between the *in vivo* and *in vitro* models. To further understand the role of DNA methylation during the establishment of uterine receptivity, a more sensitive and cell-type specific analysis of genome-wide DNA methylation is needed. Moreover, functional study utilizing the genetic mouse model will also provide critical cues.

### Histone modification and uterine receptivity

In the early 1970s, it was found that treatment with estradiol led to increased histone acetylation in the uterus of rat and mouse [92–94], suggesting that the ovarian steroid hormones might act through chromatin alterations to effect gene transcription. In fact, co-treatment with E2 and P4 increased the levels of acetylated histone H3 and H4, and this increase was enhanced by co-treatment with the HDAC inhibitor TSA [95]. Likewise, it has been shown

that HDAC inhibitors treatment can enhance the proliferative actions of E2 in mice [96]. Moreover, HDAC inhibitor was able to induce morphological and molecular changes comparable to treatment with E2 and P4 in human endometrial cells [97]. Consistently, during E2- and P4-induced stroma decidualization, many epigenetic regulators have been reported to be dynamically expressed [98]. These findings suggest that epigenetic regulation is involved in uterine events controlled by E2 or/and P4.

During the menstrual cycle, the endometrium undergoes morphological and functional changes, which are essential for the establishment of uterine receptivity and are also under the regulation of histone modification. Glycodelin, a secretory protein, is highly expressed at 10 days after ovulation (within the implantation window) in the human endometrium, and treatment with HDAC inhibitor upregulated the expression of glycodelin [99]. As we previously mentioned, HOXA10 is a transcription factor critical for endometrial receptivity. It has been demonstrated that acetylation of HOXA10 by PCAF downregulated HOXA10-mediated  $\beta$ 3-integrin (ITGB3) expression and diminished HOXA10-mediated embryo adhesiveness (Fig. 2b) [100]. These data suggest that the acetylated modification of receptivity-related genes is essential for the achievement of endometrial receptivity. Moreover, BMI1, a key component of the Polycomb repressive complex-1 (PRC1), interacts with the PR as well as the E3 ligase E6AP in a polycomb complex-independent manner and regulates the PR ubiquitination that is essential for normal P4 responsiveness during the acquisition of uterine receptivity (Fig. 2b) [101]. In addition, uterine glands are essential for embryo implantation and pregnancy, as demonstrated by implantation or pregnancy failure in several gene mutation mouse models [102, 103]. The histone H4K20me1 methyltransferase Pr-set7 has been reported to regulate gland development and thus female fertility [104].

A number of transcription factors have been demonstrated to be critical for uterine receptivity establishment, such as Hand2 and Msx1/2. Hand2 is specifically expressed in uterine stroma cells and induced by P4 signalling through PR. Hand2 mediated one of the PR's critical functions, inhibiting uterine ER signalling and epithelium proliferation, mainly through repressing the expression of the stroma-derived growth factors [105]. However, how Hand2 inhibits target gene expression in uterus is still unclear, and it was reported that Hand2 could repress target gene expression through the distal enhancer [106]. Msx1 and Msx2 are another example of transcription factors, whose function is reported to repress Wnt5a to regulate the cell polarity of luminal epithelium for blastocyst attachment [107]. In myoblast cells, Msx1 could interact with the repressive PRC2 complex to inhibit target gene expression by upregulating the H3K27Me3 [108], and

whether these histone modification regulators are involved in uterine cells requires further exploration.

### Non-coding RNA for uterine receptivity

Several miRNAs have been recently identified both in the human and mouse endometrium using next-generation sequencing [109–112]. A study using mid-secretory endometrial samples from women with repeated implantation failure discovered that miRNAs with altered expression are likely to contribute to defective endometrial receptivity in humans [113]. Moreover, Altmae et al. found that the expression of miR-30b, miR-30d, and miR-494 was closely related to human endometrial receptivity [43]. Overexpression of miR-30d in human endometrial epithelial cells demonstrates that miR-30d participates in the maintenance of DNA methylation, supporting an epigenetic role for miR-30d [114]. Ponsuksili et al. further revealed that differential expression of miRNAs and their target mRNAs in the endometrium is associated with endometrial receptivity [115]. In addition, microRNA-31 is significantly elevated in both human endometrium and serum during the window of implantation, and might act as a potential biomarker for optimum receptivity [116]. Several miRNAs, including miR-96, have been demonstrated to regulate epithelial PGR expression during the establishment of uterine receptivity in rhesus monkeys and humans (Fig. 2c), but not in rodents [117]. Cox-2, an attachment reaction marker encoded by prostaglandin endoperoxide synthase 2 (PTGS2) and expressed around the implanted blastocyst, was reported to be regulated by miR-101a and miR-199a\* at the post-transcriptional level (Fig. 2c) [112]. Moreover, miRNAs secreted within the exosome by the embryo or/and uterine epithelial cells are reported to be involved not only in the interaction between the embryo and uterus during implantation [118], but also in the potential crosstalk between the ICM and TE for coordinating the trophoblast cell migration during implantation [119]. For more detailed knowledge about miRNAs during implantation, one can refer to several recent reviews [120–122].

One previous work was conducted to identify specific miRNAs involved in hormonal regulation of normal human endometrium by E2 and P4 and revealed that miRNAs down-regulate the expression of some cell cycle genes in the endometrial epithelium during the secretory phase, thereby suppressing cell proliferation [110]. Another finding revealed that the microRNA-200a locally attenuates P4 signalling in the cervix by downregulating the PR protein level and upregulating P4 inactivating enzyme (20 $\alpha$ -HSD), leading to a different response to P4 between the cervix and uterus (Fig. 2c) [123]. These findings demonstrated that the functions of E2 and P4 are mediated or affiliated by epigenetic regulators.

Several lncRNAs were reported to be actively transcribed in the uterus during embryo implantation, such as HOXA11-AS [124], Scx-AS [125], FGF2-AS [126], and EMX2-OS [127], which are naturally occurring antisense RNAs and supposed to mediate transcriptional activation or transcriptional silencing of their respective host genes. Moreover, the steroid receptor RNA activator (SRA) gene, an intergenic ncRNA and steroid receptor co-activator, was also detected in the uterus [128]. Recently, Sigurgeirsson et al. performed comprehensive RNA sequencing of healthy human endometrium at the proliferative stage and mid-secretory stage (receptive phase), and found that 516 lncRNAs were significantly differentially expressed in the endometrium [129]. Another group also explored RNA-Seq to compare the transcriptome of the endometrium between LH + 2 (pre-receptive phase) and LH + 7 (receptive phase) and identified 2372 differentially expressed genes [130]. Using similar method, different expressed lncRNA transcripts were identified during the implantation window both in the mouse and porcine [131, 132]. Although these findings have shown that lncRNAs are associated with the acquisition of uterine receptivity, the functions, and the molecular mechanism of these lncRNAs during uterine receptivity establishment need further investigation. On the other hand, lncRNA H19, one of the first genes found to be transcribed into long non-coding RNAs, was decreased in repeated implantation failure [133]. Feng et al. reported that six lncRNAs were significantly elevated in the endometrium of RIF or recurrent miscarriage patients [134]. These data suggest that lncRNAs may be prospective biomarkers for predicting endometrial receptivity.

Moreover, circRNA-9119 was demonstrated to regulate the expression of prostaglandin endoperoxide synthase 2 (PTGS2), which is involved in the regulation of some markers of endometrial receptivity, by sponging miR-26a in ovine endometrial epithelial cells [135], suggesting that circRNAs might participate in embryo implantation. Indeed, altered circular RNA expression has been reported to be associated with repeated implantation failure in humans [136].

### Epigenetic control of stromal–decidual transformation after implantation

In mice, the embryo attachment to the uterine luminal epithelium would initiate the stroma differentiation program, known as decidualization. Decidual cells in the endometrium provide nutrients to the fetus before the establishment of maternal–fetal circulation, inhibit trophoblast cells from invading too deeply inside the uterine wall, and are involved in the immunotolerance of the allogeneic fetus. Therefore, decidualization is a pivotal step for normal embryo implantation, placental development, and successful completion of pregnancy to term. While decidualization in women occurs during the secretory phase of each menstrual cycle as well



as in pregnancy, this event only occurs during pregnancy in rodents [7]. Gene expression undergoes dramatic changes in human endometrial stromal cells (HESCs) during decidualization. These changes in gene expression are associated with changes in chromatin structure, which are partially regulated by epigenetic factors. Since the finding that histone phosphorylation and acetylation were increased upon decidualization in the mouse uterus approximately 40 years ago [92], there is increasing evidence that the epigenetic mechanisms regulate gene expression during decidualization in the endometrium.

### DNA methylation and decidualization

DNA methylation is high in the proliferative phase, with a significant decline towards the end of the secretory phase (corresponding to the stage of decidualization) [71]. Consistently, the methylation inhibitor 5-Aza-2'-deoxycytidine (AZA) induced some decidualization-like changes in endometrial stromal cells possibly via the cytoskeletal reorganization pathway, even though this decidualization was different from that induced by hormones [137]. Although it has also been demonstrated that no obvious changes in global DNA methylation are observed after decidualization is induced in humans [98], the DNMT inhibitor 5-AZA-Dc has negative effects on decidualization in mice [138, 139], suggesting that gene locus-specific changes in DNA methylation or DNA methylation maintenance play an important role during decidual development. The expression of DNMT1, DNMT3a, and DNMT3b in the endometrium fluctuates during the cycle and in differentiating HESCs [87–89, 98, 140]. Even though there is considerable discrepancy between these investigations due to different treatment recipes, cell sources, and other experimental variations, these findings further support the notion that active changes in DNA methylation are essential for the expression of decidual genes. Recently, using mice with different genetic backgrounds, decidual caps with widely and apparently randomly disturbed DNA methylation were demonstrated to be responsible for the pregnancy failure in the CBA/J X DBA/2 spontaneous abortion mouse model [141]. Combining the DNA methylation analysis and RNA-Seq in decidua tissue from women with normal pregnancy and recurrent pregnancy loss, Yu et al. identified 539 differentially methylated regions (DMRs) that are significantly correlated with gene expression [142]. Moreover, hypermethylation at the promoter region of FOXP3, which is extremely important for the development and function of Treg cells, down-regulates FOXP3 expression, resulting in a broken immune balance at the maternal–fetal interface and pregnancy failure [143]. Hand2, another critical molecule for embryo implantation and decidualization, is reported to be silenced in endometrial cancer through DNA methylation at its promoter [144].

### Histone modification and decidualization

Recent studies confirmed dynamic regulation of histone marks during the menstrual cycle in vivo [67] and upon decidualization of HESCs in culture. It has been demonstrated that trichostatin A, an inhibitor of histone deacetylation, enhanced estradiol and progesterin-induced decidualization of primary HESCs, suggesting that histone acetylation is involved in uterine decidualization [95]. Treating HESCs with TSA was also reported to increase the expression of TIMP-1 and TIMP-3 by acetylating histones in their promoters, and inhibit trophoblast invasion during implantation [145]. Retinoblastoma-binding protein 7 (Rbbp7), a core component of many histone modifications and chromatin remodelling complexes, has been demonstrated to be a potentially functional player regulating normal histone acetylation and cyclin D3 expression during decidual development [146]. Moreover, it has been shown that C/EBP $\beta$  regulates the expression of IGFBP-1 and PRL by altering the histone acetylation status of their promoters during HESC decidualization [147]. Recently, the increases of H3K27ac and H3K4me3 in both the proximal and distal regulatory regions have been found to be the main changes of histone modifications accompanying uterine decidualization, which are tightly associated the upregulated gene expression (Fig. 2d) [148]. For example, increased H3K27ac levels in the distal upstream region of IGFBP1 can significantly stimulate IGFBP1 expression in endometrial stromal cells during decidualization [149]. In addition, the down-regulation of the histone methyltransferase EZH2 was reported to contribute to epigenetic programming during the decidualization of HESCs [150].

In mouse decidual differentiation, the H3K27me3-mediated silencing of inflammatory chemokine genes can limit T cell access to the fetal–maternal interface for the establishment of immune tolerance during pregnancy [151]. An additional study demonstrated the dynamic changes of H3K27Me3 during the whole pregnancy: transcriptional silence of target genes related to parturition-inducing signals through H3K27Me3 to ensure uterine quiescence during early pregnancy; then, genome-wide demethylation occurs to derepress these genes for labor [152]. A recent report uncovered that uterus-specific deletion of *Ezh2* derail post-implantation decidualization [153]. Cbx4/Ring1B-containing PRC1, which is responsible for the ubiquitination of histone-H2A at lysine-119, was found to control decidualization through the regulation of extracellular matrix remodelling genes [154]. These findings demonstrate that different histone modifications are involved in the regulation of decidualization.

## Non-coding RNA and decidualization

During the cAMP and medroxyP4 acetate induced in vitro human uterine stromal decidualization, some of the miRNAs were found to be significantly differentially expressed [155, 156]. However, siRNA-mediated knockdown of Dicer displayed no obvious defect in decidualization marker expression, and further evidence suggested microRNA resistance in decidual cells due to the limited level of miRNA processing enzyme Ago1/2 [156]. It has also been reported that individual miRNAs regulate the stroma decidualization [157, 158]. Therefore, more specific studies are needed to explore the function of miRNAs during decidualization.

The different manners of epigenetic modification do not function independently. Different epigenetic modifiers interact with one another. In detail, the establishment of histone modifications and DNA methylation is interdependent. They not only interplay with each other, but also with regulatory proteins and non-coding RNAs [159–164]. Specifically, the crosstalk between DNA methylation and histone modifications is established by MBD proteins, which contain a methyl-CpG binding domain. MBDs specifically recognize and bind to methylated CpGs, and subsequently recruit histone deacetylases (HDACs), which remove acetyl groups from the lysine of histone tails, leading to chromatin condensation and transcriptional inactivation [159, 160]. The expression of DNMT1 and MBD2 was increased significantly by treatment with E2 in combination with P4, suggesting that the interactions between different epigenetic modifications may also exist in the endometrium.

## Epigenetic control of infertility-related uterine disease

Increasing evidence demonstrates that uterine diseases, such as endometriosis and endometritis, are closely related to subfertility or infertility [165–167]. The occurrence of these diseases and disease-induced infertility are associated with disturbed epigenetic regulation.

As mentioned above, HOXA10 has an important function in regulating endometrial development and in establishing conditions necessary for implantation. It has been reported that HOXA10 expression was decreased in the endometrium of women with endometriosis [168]. In endometriosis models established in both mouse and baboon, DNA hypermethylation of Hoxa10/HOXA10 promoters was demonstrated in eutopic endometrium [169, 170]. In humans, the HOXA10 promoter was also hypermethylated in the endometrium of women with endometriosis [80, 171]. Moreover, the overexpression of DNMT1, DNMT3A, and DNMT3B was revealed in ectopic endometrium compared with normal controls [172], further suggesting that DNA methylation might be one of the possible mechanisms underlying endometriosis.

A hypermethylated promoter of PRB is also observed in endometriosis with decreased PRB expression, which might account for P4 resistance in endometriosis since P4 functions through its receptors PRB [173]. In addition, the positive regulators for endometriosis, such as aromatase and ER $\beta$ , are both with hypomethylated promoters and display aberrant upregulation [174, 175]. In addition to DNA methylation, histone modification is also involved in the underlying mechanism of endometriosis. HDAC1 and HDAC2 are reported to be differentially expressed in endometriosis [176]. Hypoacetylation at H3K4, H3K9, and H4K16, and hypermethylation at H3K4, H3K9, and H3K27 were found in endometriosis tissues [177, 178], suggesting a possible role of histone modification in regulating gene expression in endometriosis.

In addition to endometriosis, several other common gynecologic disease, such as hydrosalpinges, polyps, and submucosal myomas, are associated with implantation defects [179]. CpG sites within the HOXA10 promoter were highly methylated [180], which is responsible for the defective HOXA10 expression [181] in these diseases. E-cadherin, the major cadherin molecule expressed in epithelial cells, is known to act as an invasion suppressor in cancer cells. It has been shown that hypermethylation in the promoter region of E-cadherin is associated with dedifferentiation and myometrial invasion of endometrial carcinoma [182]. Moreover, several studies have demonstrated that circRNAs are involved in uterine dysfunction, such as in repeated implantation failure [136].

## Conclusion

At present, although an increasing number of epigenetic factors and their expressions, as well as the regulations have been reported during the peri-implantation stage (summarized in Table 1), the systemic exploration of their physiological function and the underlying mechanism is still a major challenge in this field. Genetic mouse models, especially temporal induced genetic manipulation models in different uterine compartments, and other in vivo/ex vivo approaches combined with the newly developed genome-wide strategy must be utilized to uncover the mystery of the genetic regulation network during embryo implantation. Moreover, some epigenetic regulators such as miRNA and circRNAs, which can be stably detected in the serum and other body fluids, might act as biomarkers for the diagnosis and targets for potential treatment of embryo implantation defects [183]. In addition, both the internal environment such as uterus aging [184, 185], and external environment, such as malnutrition and endocrine disruptors [74, 186–188], have been reported to influence embryo implantation and lead to an adverse pregnancy

**Table 1** Represent epigenetic regulators for embryo implantation in the mouse and human

Epigenetic regulator	Function	References
Let-7a	Regulate the blastocyst implantation potential via its action on $\beta$ 3-Integrin and Dicer	[63, 64]
miR-30d	Upregulated during the acquisition of receptivity in the endometrium	[43]
	Overexpression induced gene expression changes at transcriptome and proteomics changes in the epithelium	[114]
	Involved in interaction between embryo and uterus as secreted by uteri and taken up by the embryo	[118]
miR-31	Elevated in both human endometrium and serum during the implantation window	[116]
miR-96	Modulate PGR expression during the establishment of uterine receptivity in rhesus monkey and human, but not in rodents	[117]
miR-101a/miR-199a*	Regulate the attachment reaction marker Cox2 expression at the post-transcriptional level	[189]
miR-200a	Locally attenuates progesterone signalling in the cervix through downregulating the PR protein level and upregulation of progesterone inactivating enzyme (20 $\alpha$ -HSD)	[123]
DNMT	Regulate the expression of implantation- and pregnancy maintenance- related genes, such as E-cadherin, Hoxa10, Muc1, FoxP3 and Hand2 were influenced through the DNA methylation	[79, 86, 143, 144]
	DNA methylation changes in endometrium were correlated with gene expression during the transition from pre-receptive to receptive phase in human	[83, 90]
	DNMT inhibitor 5-AZA-Dc inhibits decidualization in mice	[138, 139]
	Disturbed DNA methylation of decidual tissue in spontaneous abortion mouse model	[141]
	PR down-regulation through DNA methylation in Notch1 constitutive activation mouse uterus	[72]
	Altered DNA methylation states of promoters and expressions of ER $\alpha$ in endocrine disruptor endosulfan or folate deficiency mouse uterus	[73, 74]
	Hypomethylated region near CREB5 recruited transcription factors binding, such as P53 and SP1, and in turn upregulated CREB5 in recurrent pregnancy loss, which induced compromised trophoblast cell migration and apoptosis	[142]
circRNA-9119	Regulate the expression of Cox2, by sponging miR-26a in endometrial epithelial cells	[135]
Histone acetylation	HDAC inhibitors upregulated the glycodefin expression, which induced LIF expression in glandular epithelial and enhanced implantation	[99]
	Acetylation of HOXA10 diminished its transcription activity on $\beta$ 3-integrin promoter	[100]
	TSA treatment enhanced the estradiol and progestin-induced decidualization	[95]
	Rbbp7, a core component of histone acetylation, regulate the mouse decidualization	[146]
	Upregulation of decidualization marker IGFBP-1 and PRL was associated with increased histone acetylation status in their promoters	[147]
	Increased H3K27ac and H3K4me3 levels in both the proximal and distal promoter regions of decidual upregulated genes	[148]
Histone methylation	Down-regulation of Ezh2 during the human stroma cell decidualization	[150]
	H3K27me3 mediate silence of inflammatory chemokine genes in mouse decidualization	[151]
	Dynamic H3K27me3 fluctuation for pregnancy maintenance and labor	[152]
	H4K20Me1 methyltransferase Pr-set7 is indispensable for uterine gland development	[104]
Histone mono- ubiquitination	PR ubiquitination regulated by Bmi1 independent of PRC1 complex is essential for normal progesterone responsiveness in endometrium	[101]
	Cbx4/Ring1B-containing PRC1 control decidualization through the regulation of extracellular matrix remodelling gene	[154]

outcome. However, the underlying mechanism, especially from the epigenetic aspect, must be first addressed before the potential management for these disturbances.

81830045 and 81490744 to H. W. and 31600945 to J.L.) and Fundamental Research Funds for the Central Universities (20720180039 to S.K., 20720180041 to J.L.).

**Acknowledgements** This review article was supported in parts by National Key R&D program of China (2017YFC1001402 to H.W.), and the National Natural Science Foundation (81601285 to S.K.,

## References

- Dickmann Z, Noyes RW (1961) The zona pellucida at the time of implantation. *Fertil Steril* 12:310–318
- Wang H, Dey SK (2006) Roadmap to embryo implantation: clues from mouse models. *Nat Rev Genet* 7:185–199
- Zhang S, Lin H, Kong S, Wang S, Wang H, Wang H, Armant DR (2013) Physiological and molecular determinants of embryo implantation. *Mol Aspects Med* 34:939–980
- Paria BC, Huet-Hudson YM, Dey SK (1993) Blastocyst's state of activity determines the “window” of implantation in the receptive mouse uterus. *Proc Natl Acad Sci USA* 90:10159–10162
- Cha J, Sun X, Dey SK (2012) Mechanisms of implantation: strategies for successful pregnancy. *Nat Med* 18:1754–1767
- Lim HJ, Wang H (2010) Uterine disorders and pregnancy complications: insights from mouse models. *J Clin Invest* 120:1004–1015
- Gellersen B, Brosens JJ (2014) Cyclic decidualization of the human endometrium in reproductive health and failure. *Endocr Rev* 35:851–905
- Berger SL, Kouzarides T, Shiekhatar R, Shilatifard A (2009) An operational definition of epigenetics. *Genes Dev* 23:781–783
- Nelissen EC, van Montfort AP, Dumoulin JC, Evers JL (2011) Epigenetics and the placenta. *Hum Reprod Update* 17:397–417
- Herman JG, Baylin SB (2003) Gene silencing in cancer in association with promoter hypermethylation. *N Engl J Med* 349:2042–2054
- Laird PW (2003) The power and the promise of DNA methylation markers. *Nat Rev Cancer* 3:253–266
- Ooi SK, Qiu C, Bernstein E, Li K, Jia D, Yang Z, Erdjument-Bromage H, Tempst P, Lin SP, Allis CD, Cheng X, Bestor TH (2007) DNMT3L connects unmethylated lysine 4 of histone H3 to de novo methylation of DNA. *Nature* 448:714–717
- Veland N, Lu Y, Hardikar S, Gaddis S, Zeng Y, Liu B, Estecio MR, Takata Y, Lin K, Tomida MW, Shen J, Saha D, Gowher H, Zhao H, Chen T (2018) DNMT3L facilitates DNA methylation partly by maintaining DNMT3A stability in mouse embryonic stem cells. *Nucleic Acids Res* 47(1):152–167
- Jaenisch R, Bird A (2003) Epigenetic regulation of gene expression: how the genome integrates intrinsic and environmental signals. *Nat Genet* 33(Suppl):245–254
- Kim TH, Barrera LO, Zheng M, Qu C, Singer MA, Richmond TA, Wu Y, Green RD, Ren B (2005) A high-resolution map of active promoters in the human genome. *Nature* 436:876–880
- Weber M, Schubeler D (2007) Genomic patterns of DNA methylation: targets and function of an epigenetic mark. *Curr Opin Cell Biol* 19:273–280
- Branco MR, Ficiz G, Reik W (2011) Uncovering the role of 5-hydroxymethylcytosine in the epigenome. *Nat Rev Genet* 13:7–13
- Kouzarides T (2007) SnapShot: histone-modifying enzymes. *Cell* 131:822
- Turner BM (2002) Cellular memory and the histone code. *Cell* 111:285–291
- Kouzarides T (2007) Chromatin modifications and their function. *Cell* 128:693–705
- Peterson CL, Laniel MA (2004) Histones and histone modifications. *Curr Biol* 14:R546–R551
- Sims RJ 3rd, Reinberg D (2006) Histone H3 Lys 4 methylation: caught in a bind? *Genes Dev* 20:2779–2786
- Patel DJ, Wang Z (2013) Readout of epigenetic modifications. *Annu Rev Biochem* 82:81–118
- Boyer LA, Plath K, Zeitlinger J, Brambrink T, Medeiros LA, Lee TI, Levine SS, Wernig M, Tajonar A, Ray MK, Bell GW, Otte AP, Vidal M, Gifford DK, Young RA, Jaenisch R (2006) Polycomb complexes repress developmental regulators in murine embryonic stem cells. *Nature* 441:349–353
- Lee TI, Jenner RG, Boyer LA, Guenther MG, Levine SS, Kumar RM, Chevalier B, Johnstone SE, Cole MF, Isono K, Koseki H, Fuchikami T, Abe K, Murray HL, Zucker JP, Yuan B, Bell GW, Herbolsheimer E, Hannett NM, Sun K, Odom DT, Otte AP, Volkert TL, Bartel DP, Melton DA, Gifford DK, Jaenisch R, Young RA (2006) Control of developmental regulators by Polycomb in human embryonic stem cells. *Cell* 125:301–313
- Reik W (2007) Stability and flexibility of epigenetic gene regulation in mammalian development. *Nature* 447:425–432
- Boon RA, Dimmeler S (2015) MicroRNAs in myocardial infarction. *Nat Rev Cardiol* 12:135–142
- Ransohoff JD, Wei Y, Khavari PA (2018) The functions and unique features of long intergenic non-coding RNA. *Nat Rev Mol Cell Biol* 19:143–157
- Ernst C, Odom DT, Kutter C (2017) The emergence of piRNAs against transposon invasion to preserve mammalian genome integrity. *Nat Commun* 8:1411
- Ferlita A, Battaglia R, Andronico F, Caruso S, Cianci A, Purrello M, Pietro CD (2018) Non-coding RNAs in endometrial physiopathology. *Int J Mol Sci* 19(7):2120
- Koerner MV, Pauler FM, Huang R, Barlow DP (2009) The function of non-coding RNAs in genomic imprinting. *Development* 136:1771–1783
- Gendrel AV, Heard E (2014) Noncoding RNAs and epigenetic mechanisms during X-chromosome inactivation. *Annu Rev Cell Dev Biol* 30:561–580
- Bagga S, Bracht J, Hunter S, Massirer K, Holtz J, Eachus R, Pasquinelli AE (2005) Regulation by let-7 and lin-4 miRNAs results in target mRNA degradation. *Cell* 122:553–563
- Folco HD, Pidoux AL, Urano T, Allshire RC (2008) Heterochromatin and RNAi are required to establish CENP-A chromatin at centromeres. *Science* 319:94–97
- Varambally S, Cao Q, Mani RS, Shankar S, Wang X, Ateeq B, Laxman B, Cao X, Jing X, Ramnarayanan K, Brenner JC, Yu J, Kim JH, Han B, Tan P, Kumar-Sinha C, Lonigro RJ, Palanisamy N, Maher CA, Chinnaiyan AM (2008) Genomic loss of microRNA-101 leads to overexpression of histone methyltransferase EZH2 in cancer. *Science* 322:1695–1699
- Kopp F, Mendell JT (2018) Functional classification and experimental dissection of long noncoding RNAs. *Cell* 172:393–407
- Sun Q, Hao Q, Prasanth KV (2018) Nuclear long noncoding RNAs: key regulators of gene expression. *Trends Genet* 34:142–157
- Carthew RW, Sontheimer EJ (2009) Origins and mechanisms of miRNAs and siRNAs. *Cell* 136:642–655
- Bartel DP (2004) MicroRNAs: genomics, biogenesis, mechanism, and function. *Cell* 116:281–297
- Lim LP, Lau NC, Garrett-Engele P, Grimson A, Schelter JM, Castle J, Bartel DP, Linsley PS, Johnson JM (2005) Microarray analysis shows that some microRNAs downregulate large numbers of target mRNAs. *Nature* 433:769–773
- He L, Hannon GJ (2004) MicroRNAs: small RNAs with a big role in gene regulation. *Nat Rev Genet* 5:522–531
- Ghildiyal M, Zamore PD (2009) Small silencing RNAs: an expanding universe. *Nat Rev Genet* 10:94–108
- Altmae S, Martinez-Conejero JA, Esteban FJ, Ruiz-Alonso M, Stavreus-Evers A, Horcajadas JA, Salumets A (2013) MicroRNAs miR-30b, miR-30d, and miR-494 regulate human endometrial receptivity. *Reprod Sci* 20:308–317
- Gebert LFR, MacRae IJ (2018) Regulation of microRNA function in animals. *Nat Rev Mol Cell Biol* 431:350
- Bartel DP (2018) Metazoan MicroRNAs. *Cell* 173:20–51

46. Chen YG, Satpathy AT, Chang HY (2017) Gene regulation in the immune system by long noncoding RNAs. *Nat Immunol* 18:962–972
47. Fatica A, Bozzoni I (2014) Long non-coding RNAs: new players in cell differentiation and development. *Nat Rev Genet* 15:7–21
48. Yan P, Luo S, Lu JY, Shen X (2017) Cis- and trans-acting lncRNAs in pluripotency and reprogramming. *Curr Opin Genet Dev* 46:170–178
49. Hansen TB, Jensen TI, Clausen BH, Bramsen JB, Finsen B, Damgaard CK, Kjems J (2013) Natural RNA circles function as efficient microRNA sponges. *Nature* 495:384–388
50. Tatomer DC, Wilusz JE (2017) An uncharted journey for ribosomes: circumnavigating circular RNAs to produce proteins. *Mol Cell* 66:1–2
51. Lee HJ, Hore TA, Reik W (2014) Reprogramming the methylome: erasing memory and creating diversity. *Cell Stem Cell* 14:710–719
52. Lepikhov K, Zakhartchenko V, Hao R, Yang F, Wrenzycki C, Niemann H, Wolf E, Walter J (2008) Evidence for conserved DNA and histone H3 methylation reprogramming in mouse, bovine and rabbit zygotes. *Epigenetics Chromatin* 1:8
53. Lepikhov K, Walter J (2004) Differential dynamics of histone H3 methylation at positions K4 and K9 in the mouse zygote. *BMC Dev Biol* 4:12
54. Zhang B, Zheng H, Huang B, Li W, Xiang Y, Peng X, Ming J, Wu X, Zhang Y, Xu Q, Liu W, Kou X, Zhao Y, He W, Li C, Chen B, Li Y, Wang Q, Ma J, Yin Q, Kee K, Meng A, Gao S, Xu F, Na J, Xie W (2016) Allelic reprogramming of the histone modification H3K4me3 in early mammalian development. *Nature* 537:553–557
55. Liu X, Wang C, Liu W, Li J, Li C, Kou X, Chen J, Zhao Y, Gao H, Wang H, Zhang Y, Gao Y, Gao S (2016) Distinct features of H3K4me3 and H3K27me3 chromatin domains in pre-implantation embryos. *Nature* 537:558–562
56. Wu J, Huang B, Chen H, Yin Q, Liu Y, Xiang Y, Zhang B, Liu B, Wang Q, Xia W, Li W, Li Y, Ma J, Peng X, Zheng H, Ming J, Zhang W, Zhang J, Tian G, Xu F, Chang Z, Na J, Yang X, Xie W (2016) The landscape of accessible chromatin in mammalian preimplantation embryos. *Nature* 534:652–657
57. Xu Q, Xie W (2018) Epigenome in early mammalian development: inheritance, reprogramming and establishment. *Trends Cell Biol* 28:237–253
58. Ladstatter S, Tachibana K (2018) Genomic insights into chromatin reprogramming to totipotency in embryos. *J Cell Biol* 218:70–82
59. Fenelon JC, Banerjee A, Murphy BD (2014) Embryonic diapause: development on hold. *Int J Dev Biol* 58:163–174
60. Fu Z, Wang B, Wang S, Wu W, Wang Q, Chen Y, Kong S, Lu J, Tang Z, Ran H, Tu Z, He B, Zhang S, Chen Q, Jin W, Duan E, Wang H, Wang YL, Li L, Wang F, Gao S, Wang H (2014) Integral proteomic analysis of blastocysts reveals key molecular machinery governing embryonic diapause and reactivation for implantation in mice. *Biol Reprod* 90:52
61. Bulut-Karslioglu A, Biechele S, Jin H, Macrae TA, Hejna M, Gertsenstein M, Song JS, Ramalho-Santos M (2016) Inhibition of mTOR induces a paused pluripotent state. *Nature* 540:119–123
62. Scognamiglio R, Cabezas-Wallscheid N, Thier MC, Altamura S, Reyes A, Prendergast AM, Baumgartner D, Carnevalli LS, Atzberger A, Haas S, von Paleske L, Boroviak T, Worsdorfer P, Essers MA, Kloz U, Eisenman RN, Edenhofer F, Bertone P, Huber W, van der Hoeven F, Smith A, Trumpp A (2016) Myc depletion induces a pluripotent dormant state mimicking diapause. *Cell* 164:668–680
63. Cheong AW, Pang RT, Liu WM, Kottawatta KS, Lee KF, Yeung WS (2014) MicroRNA Let-7a and dicer are important in the activation and implantation of delayed implanting mouse embryos. *Hum Reprod* 29:750–762
64. Liu WM, Pang RT, Cheong AW, Ng EH, Lao K, Lee KF, Yeung WS (2012) Involvement of microRNA lethal-7a in the regulation of embryo implantation in mice. *PLoS One* 7:e37039
65. Capalbo A, Ubaldi FM, Cimadomo D, Noli L, Khalaf Y, Farcomeni A, Ilic D, Rienzi L (2016) MicroRNAs in spent blastocyst culture medium are derived from trophectoderm cells and can be explored for human embryo reproductive competence assessment. *Fertil Steril* 105(225–235):e221–e223
66. Gao L, Wu K, Liu Z, Yao X, Yuan S, Tao W, Yi L, Yu G, Hou Z, Fan D, Tian Y, Liu J, Chen ZJ, Liu J (2018) Chromatin accessibility landscape in human early embryos and its association with evolution. *Cell* 173(248–259):e215
67. Munro SK, Farquhar CM, Mitchell MD, Ponnampalam AP (2010) Epigenetic regulation of endometrium during the menstrual cycle. *Mol Hum Reprod* 16:297–310
68. Sasaki M, Kotcherguina L, Dharia A, Fujimoto S, Dahiya R (2001) Cytosine-phosphoguanine methylation of estrogen receptors in endometrial cancer. *Cancer Res* 61:3262–3266
69. Hori M, Iwasaki M, Shimazaki J, Inagawa S, Itabashi M (2000) Assessment of hypermethylated DNA in two promoter regions of the estrogen receptor alpha gene in human endometrial diseases. *Gynecol Oncol* 76:89–96
70. Sasaki M, Dharia A, Oh BR, Tanaka Y, Fujimoto S, Dahiya R (2001) Progesterone receptor B gene inactivation and CpG hypermethylation in human uterine endometrial cancer. *Cancer Res* 61:97–102
71. Ghabreau L, Roux JP, Niveleau A, Fontaniere B, Mahe C, Mokni M, Frappart L (2004) Correlation between the DNA global methylation status and progesterone receptor expression in normal endometrium, endometrioid adenocarcinoma and precursors. *Virchow Arch Int J Pathol* 445:129–134
72. Su RW, Strug MR, Jeong JW, Miele L, Fazleabas AT (2016) Aberrant activation of canonical Notch1 signaling in the mouse uterus decreases progesterone receptor by hypermethylation and leads to infertility. *Proc Natl Acad Sci USA* 113:2300–2305
73. Gao R, Ding Y, Liu X, Chen X, Wang Y, Long C, Li S, Guo L, He J (2012) Effect of folate deficiency on promoter methylation and gene expression of *Esr1*, *Cdh1* and *Pgr*, and its influence on endometrial receptivity and embryo implantation. *Hum Reprod* 27:2756–2765
74. Milesi MM, Varayoud J, Ramos JG, Luque EH (2017) Uterine ERalpha epigenetic modifications are induced by the endocrine disruptor endosulfan in female rats with impaired fertility. *Mol Cell Endocrinol* 454:1–11
75. Taylor HS, Arici A, Olive D, Igarashi P (1998) HOXA10 is expressed in response to sex steroids at the time of implantation in the human endometrium. *J Clin Invest* 101:1379–1384
76. Gui Y, Zhang J, Yuan L, Lessey BA (1999) Regulation of HOXA10 and its expression in normal and abnormal endometrium. *Mol Hum Reprod* 5:866–873
77. Satokata I, Benson G, Maas R (1995) Sexually dimorphic sterility phenotypes in *Hoxa10*-deficient mice. *Nature* 374:460–463
78. Benson GV, Lim H, Paria BC, Satokata I, Dey SK, Maas RL (1996) Mechanisms of reduced fertility in *Hoxa-10* mutant mice: uterine homeosis and loss of maternal *Hoxa-10* expression. *Development* 122:2687–2696
79. Yoshida H, Broaddus R, Cheng W, Xie S, Naora H (2006) Deregulation of the HOXA10 homeobox gene in endometrial carcinoma: role in epithelial-mesenchymal transition. *Cancer Res* 66:889–897
80. Wu Y, Halverson G, Basir Z, Strawn E, Yan P, Guo SW (2005) Aberrant methylation at HOXA10 may be responsible for its aberrant expression in the endometrium of patients with endometriosis. *Am J Obstet Gynecol* 193:371–380



81. Leu YW, Yan PS, Fan M, Jin VX, Liu JC, Curran EM, Welshons WV, Wei SH, Davuluri RV, Plass C, Nephew KP, Huang TH (2004) Loss of estrogen receptor signaling triggers epigenetic silencing of downstream targets in breast cancer. *Cancer Res* 64:8184–8192
82. Metivier R, Gallais R, Tiffoche C, Le Peron C, Jurkowska RZ, Carmouche RP, Ibberson D, Barath P, Demay F, Reid G, Benes V, Jeltsch A, Gannon F, Salbert G (2008) Cyclical DNA methylation of a transcriptionally active promoter. *Nature* 452:45–50
83. Houshdaran S, Zelenko Z, Irwin JC, Giudice LC (2014) Human endometrial DNA methylome is cycle-dependent and is associated with gene expression regulation. *Mol Endocrinol* 28:1118–1135
84. Horne AW, Lalani EN, Margara RA, White JO (2006) The effects of sex steroid hormones and interleukin-1-beta on MUC1 expression in endometrial epithelial cell lines. *Reproduction* 131:733–742
85. Meseguer M, Aplin JD, Caballero-Campo P, O'Connor JE, Martin JC, Remohi J, Pellicer A, Simon C (2001) Human endometrial mucin MUC1 is up-regulated by progesterone and down-regulated in vitro by the human blastocyst. *Biol Reprod* 64:590–601
86. Rahnama F, Thompson B, Steiner M, Shafiei F, Lobie PE, Mitchell MD (2009) Epigenetic regulation of E-cadherin controls endometrial receptivity. *Endocrinology* 150:1466–1472
87. Vincent ZL, Farquhar CM, Mitchell MD, Ponnampalam AP (2011) Expression and regulation of DNA methyltransferases in human endometrium. *Fertil Steril* 95(1522–1525):e1521
88. Yamagata Y, Asada H, Tamura I, Lee L, Maekawa R, Taniguchi K, Taketani T, Matsuoka A, Tamura H, Sugino N (2009) DNA methyltransferase expression in the human endometrium: down-regulation by progesterone and estrogen. *Hum Reprod* 24:1126–1132
89. van Kaam KJ, Delvoux B, Romano A, D'Hooghe T, Dunselman GA, Groothuis PG (2011) Deoxyribonucleic acid methyltransferases and methyl-CpG-binding domain proteins in human endometrium and endometriosis. *Fertil Steril* 95:1421–1427
90. Kukushkina V, Modhukur V, Suhorutsenko M, Peters M, Magi R, Rahmioglu N, Velthut-Meikas A, Altmae S, Esteban FJ, Vilo J, Zondervan K, Salumets A, Laisk-Podar T (2017) DNA methylation changes in endometrium and correlation with gene expression during the transition from pre-receptive to receptive phase. *Sci Rep* 7:3916
91. Maekawa R, Tamura I, Shinagawa M, Mihara Y, Sato S, Okada M, Taketani T, Tamura H, Sugino N (2019) Genome-wide DNA methylation analysis revealed stable DNA methylation status during decidualization in human endometrial stromal cells. *BMC Genomics* 20:324
92. Serra MJ, Ledford BE, Baggett B (1979) Synthesis and modification of the histones during the decidual cell reaction in the mouse uterus. *Biol Reprod* 20:214–220
93. Libby PR (1972) Histone acetylation and hormone action. Early effects of oestradiol-17beta on histone acetylation in rat uterus. *Biochem J* 130:663–669
94. Guo JZ, Gorski J (1989) Estrogen effects on modifications of chromatin proteins in the rat uterus. *J Steroid Biochem* 32:13–20
95. Sakai N, Maruyama T, Sakurai R, Masuda H, Yamamoto Y, Shimizu A, Kishi I, Asada H, Yamagoe S, Yoshimura Y (2003) Involvement of histone acetylation in ovarian steroid-induced decidualization of human endometrial stromal cells. *J Biol Chem* 278:16675–16682
96. Gunin AG, Kapitova IN, Suslonova NV (2005) Effects of histone deacetylase inhibitors on estradiol-induced proliferation and hyperplasia formation in the mouse uterus. *J Endocrinol* 185:539–549
97. Uchida H, Maruyama T, Nagashima T, Asada H, Yoshimura Y (2005) Histone deacetylase inhibitors induce differentiation of human endometrial adenocarcinoma cells through up-regulation of glycodelin. *Endocrinology* 146:5365–5373
98. Grimaldi G, Christian M, Quenby S, Brosens JJ (2012) Expression of epigenetic effectors in decidualizing human endometrial stromal cells. *Mol Hum Reprod* 18:451–458
99. Uchida H, Maruyama T, Ohta K, Ono M, Arase T, Kagami M, Oda H, Kajitani T, Asada H, Yoshimura Y (2007) Histone deacetylase inhibitor-induced glycodelin enhances the initial step of implantation. *Hum Reprod* 22:2615–2622
100. Zhu LH, Sun LH, Hu YL, Jiang Y, Liu HY, Shen XY, Jin XY, Zhen X, Sun HX, Yan GJ (2013) PCAF impairs endometrial receptivity and embryo implantation by down-regulating beta3-integrin expression via HOXA10 acetylation. *J Clin Endocrinol Metab* 98:4417–4428
101. Xin Q, Kong S, Yan J, Qiu J, He B, Zhou C, Ni Z, Bao H, Huang L, Lu J, Xia G, Liu X, Chen ZJ, Wang C, Wang H (2018) Polycomb subunit BMI1 determines uterine progesterone responsiveness essential for normal embryo implantation. *J Clin Invest* 128:175–189
102. Filant J, Spencer TE (2014) Uterine glands: biological roles in conceptus implantation, uterine receptivity and decidualization. *Int J Dev Biol* 58:107–116
103. Kelleher AM, Milano-Foster J, Behura SK, Spencer TE (2018) Uterine glands coordinate on-time embryo implantation and impact endometrial decidualization for pregnancy success. *Nat Commun* 9:2435
104. Cui T, He B, Kong S, Zhou C, Zhang H, Ni Z, Bao H, Qiu J, Xin Q, Reinberg D, Lydon JP, Lu J, Wang H (2017) PR-Set7 deficiency limits uterine epithelial population growth hampering postnatal gland formation in mice. *Cell Death Differ* 24:2013–2021
105. Li Q, Kannan A, DeMayo FJ, Lydon JP, Cooke PS, Yamagishi H, Srivastava D, Bagchi MK, Bagchi IC (2011) The antiproliferative action of progesterone in uterine epithelium is mediated by Hand2. *Science* 331:912–916
106. Barron F, Woods C, Kuhn K, Bishop J, Howard MJ, Clouthier DE (2011) Downregulation of Dlx5 and Dlx6 expression by Hand2 is essential for initiation of tongue morphogenesis. *Development* 138:2249–2259
107. Daikoku T, Cha J, Sun X, Tranguch S, Xie H, Fujita T, Hirota Y, Lydon J, DeMayo F, Maxson R, Dey SK (2011) Conditional deletion of Msx homeobox genes in the uterus inhibits blastocyst implantation by altering uterine receptivity. *Dev Cell* 21:1014–1025
108. Wang J, Kumar RM, Biggs VJ, Lee H, Chen Y, Kagey MH, Young RA, Abate-Shen C (2011) The Msx1 homeoprotein recruits polycomb to the nuclear periphery during development. *Dev Cell* 21:575–588
109. Creighton CJ, Benham AL, Zhu H, Khan MF, Reid JG, Nagaraja AK, Fountain MD, Dziadek O, Han D, Ma L, Kim J, Hawkins SM, Anderson ML, Matzuk MM, Gunaratne PH (2010) Discovery of novel microRNAs in female reproductive tract using next generation sequencing. *PLoS One* 5:e9637
110. Kuokkanen S, Chen B, Ojalvo L, Benard L, Santoro N, Pollard JW (2010) Genomic profiling of microRNAs and messenger RNAs reveals hormonal regulation in microRNA expression in human endometrium. *Biol Reprod* 82:791–801
111. Hu SJ, Ren G, Liu JL, Zhao ZA, Yu YS, Su RW, Ma XH, Ni H, Lei W, Yang ZM (2008) MicroRNA expression and regulation in mouse uterus during embryo implantation. *J Biol Chem* 283:23473–23484
112. Chakrabarty A, Tranguch S, Daikoku T, Jensen K, Furneaux H, Dey SK (2007) MicroRNA regulation of cyclooxygenase-2 during embryo implantation. *Proc Natl Acad Sci USA* 104:15144–15149

113. Revel A, Achache H, Stevens J, Smith Y, Reich R (2011) MicroRNAs are associated with human embryo implantation defects. *Hum Reprod* 26:2830–2840
114. Moreno-Moya JM, Vilella F, Martinez S, Pellicer A, Simon C (2014) The transcriptomic and proteomic effects of ectopic over-expression of miR-30d in human endometrial epithelial cells. *Mol Hum Reprod* 20:550–566
115. Ponsuksili S, Tesfaye D, Schellander K, Hoelker M, Hadlich F, Schwerin M, Wimmers K (2014) Differential expression of miRNAs and their target mRNAs in endometria prior to maternal recognition of pregnancy associates with endometrial receptivity for in vivo- and in vitro-produced bovine embryos. *Biol Reprod* 91:135
116. Kresowik JD, Devor EJ, Van Voorhis BJ, Leslie KK (2014) MicroRNA-31 is significantly elevated in both human endometrium and serum during the window of implantation: a potential biomarker for optimum receptivity. *Biol Reprod* 91:17
117. Liu JL, Liang XH, Su RW, Lei W, Jia B, Feng XH, Li ZX, Yang ZM (2012) Combined analysis of microRNome and 3'-UTRome reveals a species-specific regulation of progesterone receptor expression in the endometrium of rhesus monkey. *J Biol Chem* 287:13899–13910
118. Vilella F, Moreno-Moya JM, Balaguer N, Grasso A, Herrero M, Martinez S, Marcilla A, Simon C (2015) Hsa-miR-30d, secreted by the human endometrium, is taken up by the pre-implantation embryo and might modify its transcriptome. *Development* 142:3210–3221
119. Desrochers LM, Bordeleau F, Reinhart-King CA, Cerione RA, Antonyak MA (2016) Microvesicles provide a mechanism for intercellular communication by embryonic stem cells during embryo implantation. *Nat Commun* 7:11958
120. Galliano D, Pellicer A (2014) MicroRNA and implantation. *Fertil Steril* 101:1531–1544
121. Liu W, Niu Z, Li Q, Pang RT, Chiu PC, Yeung WS (2016) MicroRNA and Embryo Implantation. *Am J Reprod Immunol* 75:263–271
122. Liang J, Wang S, Wang Z (2017) Role of microRNAs in embryo implantation. *Reprod Biol Endocrinol* 15:90
123. Haraguchi H, Saito-Fujita T, Hirota Y, Egashira M, Matsumoto L, Matsuo M, Hiraoka T, Koga K, Yamauchi N, Fukayama M, Bartos A, Cha J, Dey SK, Fujii T, Osuga Y (2014) MicroRNA-200a locally attenuates progesterone signaling in the cervix, preventing embryo implantation. *Mol Endocrinol* 28:1108–1117
124. Chau YM, Pando S, Taylor HS (2002) HOXA11 silencing and endogenous HOXA11 antisense ribonucleic acid in the uterine endometrium. *J Clin Endocrinol Metab* 87:2674–2680
125. Arao Y, Carpenter K, Hewitt S, Korach KS (2010) Estrogen down-regulation of the *Sex* gene is mediated by the opposing strand-overlapping gene *Bop1*. *J Biol Chem* 285:4806–4814
126. Mihalich A, Reina M, Mangioni S, Ponti E, Alberti L, Vigano P, Vignali M, Di Blasio AM (2003) Different basic fibroblast growth factor and fibroblast growth factor-antisense expression in eutopic endometrial stromal cells derived from women with and without endometriosis. *J Clin Endocrinol Metab* 88:2853–2859
127. Noonan FC, Goodfellow PJ, Staloch LJ, Mutch DG, Simon TC (2003) Antisense transcripts at the *EMX2* locus in human and mouse. *Genomics* 81:58–66
128. Lanz RB, Chua SS, Barron N, Soder BM, DeMayo F, O'Malley BW (2003) Steroid receptor RNA activator stimulates proliferation as well as apoptosis in vivo. *Mol Cell Biol* 23:7163–7176
129. Sigurgeirsson B, Amark H, Jemt A, Ujvari D, Westgren M, Lundberg J, Gidlöf S (2017) Comprehensive RNA sequencing of healthy human endometrium at two time points of the menstrual cycle. *Biol Reprod* 96:24–33
130. Hu S, Yao G, Wang Y, Xu H, Ji X, He Y, Zhu Q, Chen Z, Sun Y (2014) Transcriptomic changes during the pre-receptive to receptive transition in human endometrium detected by RNA-Seq. *J Clin Endocrinol Metab* 99:E2744–E2753
131. Wang Q, Wang N, Cai R, Zhao F, Xiong Y, Li X, Wang A, Lin P, Jin Y (2017) Genome-wide analysis and functional prediction of long non-coding RNAs in mouse uterus during the implantation window. *Oncotarget* 8:84360–84372
132. Wang Y, Hu T, Wu L, Liu X, Xue S, Lei M (2017) Identification of non-coding and coding RNAs in porcine endometrium. *Genomics* 109:43–50
133. Zeng H, Fan X, Liu N (2017) Expression of H19 imprinted gene in patients with repeated implantation failure during the window of implantation. *Arch Gynecol Obstet* 296:835–839
134. Feng C, Shen JM, Lv PP, Jin M, Wang LQ, Rao JP, Feng L (2018) Construction of implantation failure related lncRNA-mRNA network and identification of lncRNA biomarkers for predicting endometrial receptivity. *Int J Biol Sci* 14:1361–1377
135. Zhang L, Liu X, Che S, Cui J, Liu Y, An X, Cao B, Song Y (2018) CircRNA-9119 regulates the expression of prostaglandin-endoperoxide synthase 2 (*PTGS2*) by sponging miR-26a in the endometrial epithelial cells of dairy goat. *Reprod Fertil Dev* 30(12):1759–1769
136. Liu L, Li L, Ma X, Yue F, Wang Y, Wang L, Jin P, Zhang X (2017) Altered circular RNA expression in patients with repeated implantation failure. *Cell Physiol Biochem* 44:303–313
137. Logan PC, Ponnampalam AP, Rahnema F, Lobie PE, Mitchell MD (2010) The effect of DNA methylation inhibitor 5-Aza-2'-deoxycytidine on human endometrial stromal cells. *Hum Reprod* 25:2859–2869
138. Gao F, Ma X, Rusie A, Hemingway J, Ostmann AB, Chung D, Das SK (2012) Epigenetic changes through DNA methylation contribute to uterine stromal cell decidualization. *Endocrinology* 153:6078–6090
139. Ding YB, Long CL, Liu XQ, Chen XM, Guo LR, Xia YY, He JL, Wang YX (2012) 5-aza-2'-deoxycytidine leads to reduced embryo implantation and reduced expression of DNA methyltransferases and essential endometrial genes. *PLoS One* 7:e45364
140. Logan PC, Ponnampalam AP, Steiner M, Mitchell MD (2013) Effect of cyclic AMP and estrogen/progesterone on the transcription of DNA methyltransferases during the decidualization of human endometrial stromal cells. *Mol Hum Reprod* 19:302–312
141. Brown LY, Bonney EA, Raj RS, Nielsen B, Brown S (2013) Generalized disturbance of DNA methylation in the uterine decidua in the CBA/J x DBA/2 mouse model of pregnancy failure. *Biol Reprod* 89:120
142. Yu M, Du G, Xu Q, Huang Z, Huang X, Qin Y, Han L, Fan Y, Zhang Y, Han X, Jiang Z, Xia Y, Wang X, Lu C (2018) Integrated analysis of DNA methylome and transcriptome identified CREB5 as a novel risk gene contributing to recurrent pregnancy loss. *EBioMedicine* 35:334–344
143. Hou W, Li Z, Li Y, Fang L, Li X, Huang J, Li X, You Z (2016) Correlation between protein expression of FOXP3 and level of FOXP3 promoter methylation in recurrent spontaneous abortion. *J Obstet Gynaecol Res* 42:1439–1444
144. Jones A, Teschendorff AE, Li Q, Hayward JD, Kannan A, Mould T, West J, Zikan M, Cibula D, Fiegl H, Lee SH, Wik E, Hadwin R, Arora R, Lemech C, Turunen H, Pakarinen P, Jacobs IJ, Salvesen HB, Bagchi MK, Bagchi IC, Widschwendner M (2013) Role of DNA methylation and epigenetic silencing of *HAND2* in endometrial cancer development. *PLoS Med* 10:e1001551
145. Estella C, Herrer I, Atkinson SP, Quinonero A, Martinez S, Pellicer A, Simon C (2012) Inhibition of histone deacetylase activity in human endometrial stromal cells promotes extracellular matrix remodelling and limits embryo invasion. *PLoS One* 7:e30508
146. He H, Kong S, Liu F, Zhang S, Jiang Y, Liao Y, Jiang Y, Li Q, Wang B, Zhou Z, Wang H, Huo R (2015) *Rbbp7* is required for uterine stromal decidualization in mice. *Biol Reprod* 93:13

147. Tamura I, Sato S, Okada M, Tanabe M, Lee L, Maekawa R, Asada H, Yamagata Y, Tamura H, Sugino N (2014) Importance of C/EBP $\beta$  binding and histone acetylation status in the promoter regions for induction of IGFBP-1, PRL, and Mn-SOD by cAMP in human endometrial stromal cells. *Endocrinology* 155:275–286
148. Tamura I, Ohkawa Y, Sato T, Suyama M, Jozaki K, Okada M, Lee L, Maekawa R, Asada H, Sato S, Yamagata Y, Tamura H, Sugino N (2014) Genome-wide analysis of histone modifications in human endometrial stromal cells. *Mol Endocrinol* 28:1656–1669
149. Tamura I, Jozaki K, Sato S, Shirafuta Y, Shinagawa M, Maekawa R, Taketani T, Asada H, Tamura H, Sugino N (2018) The distal upstream region of insulin-like growth factor-binding protein-1 enhances its expression in endometrial stromal cells during decidualization. *J Biol Chem* 293:5270–5280
150. Grimaldi G, Christian M, Steel JH, Henriot P, Poutanen M, Brosens JJ (2011) Down-regulation of the histone methyltransferase EZH2 contributes to the epigenetic programming of decidualizing human endometrial stromal cells. *Mol Endocrinol* 25:1892–1903
151. Nancy P, Tagliani E, Tay CS, Asp P, Levy DE, Erlebacher A (2012) Chemokine gene silencing in decidual stromal cells limits T cell access to the maternal-fetal interface. *Science* 336:1317–1321
152. Nancy P, Siewiera J, Rizzuto G, Tagliani E, Osokine I, Manandhar P, Dolgalev I, Clementi C, Tsirigos A, Erlebacher A (2018) H3K27me3 dynamics dictate evolving uterine states in pregnancy and parturition. *J Clin Invest* 128:233–247
153. Fang X, Ni N, Lydon JP, Ivanov I, Bayless KJ, Rijnkels M, Li Q (2019) Enhancer of Zeste 2 polycomb repressive complex 2 subunit is required for uterine epithelial integrity. *Am J Pathol* 189:1212–1225
154. Bian F, Gao F, Kartashov AV, Jegga AG, Barski A, Das SK (2016) Polycomb repressive complex 1 controls uterine decidualization. *Sci Rep* 6:26061
155. Estella C, Herrero I, Moreno-Moya JM, Quinonero A, Martinez S, Pellicer A, Simon C (2012) miRNA signature and Dicer requirement during human endometrial stromal decidualization in vitro. *PLoS One* 7:e41080
156. Shah KM, Webber J, Carzaniga R, Taylor DM, Fusi L, Clayton A, Brosens JJ, Hartshorne G, Christian M (2013) Induction of microRNA resistance and secretion in differentiating human endometrial stromal cells. *J Mol Cell Biol* 5:67–70
157. Zhang Q, Zhang H, Jiang Y, Xue B, Diao Z, Ding L, Zhen X, Sun H, Yan G, Hu Y (2015) MicroRNA-181a is involved in the regulation of human endometrial stromal cell decidualization by inhibiting Kruppel-like factor 12. *Reprod Biol Endocrinol* 13:23
158. Tochigi H, Kajihara T, Mizuno Y, Mizuno Y, Tamaru S, Kamei Y, Okazaki Y, Brosens JJ, Ishihara O (2017) Loss of miR-542-3p enhances IGFBP-1 expression in decidualizing human endometrial stromal cells. *Sci Rep* 7:40001
159. Nan X, Ng HH, Johnson CA, Laherty CD, Turner BM, Eisenman RN, Bird A (1998) Transcriptional repression by the methyl-CpG-binding protein MeCP2 involves a histone deacetylase complex. *Nature* 393:386–389
160. Jones PL, Veenstra GJ, Wade PA, Vermaak D, Kass SU, Landsberger N, Strouboulis J, Wolffe AP (1998) Methylated DNA and MeCP2 recruit histone deacetylase to repress transcription. *Nat Genet* 19:187–191
161. Lehnertz B, Ueda Y, Derijck AA, Braunschweig U, Perez-Burgos L, Kubicek S, Chen T, Li E, Jenuwein T, Peters AH (2003) Suv39 h-mediated histone H3 lysine 9 methylation directs DNA methylation to major satellite repeats at pericentric heterochromatin. *Curr Biol* 13:1192–1200
162. Vire E, Brenner C, Deplus R, Blanchon L, Fraga M, Didelot C, Morey L, Van Eynde A, Bernard D, Vanderwinden JM, Bollen M, Esteller M, Di Croce L, de Launoit Y, Fuks F (2006) The polycomb group protein EZH2 directly controls DNA methylation. *Nature* 439:871–874
163. Delcuve GP, Rastegar M, Davie JR (2009) Epigenetic control. *J Cell Physiol* 219:243–250
164. Januchowski R, Dabrowski M, Ofori H, Jagodzinski PP (2007) Trichostatin A down-regulate DNA methyltransferase 1 in Jurkat T cells. *Cancer Lett* 246:313–317
165. Barnhart K, Dunsmoor-Su R, Coutifaris C (2002) Effect of endometriosis on in vitro fertilization. *Fertil Steril* 77:1148–1155
166. Olive DL, Schwartz LB (1993) Endometriosis. *N Engl J Med* 328:1759–1769
167. Matsuzaki S, Canis M, Darcha C, Pouly JL, Mage G (2009) HOXA-10 expression in the mid-secretory endometrium of infertile patients with either endometriosis, uterine fibromas or unexplained infertility. *Hum Reprod* 24:3180–3187
168. Taylor HS, Bagot C, Kardana A, Olive D, Arici A (1999) HOX gene expression is altered in the endometrium of women with endometriosis. *Hum Reprod* 14:1328–1331
169. Lee B, Du H, Taylor HS (2009) Experimental murine endometriosis induces DNA methylation and altered gene expression in eutopic endometrium. *Biol Reprod* 80:79–85
170. Kim JJ, Taylor HS, Lu Z, Ladhani O, Hastings JM, Jackson KS, Wu Y, Guo SW, Fazleabas AT (2007) Altered expression of HOXA10 in endometriosis: potential role in decidualization. *Mol Hum Reprod* 13:323–332
171. Szczepanska M, Wirstlein P, Luczak M, Jagodzinski PP, Skrzypczak J (2010) Reduced expression of HOXA10 in the midluteal endometrium from infertile women with minimal endometriosis. *Biomed Pharmacother* 64:697–705
172. Wu Y, Strawn E, Basir Z, Halverson G, Guo SW (2007) Aberrant expression of deoxyribonucleic acid methyltransferases DNMT1, DNMT3A, and DNMT3B in women with endometriosis. *Fertil Steril* 87:24–32
173. Wu Y, Strawn E, Basir Z, Halverson G, Guo SW (2006) Promoter hypermethylation of progesterone receptor isoform B (PR-B) in endometriosis. *Epigenetics* 1:106–111
174. Izawa M, Harada T, Taniguchi F, Ohama Y, Takenaka Y, Terakawa N (2008) An epigenetic disorder may cause aberrant expression of aromatase gene in endometriotic stromal cells. *Fertil Steril* 89:1390–1396
175. Xue Q, Lin Z, Cheng YH, Huang CC, Marsh E, Yin P, Milad MP, Confino E, Reierstad S, Innes J, Bulun SE (2007) Promoter methylation regulates estrogen receptor 2 in human endometrium and endometriosis. *Biol Reprod* 77:681–687
176. Colon-Diaz M, Baez-Vega P, Garcia M, Ruiz A, Monteiro JB, Fourquet J, Bayona M, Alvarez-Garriga C, Achille A, Seto E, Flores I (2012) HDAC1 and HDAC2 are differentially expressed in endometriosis. *Reprod Sci* 19:483–492
177. Monteiro JB, Colon-Diaz M, Garcia M, Gutierrez S, Colon M, Seto E, Laboy J, Flores I (2014) Endometriosis is characterized by a distinct pattern of histone 3 and histone 4 lysine modifications. *Reprod Sci* 21:305–318
178. Colon-Caraballo M, Monteiro JB, Flores I (2015) H3K27me3 is an epigenetic mark of relevance in endometriosis. *Reprod Sci* 22:1134–1142
179. Rackow BW, Jorgensen E, Taylor HS (2011) Endometrial polyps affect uterine receptivity. *Fertil Steril* 95:2690–2692
180. Kulp JL, Mamillapalli R, Taylor HS (2016) Aberrant HOXA10 methylation in patients with common gynecologic disorders: implications for reproductive outcomes. *Reprod Sci* 23:455–463
181. Daftary GS, Taylor HS (2002) Hydrosalpinx fluid diminishes endometrial cell HOXA10 expression. *Fertil Steril* 78:577–580

182. Saito T, Nishimura M, Yamasaki H, Kudo R (2003) Hypermethylation in promoter region of E-cadherin gene is associated with tumor dedifferentiation and myometrial invasion in endometrial carcinoma. *Cancer* 97:1002–1009
183. Memczak S, Papavasileiou P, Peters O, Rajewsky N (2015) Identification and characterization of circular RNAs as a new class of putative biomarkers in human blood. *PLoS One* 10:e0141214
184. Li MQ, Yao MN, Yan JQ, Li ZL, Gu XW, Lin S, Hu W, Yang ZM (2017) The decline of pregnancy rate and abnormal uterine responsiveness of steroid hormones in aging mice. *Reprod Biol* 17:305–311
185. Woods L, Perez-Garcia V, Kieckbusch J, Wang X, DeMayo F, Colucci F, Hemberger M (2017) Decidualisation and placenta-tion defects are a major cause of age-related reproductive decline. *Nat Commun* 8:352
186. Jefferson WN, Chevalier DM, Phelps JY, Cantor AM, Padilla-Banks E, Newbold RR, Archer TK, Kinyamu HK, Williams CJ (2013) Persistently altered epigenetic marks in the mouse uterus after neonatal estrogen exposure. *Mol Endocrinol* 27:1666–1677
187. Franczak A, Zglejc K, Waszkiewicz E, Wojciechowicz B, Martyniak M, Sobotka W, Okrasa S, Kotwica G (2017) Periconceptional undernutrition affects in utero methyltransferase expression and steroid hormone concentrations in uterine flushings and blood plasma during the peri-implantation period in domestic pigs. *Reprod Fertil Dev* 29:1499–1508
188. Zglejc K, Franczak A (2017) Peri-conceptional under-nutrition alters the expression of TRIM28 and ZFP57 in the endometrium and embryos during peri-implantation period in domestic pigs. *Reprod Domest Anim* 52:542–550

**Publisher's Note** Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.