

# UC Berkeley

## UC Berkeley Previously Published Works

**Title**

On imbedding differentiable manifolds in Euclidean space

**Permalink**

<https://escholarship.org/uc/item/1jh3g3fz>

**Journal**

Annals of Mathematics, 73(3)

**ISSN**

1939-8980

**Author**

Hirsch, MW

**Publication Date**

1961-05-01

Peer reviewed

## Annals of Mathematics

---

On Imbedding Differentiable Manifolds in Euclidean Space

Author(s): Morris W. Hirsch

Source: *Annals of Mathematics*, Second Series, Vol. 73, No. 3 (May, 1961), pp. 566-571

Published by: [Annals of Mathematics](#)

Stable URL: <http://www.jstor.org/stable/1970318>

Accessed: 11/11/2014 22:26

---

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <http://www.jstor.org/page/info/about/policies/terms.jsp>

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact support@jstor.org.



*Annals of Mathematics* is collaborating with JSTOR to digitize, preserve and extend access to *Annals of Mathematics*.

<http://www.jstor.org>

## ON IMBEDDING DIFFERENTIABLE MANIFOLDS IN EUCLIDEAN SPACE\*

BY MORRIS W. HIRSCH

(Received May 25, 1960)

### 1. Introduction

Recently R. Penrose, J. H. C. Whitehead and E. C. Zeeman have obtained powerful results on rectilinear imbeddings of combinatorial manifolds in euclidean  $n$ -space  $R^n$  [4]. The purpose of the present paper is to show how their methods can be used to obtain information about differentiable imbeddings of smooth manifolds in  $R^n$ . We also use results due to J. H. C. Whitehead [6].

Our chief results are the following.

**THEOREM 4.6.** *A differentiable open  $n$ -manifold can be differentially imbedded in  $R^{2n-1}$ .*

**THEOREM 4.7.** *A differentiable open  $n$ -manifold which is parallelizable can be immersed in  $R^n$ .*

**THEOREM 4.1.** *Let  $M$  be a closed differentiable  $n$ -manifold, and  $x$  a point of  $M$ . Assume  $M$  has a smooth triangulation. If  $M$  is  $(m-1)$ -connected,  $2 \leq 2m \leq n$ , and if a neighborhood of the  $(n-m)$ -skeleton can be immersed in  $R^q$ ,  $q \geq 2n - 2m + 1$ , then  $M - x$  can be differentially imbedded in  $R^q$ .*

The power of 4.1 comes from the fact that much is known concerning the immersibility of a neighborhood of a skeleton [1]. For example, if  $M$  is  $m-1$  connected, a neighborhood of the  $(n-m)$ -skeleton is always immersible in  $R^{2n-m}$ ; if  $M$  is a  $\pi$ -manifold,  $M$  is immersible in  $R^{n+1}$ . Some applications are given in 4.2 - 4.5.

The author is indebted to the late Professor J. H. C. Whitehead for many fruitful discussions of the ideas presented here.

### 2. Terminology

We assume familiarity with the concept of a *differentiable manifold*. Any such manifold possesses a *smooth triangulation* [7], making the

---

\* This research was supported by the United States Air Force through the Air Force Office of Scientific Research of the Air Research and Development Command, under Contract No. AF49 (638)-253. Reproduction in whole or in part is permitted for any purpose of the United States Government.

manifold homeomorphic to a *combinatorial manifold*; for this last and related notions, see [4]. *Differentiable* means of class  $C^1$ . A *differentiable imbedding*  $f: M \rightarrow N$  of one differentiable manifold in another is a differentiable homeomorphism of  $M$  onto a submanifold of  $N$  such that  $f^{-1}$  is differentiable. If, in addition  $f(M) = N$ , we say  $f$  is a *diffeomorphism*. An *immersion* of one differentiable manifold  $M$  in another is a differentiable map whose jacobian matrix has rank everywhere equal to  $\dim M$ . We use  $R$  for the real line,  $R^n$  for euclidean  $n$ -space,  $R'$  for the non-negative ray. We consider  $R^{n-1}$  as contained in  $R^n$  in the usual way, and  $R^n = R^{n-1} \times R$ . We define *half- $n$ -space* to be  $R^{n-1} \times R' = F^n \subset R^n$ . We give  $R^n$  and  $F^n$  their usual differentiable and combinatorial structures. If  $M$  is a combinatorial or differentiable manifold, we denote its boundary by  $\partial M$ . Thus  $\partial F^n = R^{n-1}$ . If  $M$  is a combinatorial manifold, an  *$n$ -element in  $M$*  is the image of an  $n$ -simplex in  $M$  under a piecewise linear imbedding. If  $M$  is a differentiable manifold, an  *$n$ -disk in  $M$*  is the image in  $M$  of the unit ball of  $R^n$  under a differentiable imbedding. The concepts of a *combinatorial half- $n$ -space in  $M$*  and a *differentiable half- $n$ -space in  $M$*  are similarly defined, with the restriction that the latter be a closed subset of  $M$ . A manifold is *closed* if it is compact and has no boundary. It is *open* if it is not compact and has no boundary. It is *bounded* if it has a boundary. We use  $\text{int } X$  for the interior of  $X$ , and  $\text{cl } (X)$  for the closure of  $X$ .

### 3. Combinatorial results and their differentiable implications

The main results of this section are 3.3 and 3.7.

**THEOREM 3.1.** *Let  $M$  be a closed  $(m-1)$ -connected combinatorial  $n$ -manifold,  $2 \leq 2m \leq n$ . If  $U$  is a neighborhood of the  $n-m$  skeleton of rectilinear triangulation of  $M$ , there is an  $n$ -element  $E$  in  $M$  such that  $M - \text{int } E \subset U$ .*

**PROOF.** This is a corollary of Lemma 2.7 of [4].

**THEOREM 3.2.** *Let  $M$  be an open combinatorial  $n$ -manifold. There exists a subcomplex  $K$  of dimension  $n-1$  with the following property. Let  $U$  be a neighborhood of  $K$ . There exists a collection  $A_1, A_2, \dots$ , of mutually disjoint combinatorial half- $n$ -spaces in  $M$ , such that  $M - \bigcup \text{int } A_i$  is contained in  $U$ . There is a triangulation of  $M$  in which every  $A_i$  is a subcomplex.*

**PROOF.** This is essentially proved in the proof of theorem 3.2 of [6].

**THEOREM 3.3.** *Let  $M$  be a closed differentiable  $n$ -manifold which is  $m-1$  connected,  $2 \leq 2m \leq n$ . Let  $U$  be a neighborhood of the  $n-m$  skeleton*

of a smooth triangulation of  $M$ . There is an  $n$ -disk  $D$  in  $M$  such that  $M - \text{int } D \subset U$ .

PROOF. By 3.1, there exists an  $n$ -element in  $M$  (considered as a combinatorial manifold) such that  $M - \text{int } E \subset U$ . The interior of  $E$  is an open subset of  $M$  and is therefore a differentiable manifold. Since  $E$  is an  $n$ -element, there is a piecewise linear homeomorphism  $g: R^n \rightarrow \text{int } E$ . Since  $R^n$  is contractible, by a result of J. Munkres [3, 6.6], there is a diffeomorphism  $h: R^n \rightarrow \text{int } E$ . Since  $E$  is compact, there is a compact subset  $Q \subset \text{int } E$  such that  $M - Q \subset U$ . Let  $B$  be a ball in  $R^n$  so large that  $h^{-1}(Q) \subset \text{int } B$ . Then  $D = h(B)$  is the required disk.

LEMMA 3.4. *Let  $M$  be an open differentiable  $n$ -manifold with a smooth triangulation. There exists a subcomplex  $K$  of dimension  $n - 1$  with the following property. If  $U$  is a neighborhood of  $K$ , there exist a collection  $B_1, B_2, \dots$  of mutually disjoint differentiable half- $n$ -spaces in  $M$  such that  $M - \bigcup \text{int } B_i$  is contained in  $U$ . Each  $B_i$  is contained in an open set  $V_i$  of  $M$  such that  $V_i \cap V_j$  is empty for  $i \neq j$ .*

PROOF. Let  $K$  be the  $(n - 1)$ -dimensional subcomplex of  $M$  described in 3.2. Let  $U$  be a neighborhood of  $K$ . Let  $A_1, A_2, \dots$  be combinatorial half  $n$ -spaces in  $M$  as in 3.2. For each  $i$  let  $V_i$  be a neighborhood of  $A_i$  such that  $V_i \cap V_j$  is empty for  $i \neq j$ . We shall find a differentiable half- $n$ -space  $B_i \subset V_i$  such that  $V_i - \text{int } B_i \subset U$ , and doing this for  $i = 1, 2, \dots$  prove the lemma. It is enough to consider a single  $i$ . Replacing  $V_i$  by  $M, V_i \cap U$  by  $U$ , and  $A_i$  by  $A$ , we must prove

LEMMA 3.5. *Let  $M$  be an open differentiable  $n$ -manifold and  $A$  a combinatorial half- $n$ -space in  $M$ . Let  $U$  be an open set in  $M$  such that  $M - \text{int } A \subset U$ . There is a differentiable half- $n$ -space  $B$  in  $M$  such that  $M - \text{int } B \subset U$ .*

PROOF. Let  $f: F^n \rightarrow M$  be a piecewise linear imbedding such that  $f(F^n) = A$ . Let  $d$  be a metric on  $M$ . Let  $\delta: F^n \rightarrow R$  be a continuous function such that  $\delta(x, 0) = 0, \delta(x, y) > 0$  if  $y > 0$ . (Recall that  $F^n = R^{n-1} \times R'$ .) We apply [3, 6.2] and conclude that there exists a diffeomorphism  $g: \text{int } F^n \rightarrow \text{int } A$  such that  $d(g(x), f(x)) < \delta(x)$  for every  $x \in F^n$ . Using the properties of  $\delta$  and the paracompactness of  $R^{n-1} = \partial F^n$ , we choose a positive differentiable function  $\alpha: R^{n-1} \rightarrow R$  such that if  $(x, y) \in F^n$  and  $0 \leq y \leq \alpha(x)$ , then  $g(x, y) \in U$ . Define  $h: F^n \rightarrow M$  by  $h(x, y) = g(x, y + \alpha(x))$ . Then  $h$  is a differentiable imbedding and  $M - \text{int } h(F^n) \subset U$ . Thus  $B = h(F^n)$  is the required differentiable half- $n$ -space.

LEMMA 3.6. *Let  $B$  be a differentiable half- $n$ -space in an open differentiable manifold  $M$ . Let  $V$  be a neighborhood of  $B$ . There is a diffeomor-*

phism  $f: M \rightarrow M - B$  such that if  $x \in M - V$ ,  $f(x) = x$ .

PROOF. Let  $h_0: F^n \rightarrow B$  be a diffeomorphism. We may find a neighborhood  $U$  of  $F^n$  in  $R^n$  and a differentiable imbedding  $h: U \rightarrow M$  such that  $\text{cl}(h(U)) \subset V$ , and  $h|_{F^n} = h_0$ ; see e.g., [2]. The lemma follows from the easily proved fact that there is a diffeomorphism  $g: R^n \rightarrow R^n - F^n$  such that  $g(x) = x$  for  $x \in R^n - U$ .

**THEOREM 3.7.** *Let  $M$  be an open differentiable  $n$ -manifold with a smooth triangulation. Let  $U$  be a neighborhood of the  $n - 1$  skeleton. There is a differentiable imbedding  $f: M \rightarrow U$ .*

PROOF. Let  $K$  be the  $(n - 1)$ -dimensional subcomplex of  $M$  of 3.4. Let  $B_1, B_2, \dots$ , and  $V_1, V_2, \dots$  be as in 3.4, taking  $U$  of 3.7 for  $U$  of 3.4. By 3.6, for each  $i = 1, 2, \dots$ , there is a diffeomorphism  $f_i: M \rightarrow M - B_i$  which leaves  $M - V_i$  fixed. Define  $f: M \rightarrow M - \bigcup B_i$  by

$$f(x) = \begin{cases} x & \text{if } x \in M - \bigcup V_i, \\ f_i(x) & \text{if } x \in V_i. \end{cases}$$

Since the  $V_i$  are disjoint,  $f$  is well defined. Since  $M - \bigcup B_i \subset U$ ,  $f$  is as required.

It is perhaps worth noting that the analogue of 3.7 for  $M$  compact and bounded can be proved without using [6] or J. H. C. Whitehead's "regular neighborhood" theory. This can be done by successively "pushing in" the  $n$ -simplices of  $M$ , starting with those that meet  $\partial M$ , until all the  $n$ -simplices have been deformed into a neighborhood of an  $(n - 1)$ -dimensional subcomplex.

#### 4. Applications

From now on all manifolds and imbeddings will be differentiable.

**THEOREM 4.1.** *Let  $M$  be a closed  $(m - 1)$ -connected  $n$ -manifold,  $2 \leq 2m \leq n$ , with a smooth triangulation; let  $x$  be a point of  $M$ . If a neighborhood of the  $n - m$  skeleton can be immersed in  $R^q$  for  $q \geq 2n - 2m + 1$ , then  $M - x$  can be imbedded in  $R^q$ .*

PROOF. Let  $M_{n-m}$  denote the  $n - m$  skeleton. Let  $V$  be a neighborhood of  $M_{n-m}$  and  $f: V \rightarrow R^q$  an immersion. Since  $2(n - m) < q$ , there is an approximation  $g$  to  $f$  which is also an immersion and which imbeds  $M_{n-m}$ ; see Theorem 2(e) of [9]. A compactness argument shows that there is a smaller neighborhood  $U$  of  $M_{n-m}$  such that  $g: U \rightarrow R^q$  is an imbedding. Let  $D$  be an  $n$ -disk in  $M$  such that  $M - \text{int } D \subset U$ ; this is possible by 3.3. Thus  $g: M - D \rightarrow R^q$  is an imbedding. It is easy to construct a diffeomorphism  $h: M \rightarrow M$  such that  $h(x) \in \text{int } D$ . Let  $v: M - h(x) \rightarrow M - D$  be a diffeomorphism. Then  $gvh: M - x \rightarrow R^q$  is the required imbedding.

**COROLLARY 4.2.**  *$M - x$  can be imbedded in  $R^{2n-m}$ .*

**PROOF.** By 4.1, it suffices to immerse a neighborhood of  $M_{n-m}$  in  $R^{2n-m}$ . This can be done by [1, 6.2].

A  $\pi$ -manifold is one which is imbeddable in some  $R^q$  with a trivial normal bundle. It is well known that a parallelizable manifold is a  $\pi$ -manifold [8].

**COROLLARY 4.3.** *If  $M$  is as in 4.1 and if  $M - x$  is a  $\pi$ -manifold,  $M - x$  is imbeddable in  $R^{2n-2m+1}$ .*

**PROOF.** By 4.1, it suffices to immerse  $M - x$  in  $R^{2n-2m+1}$ . By 6.5 of [1],  $M - x$  is immersible in  $R^{n+1} \subset R^{2n-2m+1}$ .

**COROLLARY 4.4.**<sup>1</sup> *If  $M$  is a closed homotopy  $n$ -sphere (i.e.,  $M$  is  $n - 1$  connected),  $M - x$  is imbeddable in  $R^{n+2}$  if  $n$  is odd, and in  $R^{n+1}$  if  $n$  is even.*

**PROOF.**  $M - x$  is parallelizable and  $(m - 1)$ -connected, where  $2m = n - 1$  or  $n$ , and 4.3 applies.

The double of a bounded manifold  $M$  is the manifold  $N$  obtained by identifying two copies of  $M$  along their boundaries. We consider  $M$  as a submanifold of  $N$  in an obvious way. One can obtain information about a bounded compact manifold by applying the preceding results to the double. For example,

**THEOREM 4.5.**<sup>1</sup> *Let  $M$  be a compact bounded  $n$ -manifold whose double is  $N$ . Suppose  $M$  is contractible. Then  $M$  is imbeddable in  $R^{n+2}$  or  $R^{n+1}$ , and  $N$  in  $R^{n+3}$  or  $R^{n+2}$ , according to whether  $n$  is odd or even.*

**PROOF.** Let  $x$  be an element of  $N$ . It is easy to see that  $N$  is a homotopy  $n$ -sphere, so  $N - x$ , and hence  $M \subset N - x$ , is imbeddable in  $R^q$ , where  $q = n + 2$  if  $n$  is odd,  $q = n + 1$  if  $n$  is even. Thus  $M \subset R^q \subset R^{q+1}$ . We may deform  $M$  in  $R^{q+1}$  so that  $M \subset F^{q+1}$ ,  $M \cap R^q = \partial M$ , and the rays in  $F^{q+1}$  normal to  $R^q$  are tangent to  $M$ . The double of  $F^{q+1}$  is  $R^{q+1}$ , and the double  $N$  of  $M$  is thus naturally contained in  $R^{q+1}$ . The imbedding of  $N$  can easily be made differentiable; it is already so except along  $\partial M$ , where the tangent planes to the two halves of  $N$  coincide.

**THEOREM 4.6.** *A non-closed  $n$ -manifold  $M$  can be imbedded in  $R^{2n-1}$ .*

**PROOF.** It is enough to assume  $M$  is open. Whitney [10] proves that  $M$  can be immersed in  $R^{2n-1}$ . Reasoning as in the beginning of the proof of 4.1, we see that a neighborhood  $U$  of the  $(n - 1)$ -skeleton (of a smooth triangulation) can be imbedded. Let  $g: U \rightarrow R^{2n-1}$  be such an imbedding.

<sup>1</sup> In the case  $n \geq 5$ , these results are superseded by the recent solutions of the generalized Poincaré conjecture by Smale [12] and Stallings [11]. Using the results of either of these works, it can be shown that  $M - x$  of 4.4, and  $M$  of 4.5, can be imbedded in  $R^n$ .

By 3.7 there is an imbedding  $f: M \rightarrow U$ . Then  $gf: M \rightarrow R^{2n-1}$  is the required imbedding.

**THEOREM 4.7.** *A non-closed parallelizable  $n$ -manifold  $M$  is immersible in  $R^n$ .*

**PROOF.** Again it suffices to take  $M$  open. It follows from 5.7 of [1] that a neighborhood of the  $(n-1)$ -skeleton of a smooth triangulation can be immersed in  $R^n$ . (Although the hypothesis of the reference requires  $\dim M < n$ , this is not used in the proof if  $L$  of the reference has dimension less than  $n$ .) Applying 3.7 proves 4.7.

This was proved for  $n = 3$  by J. H. C. Whitehead [6], assuming merely orientability instead of parallelizability.

The following theorem does not seem to be in the literature. It is usually "proved" by saying, "push the singularities to infinity". It was pointed out to the author by J. H. C. Whitehead.

**THEOREM 4.8.** *If  $M$  is a non-closed  $n$ -manifold there exists a differentiable function on  $M$  without critical points.*

**PROOF.** We may assume  $M$  is open. A theorem due to M. Morse states that there exists a differentiable map  $f: M \rightarrow R$  with only isolated critical points; see e.g., [5, p. 72]. We may assume none of these points lies on the  $(n-1)$ -skeleton of a smooth triangulation of  $M$ . The conclusion follows from 3.7.

INSTITUTE FOR ADVANCED STUDY

#### BIBLIOGRAPHY

1. M. W. HIRSCH, *Immersions of manifolds*, Trans. Amer. Math. Soc., 93 (1959), 242-276.
2. J. MILNOR, On the relationship between differentiable and combinatorial manifolds, Princeton University, 1956, mimeographed.
3. J. MUNKRES, *Obstructions to the smoothing of piecewise-differentiable homeomorphisms*, Ann. of Math., 72 (1960), 521-554.
4. R. PENROSE, J. H. C. WHITEHEAD and E. C. ZEEMAN, *Imbedding of manifolds in euclidean space*, Ann. of Math., 73 (1961), 613-623.
5. R. THOM, *Les singularités des applications différentiables*, Ann. Inst. Fourier, Grenoble, XI (1956), 43-87.
6. J. H. C. WHITEHEAD, *The immersion of an open 3-manifold in euclidean 3-space*, to appear.
7. ———, *On  $C^1$ -complexes*, Ann. of Math., 41 (1940), 809-824.
8. ———, *On the homotopy type of manifolds*, Ann. of Math., 41 (1940), 825-832.
9. H. WHITNEY, *Differentiable manifolds*, Ann. of Math., 37 (1936), 645-680.
10. ———, *Singularities of a smooth  $n$ -manifold in  $2n-1$  space*, Ann. of Math., 45 (1954), 247-293.
11. J. STALLINGS, *Polyhedral homotopy-spheres*, Bull. Amer. Math. Soc., to appear.
12. S. SMALE, *The generalized Poincaré conjecture in higher dimensions*, Bull. Amer. Math. Soc., 66 (1960), 373-375.