

UCLA

UCLA Previously Published Works

Title

Neurobiological Markers of Resilience to Early-Life Adversity During Adolescence

Permalink

<https://escholarship.org/uc/item/1jw0t1hx>

Journal

Biological Psychiatry Cognitive Neuroscience and Neuroimaging, 6(2)

ISSN

2451-9022

Authors

Méndez Leal, Adriana S
Silvers, Jennifer A

Publication Date

2021-02-01

DOI

10.1016/j.bpsc.2020.08.004

Peer reviewed

Adolescent Resilience to Early Life Adversity

Neurobiological Markers of Resilience to Early Life Adversity During
Adolescence

Adriana S. Méndez Leal and Jennifer A. Silvers*

For Consideration at *Biological Psychiatry: Cognitive Neuroscience and
Neuroimaging*
Special issue on Imaging Risk and Resilience

Word count: 3,476

Figures: 1

Tables: 0

Keywords: Early life adversity, resilience, adolescence, puberty, emotion,
neurodevelopment

*Corresponding Author:

Jennifer A. Silvers, Ph.D.

Department of Psychology, University of California, Los Angeles

1285 Franz Hall, Box 951563, Los Angeles, California, 90095, USA

Email: silvers@ucla.edu

Phone: 310-206-9024

Abstract

Early life adversity (ELA) exposure (including trauma, abuse, neglect or institutional care) is a precursor to poor physical and mental health outcomes, and is implicated in 30% of adult mental illness. In recent decades, ELA research has increasingly focused on characterizing factors that confer resilience to ELA, and on identifying opportunities for intervention. In this review, we describe recent behavioral and neurobiological resilience work that suggests adolescence (a period marked by heightened plasticity, development of key neurobiological circuitry, and sensitivity to the social environment) may be a particularly opportune moment for ELA intervention. We review intrapersonal factors associated with resilience that become increasingly important during adolescence (specifically, reward processing, affective learning, and self-regulation), and describe the contextual factors (family, peers, and broader social environment) that modulate them. Additionally, we describe how the onset of puberty interacts with each of these factors, and explore recent findings that point to possible “pubertal recalibration” of ELA exposure as an opportunity for intervention. Lastly, we conclude by describing considerations and future directions for resilience research in adolescents, with a focus on understanding developmental trajectories using dimensional, holistic models of resilience.

Early life adversity (ELA) exposure is associated with poor mental and physical health outcomes, and is implicated in 30% of adult mental illness (1-3). Exposure to ELA (e.g., trauma, abuse, neglect or institutional care) is the normative experience in the United States and worldwide, with half of children reporting one or more such events (3,4). However, individuals vary greatly in their responses to ELA, and many demonstrate resilience in one or more domains following ELA exposure. Current ELA research is increasingly focused on identifying factors that promote resilience or may be targets for concurrent and retrospective intervention (5).

Adolescence is a developmental period marked by heightened plasticity, sensitivity to the social environment, and the rapid development of critical functions related to self-regulation, reward processing, and affective learning (6). Adolescence may present a unique opportunity for cultivating resilience by targeting the aforementioned domains, particularly in light of recent evidence that puberty provides a “recalibration” window for specific biological systems following ELA (7). Intervention during adolescence may be particularly beneficial given that the transitions associated with this developmental stage are normatively stressful, and the onset of anxiety, mood, and substance use disorders is most common during this period (8).

This review highlights individual and contextual factors¹ that promote resilience, which we define as positive physical and mental health outcomes following ELA (5,9). According to recent scientific consensus, resilience is not a personality trait (and not the responsibility of individuals to cultivate), but rather various domain-specific adaptations that improve post-ELA outcomes in the short or long term (9). Under this definition, resilience in a given domain arises from individual and contextual factors that contribute to adaptation to and/or recovery from exposures, and may vary over time. As a result, we operationalize resilience as distinct from the inverse of an individual’s vulnerability (9).

In neurodevelopment, resilience can manifest in reduced impact of ELA on a circuit, through neurobehavioral adaptations that promote better outcomes, or through adaptations that confer risk or benefit depending on the current context. These processes are modulated by local, community, and societal level contextual factors. Rather than providing a systematic

¹ Given our focus on neurodevelopmental effects of ELA, we use “resilience factors” to describe both characteristics that may be stable or trait-like in adulthood (e.g. cognitive emotion regulation ability or capacity) and the adaptive behaviors these traits may promote in practice (e.g. use of emotion regulation ability to reduce a response to a particular stressor), as these have yet to be fully parsed in the developmental literature (see (9) for discussion in adults).

review of all factors associated with resilience, here we highlight a subset of domains fundamental to adolescent neurodevelopment that are promising candidates for intervention: specifically, reward-processing, affective learning, and self-regulation (6). An overview of these resilience factors, the neural circuits they rely on, and candidate techniques for intervention during adolescence are depicted in Figure 1. While we focus primarily on neurobiological and behavioral resilience markers, we also highlight other systems implicated in stress neurobiology, including the immune and neuroendocrine systems. Given that this literature is still nascent, we review resilience to all ELA, but when possible identify which dimensions of ELA – threat or deprivation, for example – may benefit from a particular resilience factor (10,11). Lastly, we describe future directions for this area of research.

Individual Factors Affecting Resilience

Reward Processing

Adolescence is marked by increased behavioral and neural reward sensitivity, defined as enhanced arousal in response to and heightened motivation to seek rewards (13). While increased reward sensitivity is often described as a risk factor in adolescents (13), it also confers benefits (maximizing exploration and reward optimization, for instance; 14), and in adversity-exposed adolescents may be a source of resilience. Across species, reward processing is guided by a mesocorticolimbic dopaminergic system that includes the basal ganglia (including the ventral striatum, commonly associated with reward processing), the orbitofrontal cortex (involved in contingency representation and reward learning), ventromedial prefrontal cortex (vmPFC; linked to self-referential thinking, reward processing, and emotional learning), and limbic regions including the amygdala (implicated in affective valuation, particularly threat and reward) and the hippocampus (associated with learning and memory, alongside biological tuning of the hypothalamic-pituitary-adrenal, or HPA, axis, which coordinates stress responses) (15,16). Functional magnetic resonance imaging (fMRI) studies of adolescents indicate that early deprivation (e.g. institutional orphanage care) and trauma dampen behavioral and neural sensitivity to rewards and weaken reward-based learning (17–19). These effects are linked to anhedonia, which mediates relationships between ELA and psychopathology, social challenges, and substance abuse (17,20–24).

Conversely, heightened reward sensitivity (behavioral and neural, particularly in the striatum) predicts concurrent and longitudinal resilience for adolescents exposed to ELA, in part through reductions in anhedonia (25–28). Reward sensitivity also increases positive affect and the propensity to appraise events positively following ELA, which in turn predict better mental

and physical health (26,29,30).² The aforementioned findings suggest that reward processing interventions may support adolescent ELA resilience. Candidate interventions include positive affect treatment, which enhances reward anticipation, reward learning, and savoring of positive experiences (32), and behavioral activation therapy (currently being evaluated in maltreated adolescents), which emphasizes positive reinforcement to reduce anhedonia and enhance motivation and pleasure (33,34).

Affective Learning

Preliminary evidence from fear conditioning research in humans and non-human animals points to fear learning as a possible ELA intervention target. Converging classical conditioning work has implicated the amygdala in producing and storing fear memories, the hippocampus in context learning, and various prefrontal cortex (PFC) regions (including vmPFC) in increasingly inhibiting fear expression across adolescence (35). Across species, adolescents exposed to ELA exhibit accelerated development of fear learning behaviors and medial PFC, hippocampus and amygdala circuitry (10,36). In human adolescents, childhood maltreatment has been associated with threat discrimination difficulty and reduced amygdala and hippocampal volumes (37). By contrast, youth exposed to deprivation (specifically previously institutionalization; PI) show a positive relationship between aversive learning behavior and anxiety, though those that show stronger (i.e., more “mature”) vmPFC-hippocampus functional coupling are buffered against anxiety two years later (38). Similarly, maternally separated rats that exhibit increased amygdala-PFC functional coupling during fear learning display decreased anxiety behaviors (described in 39). However, current evidence is more mixed about whether non-affective forms of memory and learning that rely on hippocampus-PFC circuits also contribute to resilience (31). Overall, the extant data suggest that accelerated development of fear learning may confer resilience for adolescents exposed to ELA (particularly deprivation), and thus, extinction-enhancing exposures might improve outcomes in this population (34).

Self-Regulation

Successful self-regulation (including cognitive control and regulation of negative emotion) may also buffer against ELA. Emotion regulation is defined as a collection of implicit (automatic) and explicit (deliberate) strategies that modify the intensity, valence or timing of an emotional response (40), and cognitive control as mechanisms (including response inhibition, cognitive flexibility, attentional control and working memory) that facilitate overriding automatic mental or behavioral responses in accordance with one’s goals (41).

² While early threat exposure may alter related systems like the salience network, such evidence is limited, particularly in regards to adolescent resilience (31).

Although cognitive control functions emerge during childhood, they undergo protracted development during adolescence (42,43). These functions are supported by a broad network that includes fronto-parietal regions (including the dorsolateral PFC, dorsomedial PFC, the anterior cingulate cortex, and superior parietal cortex), the basal ganglia, and the thalamus (42). Neuropsychological testing, electroencephalography, and fMRI evidence suggests that exposure to both threat and deprivation diminishes behavioral and neurocognitive markers of cognitive control (44-47),³ but also that ELA-exposed youth with strong cognitive control display fewer internalizing symptoms (10,49).

While less commonly explored, emerging work suggests cognitive control may contribute to neurodevelopmental resilience during adolescence. Initial fMRI studies have shown decreased PFC recruitment during cognitive control in adolescents exposed to early threat (50-53) and deprivation (46). Though not an explicit evaluation of neural functioning during cognitive control, a large-scale structural analysis (N = 1870) found that adolescents who were resilient across social, academic, and risk-taking domains had increased gray matter volumes in multiple dorsolateral PFC regions associated with cognitive control (54). Cognitive control may also interact with other resilience factors during adolescence: one recent study found that adolescents with high amygdala threat responses and decreased striatum reward responses reported increased anxiety, unless they displayed robust dlPFC recruitment during executive control (55). ELA-exposed youth may therefore benefit from existing well-validated cognitive control interventions (e.g., task-switch and working memory training; (42).

A larger body of work has explored emotional control and regulation as sources of resilience to ELA. During adolescence, age-related improvements in implicit and explicit emotion regulation behavior are mirrored by the emergence of negative functional coupling between the amygdala and vmPFC and ventrolateral prefrontal cortex (vlPFC), respectively (56-58). Converging evidence from human lesion work, meta-analysis of fMRI emotion regulation studies, and neuroanatomical tracing in non-human primates suggests that negative PFC-amygdala functional coupling during emotional events likely reflects the PFC exerting inhibitory control over amygdala activity (59-62). Negative PFC-amygdala coupling is broadly considered the mature emotion regulation phenotype, as it emerges during the transition to adolescence and predicts both better regulation of negative affect and reduced risk for internalizing psychopathology (56,63-66). Similarly, disruption to (behavioral and neural) emotion regulation processes

³ Although some recent findings suggest cognitive control deficits following ELA may be linked to systemic barriers (e.g. low-SES) that increase risk of ELA, rather than ELA itself (48)

constitutes a transdiagnostic mechanism connecting ELA and psychopathology risk (67–70).

Although early deprivation and trauma reduce self-regulation ability and confer increased risk for psychopathology at the population level, in individuals there is marked variability in self-regulation capacity, and successful regulation appears to buffer against negative outcomes (71,72). In young adults who experienced early deprivation (e.g. low socioeconomic status), adaptive coping strategies confer benefits for both mental and physical health (e.g. reduced inflammatory activity) (73–75). A systematic review of child and adolescent studies found that emotion regulation success reduces psychopathology risk post-ELA (76), and use of adaptive emotion regulation strategies is associated with resilience to psychopathology in adolescents who experienced childhood poverty (77,78) or war (79).

Similarly, ELA-exposed individuals who display neural phenotypes associated with effective emotion regulation show greater resilience to psychopathology. This effect has primarily been observed in neuroimaging studies examining cognitive reappraisal, an explicit emotion regulation strategy that involves reinterpreting emotional events through lateral PFC regulation of amygdala responses (59). Although ELA (abuse and poverty) confers risk for weak lateral prefrontal recruitment during reappraisal (68,80), adolescents exposed to ELA who exhibit robust lateral prefrontal recruitment and attenuated amygdala reactivity during cognitive reappraisal are at decreased risk for depression (81) and anxiety (82).

While neuroimaging research on ELA and explicit emotion regulation is relatively nascent, far more work has examined interactions between ELA and implicit forms of emotion regulation, including discrimination learning, extinction, and automatic regulation of emotional threat responses. In contrast to reappraisal, which strongly recruits lateral PFC, implicit regulation processes rely heavily upon vmPFC-amygdala interactions (83). Cross-species models of caregiving deprivation suggest that ELA accelerates development of vmPFC-amygdala networks underlying implicit emotion regulation (84). For example, PI youth display negative (mature) vmPFC-amygdala functional coupling patterns during childhood, in contrast with the positive coupling observed in comparison children (38,85). Importantly, PI youth who demonstrate the mature phenotype display decreased anxiety symptoms relative to those who do not (38,85). Thus, accelerated development of vmPFC-amygdala circuitry may be an adaptive response to the need for self-regulation in the absence of a caregiver (84), given that parental presence tunes such networks in typical development (86,87).

Together, these findings highlight self-regulation training as a candidate intervention in adolescents. In particular, cognitive behavioral therapy (CBT) relies upon ingredients ascribed to both explicit (i.e.,

reframing emotional events) and implicit (i.e., gradual exposure to emotional triggers) emotion regulation (88). Trauma-focused CBT, which is well-validated in children, may promote resilience through relaxation and emotion regulation (34,89,90). Likewise, transdiagnostic treatment approaches that focus more squarely on reappraisal and relaxation⁴ may support self-regulation development (91,34). However, because such treatments target multiple domains, more work is needed to assess the utility of interventions specifically targeting emotion regulation (e.g. 92) to confer resilience.

Contextual Factors Affecting Resilience

Development of the individual resilience factors reviewed above is scaffolded by one's social context. Because adolescence confers increased sensitivity to the social environment (6), this developmental period provides a unique opportunity for the cultivation of resilience factors supported by the social context.

High-quality caregiving and a supportive family environment promote resilience following ELA. Caregivers buffer biological stress responses in childhood (93), and scaffold the development of affective (fear) learning, emotion regulation, and cognitive control (94-97), as well as the development of vmPFC-amygdala circuitry during childhood (84,87). Similarly, high-quality caregiving and attachment promote resilience during childhood and adolescence (30,98,99), although individuals vary in their tendency to demonstrate parental buffering effects to stress (86). The family environment (e.g., positive parenting, cohesion and involvement) plays a critical role in resilience and can buffer ELA-exposed adolescents against psychopathology (100). Critically, the family context during adolescence not only protects mental health concurrently – perhaps partially by diminishing sensitivity to other adversities (101) – but into adulthood as well. Indeed, one study found that a family-based parenting intervention in early adolescence was associated with improved self-regulation (and consequently, vmPFC-hippocampal functional coupling) at age 25 (102).

While far more work has examined how social support, particularly caregiving, sculpts fear behavior and vmPFC-amygdala circuitry, emerging evidence suggests positive social contexts also promote adaptive reward-motivated behavior. For example, caregiver presence increases adolescents' ventral striatal activity when making safe choices but decreases said activity during risky choices (103), and adolescents with a greater sense of family obligation display decreased striatal activity during risk taking (104). Although the presence of peers generally increases reward sensitivity in adolescents (105), in ELA-exposed youth peer effects are less well understood. While preliminary evidence suggests that as children become

⁴ e.g., the FIRST program, which promotes calmness, changing thoughts, and problem solving (91)

adolescents, peers increasingly modulate biological stress processes (106,107) and resilience factors (105,108–110), reported effects are mixed (and paradigms disproportionately employ social evaluative stressors over other stressors like shared threats). A systematic review found that peer relationships do not modulate adolescent resilience to ELA (100), although this may partially reflect social challenges experienced by ELA-exposed youth (111).

The broader social environment also affects resilience. Adolescents who belong to marginalized groups (particularly racial/ethnic minorities) often face ongoing discrimination and systemic barriers that prevent access to therapeutic resources. Such ongoing stress can affect physical and mental health in adolescence and beyond (112–114). While addressing oppression of minority and low-income groups will require large-scale sociopolitical changes, interventions that target systemic inequalities can confer great benefit. For example, regular small unconditional cash transfers to low-income families improve maternal mental health and reduce depression rates in adolescent boys (115,116). Likewise, animal models suggest that peripubertal environmental enrichment may reverse negative effects of ELA on hippocampal development, HPA-axis reactivity, cognitive functioning, and play behaviors (117–119). Additionally, interventions that target identity development may contribute to resilience in adolescents. Community-based interventions like the Strong African-American Families (SAAF) program, which supports positive parenting, body image and adolescent pride in racial identity have been shown to reduce risk-taking, depression, and conduct disorders, and improve neural, endocrine, immune, and biological aging outcomes more than a decade later (120,121,121–123). Given that identity development is a prominent developmental task during adolescence that relies on circuits underlying individual resilience factors described above (including vmPFC, dmPFC, ACC, and the striatum; 124), it may be a particularly important intervention focus (6). Community-based interventions like SAAF have the added benefit of cultivating resilience in families, which may support more sustainable resilience in youth. Parents of children exposed to ELA are disproportionately likely to have experienced adversity themselves, and community interventions may therefore foster resilience in both youth and their caregivers, potentially mitigating cross-generational adversity transmission (5).

Pubertal Stress Recalibration Hypothesis

In normative development, puberty is associated with dramatic changes in the brain, neuroendocrine system, and HPA axis (125,126). Puberty alters the function of each of these systems and their interactions – for example, circulating sex hormones and inflammatory markers predict brain function related to emotion regulation and executive control in adolescence (127,128). Accumulating cross-species work suggests that ELA

not only modulates brain and biology concurrently, but also may have ongoing or novel effects during and after puberty. In primates (including humans), trauma and psychosocial stress may even accelerate pubertal onset, particularly in girls (129–131). While the aforementioned work has focused on identifying mechanisms of risk, emerging evidence indicates puberty may offer a window of resilience for adolescents exposed to ELA (129,132,133). Recent research in internationally-adopted PI youth suggests that pubertal onset may present a stress recalibration opportunity for HPA axis functioning (specifically, cortisol reactivity to a social stressor) (7,134). These adolescents experienced extreme social deprivation during early HPA axis development, followed by high-resource environments post-adoption (135). While in this study PI children displayed blunted cortisol responses to stressors, as they progressed through puberty they began to show more normative cortisol responses, suggesting that puberty constitutes a second sensitive period for HPA development that facilitates recalibration to the current environment (7).

The onset of puberty (and thus the beginning of adolescence) may be an ideal intervention opportunity for developing the intrapersonal resilience factors reviewed above. Developmental trajectories of self-regulation appear synchronized to pubertal onset, pointing to a possible pubertal interaction: cognitive reappraisal ability develops rapidly around age 10 and increases across adolescence, and mPFC-amygdala functional coupling patterns associated with implicit emotion regulation ability shift from positive coupling during childhood to negative coupling (the adult phenotype) at approximately the same age (56–58), with similar patterns in cognitive control development (42,43). Similarly, recent developmental theory posits that because puberty initiates peak behavioral and neural reward sensitivity (136), adolescents may be particularly amenable to interventions utilizing “health promoting” rewards like prosocial behavior (132,137).

Puberty also marks a shift in environmental modulation of resilience factors. In childhood, caregiver presence reduces HPA reactivity to stressors, but this dampening diminishes after middle childhood – an effect driven primarily by pubertal onset, rather than chronological age (93,96,138). Likewise, while children display similar stress-induced cortisol responses in the presence of peers and strangers, peers increasingly modulate HPA stress responses in adolescence (although effects vary across populations) (106,107,139,140). Similarly, although parental presence scaffolds negative mPFC-amygdala coupling in children, it does not in adolescents, implying parents impact emotion regulation circuitry to a lesser degree after puberty (141). These findings motivate investigation of interactions between puberty and social context effects on resilience, which may facilitate intervention development (132).

Future Directions

Just as ELA research has embraced dimensional models of risk conferred by different features of stress (e.g., type, timing, severity, controllability) (11,142,143), so too has resilience work begun to adopt a similar approach, assessing resilience with consideration for the type and severity of ELA experienced, and jointly investigating “bottom-up” (e.g. polygenetic and epigenetic) and “top-down” resilience factors (e.g., social environment) (144). Evaluating functioning across biobehavioral systems may be particularly important for understanding resilience, as resilience in one domain may come at the expense of outcomes in another. For example, in low-SES youth, higher self-regulation predicts positive psychosocial outcomes, but also accelerated epigenetic aging (78). Similarly, a recent network analysis revealed that 10 commonly studied resilience factors were all positively associated in non-adversity-exposed adolescents, but showed more antagonistic associations in adversity-exposed adolescents (100). While recent work has probed interactions between neurobiological resilience and resilience in other systems (epigenetics, for example, see 145), there has not been systematic evaluation of neurobiological resilience factors within individuals (e.g., between accelerated development of emotion regulation circuitry as compared to fear learning circuitry; 146). Future work should evaluate the interplay between resilience factors (particularly following differing ELA exposures) to examine whether they are synergistic, antagonistic or orthogonal to each other. Lastly, these holistic approaches should assess possible pubertal recalibration of stress-related biobehavioral processes across domains (147, for example), and subsequent opportunities for tailored intervention work (7).

Future research should also probe how the biological manifestation and relative impact of specific resilience factors may change and interact across development to predict varying outcomes across the lifespan. Although evidence for sensitive periods of ELA exposure is mixed (despite indication that there may sensitive periods for the effects of caregiver deprivation), future investigations should consider that ELA may shift the timing of neural and biological sensitive periods, including puberty (148,129,38,149,150). These shifts may be adaptive in the short-term (e.g. the absence of a caregiver), but long-term effects are not well characterized. This is particularly relevant to adolescence, given the biological and social transitions that occur during this period. Lastly, parsing the contributions of ongoing and resolved sources of adversity may inform adolescent resilience research, given adolescence’s temporal proximity to childhood and the frequent co-occurrence of ELA and complex or chronic stress exposures (as with persistent systemic inequalities). Further evaluation of developmental timing effects may therefore inform resilience models and later intervention design.

Taken together, these findings point to the continued need to employ long-term, large-scale longitudinal studies that evaluate a) resilience given severity and type of ELA exposure b) resilience factors across multiple levels (genetic, immune, neurological, behavioral, and social), with an eye to pubertal effects, c) developmental trajectories for each resilience factor and d) the interplay between resilience sources across developmental stages. Information gleaned from this work may inform targeted interventions and improved understanding of resilience during development.

Acknowledgements

This work was supported by NSF Grant 1848004 awarded to JAS and NSF Graduate Research Fellowship Program Grant DGE-1650604 awarded to ASML.

Disclosures

AML and JAS have no disclosures to report.

References

1. Brown DW, Anda RF, Tiemeier H, Felitti VJ, Edwards VJ, Croft JB, Giles WH (2009): Adverse Childhood Experiences and the Risk of Premature Mortality. *Am J Prev Med* 37: 389–396.
2. Friedman EM, Karlamangla AS, Gruenewald T, Koretz B, Seeman TE (2015): Early Life Adversity and Adult Biological Risk Profiles. *Psychosom Med* 77: 176–185.
3. Kessler RC, McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky AM, et al. (2010): Childhood adversities and adult psychopathology in the WHO World Mental Health Surveys. *Br J Psychiatry* 197: 378–385.
4. McLaughlin KA, Green JG, Gruber MJ, Sampson NA, Zaslavsky AM, Kessler RC (2012): Childhood Adversities and First Onset of Psychiatric Disorders in a National Sample of US Adolescents. *Arch Gen Psychiatry* 69: 1151–1160.
5. Traub F, Boynton-Jarrett R (2017): Modifiable Resilience Factors to Childhood Adversity for Clinical Pediatric Practice. *Pediatrics* 139. <https://doi.org/10.1542/peds.2016-2569>
6. National Academies of Sciences, Engineering, and Medicine (2019): *The Promise of Adolescence: Realizing Opportunity for All Youth*. National Academies Press.
7. Gunnar MR, DePasquale CE, Reid BM, Donzella B (2019): Pubertal stress recalibration reverses the effects of early life stress in postinstitutionalized children. *Proc Natl Acad Sci* 116: 23984–23988.
8. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE (2005): Lifetime Prevalence and Age-of-Onset Distributions of DSM-IV Disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry* 62: 593–602.
9. Kalisch R, Baker DG, Basten U, Boks MP, Bonanno GA, Brummelman E, et al. (2017): The resilience framework as a strategy to combat stress-related disorders [no. 11]. *Nat Hum Behav* 1: 784–790.
10. Machlin L, Miller AB, Snyder J, McLaughlin KA, Sheridan MA (2019): Differential Associations of Deprivation and Threat With Cognitive Control and Fear Conditioning in Early Childhood. *Front Behav Neurosci* 13. <https://doi.org/10.3389/fnbeh.2019.00080>
11. McLaughlin KA, Sheridan MA (2016): Beyond Cumulative Risk: A Dimensional Approach to Childhood Adversity. *Curr Dir Psychol Sci* 25: 239–245.
12. Galván A (2013): The Teenage Brain: Sensitivity to Rewards. *Curr Dir Psychol Sci* 22: 88–93.
13. Steinberg L, Icenogle G, Shulman EP, Breiner K, Chein J, Bacchini D, et al. (2018): Around the world, adolescence is a time of heightened sensation seeking and immature self-regulation. *Dev Sci* 21: e12532.

14. Lloyd A, McKay R, Sebastian C, Balsters J (2020): Are Adolescents More Optimal Decision-Makers in Novel Environments? Examining Exploration and Learning in a Patch Foraging Paradigm. <https://doi.org/10.31234/osf.io/ypknq>
15. Haber SN, Knutson B (2010): The Reward Circuit: Linking Primate Anatomy and Human Imaging. *Neuropsychopharmacology* 35: 4–26.
16. Tottenham N, Sheridan MA (2010): A review of adversity, the amygdala and the hippocampus: a consideration of developmental timing. *Front Hum Neurosci* 3. <https://doi.org/10.3389/neuro.09.068.2009>
17. Fareri DS, Tottenham N (2016): Effects of early life stress on amygdala and striatal development. *Dev Cogn Neurosci* 19: 233–247.
18. Gerin MI, Puetz VB, Blair RJR, White S, Sethi A, Hoffmann F, *et al.* (2017): A neurocomputational investigation of reinforcement-based decision making as a candidate latent vulnerability mechanism in maltreated children. *Dev Psychopathol* 29: 1689–1705.
19. Sheridan MA, McLaughlin KA, Winter W, Fox N, Zeanah C, Nelson CA (2018): Early deprivation disruption of associative learning is a developmental pathway to depression and social problems [no. 1]. *Nat Commun* 9: 1–8.
20. Fries ABW, Pollak SD (2017): The role of learning in social development: Illustrations from neglected children. *Dev Sci* 20: e12431.
21. Goff B, Gee DG, Telzer EH, Humphreys KL, Gabard-Durnam L, Flannery J, Tottenham N (2013): Reduced nucleus accumbens reactivity and adolescent depression following early-life stress. *Neuroscience* 249: 129–138.
22. Guyer AE, Kaufman J, Hodgdon HB, Masten CL, Jazbec S, Pine DS, Ernst M (2006): Behavioral Alterations in Reward System Function: The Role of Childhood Maltreatment and Psychopathology. *J Am Acad Child Adolesc Psychiatry* 45: 1059–1067.
23. Hanson JL, Hariri AR, Williamson DE (2015): Blunted Ventral Striatum Development in Adolescence Reflects Emotional Neglect and Predicts Depressive Symptoms. *Biol Psychiatry* 78: 598–605.
24. Hanson JL, Albert D, Iselin A-MR, Carré JM, Dodge KA, Hariri AR (2016): Cumulative stress in childhood is associated with blunted reward-related brain activity in adulthood. *Soc Cogn Affect Neurosci* 11: 405–412.
25. Corral-Frías NS, Nikolova YS, Michalski LJ, Baranger D a. A, Hariri AR, Bogdan R (2015): Stress-related anhedonia is associated with ventral striatum reactivity to reward and transdiagnostic psychiatric symptomatology. *Psychol Med* 45: 2605–2617.
26. Corral-Frías NS, Nadel L, Fellous J-M, Jacobs WJ (2016): Behavioral and self-reported sensitivity to reward are linked to stress-related differences in positive affect. *Psychoneuroendocrinology* 66: 205–213.
27. Dennison MJ, Sheridan MA, Busso DS, Jenness JL, Peverill M, Rosen ML, McLaughlin KA (2016): Neurobehavioral markers of resilience to

- depression amongst adolescents exposed to child abuse. *J Abnorm Psychol* 125: 1201-1212.
28. Geschwind N, Peeters F, Jacobs N, Delespaul P, Derom C, Thiery E, et al. (2010): Meeting risk with resilience: high daily life reward experience preserves mental health. *Acta Psychiatr Scand* 122: 129-138.
 29. Steptoe A, Wardle J, Marmot M (2005): Positive affect and health-related neuroendocrine, cardiovascular, and inflammatory processes. *Proc Natl Acad Sci* 102: 6508-6512.
 30. Vantieghem MR, Gabard-Durnam L, Goff B, Flannery J, Humphreys KL, Telzer EH, et al. (2017): Positive valence bias and parent-child relationship security moderate the association between early institutional caregiving and internalizing symptoms. *Dev Psychopathol* 29: 519-533.
 31. McLaughlin KA, Weissman D, Bitrán D (2019): Childhood Adversity and Neural Development: A Systematic Review. *Annu Rev Dev Psychol* 1: 277-312.
 32. Craske MG, Meuret AE, Ritz T, Treanor M, Dour HJ (2016): Treatment for Anhedonia: A Neuroscience Driven Approach. *Depress Anxiety* 33: 927-938.
 33. Dimidjian S, Hollon SD, Dobson KS, Schmaling KB, Kohlenberg RJ, Addis ME, et al. (2006): Randomized trial of behavioral activation, cognitive therapy, and antidepressant medication in the acute treatment of adults with major depression. *J Consult Clin Psychol* 74: 658-670.
 34. McLaughlin KA, DeCross SN, Jovanovic T, Tottenham N (2019): Mechanisms linking childhood adversity with psychopathology: Learning as an intervention target. *Behav Res Ther* 118: 101-109.
 35. Callaghan BL, Meyer H, Opendak M, Van Tieghem M, Harmon C, Li A, et al. (2019): Using A Developmental Ecology Framework to Align Fear Neurobiology Across Species. *Annu Rev Clin Psychol* 15: null.
 36. Muhammad A, Carroll C, Kolb B (2012): Stress during development alters dendritic morphology in the nucleus accumbens and prefrontal cortex. *Neuroscience* 216: 103-109.
 37. McLaughlin KA, Sheridan MA, Gold AL, Duys A, Lambert HK, Peverill M, et al. (2016): Maltreatment Exposure, Brain Structure, and Fear Conditioning in Children and Adolescents [no. 8]. *Neuropsychopharmacology* 41: 1956-1964.
 38. Silvers JA, Lumian DS, Gabard-Durnam L, Gee DG, Goff B, Fareri DS, et al. (2016): Previous Institutionalization Is Followed by Broader Amygdala-Hippocampal-PFC Network Connectivity during Aversive Learning in Human Development. *J Neurosci* 36: 6420-6430.
 39. Brenhouse HC, Bath KG (2019): Bundling the haystack to find the needle: Challenges and opportunities in modeling risk and resilience following early life stress. *Front Neuroendocrinol* 54: 100768.
 40. Thompson RA (1994): Emotion Regulation: A Theme in Search of Definition. *Monogr Soc Res Child Dev* 59: 25-52.

41. Miller EK (2000): The prefrontal cortex and cognitive control [no. 1]. *Nat Rev Neurosci* 1: 59–65.
42. Karbach J, Unger K (2014): Executive control training from middle childhood to adolescence. *Front Psychol* 5. <https://doi.org/10.3389/fpsyg.2014.00390>
43. Luna B (2009): Developmental Changes in Cognitive Control through Adolescence. In: Bauer P, editor. *Advances in Child Development and Behavior*, vol. 37. JAI, pp 233–278.
44. Bos KJ, Fox N, Zeanah CH, Nelson CA (2009): Effects of early psychosocial deprivation on the development of memory and executive function. *Front Behav Neurosci* 3. <https://doi.org/10.3389/neuro.08.016.2009>
45. McDermott JM, Westerlund A, Zeanah CH, Nelson CA, Fox NA (2012): Early adversity and neural correlates of executive function: Implications for academic adjustment. *Dev Cogn Neurosci* 2: S59–S66.
46. Mueller SC, Maheu FS, Dozier M, Peloso E, Mandell D, Leibenluft E, et al. (2010): Early-life stress is associated with impairment in cognitive control in adolescence: An fMRI study. *Neuropsychologia* 48: 3037–3044.
47. Spratt EG, Friedenbergl SL, Swenson CC, LaRosa A, De Bellis MD, Macias MM, et al. (2012): The Effects of Early Neglect on Cognitive, Language, and Behavioral Functioning in Childhood. *Psychol Irvine Calif* 3: 175–182.
48. Danese A, Moffitt TE, Arseneault L, Bleiberg BA, Dinardo PB, Gandelman SB, et al. (2016): The Origins of Cognitive Deficits in Victimized Children: Implications for Neuroscientists and Clinicians. *Am J Psychiatry* 174: 349–361.
49. Lambert HK, King KM, Monahan KC, McLaughlin KA (2017): Differential associations of threat and deprivation with emotion regulation and cognitive control in adolescence. *Dev Psychopathol* 29: 929–940.
50. Blair KS, Aloi J, Crum K, Meffert H, White SF, Taylor BK, et al. (2019): Association of Different Types of Childhood Maltreatment With Emotional Responding and Response Control Among Youths. *JAMA Netw Open* 2: e194604–e194604.
51. Hart H, Lim L, Mehta MA, Curtis C, Xu X, Breen G, et al. (2018): Altered Functional Connectivity of Fronto-Cingulo-Striatal Circuits during Error Monitoring in Adolescents with a History of Childhood Abuse. *Front Hum Neurosci* 12. <https://doi.org/10.3389/fnhum.2018.00007>
52. Lim L, Hart H, Mehta MA, Simmons A, Mirza K, Rubia K (2015): Neural Correlates of Error Processing in Young People With a History of Severe Childhood Abuse: An fMRI Study. *Am J Psychiatry* 172: 892–900.
53. Lim L, Hart H, Mehta MA, Simmons A, Mirza K, Rubia K (2016): Neurofunctional Abnormalities during Sustained Attention in Severe Childhood Abuse. *PLOS ONE* 11: e0165547.

54. Burt KB, Whelan R, Conrod PJ, Banaschewski T, Barker GJ, Bokde ALW, *et al.* (2016): Structural brain correlates of adolescent resilience. *J Child Psychol Psychiatry* 57: 1287–1296.
55. Scult MA, Knodt AR, Radtke SR, Brigidi BD, Hariri AR (2019): Prefrontal Executive Control Rescues Risk for Anxiety Associated with High Threat and Low Reward Brain Function. *Cereb Cortex* 29: 70–76.
56. Gee DG, Humphreys KL, Flannery J, Goff B, Telzer EH, Shapiro M, *et al.* (2013): A Developmental Shift from Positive to Negative Connectivity in Human Amygdala–Prefrontal Circuitry. *J Neurosci* 33: 4584–4593.
57. Silvers JA, McRae K, Gabrieli JDE, Gross JJ, Remy KA, Ochsner KN (2012): Age-Related Differences in Emotional Reactivity, Regulation, and Rejection Sensitivity in Adolescence. *Emot Wash DC* 12: 1235–1247.
58. Silvers JA, Insel C, Powers A, Franz P, Helion C, Martin RE, *et al.* (2017): vIPFC–vmPFC–Amygdala Interactions Underlie Age-Related Differences in Cognitive Regulation of Emotion. *Cereb Cortex* 27: 3502–3514.
59. Buhle JT, Silvers JA, Wager TD, Lopez R, Onyemekwu C, Kober H, *et al.* (2014): Cognitive Reappraisal of Emotion: A Meta-Analysis of Human Neuroimaging Studies. *Cereb Cortex* 24: 2981–2990.
60. Ghashghaei HT, Hilgetag CC, Barbas H (2007): Sequence of information processing for emotions based on the anatomic dialogue between prefrontal cortex and amygdala. *NeuroImage* 34: 905–923.
61. Salas CE, Gross JJ, Rafal RD, Viñas-Guasch N, Turnbull OH (2013): Concrete behaviour and reappraisal deficits after a left frontal stroke: A case study. *Neuropsychol Rehabil* 23: 467–500.
62. Salas CE, Radovic D, Yuen KSL, Yeates GN, Castro O, Turnbull OH (2014): “Opening an emotional dimension in me”: Changes in emotional reactivity and emotion regulation in a case of executive impairment after left fronto-parietal damage. *Bull Menninger Clin* 78: 301–334.
63. Fitzgerald JM, Klumpp H, Langenecker S, Phan KL (2019): Transdiagnostic neural correlates of volitional emotion regulation in anxiety and depression. *Depress Anxiety* 36: 453–464.
64. Lee H, Heller AS, van Reekum CM, Nelson B, Davidson RJ (2012): Amygdala–prefrontal coupling underlies individual differences in emotion regulation. *NeuroImage* 62: 1575–1581.
65. Perlman G, Simmons AN, Wu J, Hahn KS, Tapert SF, Max JE, *et al.* (2012): Amygdala response and functional connectivity during emotion regulation: A study of 14 depressed adolescents. *J Affect Disord* 139: 75–84.
66. Tottenham N, Gabard-Durnam LJ (2017): The developing amygdala: a student of the world and a teacher of the cortex. *Curr Opin Psychol* 17: 55–60.
67. Jenness JL, Peverill M, Miller AB, Heleniak C, Robertson MM, Sambrook KA, *et al.* (2020): Alterations in neural circuits underlying emotion regulation following child maltreatment: a mechanism underlying trauma-related psychopathology. *Psychol Med* 1–10.

68. Kim P, Evans GW, Angstadt M, Ho SS, Sripada CS, Swain JE, *et al.* (2013): Effects of childhood poverty and chronic stress on emotion regulatory brain function in adulthood. *Proc Natl Acad Sci* 110: 18442–18447.
69. Peverill M, Sheridan MA, Busso DS, McLaughlin KA (2019): Atypical Prefrontal–Amygdala Circuitry Following Childhood Exposure to Abuse: Links With Adolescent Psychopathology. *Child Maltreat* 24: 411–423.
70. Weissman DG, Bitran D, Miller AB, Schaefer JD, Sheridan MA, McLaughlin KA (2019): Difficulties with emotion regulation as a transdiagnostic mechanism linking child maltreatment with the emergence of psychopathology. *Dev Psychopathol* 31: 899–915.
71. Blair C, Raver CC (2012): Individual development and evolution: Experiential canalization of self-regulation. *Dev Psychol* 48: 647–657.
72. Tottenham N (2012): Risk and Developmental Heterogeneity in Previously Institutionalized Children. *J Adolesc Health* 51: S29–S33.
73. Chen E, Miller GE, Lachman ME, Gruenewald TL, Seeman TE (2012): Protective Factors for Adults from Low Childhood Socioeconomic Circumstances: The Benefits of Shift-and-Persist for Allostatic Load. *Psychosom Med* 74: 178–186.
74. Chen E, McLean KC, Miller GE (2015): Shift-and-Persist Strategies: Associations With Socioeconomic Status and the Regulation of Inflammation Among Adolescents and Their Parents. *Psychosom Med* 77: 371–382.
75. Chen E, Miller GE (2012): “Shift-and-Persist” Strategies: Why Low Socioeconomic Status Isn’t Always Bad for Health. *Perspect Psychol Sci* 7: 135–158.
76. Compas BE, Jaser SS, Bettis AH, Watson KH, Gruhn MA, Dunbar JP, *et al.* (2017): Coping, emotion regulation, and psychopathology in childhood and adolescence: A meta-analysis and narrative review. *Psychol Bull* 143: 939–991.
77. Buckner JC, Mezzacappa E, Beardslee WR (2003): Characteristics of resilient youths living in poverty: The role of self-regulatory processes. *Dev Psychopathol* 15: 139–162.
78. Miller GE, Yu T, Chen E, Brody GH (2015): Self-control forecasts better psychosocial outcomes but faster epigenetic aging in low-SES youth. *Proc Natl Acad Sci* 112: 10325–10330.
79. Amony-P’Olak K, Garnefski N, Kraaij V (2007): Adolescents caught between fires: Cognitive emotion regulation in response to war experiences in Northern Uganda. *J Adolesc* 30: 655–669.
80. McLaughlin KA, Peverill M, Gold AL, Alves S, Sheridan MA (2015): Child Maltreatment and Neural Systems Underlying Emotion Regulation. *J Am Acad Child Adolesc Psychiatry* 54: 753–762.
81. Rodman AM, Jenness JL, Weissman DG, Pine DS, McLaughlin KA (2019): Neurobiological Markers of Resilience to Depression Following Childhood Maltreatment: The Role of Neural Circuits Supporting the Cognitive Control of Emotion. *Biol Psychiatry* 86: 464–473.

82. Herringa RJ, Burghy CA, Stodola DE, Fox ME, Davidson RJ, Essex MJ (2016): Enhanced Prefrontal-Amygdala Connectivity Following Childhood Adversity as a Protective Mechanism Against Internalizing in Adolescence. *Biol Psychiatry Cogn Neurosci Neuroimaging* 1: 326–334.
83. Fullana MA, Albajes-Eizagirre A, Soriano-Mas C, Vervliet B, Cardoner N, Benet O, *et al.* (2018): Fear extinction in the human brain: A meta-analysis of fMRI studies in healthy participants. *Neurosci Biobehav Rev* 88: 16–25.
84. Callaghan BL, Tottenham N (2016): The Stress Acceleration Hypothesis: effects of early-life adversity on emotion circuits and behavior. *Curr Opin Behav Sci* 7: 76–81.
85. Gee DG, Gabard-Durnam LJ, Flannery J, Goff B, Humphreys KL, Telzer EH, *et al.* (2013): Early developmental emergence of human amygdala-prefrontal connectivity after maternal deprivation. *Proc Natl Acad Sci* 110: 15638–15643.
86. Callaghan BL, Gee DG, Gabard-Durnam L, Telzer EH, Humphreys KL, Goff B, *et al.* (2019): Decreased amygdala reactivity to parent cues protects against anxiety following early adversity: an examination across 3-years. *Biol Psychiatry Cogn Neurosci Neuroimaging*. <https://doi.org/10.1016/j.bpsc.2019.02.001>
87. Gee DG (2016): Sensitive Periods of Emotion Regulation: Influences of Parental Care on Frontoamygdala Circuitry and Plasticity. *New Dir Child Adolesc Dev* 2016: 87–110.
88. Westen D (2000): Commentary: Implicit and Emotional Processes in Cognitive-Behavioral Therapy. *Clin Psychol Sci Pract* 7: 386–390.
89. Cohen JA, Mannarino AP, Deblinger E (2017): *Treating Trauma and Traumatic Grief in Children and Adolescents, 2nd Ed.* New York, NY, US: Guilford Press, pp xix, 356.
90. Dorsey S, McLaughlin KA, Kerns SEU, Harrison JP, Lambert HK, Briggs EC, *et al.* (2017): Evidence Base Update for Psychosocial Treatments for Children and Adolescents Exposed to Traumatic Events. *J Clin Child Adolesc Psychol* 46: 303–330.
91. Weisz J, Bearman SK, Santucci LC, Jensen-Doss A (2017): Initial Test of a Principle-Guided Approach to Transdiagnostic Psychotherapy With Children and Adolescents. *J Clin Child Adolesc Psychol* 46: 44–58.
92. Weytens F, Luminet O, Verhofstadt LL, Mikolajczak M (2014): An Integrative Theory-Driven Positive Emotion Regulation Intervention. *PLoS ONE* 9. <https://doi.org/10.1371/journal.pone.0095677>
93. Hostinar CE, Johnson AE, Gunnar MR (2015): Parent support is less effective in buffering cortisol stress reactivity for adolescents compared to children. *Dev Sci* 18: 281–297.
94. Brody GH, Murry VM, McNair L, Chen Y-F, Gibbons FX, Gerrard M, Wills TA (2005): Linking Changes in Parenting to Parent-Child Relationship Quality and Youth Self-Control: The Strong African American Families Program. *J Res Adolesc* 15: 47–69.

95. Feldman R (2007): Parent-Infant Synchrony: Biological Foundations and Developmental Outcomes. *Curr Dir Psychol Sci* 16: 340-345.
96. Moriceau S, Sullivan RM (2006): Maternal presence serves as a switch between learning fear and attraction in infancy. *Nat Neurosci* 9: 1004.
97. Morris AS, Silk JS, Steinberg L, Myers SS, Robinson LR (2007): The Role of the Family Context in the Development of Emotion Regulation. *Soc Dev* 16: 361-388.
98. Humphreys KL, Miron D, McLaughlin KA, Sheridan MA, Nelson CA, Fox NA, Zeanah CH (2018): Foster care promotes adaptive functioning in early adolescence among children who experienced severe, early deprivation. *J Child Psychol Psychiatry* 59: 811-821.
99. McLaughlin KA, Zeanah CH, Fox NA, Nelson CA (2012): Attachment security as a mechanism linking foster care placement to improved mental health outcomes in previously institutionalized children. *J Child Psychol Psychiatry* 53: 46-55.
100. Fritz J, Fried EI, Goodyer IM, Wilkinson PO, van Harmelen A-L (2018): A Network Model of Resilience Factors for Adolescents with and without Exposure to Childhood Adversity [no. 1]. *Sci Rep* 8: 1-13.
101. Schriber RA, Rogers CR, Ferrer E, Conger RD, Robins RW, Hastings PD, Guyer AE (2018): Do Hostile School Environments Promote Social Deviance by Shaping Neural Responses to Social Exclusion? *J Res Adolesc* 28: 103-120.
102. Hanson JL, Gillmore AD, Yu T, Holmes CJ, Hallowell ES, Barton AW, et al. (2019): A Family Focused Intervention Influences Hippocampal-Prefrontal Connectivity Through Gains in Self-Regulation. *Child Dev* 90: 1389-1401.
103. Guassi Moreira JF, Telzer EH (2018): Mother still knows best: Maternal influence uniquely modulates adolescent reward sensitivity during risk taking. *Dev Sci* 21: e12484.
104. Telzer EH, Fuligni AJ, Lieberman MD, Galván A (2012): Meaningful Family Relationships: Neurocognitive Buffers of Adolescent Risk Taking. *J Cogn Neurosci* 25: 374-387.
105. Smith AR, Rosenbaum GM, Botdorf MA, Steinberg L, Chein JM (2018): Peers influence adolescent reward processing, but not response inhibition. *Cogn Affect Behav Neurosci* 18: 284-295.
106. Adams RE, Santo JB, Bukowski WM (2011): The presence of a best friend buffers the effects of negative experiences. *Dev Psychol* 47: 1786-1791.
107. Calhoun CD, Helms SW, Heilbron N, Rudolph KD, Hastings PD, Prinstein MJ (2014): Relational victimization, friendship, and adolescents' hypothalamic-pituitary-adrenal axis responses to an in vivo social stressor. *Dev Psychopathol* 26: 605-618.
108. Breiner K, Li A, Cohen AO, Steinberg L, Bonnie RJ, Scott ES, et al. (2018): Combined effects of peer presence, social cues, and rewards on cognitive control in adolescents. *Dev Psychobiol* 60: 292-302.

109. Kim J, Cicchetti D (2010): Longitudinal pathways linking child maltreatment, emotion regulation, peer relations, and psychopathology. *J Child Psychol Psychiatry* 51: 706–716.
110. Weigard A, Chein J, Albert D, Smith A, Steinberg L (2014): Effects of anonymous peer observation on adolescents' preference for immediate rewards. *Dev Sci* 17: 71–78.
111. McCrory E, Ogle JR, Gerin MI, Viding E (2019): Neurocognitive Adaptation and Mental Health Vulnerability Following Maltreatment: The Role of Social Functioning. *Child Maltreat* 24: 435–451.
112. Brody GH, Chen Y-F, Murry VM, Ge X, Simons RL, Gibbons FX, *et al.* (2006): Perceived Discrimination and the Adjustment of African American Youths: A Five-Year Longitudinal Analysis With Contextual Moderation Effects. *Child Dev* 77: 1170–1189.
113. Brody GH, Lei M-K, Chae DH, Yu T, Kogan SM, Beach SRH (2014): Perceived Discrimination Among African American Adolescents and Allostatic Load: A Longitudinal Analysis With Buffering Effects. *Child Dev* 85: 989–1002.
114. Chae DH, Nuru-Jeter AM, Adler NE, Brody GH, Lin J, Blackburn EH, Epel ES (2014): Discrimination, Racial Bias, and Telomere Length in African-American Men. *Am J Prev Med* 46: 103–111.
115. Kilburn K, Thirumurthy H, Halpern CT, Pettifor A, Handa S (2016): Effects of a Large-Scale Unconditional Cash Transfer Program on Mental Health Outcomes of Young People in Kenya. *J Adolesc Health* 58: 223–229.
116. Rojas NM, Yoshikawa H, Gennetian L, Rangel ML, Melvin S, Noble K, *et al.* (2020): Exploring the experiences and dynamics of an unconditional cash transfer for low-income mothers: A mixed-methods study. *J Child Poverty* 26: 64–84.
117. Francis DD, Diorio J, Plotsky PM, Meaney MJ (2002): Environmental Enrichment Reverses the Effects of Maternal Separation on Stress Reactivity. *J Neurosci* 22: 7840–7843.
118. Laviola G, Rea M, Morley-Fletcher S, Carlo SD, Bacosi A, Simone RD, *et al.* (2004): Beneficial effects of enriched environment on adolescent rats from stressed pregnancies. *Eur J Neurosci* 20: 1655–1664.
119. Morley-Fletcher S, Rea M, Maccari S, Laviola G (2003): Environmental enrichment during adolescence reverses the effects of prenatal stress on play behaviour and HPA axis reactivity in rats. *Eur J Neurosci* 18: 3367–3374.
120. Brody GH, Chen Y, Kogan SM, Yu T, Molgaard VK, DiClemente RJ, Wingood GM (2012): Family-centered Program Deters Substance Use, Conduct Problems, and Depressive Symptoms in Black Adolescents. *Pediatrics* 129: 108–115.
121. Brody GH, Yu T, Chen E, Beach SRH, Miller GE (2016): Family-centered prevention ameliorates the longitudinal association between risky

- family processes and epigenetic aging. *J Child Psychol Psychiatry* 57: 566-574.
122. Brody GH, Gray JC, Yu T, Barton AW, Beach SRH, Galván A, *et al.* (2017): Protective Prevention Effects on the Association of Poverty With Brain Development. *JAMA Pediatr* 171: 46-52.
 123. Murry VM, Berkel C, Chen Y, Brody GH, Gibbons FX, Gerrard M (2011): Intervention Induced Changes on Parenting Practices, Youth Self-Pride and Sexual Norms to Reduce HIV-Related Behaviors Among Rural African American Youths. *J Youth Adolesc* 40: 1147-1163.
 124. Pfeifer JH, Berkman ET (2018): The Development of Self and Identity in Adolescence: Neural Evidence and Implications for a Value-Based Choice Perspective on Motivated Behavior. *Child Dev Perspect* 12: 158-164.
 125. Romeo RD (2018): The metamorphosis of adolescent hormonal stress reactivity: A focus on animal models. *Front Neuroendocrinol* 49: 43-51.
 126. Sisk CL, Romeo RD (2019): *Coming of Age: The Neurobiology and Psychobiology of Puberty and Adolescence*. Oxford University Press.
 127. Chung YS, Poppe A, Novotny S, Epperson CN, Kober H, Granger DA, *et al.* (2019): A preliminary study of association between adolescent estradiol level and dorsolateral prefrontal cortex activity during emotion regulation. *Psychoneuroendocrinology* 109: 104398.
 128. Nusslock R, Brody GH, Armstrong CC, Carroll AL, Sweet LH, Yu T, *et al.* (2019): Higher Peripheral Inflammatory Signaling Associated With Lower Resting-State Functional Brain Connectivity in Emotion Regulation and Central Executive Networks. *Biol Psychiatry* 86: 153-162.
 129. Mendle J, Leve LD, Ryzin MV, Natsuaki MN, Ge X (2011): Associations Between Early Life Stress, Child Maltreatment, and Pubertal Development Among Girls in Foster Care. *J Res Adolesc* 21: 871-880.
 130. Noll JG, Trickett PK, Long JD, Negri S, Susman EJ, Shalev I, *et al.* (2017): Childhood Sexual Abuse and Early Timing of Puberty. *J Adolesc Health* 60: 65-71.
 131. Wilson ME, Bounar S, Godfrey J, Michopoulos V, Higgins M, Sanchez M (2013): Social and emotional predictors of the tempo of puberty in female rhesus monkeys. *Psychoneuroendocrinology* 38: 67-83.
 132. Dorn LD, Hostinar CE, Susman EJ, Pervanidou P (2019): Conceptualizing Puberty as a Window of Opportunity for Impacting Health and Well-Being Across the Life Span. *J Res Adolesc* 29: 155-176.
 133. Rudolph KD, Flynn M (2007): Childhood adversity and youth depression: Influence of gender and pubertal status. *Dev Psychopathol* 19: 497-521.
 134. Quevedo K, Johnson AE, Loman ML, LaFavor TL, Gunnar M (2012): The confluence of adverse early experience and puberty on the cortisol awakening response. *Int J Behav Dev* 36: 19-28.

135. Koss KJ, Hostinar CE, Donzella B, Gunnar MR (2014): Social deprivation and the HPA axis in early development. *Psychoneuroendocrinology* 50: 1-13.
136. Doremus-Fitzwater TL, Spear LP (2016): Reward-centricity and attenuated aversions: An adolescent phenotype emerging from studies in laboratory animals. *Neurosci Biobehav Rev* 70: 121-134.
137. Steinberg L (2015): How to Improve the Health of American Adolescents. *Perspect Psychol Sci* 10: 711-715.
138. Doom JR, Hostinar CE, VanZomeren-Dohm AA, Gunnar MR (2015): The roles of puberty and age in explaining the diminished effectiveness of parental buffering of HPA reactivity and recovery in adolescence. *Psychoneuroendocrinology* 59: 102-111.
139. Doom JR, Doyle CM, Gunnar MR (2017): Social stress buffering by friends in childhood and adolescence: Effects on HPA and oxytocin activity. *Soc Neurosci* 12: 8-21.
140. Pauling SN, Doom JR, Gunnar MR (2017): Gender Differences in Buffering Stress Responses in Same-Sex Friend Dyads. Retrieved March 18, 2019, from <http://unicatalog.mediu.edu.my:8080/xmlui/handle/123456789/66205>
141. Gee DG, Gabard-Durnam L, Telzer EH, Humphreys KL, Goff B, Shapiro M, et al. (2014): Maternal Buffering of Human Amygdala-Prefrontal Circuitry During Childhood but Not During Adolescence. *Psychol Sci* 25: 2067-2078.
142. Miller AB, Sheridan MA, Hanson JL, McLaughlin KA, Bates JE, Lansford JE, et al. (2018): Dimensions of deprivation and threat, psychopathology, and potential mediators: A multi-year longitudinal analysis. *J Abnorm Psychol* 127: 160-170.
143. Cohodes EM, Kitt ER, Baskin-Sommers A, Gee DG (2020): Influences of early-life stress on frontolimbic circuitry: Harnessing a dimensional approach to elucidate the effects of heterogeneity in stress exposure. *Dev Psychobiol* n/a. <https://doi.org/10.1002/dev.21969>
144. Ioannidis K, Askelund AD, Kievit RA, van Harmelen A-L (2020): The complex neurobiology of resilient functioning after childhood maltreatment. *BMC Med* 18: 32.
145. Frodl T, Szyf M, Carballedo A, Ly V, Dymov S, Vaisheva F, et al. (2015): DNA methylation of the serotonin transporter gene (SLC6A4) is associated with brain function involved in processing emotional stimuli. *J Psychiatry Neurosci JPN* 40: 296-305.
146. Silvers JA (2020): Extinction Learning and Cognitive Reappraisal: Windows Into the Neurodevelopment of Emotion Regulation. *Child Dev Perspect* n/a. <https://doi.org/10.1111/cdep.12372>
147. King LS, Graber MG, Colich NL, Gotlib IH (2020): Associations of waking cortisol with DHEA and testosterone across the pubertal transition: Effects of threat-related early life stress. *Psychoneuroendocrinology* 115: 104651.

148. Callaghan BL, Tottenham N (2016): The Neuro-Environmental Loop of Plasticity: A Cross-Species Analysis of Parental Effects on Emotion Circuitry Development Following Typical and Adverse Caregiving. *Neuropsychopharmacology* 41: 163-176.
149. Dunn EC, Soare TW, Zhu Y, Simpkin AJ, Suderman MJ, Klengel T, *et al.* (2019): Sensitive Periods for the Effect of Childhood Adversity on DNA Methylation: Results From a Prospective, Longitudinal Study. *Biol Psychiatry* 85: 838-849.
150. Gabard-Durnam LJ, McLaughlin KA (2019): Do Sensitive Periods Exist for Exposure to Adversity? *Biol Psychiatry* 85: 789-791.

Figure

Figure 1. Schematic of resilience factors and associated neural systems and interventions. dACC, dorsal anterior cingulate cortex; dlPFC, dorsolateral prefrontal cortex; dmPFC, dorsomedial PFC; SPL, superior parietal lobule; vlPFC, ventrolateral PFC; vmPFC, ventromedial PFC; VS, ventral striatum.

<i>Resilience Factor</i>	<i>Neural System</i>	<i>Interventions</i>
<i>Reward Processing</i>		Positive Affect Treatment Behavioral Activation Therapy
<i>Affective Learning*</i>		Extinction-enhancing exposure
<i>Cognitive Control</i>		Task-switch training Working memory training
<i>Emotion Regulation</i> <i><u>Implicit:</u></i>		Cognitive behavioral therapy (CBT) Trauma-focused CBT FIRST
<i><u>Explicit:</u></i>		

Increased activation during task
 Decreased activation during task

* Primarily linked to deprivation