

# UC Irvine

## UC Irvine Previously Published Works

### Title

Using mouse and zebrafish models to understand the etiology of developmental defects in Cornelia de Lange Syndrome

### Permalink

<https://escholarship.org/uc/item/1kn4m2w9>

### Journal

American Journal of Medical Genetics Part C Seminars in Medical Genetics, 172(2)

### ISSN

1552-4868

### Authors

Kawauchi, Shimako  
Santos, Rosaysela  
Muto, Akihiko  
[et al.](#)

### Publication Date

2016-06-01

### DOI

10.1002/ajmg.c.31484

Peer reviewed

# Using Mouse and Zebrafish Models to Understand the Etiology of Developmental Defects in Cornelia de Lange Syndrome

SHIMAKO KAWAUCHI, ROSAYSELA SANTOS, AKIHIKO MUTO, MARTHA E. LOPEZ-BURKS, THOMAS F. SCHILLING, ARTHUR D. LANDER, AND ANNE L. CALOF\*

Cornelia de Lange Syndrome (CdLS) is a multisystem birth defects disorder that affects every tissue and organ system in the body. Understanding the factors that contribute to the origins, prevalence, and severity of these developmental defects provides the most direct approach for developing screens and potential treatments for individuals with CdLS. Since the majority of cases of CdLS are caused by haploinsufficiency for *NIPBL* (*Nipped-B-like*, which encodes a cohesin-associated protein), we have developed mouse and zebrafish models of CdLS by using molecular genetic tools to create *Nipbl*-deficient mice and zebrafish (*Nipbl*<sup>+/−</sup> mice, zebrafish *nipbl* morphants). Studies of these vertebrate animal models have yielded novel insights into the developmental etiology and genes/gene pathways that contribute to CdLS-associated birth defects, particularly defects of the gut, heart, craniofacial structures, nervous system, and limbs. Studies of these mouse and zebrafish CdLS models have helped clarify how deficiency for NIPBL, a protein that associates with cohesin and other transcriptional regulators in the nucleus, affects processes important to the emergence of the structural and physiological birth defects observed in CdLS: NIPBL exerts chromosome position-specific effects on gene expression; it influences long-range interactions between different regulatory elements of genes; and it regulates combinatorial and synergistic actions of genes in developing tissues. Our current understanding is that CdLS should be considered as not only a cohesinopathy, but also a “transcriptomopathy,” that is, a disease whose underlying etiology is the global dysregulation of gene expression throughout the organism. © 2016 Wiley Periodicals, Inc.

**KEY WORDS:** *Nipped-B-like* (NIPBL) gene; *Protocadherin* genes; *Hox* genes; *Shh* genes; chromatin conformation

**How to cite this article:** Kawauchi S, Santos R, Muto A, Lopez-Burks ME, Schilling TF, Lander AD, Calof AL. 2016. Using mouse and zebrafish models to understand the etiology of developmental defects in Cornelia de Lange syndrome. *Am J Med Genet Part C Semin Med Genet* 172C:138–145.

Shimako Kawauchi, Ph.D. is an Associate Project Scientist who has been a primary researcher in developing and analyzing mouse models of CdLS since she was a Postdoctoral Scholar in the Calof lab. Dr. Kawauchi's research interests are in the regulation of gene expression and pattern formation during mammalian development.

Rosaysela Santos, Ph.D., is a Postdoctoral Scholar who has been a lead researcher in developing and analyzing mouse models of CdLS since she was a graduate student in the Calof lab. Dr. Santos's research interests lie in using animal models to understand the origins of birth defects associated with developmental and genetic disorders in man.

Akihiko Muto, Ph.D., is Assistant Professor of Biological Sciences at Hiroshima University in Japan. He was formerly a Project Scientist at the University of California, Irvine, where he began his research on zebrafish models of CdLS in collaboration with Thomas Schilling, Arthur Lander, and Anne Calof. Dr. Muto's research interests are in understanding the molecular mechanisms by which chromatin structure affects gene expression, and the implications of this process for disease pathogenesis.

Martha E. Lopez-Burks, M.S., is a Staff Research Associate who specializes in molecular biology. During the past decade, her research has focused on molecular analysis of *Nipbl*-deficient mice. Currently, she leads the molecular and phenotypic analysis of limb and skeletal development in the CdLS mouse model.

Thomas F. Schilling, Ph.D., is Professor and Chair of the Department of Developmental & Cell Biology at the University of California, Irvine. Dr. Schilling has a long-standing interest in craniofacial development, with work in his lab largely focused on developmental genetics in zebrafish as a model for human structural birth defects.

Arthur D. Lander, M.D., Ph.D., is Donald Bren Professor of Developmental & Cell Biology, Director of the Center for Complex Biological Systems, and Co-Director the Center of Excellence in Research on Cornelia de Lange Syndrome at the University of California, Irvine. Dr. Lander's research uses experimental, mathematical, and computational approaches to elucidate the engineering principles that underlie human development, with a focus on fundamental problems in patterning and growth control, and the origins of complex traits.

Anne L. Calof, Ph.D., is Professor of Anatomy & Neurobiology at the University of California, Irvine. She serves as Co-Director of UCI's Center of Excellence in Research on Cornelia de Lange Syndrome and as a Theme Leader in the Center for Complex Biological Systems. Dr. Calof's research is focused on understanding how cell differentiation, tissue morphogenesis, and tissue size are regulated, both during normal development and when pathological conditions result in birth defects.

Grant sponsor: National Institutes of Health; Grant numbers: P01-HD052860; P50-GM076516; Grant sponsor: Cornelia de Lange Syndrome Foundation.

\*Correspondence to: Anne L. Calof, Department of Anatomy & Neurobiology, Center for Complex Biological Systems, 2626 Bio Sci 3, University of California, Irvine, CA 92697-2280. E-mail: [alcalof@uci.edu](mailto:alcalof@uci.edu)

DOI 10.1002/ajmg.c.31484

Article first published online 27 April 2016 in Wiley Online Library ([wileyonlinelibrary.com](http://wileyonlinelibrary.com)).

## INTRODUCTION

Cornelia de Lange syndrome (CdLS) is a birth defects disorder that affects every system in the body. Although long understood by clinicians to be a distinctive syndrome [Opitz, 1985], it was not until the discovery that the majority of individuals with CdLS carry mutations in *NIPBL* [Krantz et al., 2004; Tonkin et al., 2004], which encodes a protein that binds to the cohesin complex of chromosomal structural proteins, that CdLS was recognized as the prototypic cohesinopathy [Dorsett and Krantz, 2009]. Subsequent to the identification of *NIPBL* as the major causative gene for CdLS, researchers have gone on to identify a number of cohesin- and/or cohesin regulatory protein-encoding genes which, when mutated, give rise to the classical spectrum of features and deficits associated with CdLS: these genes include *NIPBL*, *SMC1A*, *SMC3*, *RAD21*, and *HDAC8* [Dear-dorff and Krantz, 2005]. For most of these genes, animal models have been generated, including models from *Drosophila*, *Xenopus*, zebrafish, and mouse; these model systems have been widely used as tools for understanding the etiology of CdLS related syndromes [Horsfield et al., 2012; Singh and Gerton, 2015]. In the present paper, we review studies from our group on two vertebrate animal models of CdLS: *nipbl*-morphant zebrafish and *Nipbl*-deficient mice. These two model systems have been particularly useful for understanding how *Nipbl* deficiency leads to global changes in gene expression, and how global gene expression changes act additively and synergistically to generate the structural birth defects that are the hallmarks of CdLS.

## DEVELOPMENT OF MOUSE AND ZEBRAFISH ANIMAL MODELS OF CDLS

The discovery that most individuals with CdLS carry mutations in, and are likely haploinsufficient for, *NIPBL* [Krantz et al., 2004; Tonkin et al., 2004], led to the rapid realization that understanding the etiology of CdLS

would require a deeper understanding of how the NIPBL protein, which associates with the cohesin complex of chromosomal proteins, functions in the regulation of chromatin structure and gene expression [Dorsett, 2009]. Since a *Drosophila* model of *Nipbl* deficiency already existed [Rollins et al., 1999], and increased insight into the interplay between gene expression and the origins of structural birth defects in CdLS was most likely to arise from studies of vertebrate models of the syndrome, our group decided to generate mouse and zebrafish models of CdLS. Mouse and zebrafish models of *Nipbl* deficiency recapitulate many of the important structural and functional deficits observed in individuals with CdLS, and have proven invaluable for gaining information about the molecular etiology of the syndrome.

---

***Mouse and zebrafish models of *Nipbl* deficiency recapitulate many of the important structural and functional deficits observed in individuals with CdLS, and have proven invaluable for gaining information about the molecular etiology of the syndrome.***

---


In mouse, we generated a line carrying one null allele of *Nipbl*. *Nipbl*<sup>+/-</sup> mice, created by introduction of a gene trap into Intron 1 of the *Nipbl* gene, show a high degree of perinatal lethality (~18% of mutant animals survive to weaning), are much smaller than their wildtype littermates, and have numerous structural, physiological, and gene expression defects that mimic the defects seen in CdLS (see below and Fig. 1A). The high incidence of perinatal lethality and small size of *Nipbl*<sup>+/-</sup> mice are heritable phenotypes and have been reproduced over many generations

[Kawauchi et al., 2009]. *Nipbl*-deficient zebrafish embryos (Fig. 1B) were created by using translation-blocking morpholinos (MOs) to knock down expression of the two zebrafish *nipbl* genes (*nipbla* and *nipblb*), which our group first cloned (Fig. 1C; since the two zebrafish *nipbl* genes have functions that do not overlap completely, both are knocked down for most analyses [Muto et al., 2011]). Zebrafish *nipbl* morphants die at larval stages and exhibit stunted growth: embryos begin to exhibit obvious phenotypes within 24 hr postfertilization (hpf), with growth retardation—especially of the head and tail—and deficits in heart and circulatory system becoming progressively more pronounced over time (Fig. 1B).

## ***NIPBL-DEFICIENT MICE AND ZEBRAFISH SHOW PRONOUNCED DEFECTS IN HEART AND GUT DEVELOPMENT RELATED TO CHANGES IN ENDODERMAL GENE EXPRESSION***

Heart and gut malformations are a prominent part of the pathology of CdLS. Structural heart defects are well-documented, and occur in about 30–40% of individuals with CdLS [Selicorni et al., 2009; Chatfield et al., 2012]. In our examinations of *Nipbl*<sup>+/-</sup> mice, we also noted structural defects in the heart. In particular, defects in atrial septum formation (ASD) were noted in the hearts of about half of embryos taken between E15.5 and 18.5 (Fig. 2A and B). Although no ASDs or other obvious cardiac malformations are observed in *Nipbl*<sup>+/-</sup> mice that survive to adulthood, these results strongly suggest that perinatal mortality in *Nipbl*-deficient mice may be caused by cardiac defects [Kawauchi et al., 2009]. Because of the prominence and severity of cardiac defects in *Nipbl*<sup>+/-</sup> mice, their underlying cellular and genetic etiology is being studied in detail [Santos, Kawauchi, Calof; unpublished results].

Defects in early development of internal organs are readily visualized in zebrafish embryos, so we took advantage




**Figure 1.** *Nipbl*-deficient mutant animals show growth abnormalities. (A) *Nipbl*<sup>+/-</sup> mouse is markedly smaller than wildtype (+/+) sibling at 4 weeks of age. (B) *Nipbl* *a+b* morpholino (MO) co-injected zebrafish larvae (*nipbl a/b* morphants) resemble uninjected and mixed MO controls at 18.5 hr post-fertilization (hpf), but show marked circulatory defects, pericardial edema (asterisk) and tail defects (arrowheads) by 34 hpf. Scale bar = 100  $\mu$ m. (C) Structural comparison of NIPBL orthologues among different species. *H.s.*, *Homo sapiens*, 2,804 amino acids (aa); *M.m.*, *Mus musculus*, 2,798 aa; *D.r. (a)*, *Danio rerio*, form a, 2,876 aa; *D.r. (b)*, form b, 2,381 aa; *D.m.*, *Drosophila melanogaster*, 2,077 aa; *S.c.*, *Saccharomyces cerevisiae*, 1,493 aa. Numbers indicate percent amino acid sequence identity compared to human NIPBL. Adapted from [Kawauchi et al., 2009; Muto et al., 2011].

of *nipbla/b* morphants to study the early development of heart and other visceral organs when *Nipbl* is deficient in this model. As shown in Figure 2C, the developing heart tube of zebrafish embryos is readily visualized by in situ hybridization (ISH) for *cmlc2*. Pronounced defects in early heart

development, ranging from reduced heart tube looping (“type A defects”) to cardia bifida (“type B” defects), are observed in the majority of *nipbla/b* morphants. Early development of organs such as eyes is not affected (Fig. 2C, *ath5* ISH), suggesting that heart development is particularly sensitive to

decreases in *Nipbl* levels. As development proceeds, impaired heart function is observed in *nipbla/b* morphants, accompanied by circulation defects, pericardial edema, and premature death [Muto et al., 2011].

Gastrointestinal (GI) reflux is a hallmark of CdLS. Many individuals



**Figure 2.** Heart and gut abnormalities in *Nipbl*-deficient animals. (A and B) Cryosections, stained with hematoxylin/eosin, of wildtype and *Nipbl*<sup>+/-</sup> littermate embryos at gestational day 17.5 (E17.5). A well-formed atrial septum is apparent in the wildtype heart, but absent in a significant fraction of mutants. Scale bar = 1 mm. (C) Zebrafish embryos injected with *nipbla/b* MOs show a range of defects in heart tube formation at 32 hpf, ranging from mild (type A) to severe (type B). Dashed white lines indicate heart tube, labeled by in situ hybridization (ISH) for a *cmlc2* probe (*ath5* ISH shows that eye is relatively unaffected). (D) Gut and visceral organ morphology assessed by ISH for *foxa3* in *nipbla/b* morphants at 52 hpf. Defects ranged from mild looping defects (type I) to more severe defects such as bifurcation (type II) and even absence (type III) of gut parenchyma. gu, gut tube; li, liver; pa, pancreas. (E) Frequency of gut bifurcation in *sox17* and *foxa2* single and double morphants at 52 hpf. (F) Example of severe gut bifurcation in *sox17/foxa2* double morphant embryo. Scale bar in D and F = 50  $\mu$ m. Adapted from [Kawauchi et al., 2009; Muto et al., 2011].

with CdLS die as a result of adverse events affecting the GI system, with gut obstructions and volvulus being reported frequently [Luzzani et al., 2003; Schrier et al., 2011]. We therefore examined *nipbla/b* morphant embryos for malformations of the developing gut and associated organs (pancreas, liver) using ISH for *foxa3*, which is expressed by early gut and associated endodermal derivatives. The defects observed in *nipbla/b* morphants, which range from decreases in gut size to severe patterning defects as extreme as duplications and even absence of associated organs, are shown in Figure 2D [Muto et al., 2011].

To gain insight into the genetic origins of these defects, we performed transcriptome analysis using mRNA obtained from gastrulation-stage (6-hr postfertilization [hpf]) uninjected and *nipbla/b* morphant embryos. Many small, but significant, gene expression changes were observed when *Nipbl* levels were reduced, consistent with our findings in mice (see below) and the findings of our colleagues in human CdLS cell lines [Liu et al., 2009]. In particular, genes that function as regulators of endoderm development, such as *sox17* and *foxa2*, were found to be significantly downregulated in their expression. These genes play important roles in heart and visceral organ development [Ober et al., 2003; Fukuda and Kikuchi, 2005; Pfister et al., 2011], and since our experiments indicated that they are directly affected by decreases in *Nipbl* levels, we performed further studies to determine if knockdown of one or both would result in gut defects similar to those observed in *nipbla/b* morphants. As shown in Figure 2E and F, reduction of *sox17* or *foxa2* levels alone causes gut defects in a significant fraction (~5–7%) of morphants. Importantly, when levels of both *sox17* and *foxa2* are reduced simultaneously, the fraction of morphants with gut defects increases drastically, with almost 60% of morphants displaying severe defects such as gut bifurcation (Fig. 2E and F). Thus, reductions in expression of *sox17* and *foxa2* act synergistically to reproduce the types of defects in gut development seen in *nipbla/b* morphants [Muto et al.,

2011]. These data strongly support the hypothesis that defects in GI structure/function, observed in individuals with CdLS, arise from the collective effects of multiple gene expression changes resulting from reductions in *NIPBL* levels.

### CRANIOFACIAL, NEUROANATOMICAL, AND BRAIN GENE EXPRESSION ABNORMALITIES IN *NIPBL*<sup>+/-</sup> MICE

Individuals with CdLS display distinctive craniofacial features that play an important role in clinical diagnosis [Jackson et al., 1993]. Micro-CT analysis of adult *Nipbl*<sup>+/-</sup> mice demonstrated that they exhibit microbrachycephaly and an upward deflection of the snout (“upturned nose”), two of the most distinctive features of CdLS (Fig. 3A and [Kawauchi et al., 2009]). The smaller skull size of *Nipbl*<sup>+/-</sup> mice is reflected in their smaller endocranial volume and smaller overall brain size relative to control littermates (Fig. 3B and C). Overall the brains of adult *Nipbl*<sup>+/-</sup> mice are grossly normal (aside from their reduced size), although cerebellar hypoplasia is observed (Fig. 3D), consistent with clinical reports [Hayashi et al., 1996; Ozkinay et al., 1998]. The neuroanatomical abnormalities observed in *Nipbl*<sup>+/-</sup> mice are reflected in aberrant behaviors and in impairment of neurological functions such as hearing (which is frequently impaired in CdLS [Sakai et al., 2002; Kawauchi et al., 2009]).


To gain insight into the gene expression changes that underlie the behavioral and neurological abnormalities we observe in *Nipbl*<sup>+/-</sup> mice, we performed transcriptome analysis on gestational day 13.5 (E13.5) brain RNA from large cohorts of mutant and wildtype embryos. Expression of 978 genes was found to be changed significantly in E13.5 *Nipbl*<sup>+/-</sup> brain compared to normal; interestingly, all gene expression changes (which included both increases and decreases in expression) were small, with the single greatest change being only 2.5-fold. Significant expression changes were

observed for numerous genes with obvious roles in neuronal function, including genes encoding ion channels, neuropeptides, neurotransmitter receptors, axon guidance molecules, and so on. However, expression changes for genes encoding proteins associated with chromatin structure and function, metabolism, lipoprotein binding, and many other functions and were also observed [Kawauchi et al., 2009]. These data are consistent with studies by us and others, which indicate that *Nipbl* deficiency results in global dysregulation of gene expression in many—if not all—tissues [Kawauchi et al., 2009; Liu et al., 2009; Muto et al., 2011; Muto et al., 2014; Yuan et al., 2015].

### EFFECTS OF *NIPBL* DEFICIENCY ON DEVELOPMENT AND GENE EXPRESSION IN LIMBS

Alterations in limb and digit development, ranging from mild oligodactyly to frank truncations of the arms, are a hallmark of CdLS [Jackson et al., 1993]. Interestingly, limb truncations are not seen in *Nipbl*<sup>+/-</sup> mice, in which only aberrations in ossification, formation of the olecranon process, and mild digit defects are observed (see Lopez-Burks et al. [2016]; and Kawauchi et al. [2009]). As a result, we have concentrated our studies of how *Nipbl* deficiency affects limb development on the zebrafish model system, where we hypothesized that *Nipbl* deficiency may be more effective at triggering limb defects than in the mouse.

We performed an extensive analysis of alterations in gene expression and cellular/structural changes in the development of pectoral fins (the structural homologues of mammalian forelimbs) in zebrafish morphants, which yielded a number of significant findings. As predicted, zebrafish embryos injected with *nipbla/b* MOs showed impaired pectoral fin development, which was dramatic at 76 hpf, when the limbs of morphants were 40% shorter than those of controls. Importantly, partial rescue of limb defects was observed when morphants were co-injected with *nipbla*




mRNA, indicating that effects of MOs were specific to *Nipbl* depletion (Fig. 4A and B).

Genes known to be of general importance in vertebrate limb development were examined in *nipbla/b* morphants, and many were found to be mis-regulated in their expression. Some of the most important of these were *fgf* genes (expression of *fgf4*, *fgf8a*, and *fgf16*

were altered in the fin bud apical ectodermal ridge/limb mesenchyme of morphants) and *shha*, whose expression in the fin bud zone of polarizing activity (ZPA) was reduced in morphants (Fig. 4C). Importantly, *Shh* expression was also reduced in the ZPA of E10.5 limb buds in *Nipbl*<sup>+/-</sup> mice (Fig. 4D and E), indicating that regulatory genes for limb development are similarly affected

by *Nipbl* deficiency in both zebrafish and mice. Transcriptome analysis of E10.5 *Nipbl*<sup>+/-</sup> and wildtype littermate limb buds demonstrated that approximately 1,000 genes are significantly altered in their expression when *Nipbl* is deficient, and that many changes in gene expression observed in mouse limb bud mirrored those observed in *nipbla/b* morphant zebrafish fin buds [Muto et al., 2014].

Appropriate spatial and temporal expression of *Hox* genes is important for limb development in all vertebrates [Zakany and Duboule, 2007]. Many *hox* genes show altered expression in the fin buds of *nipbla/b* morphant embryos, and expression of different *hox* genes is either up- or down-regulated in morphants [Muto et al., 2014]. Regulation of *hox* gene expression by *Nipbl* is dependent on the relative location of a given gene within its cluster (5' versus 3'), and the implications of such positional effects for understanding the mechanism(s) by which *Nipbl* levels effect global changes in gene expression is discussed below. Our analysis of gene expression changes in mouse limb buds also indicated that genes encoding some subunits of Mediator, a protein complex affecting chromatin conformation and global regulation of transcription [Yin and Wang, 2014], are altered in their expression when *Nipbl* levels are reduced (Table I in Muto et al. [2014]). Because it had been shown by Kagey et al. [2010] that Mediator, cohesin, and *Nipbl* all interact to regulate gene expression, and since other investigators showed that loss-of-function mutations in *med12* disrupt fin development in zebrafish [Rau et al., 2006], we generated *med12* morphants and studied pectoral fin development in them. Zebrafish *med12* morphants display disrupted fin development and mis-regulated *hox* gene expression similar to that observed in *nipbla/b* morphants. Importantly, when zebrafish embryos are injected with low doses of both *med12* and *nipbla/b* MOs (each of which gives only partial reductions in fin size), a strong synergistic effect is observed (Fig. 4F and G). Moreover, simultaneous reduction of *nipbls* and



*med12* have a synergistic effect on alterations in *hox* gene expression (Fig. 4H). Altogether, these data are consistent with *Nipbl* and Mediator acting in a common pathway to regulate gene expression and development in the vertebrate limb.


## NIPBL, CHROMATIN CONFORMATION, AND GLOBAL REGULATION OF GENE EXPRESSION

Despite the fact that we know that small changes in the expression of many genes contribute to the spectrum of birth defects in CdLS, we do not yet understand mechanistically how this occurs. Past and ongoing studies of *Nipbl*-deficient animal models do, however, provide a way forward. Transcriptome analyses performed on multiple *Nipbl*<sup>+/-</sup> tissues yield common findings, among them the observations that tens to hundreds of transcripts are significantly altered in a given tissue, with effects on gene expression being modest in size (the vast majority of gene expression

changes are less than two-fold). Distinctive profiles of gene expression changes for specific tissues are also evident, and have provided insights into genetic pathways that underlie defects in different tissues. One instance is the dysregulation of endodermal gene expression that is hypothesized to underlie, at least in part, the development of heart and visceral organ defects in CdLS (Fig. 2 and Muto et al. [2011]). Similarly, the depletion of fat observed in *Nipbl*<sup>+/-</sup> mice is likely attributable to dysregulation of genes regulating the adipocyte differentiation pathway: expression of genes in the pathway that regulates adipogenesis is altered in *Nipbl*<sup>+/-</sup> mouse embryonic fibroblasts (MEFs), which are themselves defective in their ability to differentiate into adipocytes [Kawauchi et al., 2009]. Our findings are consistent with the hypothesis that the severe developmental defects observed in CdLS and in *Nipbl*-deficient animal models are the consequence of the collective action of many small changes in gene expression in every tissue throughout the organism.

***Our findings are consistent with the hypothesis that the severe developmental defects observed in CdLS and in Nipbl-deficient animal models are the consequence of the collective action of many small changes in gene expression in every tissue throughout the organism.***

Are there types of genes, and/or gene loci, that are especially sensitive to *Nipbl* deficiency? One clue comes from the analysis of transcriptional effects that are shared across multiple tissues. In the *Nipbl*<sup>+/-</sup> mouse, for example, significant changes in the expression of genes in the *protocadherin-beta* (*Pcdhb*) cluster, located on chromosome 18, have been observed repeatedly in embryonic brain, MEFs, limbs, and



**Figure 5.** Position-specific effects of *Nipbl* on gene expression. **(A)** Schematic of protocadherin beta (*Pcdhb*) locus (consisting of 22 genes) on chromosome 18 in mouse. **(B)** Quantitative RT-PCR results showing changes in gene expression in 14 of the 22 *Pcdhb* genes in E17.5 *Nipbl*<sup>+/-</sup> brain compared to wildtype controls. **(C)** Representative graph showing sensitivity of *Pcdhb* gene expression levels to *Nipbl* levels in the same samples. Sensitivities and error bars were plotted on an abscissa corresponding to the location of the transcriptional start sites of each of the *Pcdhb* genes. The dashed line is a smooth polynomial fit to the data. Data indicate that sensitivity of *Pcdhb* gene expression to *Nipbl* levels is highest at the 5' and 3' ends of the cluster, lowest in the middle. **(D)** ISH demonstrating mis-expression of *hoxx* genes in *nipbla/b* morphant zebrafish embryos (over-expression: *hoxx4a*; under-expression: *hoxx8a*). Schematic diagram represents effects of reduced *nipbl* levels on *hoxx* gene expression, with 5' *hoxx* genes showing reductions in expression (red) and 3' *hoxx* genes showing expanded expression (green). Adapted from [Kawauchi et al., 2009; Muto et al., 2011].

even skin ([Kawauchi et al., 2009; Muto et al., 2011] and unpublished observations). As shown in Figure 5A and B, *Pcdhb* genes are both up- and down-regulated in their expression when *Nipbl* is deficient, with genes at the 3' and 5' ends of the cluster showing the greatest percent change in expression. An in-depth analysis, in which changes in expression of individual *Pcdhb* genes were evaluated as a function of *Nipbl* levels in the same samples, demonstrated that expression of *Pcdhb* genes closest to the 5' and 3' ends of the cluster are most sensitive to *Nipbl* levels, while expression of genes in the middle of the cluster are relatively insensitive to *Nipbl* levels (Fig. 5C). Therefore, within the *Pcdhb* gene cluster, position-specific effects of *Nipbl* on gene expression can clearly be demonstrated [Kawauchi et al., 2009]. Similarly, studies of *hoxx* gene expression during zebrafish limb development demonstrate a relationship between the positions of specific *hoxx* genes in a given chromosomal cluster and their expression changes when *Nipbl* levels are depleted (Fig. 5D and Muto et al. [2014]). Altogether, our results from studies in both mouse and zebrafish models support the hypothesis that chromosomal location is an important

factor in the sensitivity of gene expression to NIPBL levels.

***Altogether, our results from studies in both mouse and zebrafish models support the hypothesis that chromosomal location is an important factor in the sensitivity of gene expression to NIPBL levels.***

How do these topological changes in NIPBL sensitivity come about? One possibility, for which there is increasing evidence, is that *Nipbl*—together with cohesin and proteins of the Mediator complex—physically brings together gene enhancers and their promoters, resulting in global changes in chromatin architecture that bring about widespread changes in gene expression [Kagey et al., 2010; Whyte et al., 2013; Yin and Wang, 2014]. Indeed, both *Nipbl* and Mediator are important in limb development; and in the zebrafish model of CdLS, deficiencies of *nipbls* and *med12* together act synergistically to regulate

gene expression and the severity of limb defects (Fig. 4F–H and Muto et al., [2014]). Importantly, depletion of both *Nipbl* and *Med 12* have been shown in 3D-fluorescence ISH experiments to affect long-range chromosomal interactions at the *hoxxda* locus in zebrafish limb buds [Muto et al., 2014]. The importance of Mediator, cohesin, and other general transcriptional regulators as collaborators of *Nipbl* in regulating chromatin conformation and gene expression is an ongoing subject of investigation in our attempt to understand the etiology of birth defects in CdLS.

## REFERENCES

- Chatfield KC, Schrier SA, Li J, Clark D, Kaur M, Kline AD, Deardorff MA, Jackson LS, Goldmuntz E, Krantz ID. 2012. Congenital heart disease in Cornelia de Lange syndrome: Phenotype and genotype analysis. *Am J Med Genet A* 158A:2499–2505.
- Deardorff MA NS, Krantz ID. 2005 Sep 16 [Updated 2016 Jan 28]. Cornelia de Lange Syndrome. In: Pagon RA, Adam MP, Ardinger HH, GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993–2016. Available from: <http://www.ncbi.nlm.nih.gov/books/NBK1104/>
- Dorsett D. 2009. Cohesin, gene expression and development: Lessons from *Drosophila*. *Chromosome Res* 17:185–200.
- Dorsett D, Krantz ID. 2009. On the molecular etiology of Cornelia de Lange syndrome. *Ann N Y Acad Sci* 1151:22–37.



- Fukuda K, Kikuchi Y. 2005. Endoderm development in vertebrates: Fate mapping, induction and regional specification. *Dev Growth Differ* 47:343–355.
- Hayashi M, Sakamoto K, Kurata K, Nagata J, Satoh J, Morimatsu Y. 1996. Septo-optic dysplasia with cerebellar hypoplasia in Cornelia de Lange syndrome. *Acta Neuropathol (Berl)* 92:625–630.
- Horsfield JA, Print CG, Monnich M. 2012. Diverse developmental disorders from the one ring: Distinct molecular pathways underlie the cohesinopathies. *Front Genet* 3:171.
- Jackson L, Kline AD, Barr MA, Koch S. 1993. De Lange syndrome: A clinical review of 310 individuals. *Am J Med Genet* 47:940–946.
- Kagey MH, Newman JJ, Bilodeau S, Zhan Y, Orlando DA, van Berkum NL, Ebmeier CC, Goossens J, Rahl PB, Levine SS, Taatjes DJ, Dekker J, Young RA. 2010. Mediator and cohesin connect gene expression and chromatin architecture. *Nature* 467:430–435.
- Kawauchi S, Calof AL, Santos R, Lopez-Burks ME, Young CM, Hoang MP, Chua A, Lao T, Lechner MS, Daniel JA, Nussenzweig A, Kitzes L, Yokomori K, Hallgrímsson B, Lander AD. 2009. Multiple organ system defects and transcriptional dysregulation in the *Nipbl*(+/-) mouse, a model of Cornelia de Lange syndrome. *PLoS Genet* 5:e1000650.
- Krantz ID, McCallum J, DeScipio C, Kaur M, Gillis LA, Yaeger D, Jukofsky L, Wasserman N, Bottani A, Morris CA, Nowaczyk MJ, Toriello H, Bamshad MJ, Carey JC, Rappaport E, Kawauchi S, Lander AD, Calof AL, Li HH, Devoto M, Jackson LG. 2004. Cornelia de Lange syndrome is caused by mutations in *NIPBL*, the human homolog of *Drosophila melanogaster* *Nipped-B*. *Nat Genet* 36:631–635.
- Liu J, Zhang Z, Bando M, Itoh T, Deardorff MA, Clark D, Kaur M, Tandy S, Kondoh T, Rappaport E, Spinner NB, Vega H, Jackson LG, Shirahige K, Krantz ID. 2009. Transcriptional dysregulation in *NIPBL* and cohesin mutant human cells. *PLoS Biol* 7:e1000119.
- Lopez-Burks ME, Santos R, Kawauchi S, Calof AL, Lander AD. 2016. Genetic enhancement of limb defects in a mouse model of Cornelia de Lange syndrome. *Am J Med Genet Part C Semin Med Genet* (in this issue).
- Luzzani S, Macchini F, Valadè A, Milani D, Selicorni A. 2003. Gastroesophageal reflux and Cornelia de Lange syndrome: Typical and atypical symptoms. *Am J Med Genet* 119A:283–287.
- Muto A, Calof AL, Lander AD, Schilling TF. 2011. Multifactorial origins of heart and gut defects in *nipbl*-deficient zebrafish, a model of Cornelia de Lange syndrome. *PLoS Biol* 9:e1001181.
- Muto A, Ikeda S, Lopez-Burks ME, Kikuchi Y, Calof AL, Lander AD, Schilling TF. 2014. *Nipbl* and mediator cooperatively regulate gene expression to control limb development. *PLoS Genet* 10:e1004671.
- Ober EA, Field HA, Stainier DY. 2003. From endoderm formation to liver and pancreas development in zebrafish. *Mech Dev* 120:5–18.
- Opitz JM. 1985. The Brachmann-de Lange syndrome. *Am J Med Genet* 22:89–102.
- Ozkinay F, Cogulu O, Gunduz C, Levent E, Ozkinay C. 1998. A case of Brachman de Lange syndrome with cerebellar vermis hypoplasia. *Clin Dysmorphol* 7:303–305.
- Pfister S, Jones VJ, Power M, Truise GL, Khoo PL, Steiner KA, Kanai-Azuma M, Kanai Y, Tam PP, Loebel DA. 2011. *Sox17*-dependent gene expression and early heart and gut development in *Sox17*-deficient mouse embryos. *Int J Dev Biol* 55:45–58.
- Rau MJ, Fischer S, Neumann CJ. 2006. Zebrafish *Trap230/Med12* is required as a coactivator for *Sox9*-dependent neural crest, cartilage and ear development. *Dev Biol* 296:83–93.
- Rollins RA, Morcillo P, Dorsett D. 1999. *Nipped-B*, a *Drosophila* homologue of chromosomal adherins, participates in activation by remote enhancers in the cut and Ultrabithorax genes. *Genetics* 152:577–593.
- Sakai Y, Watanabe T, Kaga K. 2002. Auditory brainstem responses and usefulness of hearing aids in hearing impaired children with Cornelia de Lange syndrome. *Int J Pediatr Otorhinolaryngol* 66:63–69.
- Schrier SA, Sherer I, Deardorff MA, Clark D, Audette L, Gillis L, Kline AD, Ernst L, Loomes K, Krantz ID, Jackson LG. 2011. Causes of death and autopsy findings in a large study cohort of individuals with Cornelia de Lange syndrome and review of the literature. *Am J Med Genet A* 155A:3007–3024.
- Selicorni A, Colli AM, Passarini A, Milani D, Cereda A, Cerutti M, Maitz S, Alloni V, Salvini L, Galli MA, Ghiglia S, Salice P, Danzi GB. 2009. Analysis of congenital heart defects in 87 consecutive patients with Brachmann-de Lange syndrome. *Am J Med Genet A* 149A:1268–1272.
- Singh VP, Gerton JL. 2015. Cohesin and human disease: Lessons from mouse models. *Curr Opin Cell Biol* 37:9–17.
- Tonkin ET, Wang TJ, Lisgo S, Bamshad MJ, Strachan T. 2004. *NIPBL*, encoding a homolog of fungal *Scs2*-type sister chromatid cohesion proteins and fly *Nipped-B*, is mutated in Cornelia de Lange syndrome. *Nat Genet* 36:636–641.
- Whyte WA, Orlando DA, Hnisz D, Abraham BJ, Lin CY, Kagey MH, Rahl PB, Lee TI, Young RA. 2013. Master transcription factors and mediator establish super-enhancers at key cell identity genes. *Cell* 153:307–319.
- Yin JW, Wang G. 2014. The Mediator complex: A master coordinator of transcription and cell lineage development. *Development* 141:977–987.
- Yuan B, Pehlivan D, Karaca E, Patel N, Charng WL, Gambin T, Gonzaga-Jauregui C, Sutton VR, Yesil G, Bozdogan ST, Tos T, Koparir A, Koparir E, Beck CR, Gu S, Aslan H, Yuregir OO, Al Rubeaan K, Alnaqeb D, Alshammari MJ, Bayram Y, Atik MM, Aydin H, Geckinli BB, Seven M, Ulucan H, Fenercioglu E, Ozen M, Jhangiani S, Muzny DM, Boerwinkle E, Tuysuz B, Alkuraya FS, Gibbs RA, Lupski JR. 2015. Global transcriptional disturbances underlie Cornelia de Lange syndrome and related phenotypes. *J Clin Invest* 125:636–651.
- Zakany J, Duboule D. 2007. The role of *Hox* genes during vertebrate limb development. *Curr Opin Genet Dev* 17:359–366.