

# UC Irvine

## UC Irvine Previously Published Works

### Title

Preferential loss of dorsal-hippocampus synapses underlies memory impairments provoked by short, multi-modal stress

### Permalink

<https://escholarship.org/uc/item/1mr03859>

### Journal

Molecular Psychiatry, 19(7)

### ISSN

1359-4184

### Authors

Maras, PM  
Molet, J  
Chen, Y  
[et al.](#)

### Publication Date

2014-07-01

### DOI

10.1038/mp.2014.64

### Copyright Information


This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

## IMAGE

# Preferential loss of dorsal-hippocampus synapses underlies memory impairments provoked by short, multi-modal stress

PM Maras<sup>1</sup>, J Molet<sup>2,5</sup>, Y Chen<sup>1,5</sup>, C Rice<sup>2</sup>, SG Ji<sup>3</sup>, A Solodkin<sup>2,4</sup> and TZ Baram<sup>1,2,4</sup>



*Molecular Psychiatry* (2014) **19**, 745; doi:10.1038/mp.2014.64

The image illustrates the cascade of events that follow short multi-modal, ‘modern-life’-like stress or a less complex stress such as restraint or loud noise. These distinct pathways result in different outcomes: Whereas concurrent physical, psychological and ‘social’ stresses lasting several hours (multi-modal) impaired memory function tested 24 h later, a less complex stress (‘uni-modal’) did not. These divergent outcomes took place even though the intensities of the multi-modal and uni-modal stresses—measured by the output of stress hormones and by the activation of hypothalamic neurons—were indistinguishable. Looking at measures of network activation patterns, there was much overlap between multi-modal and less complex stresses. However, cross-correlation analyses demonstrated ventral-hippocampus–amygdala network predominance after the multimodal stress, and dorsal-hippocampus, sensory-processing network activation after the restraint stress. Analyses of the consequences of these stresses on synaptic integrity demonstrated that only multi-modal stress destroyed synapse-bearing dendritic spines in dorsal-hippocampus CA1, an area crucial for some forms of memory. Indeed, object memory, partially dependent on intact hippocampal function, was profoundly impaired only in mice experiencing multi-modal stress. For more information on this topic, please refer to the article by Maras *et al.* on pages 811–822.

<sup>1</sup>Department of Pediatrics, University of California Irvine, Irvine, CA, USA; <sup>2</sup>Department of Anatomy/Neurobiology, University of California Irvine, Irvine, CA, USA; <sup>3</sup>Medical Scientist Training Program, University of California Irvine, Irvine, CA, USA and <sup>4</sup>Department of Neurology, University of California Irvine, Irvine, CA, USA  
<sup>5</sup>These authors contributed equally to this work.