

UC San Diego

UC San Diego Previously Published Works

Title

Neurocognitive performance in family-based and case-control studies of schizophrenia.

Permalink

<https://escholarship.org/uc/item/1pf4q33z>

Journal

Schizophrenia research, 163(1-3)

ISSN

0920-9964

Authors

Gur, Ruben C
Braff, David L
Calkins, Monica E
[et al.](#)

Publication Date

2015-04-01

DOI

10.1016/j.schres.2014.10.049

Peer reviewed



Neurocognitive performance in family-based and case-control studies of schizophrenia



Ruben C. Gur^{a,*}, David L. Braff^{b,c}, Monica E. Calkins^a, Dorcas J. Dobie^{d,e}, Robert Freedman^f, Michael F. Green^{g,h}, Tiffany A. Greenwood^{b,c}, Laura C. Lazzeroniⁱ, Gregory A. Light^{b,c}, Keith H. Nuechterlein^{g,h}, Ann Olincy^f, Allen D. Radant^{d,e}, Larry J. Seidman^{j,k}, Larry J. Siever^{l,m}, Jeremy M. Silverman^{l,m}, Joyce Sprock^{b,c}, William S. Stone^{j,k}, Catherine A. Sugarⁿ, Neal R. Swerdlow^{b,c}, Debby W. Tsuang^{d,e}, Ming T. Tsuang^{b,c}, Bruce I. Turetsky^a, Raquel E. Gur^a

^a Department of Psychiatry, University of Pennsylvania, Philadelphia, PA, United States

^b Department of Psychiatry, University of California San Diego, La Jolla, CA, United States

^c VISN-22 Mental Illness, Research, Education and Clinical Center (MIRECC), VA San Diego Healthcare System, San Diego, CA, United States

^d Department of Psychiatry and Behavioral Sciences, University of Washington, Seattle, WA, United States

^e VA Puget Sound Health Care System, Seattle, WA, United States

^f Department of Psychiatry, University of Colorado Denver, Aurora, CO, United States

^g Department of Psychiatry and Biobehavioral Sciences, Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA, United States

^h VA Greater Los Angeles Healthcare System, Los Angeles, CA, United States

ⁱ Department of Psychiatry, Stanford University, Palo Alto, CA, United States

^j Department of Psychiatry, Harvard Medical School, Boston, MA, United States

^k Massachusetts Mental Health Center, Public Psychiatry Division of the Beth Israel Deaconess Medical Center, Boston, MA, United States

^l Department of Psychiatry, The Mount Sinai School of Medicine, New York, NY, United States

^m James J. Peters VA Medical Center, New York, NY, United States

ⁿ Department of Biostatistics, University of California Los Angeles School of Public Health, Los Angeles, CA, United States

ARTICLE INFO

Article history:

Received 15 June 2014

Received in revised form 19 October 2014

Accepted 21 October 2014

Available online 26 November 2014

Keywords:

Neurocognition
Schizophrenia
Family-based
Case-control
Ascertainment

ABSTRACT

Background: Neurocognitive deficits in schizophrenia (SZ) are established and the Consortium on the Genetics of Schizophrenia (COGS) investigated such measures as endophenotypes in family-based (COGS-1) and case-control (COGS-2) studies. By requiring family participation, family-based sampling may result in samples that vary demographically and perform better on neurocognitive measures.

Methods: The Penn computerized neurocognitive battery (CNB) evaluates accuracy and speed of performance for several domains and was administered across sites in COGS-1 and COGS-2. Most tests were included in both studies. COGS-1 included 328 patients with SZ and 497 healthy comparison subjects (HCS) and COGS-2 included 1195 patients and 1009 HCS.

Results: Demographically, COGS-1 participants were younger, more educated, with more educated parents and higher estimated IQ compared to COGS-2 participants. After controlling for demographics, the two samples produced very similar performance profiles compared to their respective controls. As expected, performance was better and with smaller effect sizes compared to controls in COGS-1 relative to COGS-2. Better performance was most pronounced for spatial processing while emotion identification had large effect sizes for both accuracy and speed in both samples. Performance was positively correlated with functioning and negatively with negative and positive symptoms in both samples, but correlations were attenuated in COGS-2, especially with positive symptoms.

Conclusions: Patients ascertained through family-based design have more favorable demographics and better performance on some neurocognitive domains. Thus, studies that use case-control ascertainment may tap into populations with more severe forms of illness that are exposed to less favorable factors compared to those ascertained with family-based designs.

Published by Elsevier B.V.

* Corresponding author at: Neuropsychiatry Section, Department of Psychiatry, Perelman School of Medicine, University of Pennsylvania, 3400 Spruce, Philadelphia PA, USA. Tel.: +1 215 662 2915; fax: +1 215 662 7903.
E-mail address: gur@upenn.edu (R.C. Gur).

1. Introduction

Methods of ascertainment are pivotal across biomedical research and are an important consideration in the research design. In genetic

studies, the utility and statistical approach of family-based and unrelated case-controls studies has been discussed (e. g. [Hiekkalinna et al., 2012](#)). The incorporation of endophenotypes to genetic investigations of schizophrenia (SZ) has grown significantly with neurocognitive measures ([Gur et al., 2007a, 2007b](#); [Lee et al., 2015](#)–in this issue; [Nuechterlein et al., 2015](#)–in this issue; [Stone et al., 2015](#)–in this issue) and neurophysiological measures ([Swerdlow et al., 2014](#); [Light et al., 2015](#)–in this issue; [Turetsky et al., 2015](#)–in this issue) playing key roles. Family-based designs enable testing the endophenotype criteria ([Braff et al., 2007](#); [Braff, 2015](#)–in this issue; [Gottesman and Gould, 2003](#)) and, when sufficiently powered, allow for the examination of heritability, association with the disease phenotype and co-segregation within families ([Glahn et al., 2014](#); [Greenwood et al., 2007, 2011, 2013](#)).

Several meta-analyses have reported that adult relatives of probands with SZ show intermediate deficits in neurocognitive measures including executive functions, such as working memory and attention, verbal fluency and sensori-motor speed ([Faraone et al., 2001](#); [Kremen and Hoff, 2004](#); [Sitskoorn et al., 2004](#); [Snitz et al., 2006](#)). Similar deficits have also been observed in younger relatives ([Niemi et al., 2003](#); [Seidman et al., 2006](#); [Keshavan et al., 2010](#); [Agnew-Blais and Siedman, 2013](#)). The neurocognitive domains implicated in family-based studies are similar to deficits observed in case-control studies ([Gur et al., 2001b](#)). Yet, direct evaluation of these complementary ascertainment strategies applying the same measures has not been conducted. The Penn computerized neurocognitive battery (CNB) used in the Consortium on the Genetics of Schizophrenia (COGS) provides a unique opportunity to evaluate effects of ascertainment methods—family-based (COGS-1) vs. case control (COGS-2)—with the same neurocognitive battery across the participating sites.

The CNB, developed in concert with functional neuroimaging studies ([Gur et al., 2010](#)), has been validated in healthy participants and people with SZ ([Gur et al., 2001a,b](#)) and is sensitive to the effects of age and sex ([Gur et al., 2012](#); [Irani et al., 2012](#)). The battery, which provides measures of performance accuracy and response time, was applied in three independent large-scale family-based genetic studies. The Multiplex Multigenerational Investigation of Schizophrenia (MGI; [Gur et al., 2007a](#)) reported that probands demonstrated greatest impairment relative to healthy controls, with intermediate performance of family members. Liability for SZ affected the speed–accuracy tradeoff differently for specific neurocognitive domains. Significant heritability estimates were obtained for accuracy of verbal, facial, and spatial memory and spatial and emotion processing. For speed, estimates of heritability were significant for abstraction and mental flexibility, attention, face memory, and spatial and sensorimotor processing. The results of the Project among African-Americans to Explore Risks for Schizophrenia (PAARTNERS) revealed that patients with SZ exhibited less accuracy and speed in most neurocognitive domains than their relatives, who were impaired relative to HCS in most domains. Significant heritabilities were observed for most neurocognitive domains, with the highest for accuracy of abstraction and mental flexibility, verbal memory, face memory, spatial processing, and emotion processing and for speed of attention ([Calkins et al., 2010](#)).

In COGS-1 all of the measures applied from the Penn CNB (Abstraction and Mental Flexibility, Face Memory, Spatial Memory, Spatial Processing, Sensorimotor Dexterity, and Emotion Recognition) were significantly heritable with heritability estimates ranging from 24% for Spatial Memory to 55% for Spatial Processing ([Greenwood et al., 2007](#)). These heritabilities are in the same range as the heritability of SZ itself in the COGS-1 families ([Light et al., in press](#)). Furthermore, we noted sex differences in familiarity effects with male probands' performance predictive of performance of their unaffected relatives ([Calkins et al., 2013](#)). The subsequent application of the CNB in the case-control design of COGS-2 enabled evaluation of the pattern of performance of individuals with SZ, compared to HCS, ascertained in family-based and case-control designs. We noted that in some endophenotypic measures in COGS-1 probands were less impaired than observed in

other samples of patients with SZ ([Greenwood et al., 2007](#)). The major ascertainment difference between the samples is that patients recruited for COGS-1 required the availability of parents and siblings while COGS-2 permitted participation of patients regardless of family availability ([Swerdlow et al., 2015](#)–in this issue). This difference likely affects multiple demographic characteristics related to age, education, socioeconomic status as well as severity of illness, favoring COGS-1. We hypothesized that while the profile of impairment would be similar, probands in the COGS-1 family-based ascertainment would perform better than those ascertained as cases in COGS-2.

2. Materials and methods

2.1. Participants

Details on the COGS-1 and COGS-2 samples' ascertainment, inclusion and exclusion criteria and clinical assessment are provided elsewhere in this issue ([Braff et al.](#); [Swerdlow et al.](#)). Briefly, COGS-1, a family-based design, and COG-2, a case-control design, included probands 18–65 years old who met DSM-IV criteria for schizophrenia based on established diagnostic procedures. COGS-1 required that both biological parents were available for genotyping, and that at least one full sibling, unaffected with schizophrenia, was available for endophenotyping and genotyping. Probands with one available parent but two or more available siblings, with at least one unaffected by schizophrenia, were also included, as were probands with no available parents but three or more available siblings (≥ 1 unaffected by schizophrenia). COGS-2 had the same diagnostic requirements for probands and controls as COGS-1, but the availability of family members was not required. Here we focus on COG-1 and COGS-2 patients and controls who completed the CNB testing. COGS-1 included 328 patients and 497 controls and COGS-2 included 1195 patients and 1009 controls. Demographic information is presented at the top portion of [Table 1](#). As can be seen, COGS-1 patients did not differ from their controls in age, or parental education, but had lower education and lower reading level with moderate effect sizes. COGS-2 patients were significantly older than their controls as well as less educated with lower parental education and Wide Range Achievement Test (WRAT4, [Wilkinson and Robertson, 2006](#)) scores, with effect sizes ranging from moderate to large. COGS-1 controls were younger, attained higher educational level, had higher paternal education and higher WRAT scores compared to COGS-2 controls, but all these effect sizes were small (<2 SD). COGS-1 patients were younger and had higher educational attainment, higher parental education and higher WRAT compared to COGS-2 patients and these effect sizes were moderate to large. Notably, the variances did not differ between the samples on most measures (Satterthwaite's correction was used for these *p* values).

2.2. The computerized neurocognitive battery (CNB)

The Penn CNB ([Gur et al., 2001a,b](#)) was administered in the COGS along with other candidate endophenotypes. It was abbreviated to reduce redundancy with other core endophenotypes. COGS-1 and COGS-2 CNB differed in three ways. First, for COGS-1 Degraded Stimulus CPT and CPT, identical pairs were used to cover the attention domain ([Nuechterlein et al., 2015](#)–in this issue), while in COGS-2 the Penn CPT data were also added to allow the full CNB to be represented. Second, for measuring working memory, different forms of the letter *n*-back test were used in COGS-1 and COGS-2. Third, many participants from COGS-1 did not receive the delayed recognition tests because the CNB was administered last and time limitations and fatigue attenuated the test sessions.

The CNB was administered on Macintosh computers (Apple Inc., Cupertino, California) in a fixed order and included brief standardized rest periods, for a total administration time of about 60 min. The

Table 1
Demographics, raw performance data and effect sizes (Cohen's *d*) in COGS-1 and COGS-2. Only large effects sizes were bolded (i.e., $ES > .8$).

			COGS-1 SZ			COGS-1 HCS			SZ vs. HCS		COGS-2 SZ			COGS-2 HCS			SZ vs. HCS		SZ1 vs. SZ2		HCS1 vs. HCS2		
			N	Mean	SD	N	Mean	SD	p	ES (d)	N	Mean	SD	N	Mean	SD	p	ES (d)	p	ES	p	ES	
DOMAIN	TEST (CONSTRUCT)	Age	328	34.86	11.00	497	36.20	12.67	0.0878	-0.11	1195	46.24	11.30	1009	38.62	13.18	<0.0001	0.63	<.0001	-1.01	0.0008	-0.19	
		Education	328	13.62	2.17	497	15.39	2.34	<0.0001	-0.78	1195	12.69	2.13	1009	14.97	2.22	<0.0001	-1.05	<.0001	0.43	0.0056	0.18	
		Mother education	323	14.41	2.96	496	14.00	3.04	0.0579	0.13	1130	12.01	3.88	999	13.85	3.31	<0.0001	-0.51	<.0001*	0.65	0.1292	0.05	
		Father education	320	14.95	3.38	483	14.36	3.40	0.0161	0.16	1069	11.81	4.66	979	13.77	3.91	<0.0001	-0.45	<.0001*	0.71	0.0058*	0.16	
		WRAT	328	102.23	11.13	497	107.16	10.61	<0.0001	-0.46	1195	95.65	12.80	1009	105.65	10.74	<0.0001	-0.84	<.0001	0.53	0.005	0.14	
		Abstraction/ flexibility	PCET Accuracy	321	1.79	1.15	489	2.41	0.96	<0.0001	-0.66	1145	1.36	0.88	991	2.09	0.92	<0.0001	-0.80	<.0001*	0.46	<.0001	0.35
			PCET Speed	321	3058.64	1887.20	489	2316.19	1251.25	<0.0001	0.62	1145	3517.44	1914.26	991	2480.85	1159.78	<0.0001	0.64	0.001	-0.24	0.0004	-0.14
			LNB Accuracy	297	26.20	3.80	493	28.14	2.32	<0.0001	-0.95	1107	16.40	3.38	992	18.53	1.89	<0.0001	-0.77				
		Episodic memory	(WM) Speed	297	592.79	164.72	493	537.75	129.79	<0.0001	0.43	1107	625.26	176.74	992	548.77	125.52	<0.0001	0.49	0.0219	-0.19	0.2624	-0.09
			WORD Accuracy	201	32.38	4.19	309	34.80	3.49	<0.0001	-0.69	1173	31.98	4.23	999	34.20	3.51	<0.0001	-0.57	0.1712	0.09	0.0089	0.17
(VME) Speed	201		1602.15	1126.86	309	1233.33	342.65	<0.0001	1.09	1173	2035.80	758.00	999	1532.50	336.00	<0.0001	0.84	<.0001	-0.53	<.0001	-0.89		
FACE Accuracy	333		31.36	4.13	499	34.29	3.34	<0.0001	-0.87	1171	30.36	3.96	999	33.31	3.40	<0.0001	-0.80	0.0003	0.25	<.0001	0.29		
Complex cognition	(FME) Speed	333	1928.42	955.70	498	1639.07	535.08	<0.0001	0.61	1171	2195.94	802.45	999	1791.03	435.39	<0.0001	0.61	<.0001	-0.32	<.0001*	-0.32		
	SHAPE Accuracy	328	14.07	2.57	495	15.72	2.12	<0.0001	-0.73	1146	13.44	2.30	993	15.25	2.31	<0.0001	-0.78	0.0021	0.27	0.0013	0.21		
	(SME) Speed	315	1532.40	738.90	487	1372.43	389.38	0.0004	0.41	1141	1746.36	664.24	993	1465.27	393.58	<0.0001	0.51	<.0001	-0.31	<.0001	-0.24		
	JOLO Accuracy	308	21.81	5.96	489	22.46	5.31	0.1209	-0.12	1160	18.98	6.44	1000	21.93	5.26	<0.0001	-0.50	<.0001	0.45	0.094	0.10		
Social cognition	(SPA) Speed	307	3364.79	1511.92	488	2940.26	1119.65	<0.0001	0.36	1157	4516.18	2327.27	1000	3168.83	1198.42	<0.0001	0.71	<.0001*	-0.53	0.0006	-0.19		
	ER40 Accuracy	328	30.68	4.70	497	33.69	2.73	<0.0001	-1.02	1139	30.25	4.57	993	33.35	3.05	<0.0001	-0.79	0.4343	0.09	0.0351	0.11		
Sensorimotor	(EMO) Speed	328	2803.23	1159.52	497	2037.94	584.93	<0.0001	1.35	1139	3087.86	1059.63	993	2183.37	560.07	<0.0001	1.05	0.0036	-0.26	<.0001	-0.26		
	MPRACT Speed	327	955.28	468.18	497	747.08	258.07	<0.0001	0.97	1195	1190.46	478.06	1009	805.18	190.92	<0.0001	1.03	<.0001	-0.49	<.0001	-0.27		
	TAP (SM) Speed									1176	94.09	16.73	1002	106.84	13.27	<0.0001	-0.84						
Attention	CPT Accuracy									1158	108.86	12.62	999	115.10	8.97	<0.0001	-0.56						
	(ATT) Speed									1157	533.19	69.81	999	483.92	49.57	<0.0001	0.80						

*Variances are significantly different and p values are Satterthwaite corrected.

following neurocognitive domains were assessed (Gur et al., 2010, 2012): 1) Executive Functions: abstraction and mental flexibility (Penn Conditional Exclusion Test, PCET); working memory (WM, letter n-back, 1-back and 2-back conditions). 2) Episodic Memory: word memory (Penn Word Memory Test); face memory (Penn Face Memory Test); spatial memory (Visual Object Learning Test). 3) Complex Cognition: spatial processing (Computerized Judgment of Line Orientation, JOLO). 4) Social Cognition: emotion processing (Penn Emotion Recognition Test). 5) Sensorimotor Speed: motor praxis and finger tapping test. For all but the last domain, two summary functions were calculated: accuracy of responses and speed, the median response time for correct answers.

2.3. Statistical analysis

The performance scores were transformed to their standard equivalents (*z*-scores) within each sample (COGS-1, COGS-2), based on the controls in that sample. These *z*-scores were the dependent measures in a Mixed Model analysis (SAS PROC MIXED), with Sample and Diagnosis as grouping factors and neurocognitive Domain as a within-group factor and age and parental education (average of mother's and father's) as covariates. We did not use patient's education as a covariate to avoid committing the "matching fallacy" because schizophrenia itself interferes with educational attainment and covarying or matching for it will falsely remove relevant variance (Meehl, 1970). Instead, we covaried parental education as recommended (Resnick, 1992) and widely practiced. Similarly, we did not covary for the WRAT score, as it is a measure of cognitive abilities that is highly correlated with performance on the neurocognitive battery. Again, covarying for it will remove relevant variance. The model was applied separately to the accuracy (7 domains) and speed (8 domains) scores. Mixed Model analysis was preferred to a MANOVA mainly because it can accommodate missing values while MANOVA would eliminate all subjects with missing data on any test. This would have affected mainly the first sample, where the word memory test was introduced in the middle of the COGS-1 study. Initially, the model included sex as a grouping factor, but since the main effects have been well established (e.g. better performance of males on spatial tests and of females on memory and emotion processing) and there were no interactions by cohort or diagnostic group, it was dropped from subsequent analyses. To examine the association between neurocognitive measures and clinical status, we correlated performance on the neurocognitive domains with clinical ratings on the Scale for the

Assessment of Negative Symptoms (SANS, Andreasen, 1984a), the Scale of Assessment of Positive Symptoms (SAPS, Andreasen, 1984b) and the Global Assessment of Functioning (GAF, Hall and Parks, 1995).

3. Results

3.1. Performance comparisons on the computerized neurocognitive battery

The means and standard deviations, as well as *p* values for Student's *t*-tests and effect sizes comparing patients and controls within each sample and between samples are presented in the bottom portion of Table 1. The *z*-scores of patients compared to their respective control groups are illustrated in Fig. 1.

As can be seen in Table 1, both COGS-1 patients and COGS-2 patients are impaired relative to their respective controls, although the effect sizes range from small to moderate in COGS-1, with six large effect sizes, while in COGS-2 all effect sizes are at least moderate and seven of them large. Notably, COGS-1 controls outperformed COGS-2 controls on all domains, but these effect sizes are generally small with only one large effect (speed of word memory). COGS-1 patients outperformed COGS-2 patients on most domains with effect sizes ranging from small to moderate.

The Mixed Model analysis on the accuracy scores showed main effects for sample, $F(1, 17,000) = 30.32, p < 0.0001$, Diagnostic group, $F(1, 17,000) = 414.85, p < 0.0001$, and Domain, $F(6, 17,000) = 18.46, p < 0.0001$. Also significant were the two-way interactions of Sample \times Diagnosis, $F(1, 17,000) = 10.17, p = 0.0014$, Sample \times Domain, $F(6, 17,000) = 3.61, p = 0.0014$, and Diagnosis \times Domain, $F(6, 17,000) = 18.52, p < 0.0001$. The three-way interaction of Sample \times Diagnosis \times Domain was also significant, $F(6, 17,000) = 3.18, p = 0.004$. The Mixed Model analysis on the speed scores showed main effects of sample, $F(1, 20,000) = 10.29, p = 0.0013$, Diagnostic group, $F(1, 20,000) = 197.24, p < 0.0001$ and Domain, $F(7, 20,000) = 65.89, p < 0.0001$. Also significant were the two-way interactions of Sample \times Domain, $F(7, 20,000) = 16.34, p < 0.0001$, and Diagnosis \times Domain, $F(7, 20,000) = 66.02, p < 0.0001$, but not Sample \times Diagnosis, $F(1, 20,000) = 0.02, p = 0.8897$. The three-way interaction of Sample \times Diagnosis \times Domain was also significant, $F(7, 20,000) = 16.34, p < 0.0001$.

Decomposing the three-way interactions for accuracy indicated that COGS-1 patients were less impaired than COGS-2 counterparts,




Fig. 1. Neurocognitive performance of COG-1 and COGS-2 participants: Mean ($\pm 95\%$ Confidence Interval) accuracy (left panel) and speed (right panel) of patients are presented in *z*-scores compared to the respective control groups (*z*-scores for response times are inverted for speed so that higher scores always reflects better performance). SZ = Schizophrenia, HCS = Healthy Comparison Subjects; ABF = Abstraction and Mental Flexibility, ATT = Attention, WM = Working Memory, VME = Word Memory, FME = Face Memory, SME = Spatial Memory, SPA = Spatial Processing, EMO = Emotion Identification, SM = Sensori-Motor, MOT = Motor Speed.

compared to their respective controls, in abstraction and mental flexibility ($p < 0.0001$) and in spatial processing ($p < 0.0001$). For speed, the three-way interaction reflected differentially greater impairment in COGS-2 patients for verbal memory ($p < 0.0001$), spatial processing ($p < 0.0001$) and sensorimotor speed ($p < 0.0001$).

3.2. Correlations of performance with clinical status

COGS-1 patients had less severe negative and positive symptoms and better functioning (Table 2). Significant correlations in the expected directions, namely negative for symptom severity and positive for GAF, were obtained in both cohorts (Table 3). They were comparable in the two cohorts for GAF, ranging generally from 0.3 to 0.47. The clinical severity ratings for COGS-1 correlated significantly with neurocognitive performance, with greater severity associated with poorer performance. These correlations were of similar magnitude for SANS and SAPS, hovering around -0.3 and as high as -0.5 for emotion processing speed. For COGS-2, the correlations between symptom severity and neurocognitive performance were considerably attenuated, hovering around -0.1 and significant for SANS but only few for SAPS.

4. Discussion

The Consortium on the Genetics of Schizophrenia (COGS) allowed for the comparison of neurocognitive performance deficits between individuals with schizophrenia ascertained through a family-based sampling (COGS-1) and those ascertained through case-control sampling (COGS-2) with their differing ascertainment strategies as discussed above and in this issue (cf. Swerdlow et al., 2015–in this issue). The results indicated very similar neurocognitive deficit profiles, for COGS-1 and COGS-2 schizophrenia patients, strongly supporting the sensitivity of the neurocognitive battery to deficits characteristic of schizophrenia. Against this similarity of profiles, the results generally supported the hypothesis that patients ascertained through family sampling are less impaired than those ascertained as cases in a case-control design. The difference between the two groups of patients was robust even when controlled for age and parental education. The effect sizes comparing patients to controls were in the small to moderate range in the COGS-1 sample and in the moderate to large range in the COGS-2 sample reflecting the greater deficits in COGS-2 patients. Thus, studies using a case-control design that does not require availability of family and therefore is likely to include older and more chronic patients should expect patients with greater neurocognitive impairment than studies where family is engaged.

The significant diagnosis by domain by sample interactions indicate that the case-control ascertained sample did not perform more poorly to the same extent in all domains. The biggest difference between the samples was in spatial processing, both for accuracy and for speed, and for sensorimotor speed. Spatial processing was the only measure of complex cognition included in this iteration of the Penn CNB (Gur et al., 2010; Moore et al., in press). It represents temporo-parietal functioning and it loads heavily on general intellectual abilities. Lower performance on this test is consistent with lower IQ estimates based on the WRAT scores. However, the COGS-2 sample was also differentially

Table 2
Clinical measures of patients in COGS-1 and COGS-2.

	COGS-1			COGS-2			COGS-1 vs. COGS-2	
	N	Mean	SD	N	Mean	SD	p	ES
SANS	461	4.09	5.68	1407	11.03	5.66	<.0001	-1.22
SAPS	461	2.40	3.87	1405	6.92	4.02	<.0001	-1.14
GAF	477	69.73	22.09	2460	62.08	22.72	<.0001	0.34

Table 3
Correlations between clinical measures and neurocognitive performance in COGS-1 and COGS-2.

	Accuracy				Speed														
	ABF	ATT	WM	VME	FME	SME	SPA	EMO	ABF	ATT	WM	VME	FME	SME	SPA	EMO	SM	MOT	
COGS-1																			
SANS	-0.303	NA	-0.366	-0.287	-0.403	-0.405	-0.126	-0.361	-0.294	NA	-0.240	-0.365	-0.189	-0.219	-0.268	-0.550	-0.470	NA	NA
SAPS	-0.265	NA	-0.351	-0.251	-0.321	-0.376	-0.117	-0.335	-0.328	NA	-0.185	-0.257	-0.173	-0.227	-0.220	-0.514	-0.493	NA	NA
GAF	0.349	NA	0.356	0.313	0.403	0.375	0.083	0.371	0.291	NA	0.247	0.348	0.185	0.189	0.207	0.515	0.402	NA	NA
COGS-2																			
SANS	-0.107	-0.148	-0.113	-0.111	-0.083	-0.105	-0.068	-0.128	-0.073	-0.118	-0.120	-0.129	-0.108	-0.133	-0.090	-0.140	-0.153	-0.145	-0.145
SAPS	-0.117	-0.077	-0.096	-0.086	-0.070	-0.098	-0.049	-0.041	-0.034	0.003	-0.040	-0.047	-0.013	-0.056	-0.050	-0.056	-0.059	-0.019	-0.019
GAF	0.384	0.275	0.368	0.286	0.367	0.373	0.231	0.370	0.317	0.364	0.256	0.391	0.308	0.283	0.343	0.473	0.464	0.380	0.380

ABF = Abstraction and Mental Flexibility, ATT = Attention, WM = Working Memory, VME = Verbal Memory, FME = Face Memory, SME = Spatial Memory, SPA = Spatial Processing, EMO = Emotion Identification, SM = Sensori-Motor, MOT = Motor Speed.

impaired in working memory suggesting that executive functioning may likewise play a role in their neurocognitive dysfunction.

Two tests were added to COGS-2 that were not included in COGS-1. The CPT in COGS-2 produced moderate effect size (-0.56 SDs) for accuracy and a large effect size (0.80 SD) for speed (see also Nuechterlein et al., 2015—in this issue). The other test that was not used in COGS-1 was finger tapping, which assessed motor-speed. This test produced a large effect size of -0.84 SD. The sensitivity of finger tapping to deficits in schizophrenia has been reported (e.g., Da Silva et al., 2012). The present study suggests that both attention and motor speed are sensitive measures that should be included in neurocognitive batteries assessing deficits in schizophrenia.

As with neurocognitive performance, symptom severity was greater and functioning was poorer in patients from COGS-2 compared to COGS-1. Neurocognitive measures were significantly correlated with these clinical parameters in both samples. It is notable that the correlations were nearly identical in both samples for GAF, but differed for severity of symptoms. The lower correlations for SAPS than for SANS are consistent with previous studies, most using the case-control design (Nuechterlein et al., 2011). Our finding that they are of similar magnitude for SAPS and SANS in COGS-1 suggests that positive symptoms may have more adverse effect on functioning of younger patients who are still engaged with family. The higher correlations with symptoms in COGS-1 are not explained by greater variance of symptom severity or performance since these did not generally differ between the samples. Thus, studies that ascertain patients in a case-control design are more likely to find lower correlations between symptom severity and neurocognitive performance, with a magnitude especially low for positive symptoms. When such designs are used in treatment trials, they are therefore likely to underestimate the potential impact of improved neurocognitive performance on symptoms and functioning.

The study has several limitations and caveats. Most importantly, two tests were missing from one of the samples and therefore no comparable data were available. Secondly, the samples differed on several demographic variables including age and education. While these factors were controlled for *a posteriori* in the analyses, it is possible they relate to other factors associated with illness that may have affected the results. Medications can be related to cognitive performance, especially motor speed and these effects have not been evaluated in the present study.

These caveats notwithstanding, the present study offers robust support, with definitively large samples of carefully diagnosed and multi-site quality assured testing, to the presence in schizophrenia of a specific pattern of cognitive deficits that is related to dysfunctional brain systems (Roalf et al., 2014). The results also support the importance of ascertainment strategy (e.g. family-based vs. case-control ascertainment) and the hypothesis that family-based ascertainment will produce samples that are less neurocognitively impaired. This finding of large CNB case-control detected deficits offers a solid neurobiologically informed platform for genomic studies to follow in the large, well-characterized COGS-2 schizophrenia sample (Braff, 2015—in this issue).

Role of funding source

This work was supported by collaborative R01 grants from the National Institute of Mental Health. COGS-1 and COGS-2: MH065571 UCSD, MH065707 UCLA, MH065554 MSSM, MH065578 Penn, MH065558 Washington; COGS-1 Only: MH065588 Colorado, MH065562 Harvard; Seidman and Stone are on a subcontract for COGS-2; COGS-2 Only: MH86135 Stanford.

Contributors

All authors significantly contributed to study design, data collection and reviewed and approved this manuscript.

Conflict of interest

Dr. Green has been a consultant to AbbVie, Biogen, DSP, EnVivo/Forum and Roche, and he is on the scientific advisory board of Mnemosyne. He has received research funds from Amgen. Dr. Lazzaroni is an inventor on a patent application filed by Stanford University on genetic polymorphisms associated with depression. Dr. Light has served as a consultant for Astellas, Forum, and Neuroverse. Dr. Nuechterlein has received unrelated research support

from Janssen Scientific Affairs, Genentech, and Brain Plasticity, Inc., and has consulted to Genentech, Otsuka, Janssen, and Brain Plasticity, Inc. Dr. Swerdlow has been a consultant for Genco Sciences, Ltd. All other authors declare that they have no conflict of interest.

Acknowledgments

We thank the research staff who administered the CNB, Allison Mott CNB Implementation Manager, University of Pennsylvania and the patients, families and healthy volunteers who participated in the study.

References

- Agnew-Blais, J., Siedman, L.J., 2013. Neurocognition in youth adults under age 30 at familial risk for schizophrenia: a quantitative and qualitative review. *Cogn. Neuropsychiatry* 18 (1–2), 44–82.
- Andreasen, N.C., 1984a. Scale for the Assessment of Negative Symptoms (SANS). University of Iowa, Iowa City, Iowa.
- Andreasen, N.C., 1984b. Scale for the Assessment of Positive Symptoms (SAPS). University of Iowa, Iowa City, Iowa.
- Braff, D.L., 2015. The importance of endophenotypes in schizophrenia research: past, present and future. *Schizophr. Res.* 163 (1–3), 1–8 (in this issue).
- Braff, D.L., Freedman, R., Schork, N.J., Gottesman, I.I., 2007. Deconstructing schizophrenia: an overview of the use of endophenotypes in order to understand a complex disorder. *Schizophr. Bull.* (33), 21–32.
- Calkins, M.E., Tepper, P., Gur, R.C., Ragland, J.D., Klei, L., Wiener, H.W., Richard, J., Savage, R.M., Allen, T.B., O'Jile, J., Devlin, B., Kwentus, J., Aliyu, M.H., Bradford, L.D., Edwards, N., Lyons, P.D., Nimgaonkar, V.L., Santos, A.B., Go, R.C., Gur, R.E., 2010. Project among African-Americans to explore risks for schizophrenia (PAARTNERS): evidence for impairment and heritability of neurocognitive functioning in families of schizophrenia patients. *Am. J. Psychiatry* 167 (4), 459–472.
- Calkins, M.E., Ray, A., Gur, R.C., Freedman, R., Green, M.F., Greenwood, T.A., Light, G.A., Nuechterlein, K.H., Olincy, A., Radant, A.D., Seidman, L.J., Siever, L.J., Silverman, J.M., Stone, W.S., Sugar, C., Swerdlow, N.R., Tsuang, D.W., Tsuang, M.T., Turetsky, B.I., Braff, D.L., Lazzaroni, L.C., Gur, R.E., 2013. Sex differences in familiarity effects on neurocognitive performance in schizophrenia. *Biol. Psychiatry* 73 (10), 976–984.
- Da Silva, F.N., Irani, F., Richard, J., Bressinger, C.M., Bilker, W.B., Gur, R.E., Gur, R.C., 2012. More than just tapping: index finger-tapping measures procedural learning in schizophrenia. *Schizophr. Res.* 137 (1–3), 234–240.
- Faraone, S.V., Green, A., Seidman, L.J., Tsuang, M.T., 2001. "Schizotaxia": clinical implications and new directions for research. *Schizophr. Bull.* 27 (1), 1–18.
- Glahn, D.C., Knowles, E.E., McKay, D.R., Sproueten, E., Raventos, H., Blangero, J., Gottesman, I.I., Almasy, L., 2014. Arguments for the sake of endophenotypes: examining common misconceptions about the use of endophenotypes in psychiatric genetics. *Am. J. Med. Genet. B Neuropsychiatr. Genet.* 165B (2), 122–130.
- Gottesman, I.I., Gould, T.D., 2003. The endophenotype concept in psychiatry: etymology and strategic intentions. *Am. J. Psychiatry* 160 (4), 636–645.
- Greenwood, T.A., Braff, D.L., Light, G.A., Cadenhead, K.S., Calkins, M.E., Dobie, D.J., Freedman, R., Green, M.F., Gur, R.E., Gur, R.C., Mintz, J., Nuechterlein, K.H., Olincy, A., Radant, A.D., Seidman, L.J., Siever, L.J., Silverman, J.M., Stone, W.S., Swerdlow, N.R., Tsuang, D.W., Tsuang, M.T., Turetsky, B.I., Schork, N.J., 2007. Initial heritability analyses of endophenotypic measures for schizophrenia: the Consortium on the Genetics of Schizophrenia. *Arch. Gen. Psychiatry* 64 (11), 1242–1250.
- Greenwood, T.A., Lazzaroni, L.C., Murray, S.S., Cadenhead, K.S., Calkins, M.E., Dobie, D.J., Green, M.F., Gur, R.E., Gur, R.C., Hardiman, G., Kelsoe, J.R., Leonard, S., Light, G.A., Nuechterlein, K.H., Olincy, A., Radant, A.D., Schork, N.J., Seidman, L.J., Siever, L.J., Silverman, J.M., Stone, W.S., Swerdlow, N.R., Tsuang, D.W., Turetsky, B.I., Freedman, R., Braff, D.L., 2011. Analysis of 94 candidate genes and 12 endophenotypes for schizophrenia from the Consortium on the Genetics of Schizophrenia. *Am. J. Psychiatry* 168 (9), 930–946.
- Greenwood, T.A., Swerdlow, N.R., Gur, R.E., Cadenhead, K.S., Calkins, M.E., Dobie, D.J., Freedman, R., Green, M.F., Gur, R.C., Lazzaroni, L.C., Nuechterlein, K.H., Olincy, A., Radant, A.D., Ray, A., Schork, N.J., Seidman, L.J., Siever, L.J., Silverman, J.M., Stone, W.S., Sugar, C.A., Tsuang, D.W., Tsuang, M.T., Turetsky, B.I., Light, G.A., Braff, D.L., 2013. Genome-wide linkage analyses of 12 endophenotypes for schizophrenia from the Consortium on the Genetics of Schizophrenia. *Am. J. Psychiatry* (170), 521–532.
- Gur, R.C., Ragland, J., Moberg, P., Turner, T., Bilker, W.B., Kohler, C., Siegel, S.J., Gur, R.E., 2001a. Computerized neurocognitive scanning: I. Methodology and validation in healthy people. *Neuropsychopharmacology* 25 (5), 766–776.
- Gur, R.C., Ragland, J.D., Moberg, P.J., Bilker, W.B., Kohler, C., Siegel, S.J., Gur, R.E., 2001b. Computerized neurocognitive scanning: II. The profile of schizophrenia. *Neuropsychopharmacology* 25 (5), 777–788.
- Gur, R.E., Nimgaonkar, V.L., Almasy, L., Calkins, M.E., Ragland, J.D., Pogue-Geile, M.F., Kanes, S., Blangero, J., Gur, R.C., 2007a. Neurocognitive endophenotypes in a multiplex multigenerational family study of schizophrenia. *Am. J. Psychiatry* 164 (5), 813–819.
- Gur, R.E., Calkins, M.E., Gur, R.C., Horan, W.P., Nuechterlein, K.H., Seidman, L.J., Stone, W.S., 2007b. The Consortium on the Genetics of Schizophrenia: neurocognitive endophenotypes. *Schizophr. Bull.* 33 (1), 49–68.
- Gur, R.C., Richard, J., Hughett, P., Calkins, M.E., Macy, L., Bilker, W.B., Bressinger, C., Gur, R.E., 2010. A cognitive neuroscience-based computerized battery for efficient measurement of individual differences: standardization and initial construct validation. *J. Neurosci. Methods* 187 (2), 254–262.
- Gur, R.C., Richard, J., Calkins, M.E., Chiavacci, R., Hansen, J.A., Bilker, W.B., Loughead, J., Connolly, J.J., Qiu, H., Mentch, F.D., Abou-Sleiman, P.M., Hakonarson, H., Gur, R.E., 2012. Age group and sex differences in performance on a computerized neurocognitive battery in children age 8–21. *Neuropsychology* 26 (2), 251–265.

- Hall, R.C., Parks, J., 1995. The modified global assessment of functioning scale: addendum. *Psychosomatics* 36 (4), 416–417.
- Hiekkaalinn, T., Goering, H.H., Lambert, B., Weiss, K.M., Norrgrann, P., Schaëffer, A.A., Terwilliger, J.D., 2012. On the statistical properties of family-based association tests in datasets containing both pedigrees and unrelated case-control samples. *Eur. J. Hum. Genet.* (20), 217–223.
- Irani, F., Brensinger, C.M., Richard, J., Calkins, M.E., Moberg, P.J., Bilker, W., Gur, R.E., Gur, R.C., 2012. Computerized neurocognitive test performance in schizophrenia: a lifespan analysis. *Am. J. Geriatr. Psychiatry* 20 (1), 41–52.
- Keshavan, M., Kulkarni, S., Bhojraj, T., Francis, A., Diwadkar, V., Montrose, D., Seidman, L.J., Sweeney, J., 2010. Premorbid cognitive deficits in young relatives of schizophrenia patients. *Front. Hum. Neurosci.* 62 (3), 1–14.
- Kremen, W., Hoff, A., 2004. Neurocognitive deficits in biological relatives of individuals with schizophrenia. In: Stone, W., Faraone, S., Tsuang, M.M. (Eds.), *Early Clinical Intervention and Prevention in Schizophrenia*. Humana Press, Totowa, NJ, pp. 133–154.
- Lee, J., Green, M.F., Calkins, M.E., Greenwood, T.A., Gur, R.E., Gur, R.C., Lazzaroni, L.C., Light, G.A., Nuechterlein, K.H., Radant, A.D., Seidman, L.J., Siever, L.J., Silverman, J.M., Sprock, J., Stone, W.S., Sugar, C.A., Swerdlow, N.R., Tsuang, D.W., Tsuang, M.T., Turetsky, B.I., Braff, D.L., 2015. Verbal working memory in schizophrenia: the moderating role of smoking status and antipsychotic medications. *Schizophr. Res.* 163 (1–3), 24–31 (in this issue).
- Light, G.A., Greenwood, T.A., Swerdlow, N.R., Calkins, M.E., Freedman, R., Green, M.F., Gur, R.E., Gur, R.C., Lazzaroni, L.C., Nuechterlein, K.H., Olincy, A., Radant, A.D., Seidman, L.J., Siever, L.J., Silverman, J.M., Sprock, J., Stone, W.S., Sugar, C.A., Tsuang, D.W., Tsuang, M.T., Turetsky, B.I., Braff, D.L., 2014. Comparison of the heritability of schizophrenia and endophenotypes in the COGS-1 family study. *Schizophr. Bull.* (in press).
- Light, G.A., Swerdlow, N.R., Thomas, M.L., Calkins, M.E., Green, M.F., Greenwood, T.A., Gur, R.E., Gur, R.C., Lazzaroni, L.C., Nuechterlein, K.H., Pela, M., Radant, A.D., Seidman, L.J., Sharp, R.F., Siever, L.J., Silverman, J.M., Sprock, J., Stone, W.S., Sugar, C.A., Tsuang, D.W., Tsuang, M.T., Braff, D.L., Turetsky, B.I., 2015. Validation of mismatch negativity and P3a for use in multi-site studies of schizophrenia: characterization of demographic, clinical, cognitive, and functional correlates in the COGS-2. *Schizophr. Res.* 163 (1–3), 63–72 (in this issue).
- Meehl, P.E., 1970. Nuisance variables and the ex post facto design. In: Radner, M., Winokur, S. (Eds.), *Minnesota Studies in the Philosophy of Science*. University of Minnesota Press, Minneapolis, pp. 373–402.
- Moore, T.M., Reise, S.T., Gur, R.E., Hakonarson, H., Gur, R.C., 2014. Psychometric properties of the Penn Computerized Neurocognitive Battery. *Neuropsychology* (in press) [Epub ahead of print].
- Niemi, L., Suvisaari, J., Tuulio-Henriksson, A., Lonnqvist, J., 2003. Childhood developmental abnormalities in schizophrenia: evidence from high-risk studies. *Schizophr. Res.* 60 (2–3), 239–258.
- Nuechterlein, K.H., Subotnik, K.L., Green, M.F., Ventura, J., Asarnow, R.F., Gitlin, M.J., Yee, C.M., Gretchen-Doorly, D., Mintz, J., 2011. Neurocognitive predictors of work outcome in recent-onset schizophrenia. *Schizophr. Bull.* 37 (Suppl. 2), S33–S40.
- Nuechterlein, K.H., Green, M.F., Calkins, M.E., Greenwood, T.A., Gur, R.E., Gur, R.C., Lazzaroni, L.C., Light, G.A., Radant, A.D., Seidman, L.J., Siever, L.J., Silverman, J.M., Sprock, J., Stone, W.S., Sugar, C.A., Swerdlow, N.R., Tsuang, D.W., Tsuang, M.T., Turetsky, B.I., Braff, D.L., 2015. Attention/vigilance in schizophrenia: performance results from a large multi-site study of the Consortium on the Genetics of Schizophrenia (COGS). *Schizophr. Res.* 163 (1–3), 38–46 (in this issue).
- Resnick, S.M., 1992. Matching for education in studies of schizophrenia. *Arch. Gen. Psychiatry* 49 (3), 246.
- Roalf, D.R., Ruparel, K., Gur, R.E., Bilker, W., Gerraty, R., Elliott, M.A., Gallagher, R.S., Almasy, L., Pogue-Geile, M.F., Prasad, K., Wood, J., Nimgaonkar, V.L., Gur, R.C., 2014. Neuroimaging predictors of cognitive performance across a standardized neurocognitive battery. *Neuropsychology* 28 (2), 161–176.
- Seidman, L.J., Giuliano, A.J., Smith, C.W., Stone, W.S., Glatt, S.J., Meyer, E., Faraone, S.V., Tsuang, M.T., Cornblatt, B., 2006. Neuropsychological functioning in adolescents and young adults at genetic risk for schizophrenia and affective psychoses: results from the Harvard and Hillside Adolescent High Risk Studies. *Schizophr. Bull.* 32 (3), 507–524.
- Sitskoorn, M.M., Aleman, A., Ebisch, S.J., Appels, M.C., Kahn, R.S., 2004. Cognitive deficits in relatives of patients with schizophrenia: a meta-analysis. *Schizophr. Res.* 71 (2–3), 285–295.
- Snitz, B.E., Macdonald III, A.W., Carter, C.S., 2006. Cognitive deficits in unaffected first-degree relatives of schizophrenia patients: a meta-analytic review of putative endophenotypes. *Schizophr. Bull.* 32 (1), 179–194.
- Stone, W.S., Braff, D.L., Calkins, M.E., Green, M.F., Greenwood, T.A., Gur, R.E., Gur, R.C., Lazzaroni, L.C., Light, G.A., Nuechterlein, K.H., Radant, A.D., Siever, L.J., Silverman, J.M., Sprock, J., Sugar, C.A., Swerdlow, N.R., Tsuang, D.W., Tsuang, M.T., Turetsky, B.I., Seidman, L.J., 2015. California Verbal Learning Test-II performance in schizophrenia as a function of ascertainment strategy: Comparing the first and second phases of the Consortium on the Genetics of Schizophrenia (COGS). *Schizophr. Res.* 163 (1–3), 32–37 (in this issue).
- Swerdlow, N.R., Light, G.A., Sprock, J., Calkins, M.E., Green, M.F., Greenwood, T.A., Gur, R.E., Gur, R.C., Lazzaroni, L.C., Nuechterlein, K.H., Radant, A.D., Ray, A., Seidman, L.J., Siever, L.J., Silverman, J.M., Stone, W.S., Sugar, C.A., Tsuang, D.W., Tsuang, M.T., Turetsky, B.I., Braff, D.L., 2014. Deficient prepulse inhibition in schizophrenia detected by the multi-site Consortium on the Genetics in Schizophrenia. *Schizophr. Res.* (152), 503–512.
- Swerdlow, N.R., Gur, R.E., Braff, D.L., 2015. Consortium on the Genetics of Schizophrenia (COGS) assessment of endophenotypes for schizophrenia: an introduction to this special issue of Schizophrenia Research. *Schizophr. Res.* 163 (1–3), 9–16 (in this issue).
- Turetsky, B.I., Dress, E.M., Braff, D.L., Calkins, M.E., Green, M.F., Greenwood, T.A., Gur, R.E., Gur, R.C., Lazzaroni, L.C., Nuechterlein, K.H., Radant, A.D., Seidman, L.J., Siever, L.J., Silverman, J.M., Sprock, J., Stone, W.S., Sugar, C.A., Swerdlow, N.R., Tsuang, D.W., Ming, T., Tsuang, M.T., Light, G.A., 2015. The utility of P300 as a schizophrenia endophenotype and predictive biomarker: Clinical and socio-demographic modulators in COGS-2. *Schizophr. Res.* 163 (1–3), 53–62 (in this issue).
- Wilkinson, G.S., Robertson, G.J., 2006. Psychological Assessment Resources. In: Lutz, F.L. (Ed.), *Wide Range Achievement Test, Fourth edition. WRAT4 Introductory Kit*.