UC San Diego

Research Summaries

Title

Combating Arundo Donax and Other Nuisance Grasses

Permalink

https://escholarship.org/uc/item/1q65352d

Author

Wijte, Antonia

Publication Date

2002



COASTAL OCEAN RESEARCH

R/CZ-151: 3.1.1998–2.28.2001 and R/CZ-163: 10.1.1999–9.30.2002 Combating *Arundo Donax* and Other Nuisance Grasses

Antonia Wijte

California State University, Long Beach

Background

rundo donax is a canelike, towering weed that
has gained notoriety for its
hearty resistance to chemical
spraying and remarkable ability to
become the dominant plant in lowlying areas such as the Santa Ana,
Santa Margarita, Santa Clara, and
Tijuana river banks.

By displacing native plants, Arundo and other invasive grasses, such as Spartina alterniflora, not only dramatically alter wild landscapes but also displace animal life. The federally protected least Bell's vireo, for instance, cannot nest in Arundo.

The Project

Dr. Antonia Wijte of California State University at Long Beach has conducted studies that show *Arundo* is sensitive to the timing of when herbicides such as Rodeo are applied. To understand why, it helps to know a little bit about how the plant stores energy and where. Normally, the plant's vascular system stores food throughout the plant—in its leaves, stems and roots—until the plant runs out of nitrogen, a nutrient needed to make nucleic acids, proteins, and enzymes. When this happens, the plant no longer has the nitrogen it needs for cell division and so begins to transport food to root-like structures called rhizomes. Since Arundo will not die unless its rhizomes are killed, the ideal time to apply herbicide occurs when the plant's nitrogen levels are depleted and food is being transported into the rhizomes.

Application

A goal of this project was to

examine whether it is possible to measure nitrogen and carbon levels in Arundo leaves and then to use these measurements to optimize the timing of herbicide application. Dr. Wijte is conducting similar experiments with two other invasive rhizomatous grasses, Spartina alterniflora and **Phragmites** australis.



Graduate student Christiana Chen of Cal State Long Beach sprays herbicide on *Arundo* in the Santa Margarita River in Camp Pendleton. Sea Grant research has shown that the optimal time to apply herbicide occurs when *Arundo*'s vascular system is transporting food into its root-like structures. Photo: Antonia Wijte.



Dr. Antonia Wijte cuts stands of *Arundo*. Photo: California State Long Beach.

At present, she has funding for an outreach program in which resource managers are encouraged to send leaf samples to her lab for nitrogen and carbon analysis. Based on this analysis, she will recommend whether and when to apply herbicide. Instructions on how to take leaf samples can be found at her website.* Field comparisons have shown that leaf-nutrient analyses significantly improve eradication efforts.

Cooperating Organizations

Camp Pendleton Marine Corps Base
Center for Natural Lands
Management
Mission Resource Conservation
District
San Diego County
San Gabriel Mountains Regional
Conservancy

Presentations

Peck, G.W., and A.H.B.M. Wijte. "Temporal variation in the nitrogen carbon ratio in *Arundo donax* L.: Herbicide effectiveness." 15th Biennial International Conference of the Estuarine Research Federation, New Orleans, Louisiana, 1999. Poster presentation.

Motamed, E.R., and A.H.B.M. Wijte. "Optimizing the timing of mechanical eradication of *Arundo donax* L," 15th Biennial International Conference of the Estuarine Research Federation, New Orleans, Louisiana, 1999. Poster presentation.

Trainees

Peck, George W., M.S. in Biology, California State University, Long Beach, 1999.

Chen, Christiana, M.S. in Biology, California State University, Long Beach, anticipated 2002. For more information:

Dr. Antonia Wijte
Department of Biological Sciences
California State University, Long Beach

Tel.: (562) 985-4917 Email: wijte@csulb.edu

*Website: www.csulb.edu/~wijte/

wpbl/arundo.html

PUB. NO. CSG-CZ-02-005 JANUARY 2002

California Sea Grant is a statewide, multiuniversity program of marine research, education, and outreach activities, administered by the University of California. Sea Grant-sponsored research contributes to the growing body of knowledge about our coastal and ocean resources and, consequently, to the solution of many marine-related problems facing our society. Through its Marine Extension Program, Sea Grant transfers information and technology developed in research efforts to a wide community of interested parties and actual users of marine information and technology, not only in California but throughout the nation. Sea Grant also supports a broad range of educational programs so that our coastal and ocean resources can be understood and used judiciously by this and future generations.

The national network of Sea Grant programs is a unique partnership of public and private sectors, combining research, education, and technology transfer for public service and dedicated to meeting the changing environmental and economic needs in our coastal, ocean, and Great Lakes regions.

This work is sponsored in part by a grant from the National Sea Grant College Program, National Oceanic and Atmospheric Administration, U.S. Department of Commerce, under grant number NA06RG0142, Project number A/P-1. The views expressed herein are those of the author and do not necessarily reflect the views of NOAA or any of its sub-agencies. The U.S. Government is authorized to reproduce and distribute for governmental purposes.

Phone: (858) 534-4440 Fax: (858) 453-2948 Web site: http://www-csgc.ucsd.edu