

# Lawrence Berkeley National Laboratory

## LBL Publications

### Title

NEGATIVE PION PRODUCTION IN RELATIVISTIC HEAVY ION COLLISIONS

### Permalink

<https://escholarship.org/uc/item/1qf5g8g6>

### Author

Fung, S.Y.

### Publication Date

1978

NEGATIVE PION PRODUCTION IN  
RELATIVISTIC HEAVY ION COLLISIONS

S. Y. Fung, W. Gorn, G. P. Kiernan, F. F. Liu,  
J. J. Lu, Y. T. Oh, J. Ozawa, R. T. Poe,  
L. Schroeder and H. Steiner

January 1978

RECEIVED  
LAWRENCE  
BERKELEY LABORATORY


MAR 14 1978

LIBRARY AND  
DOCUMENTS SECTION

Prepared for the U. S. Department of Energy  
under Contract W-7405-ENG-48

**TWO-WEEK LOAN COPY**

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 5716*



## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

NEGATIVE PION PRODUCTION IN  
RELATIVISTIC HEAVY ION COLLISIONS

S. Y. Fung, W. Gorn, G. P. Kiernan, F. F. Liu<sup>†</sup>, J. J. Lu,  
Y. T. Oh<sup>‡</sup>, J. Ozawa<sup>§</sup>, R. T. Poe

Department of Physics  
University of California, Riverside, California 92521

and

L. Schroeder and H. Steiner

Lawrence Berkeley Laboratory  
University of California  
Berkeley, California 94720

ABSTRACT

Negative pion production in relativistic heavy ion collisions has been studied in a triggered streamer chamber experiment with projectile nuclei  $^{12}\text{C}$  and  $^{40}\text{Ar}$  in the energy range 0.4 GeV/N - 2.1 GeV/N incident on targets LiH, NaF, BaI<sub>2</sub>, and Pb<sub>3</sub>O<sub>4</sub>. Negative pion yields, multiplicity distributions, and angular and momentum distributions are reported.

- 
- <sup>†</sup> Permanent Address: Physics Dept., California State College,  
San Bernardino, CA 92407
- <sup>‡</sup> Present Address: Physics Dept., University of Hawaii,  
Hilo, HI 96720
- <sup>§</sup> Present Address: Lawrence Livermore Laboratory,  
University of California, Livermore,  
CA 94550

In the study of relativistic heavy ion collisions, much interest has been focused on the production of pions. There are theoretical speculations<sup>(1-5)</sup> about the occurrence of exotic phenomena such as nuclear collective effects, shock waves, pion condensates, or quark matter production which may involve pion production significantly greater than what is expected from an aggregate of individual nucleon-nucleon collisions. The study of pion production in heavy ion collisions may also shed light on less exotic but equally important questions such as the multiple scattering and thermalization processes in nuclear matter.

We report the results of an experimental study of negative pion production using the LBL streamer chamber. The streamer chamber is particularly well suited for this kind of experiment because it can be selectively triggered, it has high efficiency for multitrack events over a  $4\pi$  solid angle, and it can yield accurate angle and momentum information on all charged tracks. The chamber, which has a volume of 127 cm x 61 cm x 40.6 cm, was operated in a magnetic field of  $\sim 14$  kgauss. The targets were located inside the chamber. They were typically quite thin (0.32 cm) in order to minimize effects due to multiple scattering and secondary interactions.

We present here results from twenty sets of runs with  $^{40}\text{Ar}$  beams at 0.4 GeV/N, 0.9 GeV/N, and 1.8 GeV/N, and  $^{12}\text{C}$  beams at 0.4 GeV/N and 2.1 GeV/N, incident on targets LiH, NaF, BaI<sub>2</sub>, and Pb<sub>3</sub>O<sub>4</sub>. These targets, rather than pure elements, were

chosen because they were compatible with the electrical operation of the chamber. Each run consists of  $\sim 2000$  pictures where the chamber was triggered in the "inelastic mode", a trigger setting which rejects most non-interacting events but is relatively unbiased for inelastic events. This triggering mode, which will be described in more detail elsewhere, is based on pulse height in a counter inside the streamer chamber immediately downstream of the production target. It subtended an angle of  $73.8$  msr at the target. Whenever it had a pulse height smaller than that produced by a beam particle, we triggered the chamber. We estimate that this trigger condition causes us to lose no more than 10-15% of the inelastic interactions. These lost events tend to leave the projectile almost intact and are most likely due to peripheral processes which involve small pion multiplicities. Consequently, our partial cross sections for the production of zero, one, and two pions are probably just slightly low.

Scanning for negative tracks yields predominantly negative pions. Contamination consists of electrons mainly from pair production. Most of these electrons are detected during scanning and the final overall electron contamination is estimated to be less than 1%. Measurement of the negative pion tracks is carried out on standard Micrometric scanner-digitizer tables and the subsequent geometric reconstruction is performed using a modified version of TVGP. The use of thin targets enables us

to extend the low momentum cutoff of the observed pion spectrum to around 50 MeV/c.

We present, in Table I, the average negative pion production  $\langle N_{\pi^-} \rangle$  per (inelastic) interaction for the twenty runs. Also shown is the ratio of average negative pion production  $\langle N_{\pi^-} \rangle$  to average number of charged fragments  $\langle N \rangle$ . We observe that this ratio rises sharply as the incident energy increases, but that the ratio is roughly independent of the projectile or the target. A recent emulsion study by McNulty et al.<sup>6</sup> reported the observation of copious pion production for heavy ion collisions in the energy range of 275 MeV/N and below. Our far lower  $\pi^-$  production observed at higher energy (400 MeV/N) is difficult to reconcile with their result.

Negative pion multiplicity distributions for four high energy runs are presented in Figure 1:  $^{40}\text{Ar}$  incident on LiH and  $\text{Pb}_3\text{O}_4$  targets at 1.8 GeV/N in Fig. 1a, and  $^{12}\text{C}$  incident on the same targets at 2.1 GeV/N in Fig. 1b. In Figures 2a and b we present two-dimensional scatter plots in the momentum- $\cos\theta$  plane and their projections: histograms of the transverse and total momenta as well as angular distributions for the produced negative pions.

We note the following qualitative features in our data: (1) the  $p_{\perp}$  distribution of the produced pions is independent of pion multiplicity; (2) the momentum (or energy) distribution is independent of pion multiplicity; (3) the pion multiplicity tends to be proportional to the total multiplicity of all charged particles

with a proportionality constant which depends on bombarding energy. This constant is the same as that given in Table 1 for the ratio  $\langle N_{\pi^-} \rangle / \langle N \rangle$ ; (4) when the data in Figure 1 are replotted in terms of the variables  $\langle N_{\pi^-} \rangle \frac{\sigma_{N_{\pi^-}}}{\sigma_{inel}}$  vs.  $N_{\pi^-} / \langle N_{\pi^-} \rangle$ , the observed distributions are quite similar for all target/projectile combinations at all but the lowest bombarding energies (see Fig. 3). (5) The pion momentum distributions observed in this experiment are consistent with the inclusive measurements of Nagamiya, et al. (7) in those kinematical domains where comparisons are possible. (6) The recently reported pion multiplicity distributions of Jakobsson, et al. (8) do not agree very well with the results reported here. (7) It is difficult to make meaningful comparisons between our data and those of the recent higher energy experiments involving pion production by 18 GeV alpha particles on various nuclear targets reported by Baldin. (9) Our multiplicity distributions have more low multiplicity events than are observed in the Dubna experiments when the average multiplicities are the same. However, in our experiment the energies are sufficiently low that kinematic threshold effects may be important. The dependence of our multiplicity distributions on the atomic numbers of projectile and target does not seem to have any of the simple forms suggested by Bialas, et al. (10) using the concept of "wounded" nucleons. Here again kinematic and reabsorption effects may be important.

Theoretical attempts to explain pion production in



nucleus-nucleus collisions have just started to be made; e.g. see the following papers by Vary<sup>(11)</sup>, and Kauffmann and Gyulassy<sup>(12)</sup>. It is our hope that the experimental data presented here will stimulate further theoretical activity and will provide useful new information for the development of meaningful theoretical models.

We wish to express our gratitude to the Bevalac staff, and in particular to James P. Brannigan for his untiring efforts to make the LBL streamer chamber a reliable facility. We are grateful to the scanning staff at Riverside. This work was supported by the U.S. Department of Energy.

## References

1. G. Chapline, M. Johnson, E. Teller and M. Weiss, Phys. Rev. D 8, 4302 (1973).
2. W. Scheid, H. Muller, and W. Greiner, Phys. Rev. Lett. 32, 741 (1974).
3. M. Sobel, P. J. Siemens, J. P. Bondorf, and H. A. Bethe, Nuclear Physics A251, 502 (1975).
4. Y. Kitazoe, M. Sano, and H. Toki, Nuovo Cimento Letters 13, 139 (1975).
5. G. Chapline and A. Kerman, to be published.
6. P. J. McNulty, G. E. Farrell, R. C. Filz, W. Schimmerling, and K. G. Vosburgh, Phys. Rev. Lett. 38, 1519 (1977).
7. S. Nagamiya, I. Tanihata, S. Schnetzer, L. Anderson, W. Brückner, O. Chamberlain, G. Shapiro, and H. Steiner, "Proton and Pion Spectra at Large Angles in Relativistic Heavy-Ion Collisions", LBL-6770.
8. B. Jakobsson, R. Kullberg, I. Otterlund, A. Ruiz, J. M. Bolta, and E. Higon, "A Search for Collective Pion Production in 2 GeV/N Heavy Ion Reactions", LUNFD6(NFFK-7007).
9. A. M. Baldin, Contributed paper at the VIIth International Conference on High-Energy Physics and Nuclear Structure, Zurich (1977).
10. A. Bialas, M. Bleszynski and W. Czyz, Nucl. Phys. B111, 461 (1976).

11. J.P. Vary, "Multiple Collision Model for Pion Production in Relativistic Nucleus-Nucleus Collisions", submitted to Phys. Rev. Lett.
12. S. K. Kauffmann and M. Gyulassy, "Pion Multiplicities in Heavy Ion Collisions", submitted to Phys. Rev. Lett.

Figure Captions.

Table 1. Average number of negative pions produced per (inelastic) interaction,  $\langle N_{\pi^-} \rangle$ , and the ratio of average number of negative pions produced to the average number of charged fragments,  $\langle N \rangle$ , for 20 beam-target combinations.

Figure 1. Negative pion multiplicity for (a) 1.8 GeV/N  $^{40}\text{Ar}$  and (b) 2.1 GeV/N  $^{12}\text{C}$  beams incident on LiH and  $\text{Pb}_3\text{O}_4$ .

Figure 2. Momentum and angular distributions of negative pions in the laboratory system for (a) 1.8 GeV/N  $^{40}\text{Ar}$  and (b) 2.1 GeV/N  $^{12}\text{C}$  incident on  $\text{Pb}_3\text{O}_4$ :  $\cos\theta$  vs. total momentum with projections, together with transverse momentum.

Figure 3. Plot of  $\langle n_{\pi^-} \rangle \sigma_{\pi^-} / \sigma_{\text{inel}} N_{\pi^-}$  vs.  $N_{\pi^-} / \langle n_{\pi^-} \rangle$  on multiplicity distribution of negative pions in 20 beam-target combinations.

Table I

Projectile	Projectile Energy (GeV/N)	TARGET							
		LiH		NaF		BaI <sub>2</sub>		Pb <sub>3</sub> O <sub>4</sub>	
		$\langle N_{\pi^-} \rangle / \text{int.}$	$\langle N_{\pi^-} \rangle / \langle N \rangle$	$\langle N_{\pi^-} \rangle / \text{int.}$	$\langle N_{\pi^-} \rangle / \langle N \rangle$	$\langle N_{\pi^-} \rangle / \text{int.}$	$\langle N_{\pi^-} \rangle / \langle N \rangle$	$\langle N_{\pi^-} \rangle / \text{int.}$	$\langle N_{\pi^-} \rangle / \langle N \rangle$
<sup>40</sup> Ar	0.4	0.043 ± 0.022	0.006 ± 0.003	0.034 ± 0.015	0.003 ± 0.001	0.107 ± 0.019	0.007 ± 0.001	0.099 ± 0.020	0.007 ± 0.001
	0.9	0.31 ± 0.04	0.033 ± 0.004	0.60 ± 0.09	0.044 ± 0.005	0.87 ± 0.09	0.045 ± 0.003	0.92 ± 0.88	0.045 ± 0.003
	1.8	0.97 ± 0.05	0.08 ± 0.003	1.91 ± 0.12	0.104 ± 0.004	3.27 ± 0.18	0.106 ± 0.003	3.27 ± 0.15	0.104 ± 0.002
<sup>12</sup> C	0.4	0.02 ± 0.01	0.004 ± 0.002	0.038 ± 0.013	0.006 ± 0.002	0.078 ± 0.014	0.010 ± 0.002	0.066 ± 0.014	0.009 ± 0.002
	2.1	0.68 ± 0.07	0.095 ± 0.008	1.03 ± 0.08	0.108 ± 0.006	1.91 ± 0.17	0.110 ± 0.006	1.79 ± 0.16	0.101 ± 0.006









FIGURE 1



1.8 GeV/N <sup>40</sup>Ar on Pb<sub>3</sub>O<sub>4</sub>



2.1 GeV/N <sup>12</sup>C on Pb<sub>3</sub>O<sub>4</sub>

FIGURE 2




Fig. 3



This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

TECHNICAL INFORMATION DEPARTMENT  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720