

# UC Davis

## UC Davis Previously Published Works

### Title

A quantitative Doignon-Bell-Scarf theorem

### Permalink

<https://escholarship.org/uc/item/1rk725c0>

### Journal

Combinatorica, 37(3)

### ISSN

0209-9683

### Authors

Aliev, Iskander  
Bassett, Robert  
De Loera, Jesús A  
et al.

### Publication Date

2017-06-01

### DOI

10.1007/s00493-015-3266-9

Peer reviewed

# A Quantitative Doignon-Bell-Scarf Theorem

Iskander Aliev, Robert Bassett, Jesús A. De Loera, and Quentin Louveaux

<sup>1</sup> Cardiff University, UK,

AlievI@cardiff.ac.uk,

<sup>2</sup> University of California, Davis

rbassett@math.ucdavis.edu

deloera@math.ucdavis.edu

<sup>3</sup> Université de Liège, Belgium

q.louveaux@ulg.ac.be

**Abstract.** The famous Doignon-Bell-Scarf theorem is a Helly-type result about the existence of integer solutions to systems of linear inequalities. The purpose of this paper is to present the following quantitative generalization: Given an integer  $k$ , we prove that there exists a constant  $c(n, k)$ , depending only on the dimension  $n$  and  $k$ , such that if a bounded polyhedron  $\{x \in \mathbb{R}^n : Ax \leq b\}$  contains exactly  $k$  integer points, then there exists a subset of the rows, of cardinality no more than  $c(n, k)$ , defining a polyhedron that contains exactly the same  $k$  integer points. In this case  $c(n, 0) = 2^n$  as in the original case of Doignon-Bell-Scarf for infeasible systems of inequalities. We work on both upper and lower bounds for the constant  $c(n, k)$  and discuss some consequences, including a Clarkson-style algorithm to find the  $l$ -th best solution of an integer program with respect to the ordering induced by the objective function.

## 1 Introduction

In a *Helly-type theorem*, there is a family of objects  $F$ , a property  $P$  and a constant  $\mu$  such that if every subfamily of  $F$  with  $\mu$  elements has property  $P$ , then the entire family has property  $P$ . This topic expands a large literature, we recommend [7, 8, 14, 18, 29] and the references there for a glimpse of this fertile subject. The classical theorem of Helly deals with the case when  $F$  is a finite family of convex sets in  $\mathbb{R}^n$ , the constant  $\mu$  is  $n + 1$ , and the property  $P$  is having a non-empty intersection. One of the most famous Helly-type theorems, due to its many applications in the theory of integer programming and computational geometry of numbers, is the 1973 theorem of Doignon [17], later reproved by Bell and Scarf [10, 26]. For an arbitrary matrix  $A \in \mathbb{R}^{m \times n}$  and a set  $S \subseteq \{1, \dots, m\}$ , we denote  $A_S \in \mathbb{R}^{|S| \times n}$  as the submatrix of  $A$  with row-indices in  $S$ .

**Theorem** *Let  $A$  be a  $m \times n$  real matrix and  $b$  a vector in  $\mathbb{R}^m$ . If the set of integer points  $\{x \in \mathbb{Z}^n : Ax \leq b\}$  is empty, then there is a subset  $S$  of the rows of  $A$ , of cardinality no more than  $2^n$ , with the property that the set  $\{x \in \mathbb{Z}^n : A_S x \leq b_S\}$  is also empty.*

It should be noted that still during the 1970's, in [22] Hoffman created an abstract framework that contains Helly's original theorem as well as the Doignon-Bell-Scarf results. Here our key contribution is to prove a weighted or quantitative generalization of Doignon-Bell-Scarf's theorem, one that is close in spirit to the quantitative versions of Helly's theorem of Bárány, Katchalski and Pach [7, 8]:

**Theorem 1.** *Given  $n, k$  two non-negative integers there exists a universal constant  $c(n, k)$ , depending only on  $n$  and  $k$ , such that for any bounded polyhedron  $P = \{x \in \mathbb{R}^n : Ax \leq b\}$  with exactly  $k$  integer points, there is a subset  $S$  of the rows of  $A$ , of cardinality no more than  $c(n, k)$ ,*

with the property that the polyhedron  $\{x \in \mathbb{R}^n : A_S x \leq b_S\}$  has exactly the same  $k$  integer points as  $P$ .

We refer to  $c(n, k)$  as the minimal such constant. The upper bound we prove in this paper is  $c(n, k) \leq \lceil 2(k+1)/3 \rceil 2^n - 2 \lfloor 2(k+1)/3 \rfloor + 2$ .

Our proof of this theorem (presented in the first section of this paper) generalizes the proof of the Doignon-Bell-Scarf theorem given in [10, 27] with some counting twists. Our new bound in Theorem 1 is a considerable improvement on the bound we presented in [1] (where the existence of the universal constant  $c(n, k)$  was first announced). The new proof is presented in Section 2.

We were also able to obtain lower bounds for the case  $k = 1$  proving that our upper bound is in fact the exact value of  $c(n, 1)$ . This is the content of the following theorem. An explicit construction of the polytope in question will be presented in Section 3 of our paper:

**Theorem 2.** *There exists a polytope  $P$  in  $\mathbb{R}^n$  that has exactly one interior integer point,  $2(2^n - 1)$  facets and one integer point in the relative interior of each facet. Thus all inequalities in  $P$  are necessary in the sense that the removal of any inequality from  $P$  results in the inclusion of at least one additional integer point in the interior of  $P$ . As a consequence, the upper bound of Theorem 1 is tight for  $k = 1$  and thus  $c(n, 1) = 2(2^n - 1)$ .*

This implies that the upper bound given in Theorem 1 for  $k = 1$  is tight, i.e.  $c(n, 1) = 2(2^n - 1)$ . Also observe that the upper bounds for  $c(n, 2)$  and  $c(n, 1)$  are equal in Theorem 1. It is an interesting problem to find exact values of  $c(n, k)$  for  $k \geq 2$ . Given the available data, a natural conjecture is that our bound for  $c(n, 2)$  is also tight, but we also know that for  $c(2, k)$  the bound is not tight for  $k \geq 3$ .

Finally, in Section 4 of the paper we discuss some consequences of Theorem 1. Using standard procedures we extend the version for half-spaces to families of convex sets and present some quantitative corollaries of Theorem 1. K. Clarkson was the first to notice that using Doignon-Bell-Scarf's theorem (or any similar Helly-type intersection result) one can obtain a probabilistic algorithm for integer linear programming [12]. Clarkson's probabilistic algorithm works in great generality for so called *violator spaces* [20]. The remarkable fact is that the running time for the algorithm is *linear* in the number of constraints defining the problem, and *subexponential* in the dimension of the problem. In this way, when the dimension of the problem is small, then these are the most efficient methods to solve a large family of optimization problems. Using violator spaces we present an application of our quantitative Doignon-Bell-Scarf theorem to the problem of finding the  $l$  best integer feasible points of an integer program.

We conclude with the remark that our results are part of the fertile and classical study of convex polyhedra with fixed number of (interior) lattice points; a topic that plays an important role in algebra, discrete geometry, and optimization. Indeed, there has been a lot of work, going back to classical results of Minkowski and van der Corput, to show that the volume of a lattice polytope  $P$  with  $k = \text{card}(\text{int } P \cap \mathbb{Z}^n) \geq 1$  is bounded above by a constant that only depends on  $n$  and  $k$  (see e.g., [23, 25]). Similarly, the supremum of the possible number of points of  $\mathbb{Z}^n$  in a lattice polytope in  $\mathbb{R}^n$  containing precisely  $k$  points of  $\mathbb{Z}^n$  in its interior, can be bounded by a constant that only depends on  $n$  and  $k$ . Such results play an important role in the theory of toric varieties and the structure of lattice polyhedra (see e.g., [9] and the references therein). In optimization it was recently shown in [5, 6] how lattice-free polyhedra can be applied to the generation of cutting planes (see also [13]). One can think of lattice-free polyhedra as dual certificates of the infeasibility of a system of the form  $\{x \in \mathbb{Z}^n : Ax = b, Cx \leq d\}$  [4]. These convex bodies play an important role in the modern theory of cutting planes [11, 16]. There has been some interesting hybrid variations of Helly and Doignon-Bell-Scarf in [3, 22].

## 2 A Generalization of Doignon-Bell-Scarf's theorem

In this section we will prove Theorem 1. The constant  $c(n, k)$  we provide is  $\lceil 2(k+1)/3 \rceil 2^n - 2\lceil 2(k+1)/3 \rceil + 2$ . This is an improvement from the constant presented in an earlier conference version of this paper [1].

[*Proof of Theorem 1*]

The proof proceeds by contradiction. We choose a counterexample system with minimal number  $m$  of linear inequalities,

$$a_1x \leq \beta_1, \dots, a_mx \leq \beta_m, x \in \mathbb{R}^n. \quad (1)$$

This counterexample system (1) defines a polytope with exactly  $k$  integer solutions and  $m \geq \lceil 2(k+1)/3 \rceil 2^n - 2\lceil 2(k+1)/3 \rceil + 3$ . Again, remember that  $m$  is smallest possible among all counterexamples to Theorem 1. The minimality of the chosen counterexample implies that if we delete any of the constraints in (1), the remaining system has at least  $k+1$  integer solutions (one is not able to remove any of the constraints otherwise that would produce a smaller counterexample). This says that there exist integer vectors  $x_1, \dots, x_m$  such that  $x_j$  violates  $a_jx \leq \beta_j$ , but satisfies all other inequalities in (1). Consider the polyhedron  $Q = \{x \in \mathbb{R}^n : a_1x \leq a_1x_1, \dots, a_mx \leq a_mx_m\}$ . The polyhedron  $Q$  is bounded. To see this, it is enough to notice that the recession cone of a polyhedron  $\{x \in \mathbb{R}^n : A'x \leq b'\}$  is independent on the right hand side vector  $b'$  (see e.g. Section 8.2 in [27]). Since  $Q$  has the same normal vectors as the original system (1) which was bounded then so is  $Q$ . Consider the set of integer points

$$G = Q \cap \mathbb{Z}^n.$$

Since the polyhedron  $Q$  is bounded, the set  $G$  is finite. Now we will associate a new system to (1) in a special critical shape constructed by shifting its hyperplanes. Consider the set  $\Gamma \subset \mathbb{R}^m$  of the vectors  $(\gamma_1, \dots, \gamma_m)$  such that

$$\gamma_j \geq \min\{a_jz : z \in G, a_jz > \beta_j\} \quad (2)$$

and

$$\text{the system } a_1x < \gamma_1, \dots, a_mx < \gamma_m \text{ has exactly } k \text{ integer solutions in } G. \quad (3)$$

The set  $\Gamma$  is nonempty as we can take the equality in (2). Next notice that (3), together with the lower bounds on the  $\gamma_i$ , implies that any integer solution of the system (1) remains feasible for the system  $a_1x < \gamma_1, \dots, a_mx < \gamma_m$  for  $\gamma \in \Gamma$ . Thus, for all  $\gamma \in \Gamma$ ,  $a_1x < \gamma_1, \dots, a_mx < \gamma_m$  share exactly the same  $k$  integer solutions as (1).

Observe also that the set  $\Gamma$  is bounded. Otherwise,  $\gamma_j$  for some  $j$  grows arbitrarily large. In particular  $\gamma_j$  is much larger than  $a_jx_j$ , for the point  $x_j \in G$  we associated with the constraint  $a_jx \leq \beta_j$  in (1). Note  $x_j$  is by construction not a solution of system (3). But by the size of  $\gamma_j$ , the point  $x_j$  satisfies all the inequalities  $a_1x < \gamma_1, \dots, a_mx < \gamma_m$ . Hence,  $x_j$  is an additional integer feasible point for (3), which is a contradiction.

*Claim.* There is a point  $(\nu_1, \dots, \nu_m) \in \Gamma$  such that

$$\text{for each } j = 1, \dots, m \text{ there exists } y_j \in G \text{ so that } a_jy_j = \nu_j \text{ and } a_iy_j < \nu_i \text{ (} i \neq j \text{)}. \quad (4)$$

Before a formal proof of the Claim there is an intuitive justification for it. Since the set  $G$  is finite, we can define the numbers  $\nu_1, \dots, \nu_m$  satisfying the Claim by changing the right-hand side of the inequalities one by one (i.e., shifting the hyperplanes), while controlling each time how many integer points are within the new polyhedron.

*Proof of Claim:* Take any point  $(\nu_1, \dots, \nu_m) \in \Gamma$  and suppose that for some  $j$  this property does not hold. Consider

$$\nu'_j = \sup\{\nu : (\nu_1, \dots, \nu_{j-1}, \nu, \nu_{j+1}, \dots, \nu_m) \in \Gamma\}. \quad (5)$$

The supremum in (5) is finite as the set  $\Gamma$  is bounded. If  $(\nu_1, \dots, \nu_{j-1}, \nu'_j, \nu_{j+1}, \dots, \nu_m) \notin \Gamma$  then, by (3), for any  $\delta > 0$  there should exist a point  $z \in G$  such that  $\nu'_j - \delta \leq a_j z < \nu'_j = a_j y_j$ . This is impossible as  $G$  is finite. Consequently,  $(\nu_1, \dots, \nu_{j-1}, \nu'_j, \nu_{j+1}, \dots, \nu_m) \in \Gamma$ . Observe that there should exist  $y_j \in G$  with  $a_j y_j = \nu'_j$  and  $a_i y_j < \nu'_i (i \neq j)$ . Otherwise  $(\nu_1, \dots, \nu_{j-1}, \nu'_j + \epsilon, \nu_{j+1}, \dots, \nu_m) \in \Gamma$  for some  $\epsilon > 0$  as  $G$  is a finite set. Hence we can replace  $\nu_j$  by  $\nu'_j$ . After at most  $m$  such replacements we will construct a point satisfying (4).  $\square$

By the established Claim, the convex hull  $H = \text{conv}(\{y_1, \dots, y_m\})$  contains at most  $k$  integer points distinct from  $y_1, \dots, y_m$ .

The property of the set  $\{y_1, \dots, y_m\}$  expressed by (4) is very important and as we will use it several times later, we formally name it.

**Definition 1.** Let  $X$  be a finite set in  $\mathbb{Z}^n$ . We say that  $X$  satisfies the support hyperplane property if for every  $y \in X$ , there exists a half-space  $f^T x \leq g$  such that  $f^T y = g$  and  $f^T z < g$  for every  $z \in X, z \neq y$ . Furthermore, we say that the inequality  $f^T x \leq g$  fulfills the support hyperplane property for  $y$ .

Observe that the support hyperplane property is equivalent to saying that all members of  $X$  are vertices of  $\text{conv}(X)$ .

The application of the following four lemmas directly gives the desired contradiction and finishes the proof of the Theorem:

**Lemma 1.** Consider a finite set  $X \subset \mathbb{Z}^n$  with  $|X| \geq 2^n + 1$  that satisfies the support hyperplane property. Then there exists a point  $z \in (\text{conv}(X) \cap \mathbb{Z}^n) \setminus X$ .

*Proof of Lemma:* Since  $|X| \geq 2^n + 1$ , by the pigeonhole principle there exist  $y_1, y_2 \in X$  with  $y_1 \neq y_2$  and  $y_1 \equiv y_2 \pmod{2}$ . Therefore  $z = \frac{1}{2}(y_1 + y_2) \in \mathbb{Z}^n$ . Since  $X$  satisfies the support hyperplane property, we conclude that  $z \notin X$ .

**Lemma 2.** Consider a finite set  $X \subset \mathbb{Z}^n$  that satisfies the support hyperplane property. Suppose there exists a point  $z \in (\text{conv}(X) \cap \mathbb{Z}^n) \setminus X$ . Then there exist disjoint subsets  $X_1, X_2 \subseteq X$  with  $X_1 \cup X_2 = X$  such that both  $X_1 \cup \{z\}$  and  $X_2 \cup \{z\}$  satisfy the support hyperplane property.

*Proof of Lemma:* There exists a hyperplane defined by the equation  $\bar{f}^T x = \bar{g}$  such that  $\bar{f}^T z = \bar{g}$  and the equality does not hold for any other member of  $X$ . We can split the other members of  $X$  into two disjoint sets  $X_1 = X \cap \{x \in \mathbb{R}^n : \bar{f}^T x < \bar{g}\}$  and  $X_2 = X \cap \{x \in \mathbb{R}^n : \bar{f}^T x > \bar{g}\}$ .

The result follows since for every  $x \in X_i$ , the inequality fulfilling the support hyperplane property for  $x$  in  $X$  fulfills the support hyperplane property for  $x$  in  $X_i \cup \{z\}$ . The inequality  $\bar{f}^T x \leq \bar{g}$  (respectively  $\bar{f}^T x \geq \bar{g}$ ) fulfills the support hyperplane property for  $z$  in  $X_1 \cup \{z\}$  (respectively in  $X_2 \cup \{z\}$ ).  $\square$

**Lemma 3.** Consider a finite set  $X \subset \mathbb{Z}^n$  that satisfies the support hyperplane property. Suppose  $z_1, z_2 \in \text{conv}(X) \cap \mathbb{Z}^n$  with  $z_1, z_2 \notin X$  and  $z_1 \neq z_2$ . There exist disjoint subsets  $X_1, X_2 \subseteq X$  with  $|X_1| + |X_2| \geq |X| - 2$  such that both  $X_1 \cup \{z_1, z_2\}$  and  $X_2 \cup \{z_1, z_2\}$  satisfy the support hyperplane property.

*Proof of Lemma:* Since  $X$  satisfies the support hyperplane property, the line  $L$  spanned by  $z_1, z_2$  intersects  $X$  in at most two points. Consequently, there exists a hyperplane  $\bar{f}^T x = \bar{g}$  such that  $\bar{f}^T z_1 = \bar{f}^T z_2 = \bar{g}$  and the equality holds for at most two points of  $X$  (precisely for the points in  $X \cap L$ ). We can split the points of the set  $Y = X \setminus (X \cap L)$  into two disjoint sets  $X_1 = Y \cap \{x \in \mathbb{R}^n : \bar{f}^T x < \bar{g}\}$  and  $X_2 = Y \cap \{x \in \mathbb{R}^n : \bar{f}^T x > \bar{g}\}$ . Clearly,  $|X_1| + |X_2| \geq |X| - 2$ .

Let us now show that  $X_1 \cup \{z_1, z_2\}$  and  $X_2 \cup \{z_1, z_2\}$  satisfy the support hyperplane property. First, for every  $x \in X_i$ , the inequality fulfilling the support hyperplane property for  $x$  in  $X$  fulfills the support hyperplane property for  $x$  in  $X_i \cup \{z_1, z_2\}$ . Next, both points  $z_1, z_2$  are on the hyperplane  $\bar{f}^T x = \bar{g}$  and this hyperplane does not intersect either of the finite sets  $X_1, X_2$ . Thus applying a sufficiently small rotation of the hyperplane  $\bar{f}^T x = \bar{g}$  around  $z_1$  we can obtain a hyperplane  $\bar{f}_1^T x = \bar{g}_1$  such that  $\bar{f}_1^T z_1 = \bar{g}_1$  and  $\bar{f}_1^T y < \bar{g}_1$  for any point  $y \in X_1 \cup \{z_2\}$ , so that the constructed hyperplane fulfills the support hyperplane property for  $z_1$  in  $X_1 \cup \{z_1, z_2\}$ . In the same manner we can construct the hyperplanes fulfilling the support hyperplane property for  $z_1$  in  $X_2 \cup \{z_1, z_2\}$  and for  $z_2$  in  $X_i \cup \{z_1, z_2\}$ ,  $i = 1, 2$ .  $\square$

**Lemma 4.** *Let  $n \geq 2$  and  $k$  be natural numbers. Consider a finite set  $X \subset \mathbb{Z}^n$  and assume that it satisfies the support hyperplane property. If the cardinality  $|X| \geq k2^n - 2k + 3$ , then there exist at least  $\lfloor 3k/2 \rfloor$  different integer points in  $\text{conv}(X) \setminus X$ .*

*Proof of Lemma:* We will proceed by induction on  $k$ . For  $k = 1$ , the result follows from Lemma 1.

We now assume that the result is true up to  $k - 1$  and prove it for  $k \geq 2$ . Assume that  $|X| \geq k2^n - 2k + 3$ . We have  $|X| \geq 2^n + 1$  and therefore by Lemma 1, there exists a point  $z_1 \in \mathbb{Z}^n$  in  $\text{conv}(X) \setminus X$ .

By Lemma 2, applied to  $X$  and  $z_1$ , there exist disjoint subsets  $Y_1, Y_2 \subseteq X$  with  $Y_1 \cup Y_2 = X$  such that both  $Y_1 \cup \{z_1\}$  and  $Y_2 \cup \{z_1\}$  satisfy the support hyperplane property.

Assume without loss of generality that  $|Y_2| \geq |Y_1|$ . Since the sets  $Y_1, Y_2$  are disjoint,  $Y_2$  has cardinality at least  $\lceil \frac{|X|}{2} \rceil \geq 2^n$  (note that  $k \geq 2$ ). By Lemma 1, applied to  $Y_2 \cup \{z_1\}$ , there exists a point  $z_2 \in \mathbb{Z}^n$  in  $\text{conv}(Y_2 \cup \{z_1\}) \setminus (Y_2 \cup \{z_1\})$ . In particular,  $z_1 \neq z_2$ . Replacing, if necessary,  $z_1, z_2$  by adjacent integer points in  $\text{conv}(\{z_1, z_2\})$ , we may also assume that  $\text{card}(\text{conv}(\{z_1, z_2\}) \cap \mathbb{Z}^n) = 2$ .

Next, by Lemma 3, applied to  $X$  and the constructed points  $z_1, z_2$ , there exist disjoint subsets  $X_1, X_2 \subseteq X$  with  $|X_1| + |X_2| \geq |X| - 2$  such that both  $X_1 \cup \{z_1, z_2\}$  and  $X_2 \cup \{z_1, z_2\}$  satisfy the support hyperplane property. It follows from the proof of Lemma 3 that if one of the sets, say  $X_1$ , is empty, then  $|X_2 \cup \{z_1, z_2\}| = |X|$ . In this case it is enough to prove the lemma for the set  $X$  replaced by the set  $X_2 \cup \{z_1, z_2\}$ . Since the set  $\text{conv}(X) \cap \mathbb{Z}^n$  is finite, after a finite number of replacements we obtain the nonempty sets  $X_1, X_2$ .

Assume now without loss of generality  $0 < |X_1| \leq |X_2|$ . We will consider the following two cases. First, suppose that there exists  $l \in \mathbb{Z}$  with  $1 \leq l \leq k$  such that

$$(l-1)2^n - 2(l-1) + 1 \leq |X_1| \leq l2^n - 2l. \quad (6)$$

Since  $X_1$  and  $X_2$  are disjoint subsets of  $X$  with  $|X_1| + |X_2| \geq |X| - 2$  and since  $|X| \geq k2^n - 2k + 3$ , it follows from the upper bound in (6) that

$$\begin{aligned} |X_2| &\geq k2^n - 2k + 1 - l2^n + 2l \\ &= (k-l)2^n - 2(k-l) + 1. \end{aligned} \quad (7)$$

Thus using (6), we have  $|X_1 \cup \{z_1, z_2\}| \geq (l-1)2^n - 2(l-1) + 3$  which implies from the induction hypothesis (note that  $l \leq k$ ) that there are  $\lfloor 3(l-1)/2 \rfloor$  additional integer points in  $\text{conv}(X_1 \cup \{z_1, z_2\}) \setminus (X_1 \cup \{z_1, z_2\})$ . Using (7), we have  $|X_2 \cup \{z_1, z_2\}| \geq (k-l)2^n - 2(k-l) + 3$  which, from

the induction hypothesis (note that  $l \geq 1$ ) implies that there are  $\lfloor 3(k-l)/2 \rfloor$  additional integer points in  $\text{conv}(X_2 \cup \{z_1, z_2\}) \setminus (X_2 \cup \{z_1, z_2\})$ . The result follows since, counting  $z_1, z_2$ , we have provided  $\lfloor 3(l-1)/2 \rfloor + \lfloor 3(k-l)/2 \rfloor + 2 \geq \lfloor 3k/2 \rfloor$  different integer points in  $\text{conv}(X) \setminus X$ . Suppose now that  $|X_1| \geq k2^n - 2k + 1$ . Then the set  $X_1 \cup \{z_1, z_2\}$  has cardinality at least  $k2^n - 2k + 3$ . Hence, similarly to the previous case, the result follows from the induction hypothesis.  $\square$

It should be emphasized here that Lemma 4 is of independent interest for the theory of lattice points in convex lattice polytopes. It can be restated as follows.

**Corollary 1.** *Let  $n \geq 2$  and  $k$  be natural numbers. Consider a convex lattice polytope  $P$  with the set of vertices  $X \subset \mathbb{Z}^n$ . If the cardinality  $|X| \geq k2^n - 2k + 3$ , then there exist at least  $\lfloor 3k/2 \rfloor$  different lattice points in  $P \setminus X$ .*

If we go back to the proof of Theorem 1, recall that existence of the counterexample system (1) implies the existence of  $m$  integer points  $y_1, \dots, y_m$  satisfying the support hyperplane property, with at most  $k$  other integer points in their convex hull  $H$  and with  $m \geq \lceil 2(k+1)/3 \rceil 2^n - 2\lceil 2(k+1)/3 \rceil + 3$ .

Applying Lemma 4, we conclude that there must be at least

$$\left\lfloor \frac{3}{2} \left\lceil \frac{2(k+1)}{3} \right\rceil \right\rfloor \quad (8)$$

other integer points in  $H$ . Observe that (8) is equal to  $k+1$  if  $k$  is congruent to 0 or 2 modulo 3 and equal to  $k+2$  if  $k$  is congruent to 1 modulo 3. Hence there are at least  $k+1$  other integer points in  $H$  which is the desired contradiction.  $\square$

### 3 Lower Bound Constructions for $k = 1$

In page 235 of [27], an example is given that shows that the bound  $2^n$  given by the Doignon-Bell-Scarf theorem is tight. In this section, we present a construction of a polytope  $P$  that shows that our upper bound for  $k = 1$  from Theorem 1 is tight. This example, together with the verification of its properties, establish Theorem 2.

For notational convenience, given a set of natural numbers  $N$ , let  $l_N := \min_{i \in N} i$  denote its least element. We define the polyhedron as follows

$$P = \{x \in \mathbb{R}^n : \sum_{i=1}^{j-1} \frac{1}{2^i} x_i + x_j + \sum_{i=j+1}^n \frac{1}{2^{i-1}} x_i \leq 1 \quad j = 1, \dots, n, \quad (9)$$

$$- \sum_{i=1}^{j-1} \frac{1}{2^i} x_i - x_j - \sum_{i=j+1}^n \frac{1}{2^{i-1}} x_i \leq 1 \quad j = 1, \dots, n, \quad (10)$$

$$- \frac{1}{|N|} x_{l_N} + \sum_{i \in N, i \neq l_N} \frac{1}{|N|} x_i - \sum_{i \notin N} \frac{1}{|N|^n} x_i \leq 1 \quad \forall N \subseteq \{1, 2, \dots, n\}; |N| \geq 2, \quad (11)$$

$$+ \frac{1}{|N|} x_{l_N} - \sum_{i \in N, i \neq l_N} \frac{1}{|N|} x_i + \sum_{i \notin N} \frac{1}{|N|^n} x_i \leq 1 \quad \forall N \subseteq \{1, 2, \dots, n\}; |N| \geq 2 \quad \}. \quad (12)$$

The rationale behind the construction of the polyhedron  $P$  is the following. First it is constructed in such a way that 0 is the only integer point in its interior. Both of the inequalities (9)-(10) are tight at a unit vector  $\pm e_i$  and exclude some integer points from  $\{-1, 0, 1\}^n$ . All inequalities (11)-(12) are all tight at exactly one of the remaining integer points of  $\{-1, 0, 1\}^n$ .

We will prove Theorem 2 through Lemmas 5 to 7. Lemma 5 proves that the only valid integer points of  $P$  are in  $\{-1, 0, 1\}^n$ . Lemma 6 uses Lemma 5 to provide a necessary and sufficient condition of feasibility of integer points. Lastly, Lemma 7 shows that each inequality defining  $P$  is necessary and contains exactly one tight integer point in its relative interior. Because we have used only rational data the polyhedron is in fact bounded, thus a polytope. This is the case because if unbounded, its recession cone would contain a rational direction which would force infinitely many points inside.

**Lemma 5.** *If  $y \in \mathbb{Z}^n$  has at least an index  $j$  such that  $|y_j| \geq 2$ , then  $y \notin P$*

*Proof.* Consider  $y \in P \cap \mathbb{Z}^n$ . Assume by contradiction that  $k$  is the largest index with  $|y_k| \geq 2$ . We prove the case  $y_k \geq 2$ . The negative case is symmetric and omitted.

Add twice Inequality (9) with  $j = k$  to Inequality (10) with  $j = 1$  which yields

$$\left(2 - \frac{1}{2^{k-1}}\right) y_k + \sum_{i=k+1}^n \frac{1}{2^{i-1}} y_i \leq 3. \quad (13)$$

Since, we have assumed that  $|y_i| \leq 1$  for all indices  $i \geq k+1$ , we can bound

$$-\frac{1}{2^{k-1}} < \sum_{i=k+1}^n \frac{1}{2^{i-1}} y_i < \frac{1}{2^{k-1}}. \quad (14)$$

Using (13) and (14), we conclude that

$$y_k < \frac{3 \cdot 2^{k-1} + 1}{2^k - 1}. \quad (15)$$

For  $k \geq 3$ , this provides a contradiction since the right-hand-side of (15) can be shown to be smaller than 2. For  $k = 1$  or  $k = 2$ , this yields  $|y_k| \leq 2$ . Observe though that if  $k = 1$ , then the inequalities (9) and (10) with  $j = 1$  together with the fact that, using  $|y_i| \leq 1$  for  $i \geq 2$ ,  $|\sum_{i=2}^n \frac{1}{2^{i-1}} y_i| < 1$  yield  $|y_1| \leq 1$ .

To finish the proof, there still remains to consider the case  $k = 2$  i.e.  $y_2 = 2$ . If  $y_1$  is nonnegative,  $y$  violates (9) with  $j = 2$ . If  $y_1$  is negative,  $y$  violates (11) with  $N = \{1, 2\}$ .  $\square$

**Definition 2.** *Given a point  $y \in \mathbb{Z}^n \setminus \{0\}$ , let  $l(y)$  be the least nonzero index of  $y$ , i.e.  $y_i = 0$  for all  $i < l(y)$ .*

**Lemma 6.** *Let  $y \in \{-1, 0, 1\}^n$ . Then  $y$  is in  $P$  if and only if one of the following is true*

- (i)  $y$  is the origin
- (ii)  $y_{l(y)} = 1$  and  $y_i \in \{-1, 0\}$ , for all  $i \geq l(y) + 1$
- (iii)  $y_{l(y)} = -1$  and  $y_i \in \{1, 0\}$ , for all  $i \geq l(y) + 1$ .

*Proof.* We first prove that if  $y \in \{-1, 0, 1\}^n$  is feasible then it must satisfy one of the three conditions. Assume therefore that  $y \in \{-1, 0, 1\}^n$  is feasible. The point  $y = 0$  is trivially feasible (option (i)). If  $y$  is not the origin, there must be some  $y_j \neq 0$ . Assume that  $y_{l(y)} = 1$ . If there is a  $k \neq l(y)$  with  $y_k = 1$ , then  $y$  violates Inequality (9) with  $j = k$

$$\sum_{i=1}^{k-1} \frac{1}{2^i} x_i + x_k + \sum_{i=k+1}^n \frac{1}{2^{i-1}} x_i \leq 1$$



so  $y$  satisfies (ii). The case that  $y_{l(y)} = -1$  is symmetric and omitted and leads to option (iii).

Assume  $y$  satisfies one of the three conditions, we want to prove that it is feasible. Obviously, if  $y$  is the origin, it is feasible. Assume that  $y$  satisfies (ii), the other case is symmetric and omitted here. We now prove that all inequalities are satisfied by  $y$ .

First consider (9). The term with  $x_{l(y)}$  is less or equal to 1 whereas the remaining part of the summation is nonpositive which makes the left-hand-side of (9) smaller or equal to 1.

Consider now (10). If  $j \leq l(y)$ , the term with  $x_{l(y)}$  is nonpositive whereas the sum of the remaining terms is less or equal to 1 which proves that the inequality is satisfied. If  $j \geq l(y) + 1$ , let us denote the inequality as  $\sum_{i=1}^n \alpha_i x_i \leq 1$ . Observe that  $\alpha_{l(y)} = -\frac{1}{2^{l(y)}}$ ,  $\alpha_j = -1$  and  $0 > \alpha_i \geq -\frac{1}{2^{l(y)+1}}$  for all  $i \geq l(y) + 1, i \neq j$ . Therefore  $\alpha_{l(y)} y_{l(y)} + \sum_{i \geq l(y), i \neq j} \alpha_i x_i \leq 0$  and  $\alpha_j x_j \leq 1$  which makes the left-hand-side of (10) smaller or equal to 1.

Consider Inequality (11). First observe that the second sum of (11) is bounded from above by  $1/|N|$ . Concerning the first two terms, we distinguish two cases. If  $l(y) \in N, l_N \neq l(y)$ , it implies that  $y_{l_N} = 0$  using condition (ii), and bounds  $-\frac{1}{|N|} x_{l_N} + \sum_{i \in N, i \neq l_N} \frac{1}{|N|} x_i \leq \frac{1}{|N|}$ . Otherwise  $-\frac{1}{|N|} x_{l_N} \leq \frac{1}{|N|}$  and  $\sum_{i \in N, i \neq l_N} \frac{1}{|N|} x_i \leq 0$  which implies that in both cases, the left-hand-side of (11) is bounded from above by 1.

Consider Inequality (12). We distinguish two cases. In the first case, we assume that  $l(y) = l_N$ . This implies that first term of (12) equals  $1/|N|$ , the first sum is bounded from above by  $(|N| - 1)/|N|$  as it contains  $|N| - 1$  terms and the last sum is nonpositive using condition (ii). Therefore the left-hand-side of (12) is bounded from above by 1. In the second case, we assume that  $l(y) \neq l_N$ . Therefore the first term of (12) is bounded from above by 0, the first sum is bounded from above by  $(|N| - 1)/|N|$  and the second sum is bounded from above by  $1/|N|^n$  and the result follows.  $\square$

**Lemma 7.** *Each of the  $2(2^n - 1)$  inequalities in  $P$  is necessary, i.e., the removal of any inequality from  $P$  results in the inclusion of at least one additional integer point in the interior of  $P$ .*

*Proof.* We will show the lemma by proving that each facet of  $P$  contains exactly one integer feasible point in its relative interior.

Consider an inequality of type (9). Observe that for any point satisfying conditions (ii) or (iii) of Lemma 6,  $\sum_{i=1}^{j-1} \frac{1}{2^i} x_i + \sum_{i=j+1}^n \frac{1}{2^{i-1}} x_i < 1$ . Therefore to make (9) tight we need  $x_j = 1$  which implies  $\sum_{i=1}^{j-1} \frac{1}{2^i} x_i + \sum_{i=j+1}^n \frac{1}{2^{i-1}} x_i = 0$ . Since the coefficients are in a geometric progression, this in turn implies  $x_i = 0$  for all  $i \neq j$ . We have therefore proven that the unit vectors are tight for all inequalities of type (9). By symmetry, for each inequality of type (10), only  $-e_j$  (where  $e_j$  denotes the  $j^{\text{th}}$  unit vector) is tight, integer and valid.

Consider an inequality of type (11). Observe that for any point satisfying conditions (ii) or (iii) of Lemma 6,  $-\sum_{i \notin N} \frac{1}{|N|^n} x_i < \frac{1}{|N|}$ . Therefore to make (11) tight, we need  $-\frac{1}{|N|} x_{l_N} + \sum_{i \in N, i \neq l_N} \frac{1}{|N|} x_i = 1$  which implies  $x_{l_N} = -1$  and  $x_i = 1$  for all  $i \in N \setminus l_N$  and  $x_i = 0$  for  $i \notin N$ . Symmetrically, for each inequality of type (12), only  $x_{l_N} = 1, x_i = -1$  for all  $i \in N \setminus l_N$  and  $x_i = 0$  for  $i \notin N$  is tight, integer and valid.

By observing that all points that were shown to be tight for the facet-defining inequalities of  $P$  are all different, the result follows.

Lastly, the fact that there are  $2(2^n - 1)$  planes follows from the fact that they are in bijection with double the number of nonempty subsets of  $\{1, \dots, n\}$ .  $\square$

In this section, we have dealt with the case  $k = 1$  and proven that the upper bound given in Theorem 1 is tight. Since the upper bound for  $k = 2$  matches that for  $k = 1$ , it is natural to conjecture that the bound is tight for  $k = 2$  as well. We know that for  $c(2, k)$  the bound is not tight for  $k \geq 3$ . We also believe that it is not tight for  $n, k \geq 3$  and can further be improved.

## 4 Consequences and variations of Theorem 1

In this section, we discuss variations and consequences of Theorem 1.

Let us begin by remarking that while replacing the  $=$  by  $\geq k$  in the statement of Theorem 1 gives a very easy-to-prove result, nevertheless one can state a more surprising corollary of Theorem 1 that involves estimations of the number of integer points and resembles more the quantitative Helly theorem of [7, 8]:

**Corollary 2.** *There exists a universal constant  $c(n, k)$  such that, given any system of inequalities  $\{x \in \mathbb{R}^n : Ax \leq b\}$ , if every subset of the constraints of cardinality  $c(n, k)$  has more than  $k$  integer solutions, then the entire system of inequalities must have more than  $k$  integer solutions.*

One natural question is how to rephrase Theorem 1 for convex sets rather than systems of linear inequalities

**Lemma 8.** *Given an infinite collection of convex sets  $(X_i)_{i \in \Lambda}$ ,  $X_i \subseteq \mathbb{R}^n$  such that there is some index  $\tilde{r}$  with  $X_{\tilde{r}}$  bounded and  $\bigcap_{i \in \Lambda} X_i$  contains exactly  $k$  integer points, there is a finite subcollection of these convex sets whose intersection contains these  $k$  integer points and no others.*

*Proof.* Consider the set  $\bigcap_{i \in \Lambda} X_i$ . Since this is a subset of  $X_{\tilde{r}}$ , the intersection is properly contained in a hypercube  $B$  with integer vertices. Consider the set

$$S = \{x \in B \setminus \bigcap_{i \in \Lambda} X_i : x \in \mathbb{Z}^n\}.$$

Since  $B$  bounds this set,  $S$  is finite. Note that  $S$  is non-empty otherwise  $B = \bigcap_{i \in \Lambda} X_i$ , which means that the (finitely many)  $X_i$  that contain the facets of  $B$  form the desired finite subcollection. For each  $x \in S$ , define  $P_x = \{X_i : x \notin X_i\}$ . Using the axiom of choice (if  $P_x$  is uncountable, else enumerate and pick the  $X_i$  of least index), pick an element  $X_{i(x)}$  in  $P_x$  for each  $x \in S$ .

Let  $I$  be the set of indices  $I := \{i(x) : x \in S\} \cup \{\tilde{r}\}$ . Note that since  $S$  is finite, then  $I$  is finite.

Claim:  $\bigcap_{i \in I} X_i$  contains the same integer points as  $\bigcap_{i \in \Lambda} X_i$  and no others.

Obviously  $\bigcap_{i \in \Lambda} X_i \subseteq \bigcap_{i \in I} X_i$  since it is the intersection of more sets. For the reverse containment, assume for contradiction that there is an integer point  $y \in \bigcap_{i \in I} X_i$  such that  $y \notin \bigcap_{i \in \Lambda} X_i$ . Obviously  $y \in B$ , since  $\tilde{r} \in I$ . By construction,  $I$  contains at least one index  $i(x)$  such that  $X_i(x)$  excludes  $y$ , so  $y \notin \bigcap_{i \in I} X_i$ . This is a contradiction. Thus

$$\left( \bigcap_{i \in I} X_i \right) \cap \mathbb{Z}^n \subseteq \bigcap_{i \in \Lambda} X_i,$$

so  $(X_i)_{i \in I}$  is the desired finite subcollection of our original convex sets.  $\square$

**Theorem 3.** *Let  $n, k$  be positive integers. There is a universal constant  $c(n, k)$ , depending only on the dimension  $n$  and  $k$ , such that, for any collection  $(X_i)_{i \in \Lambda}$  of closed convex sets in  $\mathbb{R}^n$ , where at least one of the sets is bounded and exactly  $k$  integer points are in  $\bigcap_{i \in \Lambda} X_i$ ; then there is a subcollection of size less than or equal to  $c(n, k)$  with exactly the same  $k$  integer points in their intersection.*

*Proof.* By Lemma 8, it suffices to consider a finite subcollection  $\mathcal{A} \subset \Lambda$  of indices, where at least one of the  $X_i, i \in \mathcal{A}$  is bounded, say without loss of generality  $X_1$ . Since  $X_1$  is bounded, there is a hypercube  $B$  with integer vertices that bounds it. Let  $U$  be the set of  $2n$  hyperplanes that

determine  $B$ . For each of the integer points  $y$  in  $B \setminus \bigcap_{i \in \mathcal{A}} X_i$ , there is an  $X_\alpha$  such that  $y \notin X_\alpha$ . It follows that there is a supporting hyperplane  $v_y$  of  $X_\alpha$  which is violated by  $y$ . Let

$$S = \bigcup_{y \in (B \cap \mathbb{Z}^n) \setminus \bigcap_{i \in \mathcal{A}} X_i} \{v_y\} \cup U.$$

Note that  $S$  is finite. Let  $P$  be the polytope determined by the constraints in  $S$ . Then  $P$  contains the  $k$  integer points in  $\bigcap_{i \in \mathcal{A}} X_i$ , because the hyperplanes were selected from  $B$  and the  $v_y$ s.

For the reverse containment, assume for a contradiction that there is an integer point  $y$  that is in  $P$ , but not in  $\bigcap_{i \in \mathcal{A}} X_i$ . Obviously,  $y \in B$ . But by construction,  $S$  contains a constraint which violates  $y$ , so that  $y$  is not in the polytope determined by  $S$ .

By Theorem 1,  $P$  has at most  $c(n, k)$  necessary hyperplanes, i.e. there is a subset of the constraints in  $S$  of size no more than  $c(n, k)$  that yield a polyhedron  $P_{c(n, k)}$  which contains the  $k$  original integer points and no others. By construction, the hyperplanes in  $P_{c(n, k)}$  have a well-defined identification with the  $X_i$ 's. Namely, if a hyperplane  $v \in S$  comes from  $B$ , then it is identified with  $X_1$ . Otherwise,  $v$  is identified with an arbitrary chosen  $X_\alpha$  from a finite number of convex set used in the construction of  $v$ . Call this identification  $\phi$ . The image of the hyperplanes in  $P_{c(n, k)}$  under  $\phi$  is a subcollection  $(X_i)_{i \in \mathcal{F}}$  which contains the  $k$  original integer points and no others, by construction.  $\square$

As a final application of Theorem 1 we are interested in using it within the theory of violator spaces, and then a Clarkson-type randomized algorithm, to compute the best, 2nd best, ...,  $l$ -th best solutions to a given integer linear optimization problem. We call these points the *l-best solutions of an ILP*. The literature on this problem is quite extensive and established (see e.g., [21, 24]). As we will see the resulting algorithm will be linear on the number of constraints, when  $l$  and the dimension are fixed constants.

Let us remember the basics of this theory. In the years since Clarkson wrote his well-known paper [12], several researchers observed that his algorithm works for optimization problems that fit certain abstract structures. This applies for LP-type problems e.g., in [2, 28]. More recently Gärtner et al [20] proved that in fact Clarkson's algorithm works in greatest generality for the so called *violator spaces*. Essentially, a violator space is an optimization problem in which we have a finite set of constraints or elements  $H$  and a function that given any subset of constraints  $G$ , indicates which other constraints in  $H \setminus G$  *violate* the feasible solutions to  $G$ . If one has a violator space structure, the optimal solution of the problem can be computed via a randomized method whose running time is *linear* in the number of constraints defining the problem, and *subexponential* in the dimension of the problem. Thus when  $l$  and the dimension of the problem are constant, it gives a polynomial-time method. We recall here the necessary definitions and properties of violator spaces and see that they fit the problem at hand.

**Definition 3.** A violator space is a pair  $(H, V)$  where  $H$  is a finite set and  $V$  is a mapping  $2^H \rightarrow 2^H$  such that the following two conditions hold.

- **Consistency:**  $G \cap V(G) = \emptyset$  for all  $G \subseteq H$ ,
- **Locality:** For all  $F \subseteq G \subseteq H$ , where  $G \cap V(F) = \emptyset$ , we have  $V(G) = V(F)$ .

In our case,  $H$  is the set of linear inequality constraints of an integer linear program  $IP(H) = \min \{c^T x \mid a^{(i)} x \leq b^{(i)}, i \in H, x \in \mathbb{Z}^n\}$ . For every  $G \subseteq H$ , we consider the IP defined using a subset of the constraints  $IP(G) = \min \{c^T x \mid a^{(i)} x \leq b^{(i)}, i \in G, x \in \mathbb{Z}^n\}$ .

We define the violator set  $V(G)$  as the set of inequalities  $h \in H$  such that the  $l$  best solutions of  $IP(G)$  are not identically equal to the  $l$  best solutions of  $IP(G \cup \{h\})$ . Note that we need to

have a total ordering on the possible feasible solutions of  $IP(G)$ . We therefore provide each integer program with a universal tie-breaking rule, for instance, using lexicographic ordering. Assume that  $IP(G)$  has at least  $l$  different feasible solutions. Define  $\mathbf{x}_l(G)$  to be the  $l$ -tuple consisting of the  $l$  best integer points in  $IP(G)$  under this ordering. For our application we say that a constraint  $h \in H$  is in  $V(G)$  if  $\mathbf{x}_l(G) \neq \mathbf{x}_l(G \cup \{h\})$ . If we assume that  $IP(G)$  has less than  $l$  feasible solutions, we define  $V(G)$  as being the empty set.

We call the pair  $(H, V)$  defined above as an  $l$ -th best IP.

**Lemma 9.** *The  $l$ -th best IP is a violator space.*

*Proof.* We need to check that the two conditions presented in Definition 3 are satisfied. The consistency condition is clearly satisfied.

Assume that  $F \subseteq G \subseteq H$  and  $G \cap V(F) = \emptyset$ . To show locality we want to show that  $V(F) = V(G)$ . We first consider the case where  $IP(F)$  has less than  $l$  feasible solutions. Then  $V(F) = \emptyset$ . Obviously  $IP(G)$  has less feasible solutions than  $IP(F)$  since  $IP(G)$  includes already all constraints of  $IP(F)$ . Hence  $V(G) = \emptyset = V(F)$ .

In the following, we assume that  $IP(F)$  has at least  $l$  feasible solutions. For the containment  $V(F) \subseteq V(G)$  note that  $V(F)$  can be alternatively characterized as the set of constraints that violate or cut at least one  $x_i \in \mathbf{x}_l(F)$ . These constraints cannot be in  $G$ , by the assumption that  $G \cap V(F) = \emptyset$ . It follows that the set of constraints that remove any of the  $x_i \in \mathbf{x}_l(F)$  cannot be in  $G$ , so that the same points  $x_i \in \mathbf{x}_l(G)$  for  $i = 1, \dots, l$ . Thus any constraint  $m \in V(F)$  is in  $V(G)$ , so that  $V(F) \subseteq V(G)$ . For the containment  $V(G) \subseteq V(F)$ , observe that when  $m \in G$ , then  $m \notin V(F)$  (because  $G \cap V(F) = \emptyset$ ), thus  $\mathbf{x}_l(F \cup \{m\}) = \mathbf{x}_l(F)$ . In other words, we can add any constraint in  $G \setminus F$  to  $F$  without changing  $\mathbf{x}_l(F)$ . Add all those constraints  $m \in G \setminus F$ . We then have  $\mathbf{x}_l(F) = \mathbf{x}_l(G)$  since each additional constraint left the  $l$ -tuple unaltered. Because  $\mathbf{x}_l(F) = \mathbf{x}_l(G)$  any constraint in  $V(G)$ , also violates at least one of the  $x_i \in \mathbf{x}_l(F)$ , so that  $V(G) \subseteq V(F)$ .  $\square$

Before we can apply the theory of violator spaces of [20] we need two more ingredients. First, just like a linear programming optimum is defined by a basis, we need to have a notion of basis for our optimal solutions.

**Definition 4.** *Given a violator space  $(H, V)$ , we say that  $B \subseteq H$  is a basis if for all proper subsets  $F \subset B$  we have  $B \cap V(F) \neq \emptyset$ . For  $G \subseteq H$ , a basis of  $G$  is a minimal subset  $B$  of  $G$  that is a basis and such that  $V(B) = V(G)$ .*

*The combinatorial dimension of a violator space  $(H, V)$  is the maximal cardinality of a basis.*

Intuitively a basis is a minimal subset of constraints with the same optimal value as the whole set. Now we see that the size of a basis is bounded by the constant of Theorem 1.

**Lemma 10.** *The combinatorial dimension of  $l$ -th best IP with  $n$  variables is bounded by the constant  $c(n, l)$ .*

*Proof.* Consider a set of constraints  $H$  and the corresponding  $IP(H)$ . First assume that  $IP(H)$  has  $\bar{l} < l$  feasible solutions. Then there exists a subset of the constraints  $G \subseteq H$  with cardinality at most  $c(n, \bar{l})$  that has the same number of solutions and is therefore a basis. Observe that  $c(n, \bar{l}) \leq c(n, l)$ .

Assume now that  $IP(H)$  has at least  $l$  feasible solutions. Considering the set of constraints  $\{c^T x \leq c^T x_l\} \cup H$ , where  $x_l$  is the  $l$ -th best value in  $\mathbf{x}_l(H)$ , and applying Theorem 1, we conclude that there exists a subset of the constraints of cardinality at most  $c(n, l)$  that define the same  $l$  feasible solutions. Getting rid of  $\{c^T x \leq c^T x_l\}$ , we conclude that the size of the largest basis of  $H$  is  $c(n, l)$ .  $\square$

The second ingredient is that we need to have a way to answer the following query in polynomial time in fixed dimension for subsets of size smaller than the combinatorial dimension.

**Primitive query:** Given  $G \subset H$  and  $h \in H \setminus G$ , decide whether  $h \in V(G)$ .

The reason we need to answer this query via a black-box method is because then, using Theorem 27 in Section 4 of [20], we obtain the following result

**Lemma 11.** *A basis of  $H$  in a violator space  $(H, V)$  (and thus an optimal solution to the problem) can be found by calling the algorithm that solves the primitive query an expected*

$$O(c(n, l)|H| + c(n, l)^{O(c(n, l))})$$

*number of times.*

The primitive query originally provided by Clarkson in the case of regular integer programming was Lenstra's IP algorithm in fixed dimension. For our problem, when we now look for the  $l$ -th best solution, the primitive query can be answered by calling  $l$  times an algorithm for IP in fixed dimension. Given an integer program in fixed dimension  $n$ , a fixed number of constraints  $q$  and a maximum bit size of the data of  $s$ , Eisenbrand provided in [19] an algorithm that finds an optimal solution in  $\mathcal{O}(s)$  operations. Our primitive query calls a maximum of  $l$  times Eisenbrand's algorithm in fixed dimension  $n$  and constant number of constraints  $c(n, l)$  to answer the question. We make this explicit in the following lemma:

**Lemma 12.** *Given fixed positive integers  $q, n$  and  $l$  and an integer program  $\min\{c^T x : Ax \leq b, x \in \mathbb{Z}^n, 0 \leq x \leq u\}$  with  $q$  constraints and a varying maximum encoding length of  $s$  bits for the data, it is possible to determine in  $\mathcal{O}(ls)$  operations the  $l$  best solutions to the integer program where ties are broken using lexicographic order.*

*Proof.* We use the global bound  $u$  on the variables, define  $\bar{c} := u^n c + \sum_{i=1}^n u^{n-i} e_i$  and set up the auxiliary integer program  $\min\{\bar{c}^T x : Ax \leq b, x \in \mathbb{Z}^n, 0 \leq x \leq u\}$ . Observe that by solving the latter, we have the guarantee to find the best integer point with respect to the objective function breaking ties lexicographically. Denoting  $\bar{x}^{(1)}$  the optimal solution to the above integer program, we can now find the second best integer point by solving  $\min\{\bar{c}^T x : Ax \leq b, x \in \mathbb{Z}^n, \bar{c}^T x \geq \bar{c}^T \bar{x}^{(1)} + 1, 0 \leq x \leq u\}$ . If we denote by  $\bar{x}^{(i)}$  the  $i^{\text{th}}$  best point, we obtain the  $(i+1)^{\text{th}}$  best point by solving a similar integer program with the additional constraint  $\bar{c}^T x \geq \bar{c}^T \bar{x}^{(i)} + 1$ .

This shows that we are able to compute the  $l$  best solutions to an integer program by successively solving  $l$  integer programs in dimension  $n$  with  $q+1$  constraints and with bit size  $ns$  which is of order  $s$  when  $n$  is fixed. The result follows by applying Eisenbrand's algorithm to all integer programs.  $\square$

We finally arrive at the key complexity consequence of this section:

**Corollary 3.** *Given fixed positive integer constants  $n$  and  $l$ , an integer  $m \times n$  matrix  $A$ , and the integer linear program*

$$\begin{aligned} & \min c^T x \\ & \text{subject to } Ax \leq b \\ & x \in \mathbb{Z}^n \end{aligned}$$

*of maximum bit-size  $s$  for the coefficients in  $A$ , then the  $l$ -th best solution can be computed in a expected*

$$O((ls)(c(n, l)m + c(n, l)^{O(c(n, l))}))$$

*number of operations.*

We conclude by noting that using the theory of rational generating functions (see [15] for an introduction) one can also prove a similar result with some worse complexity.

## Acknowledgements

The research of the third author was partially supported by a UCMEXUS project grant and by the Institute for Mathematics and its Applications with funds provided by the National Science Foundation. The research was carried out during the fourth author's sabbatical visit at UC Davis supported by the Belgian Science Foundation (FRS-FNRS) and UC Davis. The authors are truly grateful to the anonymous referees for their excellent detailed comments and corrections that greatly improved the quality of this paper. We are also grateful to Robert Hildebrand, Luis Montejano, Timm Oertel, Deborah Oliveros, János Pach, and Edgardo Roldán Pensado for their very useful comments.

## References

1. I. Aliev, J.A. De Loera, and Q. Louveaux, "Integer Programs with Prescribed Number of Solutions and a Weighted Version of Doignon-Bell-Scarf's Theorem", in *Proceedings of Integer Programming and Combinatorial Optimization, 17th International IPCO Conference*, Bonn Germany, June, 2014.
2. N. Amenta, "Helly-type theorems and generalized linear programming", *Discrete and Computational Geometry* 12: (1994), 241–261.
3. G. Averkov and R. Weismantel, "Transversal numbers over subsets of linear spaces". *Adv. Geom.* 12 (2012), no. 1, 19–28.
4. K. Andersen, Q. Louveaux, and R. Weismantel. "Certificates of linear mixed integer infeasibility", *Operations Research Letters*, 36 (2008), 734–738.
5. K. Andersen, Q. Louveaux, and R. Weismantel. "An analysis of mixed integer linear sets based on lattice point free convex sets", *Math of Operations Research*, 35 (2010), 233–256.
6. K. Andersen, Q. Louveaux, R. Weismantel, and L. Wolsey. "Inequalities from two rows of the simplex tableau", in *M. Fischetti & D. P., Williamson (Eds.) Integer Programming and Combinatorial Optimization, 12th International IPCO Conference, Ithaca, NY, USA, June 25-27, 2007, Proceedings, Lecture Notes in Computer Science* 4513, 1–15.
7. I. Bárány, M. Katchalski, and J. Pach. "Quantitative Helly-type theorems", *Proc. Amer. Math. Soc.*, 86 (1982), 109–114.
8. I. Bárány, M. Katchalski, and J. Pach. "Helly's theorem with volumes." *Amer. Math. Monthly*, 91 (6):362–365, 1984.
9. A. Barvinok and J. Pommersheim. "An algorithmic theory of lattice points in polyhedra". *New perspectives in algebraic combinatorics, Math. Sci. res. Inst. Publ.*, 38, Cambridge Univ. Press, Cambridge, (1999), 91-147.
10. D.E. Bell, "A theorem concerning the integer lattice". *Studies in Applied Mathematics*, 56(1), (1977), 187–188.
11. V. Borozan and G. Cornuéjols, "Minimal valid inequalities for integer constraints". *Math. Oper. Res.* 34 (2009), no. 3, 538–546.
12. K.L. Clarkson, "Las Vegas algorithms for linear and integer programming when the dimension is small", *Journal of the ACM* (1995), 42 (2), 488–499.
13. M. Conforti, G. Cornuéjols, and G. Zambelli, "Corner polyhedron and intersection cuts". *Surveys in Operations Research and Management Science*, 16 (2011), 105–120.
14. L. Danzer, B. Grünbaum, and V. Klee, "Helly's theorem and its relatives", in 1963 Proc. Sympos. Pure Math., Vol. VII pp. 101180 Amer. Math. Soc., Providence, R.I.
15. J.A. De Loera, R. Hemmecke, M. Köppe, "Algebraic and geometric ideas in the theory of discrete optimization" *MOS-SIAM Series on Optimization, 14. Society for Industrial and Applied Mathematics (SIAM)*, Philadelphia, PA; Mathematical Optimization Society, Philadelphia, PA, 2013. xx+322 pp.

16. S. Dey and L. Wolsey, “Constrained infinite group relaxations of MIPs”. *SIAM J. Optim.* 20 (2010), no. 6, 2890–2912.
17. J-P. Doignon, “Convexity in cristallographical lattices”, *Journal of Geometry* 3.1 (1973), 71–85.
18. J. Eckhoff, “Helly, Radon, and Carathéodory type theorems” in *Handbook of convex geometry*, Vol. A, B, 389448, North-Holland, Amsterdam, 1993.
19. F. Eisenbrand, “Fast integer programming in fixed dimension”, *Algorithms-ESA* (2003), 196–207.
20. B. Gärtner, J. Matoušek, L. Rüst, and P Škovroň. “Violator spaces: structure and algorithms”, *Discrete Applied Mathematics*, 156 (11): (2008), 2124–2141.
21. H.W. Hamacher and M. Queyranne, “K best solutions to combinatorial optimization problems”. *Ann. Oper. Res.* 4 (1985), no. 1-4, 123–143.
22. A.J. Hoffman, “Binding constraints and Helly numbers” in Proceedings of Second International Conference on Combinatorial Mathematics (New York, 1978), pp. 284288, *Ann. New York Acad. Sci.*, 319, New York Acad. Sci., New York, 1979.
23. J.C. Lagarias and G.M. Ziegler, “Bounds for lattice polytopes containing a fixed number of interior points in a sublattice”. *Canad. J. Math.* 43, no. 5, (1991), 1022–1035.
24. E.L. Lawler, “A procedure for computing the K-best solutions to discrete optimization problems and its application to the shortest path problem”. *Management Sci.* 18 (1971/72), 401–405.
25. O. Pikhurko, “Lattice points in lattice polytopes”. *Mathematika* 48, no. 1-2, (2003), 15–24.
26. H.E. Scarf, “An observation on the structure of production sets with indivisibilities”, *Proceedings of the National Academy of Sciences* 74.9 (1977), 3637–3641.
27. A. Schrijver, “Theory of linear and integer programming” *Wiley-Interscience Series in Discrete Mathematics* A Wiley-Interscience Publication. John Wiley & Sons, Ltd., Chichester, 1986. xii+471 pp.
28. M. Sharir, E. Welzl, “A combinatorial bound for linear programming and related problems”, in *Proceedings of 9th Annual Symposium on Theoretical Aspects of Computer Science (STACS), Lecture Notes in Computer Science* 577, Springer-Verlag, (1992), 567–579.
29. R. Wenger, “Helly-type theorems and geometric transversals” in *Handbook of discrete and computational geometry*, Handbook of discrete and computational geometry. Second edition. Edited by Jacob E. Goodman and Joseph O’Rourke. Discrete Mathematics and its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL, 2004. xviii+1539 pp.