

# UCLA

## UCLA Previously Published Works

### Title

Participant-derived cell line transcriptomic analyses and mouse studies reveal a role for ZNF335 in plasma cholesterol statin response

### Permalink

<https://escholarship.org/uc/item/1s63c192>

### Journal

Genome Medicine, 16(1)

### ISSN

1756-994X

### Authors

Theusch, Elizabeth

Ting, Flora Y

Qin, Yuanyuan

et al.

### Publication Date

2024

### DOI

10.1186/s13073-024-01366-9

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>


Peer reviewed

RESEARCH

Open Access



# Participant-derived cell line transcriptomic analyses and mouse studies reveal a role for ZNF335 in plasma cholesterol statin response

Elizabeth Theusch<sup>1\*</sup> , Flora Y. Ting<sup>1</sup>, Yuanyuan Qin<sup>1</sup>, Kristen Stevens<sup>1</sup>, Devesh Naidoo<sup>1</sup>, Sarah M. King<sup>1</sup>, Neil V. Yang<sup>1</sup>, Joseph Orr<sup>1</sup>, Brenda Y. Han<sup>2</sup>, Jason G. Cyster<sup>2</sup>, Yii-Der I. Chen<sup>3</sup>, Jerome I. Rotter<sup>3</sup>, Ronald M. Krauss<sup>1,4</sup> and Marisa W. Medina<sup>1\*</sup>

## Abstract

**Background** Statins lower circulating low-density lipoprotein cholesterol (LDLC) levels and reduce cardiovascular disease risk. Though highly efficacious in general, there is considerable inter-individual variation in statin efficacy that remains largely unexplained.

**Methods** To identify novel genes that may modulate statin-induced LDLC lowering, we used RNA-sequencing data from 426 control- and 2  $\mu$ M simvastatin-treated lymphoblastoid cell lines (LCLs) derived from European and African American ancestry participants of the Cholesterol and Pharmacogenetics (CAP) 40 mg/day 6-week simvastatin clinical trial (ClinicalTrials.gov Identifier: NCT00451828). We correlated statin-induced changes in LCL gene expression with plasma LDLC statin response in the corresponding CAP participants. For the most correlated gene identified (*ZNF335*), we followed up in vivo by comparing plasma cholesterol levels, lipoprotein profiles, and lipid statin response between wild-type mice and carriers of a hypomorphic (partial loss of function) missense mutation in *Zfp335* (the mouse homolog of *ZNF335*).

**Results** The statin-induced expression changes of 147 human LCL genes were significantly correlated to the plasma LDLC statin responses of the corresponding CAP participants in vivo (FDR = 5%). The two genes with the strongest correlations were zinc finger protein 335 (*ZNF335* aka *NIF-1*,  $\rho = 0.237$ , FDR-adj  $p = 0.0085$ ) and CCR4-NOT transcription complex subunit 3 (*CNOT3*,  $\rho = 0.233$ , FDR-adj  $p = 0.0085$ ). Chow-fed mice carrying a hypomorphic missense (R1092W; aka bloto) mutation in *Zfp335* had significantly lower non-HDL cholesterol levels than wild-type C57BL/6J mice in a sex combined model ( $p = 0.04$ ). Furthermore, male (but not female) mice carrying the *Zfp335*<sup>R1092W</sup> allele had significantly lower total and HDL cholesterol levels than wild-type mice. In a separate experiment, wild-type mice fed a control diet for 4 weeks and a matched simvastatin diet for an additional 4 weeks had significant statin-induced reductions in non-HDL cholesterol ( $-43 \pm 18\%$  and  $-23 \pm 19\%$  for males and females, respectively). Wild-type male (but

\*Correspondence:

Elizabeth Theusch  
Elizabeth.Theusch@ucsf.edu  
Marisa W. Medina  
Marisa.Medina@ucsf.edu

Full list of author information is available at the end of the article



© The Author(s) 2024. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

not female) mice experienced significant reductions in plasma LDL particle concentrations, while male mice carrying *Zfp335*<sup>R1092W</sup> allele(s) exhibited a significantly blunted LDL statin response.

**Conclusions** Our in vitro and in vivo studies identified *ZNF335* as a novel modulator of plasma cholesterol levels and statin response, suggesting that variation in *ZNF335* activity could contribute to inter-individual differences in statin clinical efficacy.

**Keywords** Statin, Cholesterol, RNA-sequencing, Lymphoblastoid cell lines, Mice, Zinc finger protein 335, Lipoprotein, Gene expression

## Background

Elevated plasma low-density lipoprotein cholesterol (LDLC) levels are a major risk factor for cardiovascular disease (CVD), the leading cause of death worldwide. Statins lower plasma LDLC levels and CVD risk by inhibiting cholesterol biosynthesis and stimulating receptor-mediated LDL uptake from the blood. Although statins are effective for primary and secondary prevention of atherosclerotic CVD [1], not everyone receives the same lipid-lowering benefit from statin therapy, and many individuals still experience CVD events while taking statins. Much of the inter-individual variation in statin efficacy remains unexplained to date.

Previous studies have identified some factors (e.g., age, smoking status, and race/ethnicity) associated with LDLC statin response [2]. In addition, genome-wide association studies (GWAS) have identified a handful of genetic loci associated with LDLC response [3–6], but together these only explain about 5% of LDLC response variation [5]. Though expanding sample sizes of participants with suitable pre- and on-drug phenotype data could bolster power for statin response GWAS, other approaches beyond GWAS could also be used to identify genetic factors that influence statin efficacy.

Statins elicit a strong transcriptional response via regulation of sterol response element binding factor 2 (SREBF2, aka SREBP2) [7] and other factors, so we and others have studied statin-induced gene expression changes to reveal novel genes involved in the modulation of clinical statin response [8–11]. To do this, we have used statin clinical trial participant-derived lymphoblastoid cell lines (LCLs) as an in vitro genetic model in which to study statin response transcriptomics. Here, we conducted a large-scale study correlating in vitro gene expression statin response with in vivo LDLC statin response to identify candidate statin efficacy modulators. We then validated a top candidate gene in vivo in a mouse model.

## Methods

### CAP study participants

This study used clinical and demographic information and cell lines derived from 426 participants in the Cholesterol and Pharmacogenetics (CAP) 40 mg/day 6-week simvastatin clinical trial [2] (ClinicalTrials.gov Identifier: NCT00451828). Written informed consent was obtained from all participants, and the study was approved by the institutional review boards of the University of California, San Francisco and the University of California, Los Angeles. Plasma lipids were measured in two baseline (pre-statin) blood samples and two on-statin blood samples collected 4 and 6 weeks after statin initiation, and the two values were averaged in each case to obtain a single baseline and on-statin value per participant. LDLC values were calculated from total cholesterol, HDL-cholesterol, and triglyceride levels using the Friedewald formula [12]. Characteristics of this CAP subset are shown in Table 1. In addition, lipoprotein particle profiles were measured and analyzed in the second baseline sample and the 6-week on-statin sample from most CAP participants ( $n = 803$ ) using ion mobility (described below).

### Lymphoblastoid cell line statin exposure

CAP lymphoblastoid cell lines (LCLs) were established by immortalizing blood-derived B lymphocytes through Epstein-Barr virus (EBV) transformation, cultured in RPMI 1640 with 10% FBS, and exposed to 2  $\mu$ M

**Table 1** Characteristics of 426 CAP participants with LCL RNA-seq data

Sex	215 F, 211 M
Self-reported race	162 AfrAm, 264 white
Age (years)	54.2 $\pm$ 12.3
BMI (kg/m <sup>2</sup> )	28.7 $\pm$ 5.9
% Smokers	17.8%
Baseline LDLC (mg/dl)	133.1 $\pm$ 34.0
On-statin LDLC (mg/dl)	77.0 $\pm$ 23.8
% Change LDLC	–41.9 $\pm$ 11.4%

Values are mean  $\pm$  SD

activated simvastatin or sham control buffer for 24 h in batches  $\leq 12$  cell lines, similar to previously described [13]. RNA was extracted from homogenized cellular lysates and RNA integrity was measured on an Agilent Bioanalyzer, as previously described [14].

### RNA-sequencing and analysis

PolyA-selected, strand-specific [15] paired-end RNA-seq libraries were prepared, and fragments were sequenced with 100–101 bp read lengths on Illumina machines at the University of Washington Northwest Genomics Center or the Baylor College of Medicine Human Genome Sequencing Center, as previously described [14, 16]. Sequences were aligned to the human (hg19) and EBV (NC\_007605) genomes using Tophat v2.0.4 [17] with Ensembl v67 [18] and EBV [19] transcriptome annotations, and sequence fragments falling within annotated genes were counted using HTSeq [20] as previously described [14, 16]. DESeq2 was used to adjust for library size and to variance stabilize gene expression levels using a  $\log_2$ -based transformation [21], and statin-induced changes in gene expression levels were calculated by subtracting expression levels in control-treated LCLs from the statin-treated expression levels of the same cell line.

After adjusting CAP participants' in vivo plasma LDLC statin responses [ $\ln(\text{on-statin LDLC}) - \ln(\text{baseline LDLC})$ ] for age, race/ethnicity, and smoking status using linear regression, the resulting residuals were correlated to statin-induced changes in gene expression in the corresponding LCLs using Spearman's rank-order correlation.  $p$  values were adjusted for multiple testing using the Benjamini–Hochberg false discovery rate (FDR) approach [22].

### Mouse studies

All mice in this study were housed under a 12-h light/dark cycle and had free access to food and water except when fasted as noted. Mice were fed a general grain-based chow diet prior to study initiation (irradiated Teklad global 18% protein rodent diet from Envigo, product 2918). C57BL/6J mice carrying an N-ethyl-N-nitrosourea (ENU)-induced hypomorphic missense mutation (R1092W, aka “bloto”) in *Zfp335* (available from JAX as stock #025853) were generated previously [23] and were backcrossed to C57BL/6J wild-type mice for many generations. For the first study, animals were fed the general chow diet until they were fasted for 4–6 h and sacrificed at 13 weeks of age. Blood was collected via cardiac puncture, and plasma was isolated. Plasma lipid levels and lipoprotein particle concentrations were measured using Liasys and ion mobility, respectively, described below. Growth curve analyses were performed using weekly weights obtained after weaning, and growth curves of

mice with different *Zfp335* genotypes were compared within sexes. Results were analyzed statistically and adjusted for multiple testing using the CGGC (Compare Groups of Growth Curves) method with 10,000 permutation tests [24, 25].

In our second study, mice were transitioned to the AIN-76A purified rodent diet (ResearchDiets product D10001i) at 13 weeks of age, which contains approximately 5% fat and no added cholesterol, and weighed weekly for the duration of the study. After mice were fed the control purified diet for 4 weeks, they were transitioned to a matched diet with the addition of 1 g simvastatin/kg diet (9.77% active; ResearchDiets product D14060903i) for 4 weeks. Plasma was obtained from mice at 15, 17, and 19 weeks of age using non-terminal retro-orbital bleeds after a 4–6-h fast. Mice were sacrificed at 21 weeks of age after a 4–6-h fast, and plasma was isolated from blood collected via cardiac puncture as in the first study.

Mouse studies were approved by the Children's Hospital & Research Center Oakland and the University of California, San Francisco Institutional Animal Care and Use Committees.

### Plasma lipid analysis

Total cholesterol (TC) and high-density lipoprotein cholesterol (HDLC) were measured in human and mouse plasma samples on an AMS Liasys 330 Clinical Chemistry Analyzer, similar to previously described [26, 27]. Mouse non-HDL cholesterol (non-HDLC) levels were calculated by subtracting HDL from total cholesterol levels. In the chow-fed mouse experiment, HDLC was not successfully measured in samples from one male heterozygote mouse, and 1 female *Zfp335*<sup>R1092W</sup> homozygote was a statistical (Tukey) outlier for HDLC (low) and non-HDLC (high) and was excluded from analyses. Plasma samples from 4 mice (1 male wild-type, 1 male heterozygote, 1 female heterozygote, and 1 male *Zfp335*<sup>R1092W</sup> homozygote) in the statin response experiment for which non-HDLC was calculated to be  $\leq 0$  and/or fasting total cholesterol levels were abnormally low ( $\leq 50$  mg/dl) were Tukey outliers and were excluded from subsequent analyses. Additive relationships were tested for association using linear regression, and comparisons between wild-type and mutant homozygotes were conducted statistically using  $t$ -tests.

Ion mobility (IM) was used to measure human CAP participant and mouse plasma lipoprotein particle concentrations as a function of particle size, as previously described [28, 29]. Of the 25 male and 33 female mice for which IM data was generated before and on the statin diet, nine were statistically significant LDL particle concentration outliers (Tukey) and excluded from

subsequent IM data analysis, leaving data from 22 male and 27 female mice. Specifically, five mice (1 female wild-type, 1 female heterozygote, 1 female *Zfp335*<sup>R1092W</sup> homozygote, and 2 male wild-type) had abnormally high pre-statin LDL particle concentrations, three mice (1 female wild-type and 2 female heterozygotes) had abnormally high on-statin LDL particle concentrations, and one (male wild-type) mouse had abnormally low on-statin LDL particle concentrations. Data was binned into lipoprotein subfraction categories based on particle diameter definitions previously determined using human studies and comparisons with other methods [29]. Two-way ANOVA with Tukey’s multiple comparisons test was used to compare statin-induced changes in LDL subfraction particle concentrations across mouse genotype categories.

**Results**

**Transcriptomic statin response in LCLs**


Statin induced a robust transcriptional response in the 426 CAP LCLs, in agreement with our previous studies [9, 14] (Fig. 1, Table S1). Of the 14,028 well-expressed human and EBV genes tested, the expression level of 9092 genes (65%) changed significantly with statin treatment (paired *t*-test, FDR-adjusted *p* < 0.0001, 4671 upregulated and 4421 downregulated), though the degree of change was modest for most genes. As expected, many

of the most robustly upregulated genes were known SREBF2 target genes.

**Correlation of in vitro gene expression statin response with in vivo LDLC statin response**

To identify genes that may modulate plasma LDLC statin response, we correlated transcriptome-wide statin-induced changes in LCL gene expression in vitro with plasma LDLC statin responses of the corresponding CAP clinical trial participants from whom the LCLs were derived. At a false discovery rate of 5%, 147 of 13,955 human genes tested had statin-induced expression changes that were significantly correlated with in vivo LDLC statin response (Table S2). The top 10 genes, which all had an FDR-adjusted *p* < 0.03, are shown in Table 2.

The top two genes, zinc finger protein 335 (*ZNF335*) and CCR4-NOT transcription complex subunit 3 (*CNOT3*), appeared to be most worthwhile for follow-up due to the significance of their correlation relationships relative to those of other genes after multiple testing adjustment. *ZNF335* (aka NIF-1) has been shown to physically interact with two other members of the CCR4-NOT complex, *CNOT6* and *CNOT9* [30], suggesting that *ZNF335* and *CNOT3* may be functionally related. *ZNF335* and *CNOT3* also exhibited similar correlation directionality (Fig. 2). For both genes, statin decreased expression in most LCLs. In addition, the




**Fig. 1** Volcano plot illustrating gene expression statin response in 426 CAP LCLs. Approximate log<sub>2</sub> fold change is 2<sup>(variance stabilized statin gene expression level – variance stabilized control gene expression level)</sup> calculated for each cell line and averaged across cell lines

**Table 2** Genes with LCL expression changes most correlated to plasma LDLC statin response

Gene symbol	Gene name	Chr	Rho	<i>p</i>	Q
<i>ZNF335</i>	Zinc finger protein 335	20	0.237	7.8E-07	0.0085
<i>CNOT3</i>	CCR4-NOT transcription complex subunit 3	19	0.233	1.2E-06	0.0085
<i>PML</i>	PML nuclear body scaffold	15	0.218	5.4E-06	0.0246
<i>SYVN1</i>	Synoviolin 1	11	0.216	7.0E-06	0.0246
<i>ANKRD46</i>	Ankyrin repeat domain 46	8	-0.208	1.6E-05	0.0293
<i>EIF4E</i>	Eukaryotic translation initiation factor 4E	4	-0.206	1.9E-05	0.0293
<i>MED15</i>	Mediator complex subunit 15	22	0.206	1.9E-05	0.0293
<i>TCF3</i>	Transcription factor 3	19	0.205	2.1E-05	0.0293
<i>OTUD6B</i>	OTU deubiquitinase 6B	8	-0.205	2.1E-05	0.0293
<i>AASDH</i>	Aminoadipate-semialdehyde dehydrogenase	4	-0.205	2.1E-05	0.0293

Rho is the Spearman correlation coefficient and Q is the FDR-adjusted *p* value




**Fig. 2** Correlation of *ZNF335* and *CNOT3* LCL gene expression statin responses with in vivo plasma LDLC statin responses of the corresponding donors (A–B) and with each other (C)

LCLs that exhibited greater statin-induced reductions in gene expression tended to be derived from participants who experienced larger statin-induced reductions in plasma LDLC in the CAP clinical trial (Fig. 2A and B). Though no other populations were available for replication of these findings, we did see good agreement between the two independent experimental batches [(batch 1:  $N=149$ ; *ZNF335* rho=0.275,  $p=7.0 \times 10^{-4}$ ; *CNOT3* rho=0.316,  $p=8.6 \times 10^{-5}$ ); (batch 2:  $N=277$ ; *ZNF335* rho=0.200,  $p=8.0 \times 10^{-4}$ ; *CNOT3* rho=0.154,  $p=0.010$ )]. Notably, the statin-induced changes in *CNOT3* and *ZNF335* were themselves highly correlated (rho=0.625,  $p=1.4 \times 10^{-47}$ , Fig. 2C). Other genes in the CCR4-NOT complex did not have gene expression statin responses that were significantly correlated to plasma LDLC response after adjustment for multiple testing, and their non-significant correlations with plasma LDLC response were negative, displaying opposite directionality to the positive *CNOT3* and *ZNF335* correlations (Table S2).

#### Plasma cholesterol levels in *Zfp335* hypomorphic mice

In mice, knocking out *Cnot3* or *Zfp335* (the murine *ZNF335* homolog) results in embryonic lethality [31, 32]. Previous studies have shown that *Cnot3*<sup>+/-</sup> (haplodeficient) mice have altered lipid and glucose metabolism [33], and liver-specific *Cnot3* knockout (*Cnot3*LKO) mice have altered lipid metabolism, among other deficits [34]. While *ZNF335* has been reported to play a role in neurogenesis [32], to date the effects of *Zfp335* disruption on lipid and cholesterol metabolism have not previously been studied. We chose to investigate plasma cholesterol levels and statin response in available mice carrying an ENU-induced hypomorphic missense mutation in *Zfp335* (R1092W aka *bloto*) that was previously studied in the context of T lymphocyte maturation [23].

We first compared fasting plasma cholesterol levels of 13-week-old *Zfp335*<sup>R1092W/R1092W</sup>, *Zfp335*<sup>R1092W/+</sup>, and wild-type mouse littermates fed a grain-based chow diet. In male mice, increasing copies of *Zfp335*<sup>R1092W</sup> mutations significantly reduced total cholesterol (TC) and HDL cholesterol (HDL) levels (Fig. 3, additive  $p=0.01$



**Fig. 3** 13-week old chow-fed mouse plasma **A** total, **B** HDL, and **C** non-HDL cholesterol levels split by sex and *Zfp335* genotype. Male sample sizes were  $N=12, 15$  (16 for TC), and 10, and female sample sizes were  $N=17, 15$ , and 5 (6 for TC) for wild-type, heterozygotes, and homozygous *Zfp335*<sup>R1092W</sup>, respectively. Values are mean  $\pm$  SD. \* $p < 0.05$ , \*\*\* $p < 0.001$

and  $p=0.0002$ , respectively). As expected based on previous studies, female mice had substantially lower HDL (and total) cholesterol levels than male mice across all *Zfp335* genotype categories. However, associations of *Zfp335* genotype with plasma TC or HDLC were not observed in female mice, with sex\**Zfp335* genotype interactions of  $p=0.15$  and  $p=0.015$ , respectively. Since we observed a significant sex interaction for HDLC levels in the mice, we went back to our CAP participant data and observed a modest association of *ZNF335* levels in control-treated LCLs with CAP baseline HDLC in men ( $p=0.048$ ) but no association in women, with a weak sex\**ZNF335* interaction ( $p=0.09$ , Fig. S1). The directionality of the correlation in men seems consistent with the male mice as well, since lower human LCL gene expression and more copies of the mouse hypomorphic *Zfp335* allele would both be presumed to have less protein activity. There was no correlation of baseline HDLC with control-treated LCL *ZNF335* levels in a sex-combined model, and there was no correlation of baseline LDLC with control-treated LCL *ZNF335* levels in either sex subset or a sex-combined model.


In contrast to HDLC, there were not significant sex differences in the relationship between mouse *Zfp335* genotype and non-HDL cholesterol (TC – HDLC). *Zfp335* genotype was significantly correlated with non-HDL cholesterol in a sex-combined model (additive  $p=0.04$ ) with sex included as a covariate. Though not statistically significant, male mice with one or two copies of the *Zfp335*<sup>R1092W</sup> mutation had lower mean levels of non-HDL cholesterol compared to wild-type littermates. In female mice, the strongest relationship of genotype with cholesterol levels was for non-HDL cholesterol, with a trend for reduced levels with increasing numbers of *Zfp335*<sup>R1092W</sup> mutations (additive  $p=0.08$ ).

### Weight and growth curves

Since we observed that *Zfp335*<sup>R1092W/R1092W</sup> mice appeared smaller on average than their littermates and this phenotype had not been documented for this mutant mouse strain before, we measured their growth curves from 4 to 11 weeks of age on a grain-based chow diet (Fig. S2). Both male (adj.  $p=0.0036$ ) and female (adj.  $p=0.036$ ) *Zfp335*<sup>R1092W/R1092W</sup> mice had significantly reduced growth curves compared to their wild-type littermates. For males (which had a larger sample size than females), *Zfp335*<sup>R1092W/+</sup> animals had growth curves that were intermediate between *Zfp335*<sup>R1092W/R1092W</sup> and *Zfp335*<sup>+/+</sup> males (heterozygote vs. *Zfp335*<sup>R1092W/R1092W</sup> adj.  $p=0.0021$ , heterozygote vs. wild-type adj.  $p=0.0498$ ), with females trending similarly (adj.  $p=0.098$  for both comparisons).

### LDL statin response in *Zfp335* hypomorphic mice

We next wanted to determine if statin efficacy differed between wild-type mice and mice carrying *Zfp335*<sup>R1092W</sup> mutations. To do this, we first fed mice a defined “control” diet (AIN-76A) for 4 weeks, starting at 13 weeks of age. We then transitioned mice to a matched diet supplemented with simvastatin for 4 weeks. We first measured statin-induced changes in non-HDL cholesterol as a crude proxy for mouse LDLC statin response, though non-HDL cholesterol also includes cholesterol found in other lipoprotein particles. Pre-statin non-HDL cholesterol levels (the average of the 15- and 17-week samples; 2 and 4 weeks on AIN-76A diet) were associated with *Zfp335* genotype in male animals alone (additive  $p=0.022$ , Fig. 4A) and in a sex-combined model with sex included as a covariate (additive  $p=0.029$ ), with a similar but non-significant trend in female animals alone (Fig. 4B).



**Fig. 4** Plasma non-HDL cholesterol levels in **A** male and **B** female mice before and during statin diet feeding and **C** statin-induced changes in non-HDL cholesterol split by sex and *Zfp335* genotype. Male sample sizes were  $N=7$ , 15, and 4 and female sample sizes were  $N=12$ , 11, and 5 for wild-type, heterozygotes, and homozygous *Zfp335*<sup>R1092W</sup>, respectively. Values are mean  $\pm$  SD. \* $p < 0.05$

Non-HDL cholesterol levels were reduced after 4 weeks of statin treatment in wild-type mice of both sexes, with wild-type male mice experiencing greater (inter-sex comparison  $p < 0.05$ ) statin-induced reductions in non-HDL cholesterol (Fig. 4A and C,  $-43 \pm 18\%$ , paired  $t$ -test  $p = 0.0016$ ,  $N = 7$ ) than wild-type females (Fig. 4B and C,  $-23 \pm 19\%$ , paired  $t$ -test  $p = 0.0020$ ,  $N = 12$ ). Female *Zfp335*<sup>R1092W/R1092W</sup> mice ( $N = 5$ ) had a significantly blunted non-HDL cholesterol statin response compared to their wild-type littermates (Fig. 4C,  $p = 0.02$ ,  $t$ -test), but this trend was not as strong in male animals.


#### Lipoprotein subfraction statin response

Non-HDL cholesterol is a broad category that includes cholesterol content from lipoprotein particles of various sizes and types. Since LDL cholesterol response was the phenotype for which we saw a correlation with *ZNF335* gene expression statin response in humans, we next performed a more detailed analysis of lipoprotein particles in the LDL size range. We therefore compared the ion mobility-based lipoprotein profiles from wild-type mice before and after they were fed 4 weeks of statin diet (Fig. 5 and Fig. S3). As has been well documented previously, most mouse lipoproteins were in the HDL size range, with much lower concentrations in the size ranges spanning LDL and intermediate density lipoproteins (IDL). Since the lipoprotein subfraction size categorization we used is based on human data, it may not always accurately reflect the type(s) of particles in the corresponding size intervals in mice. In mice, particles in the human very small LDL (designated LDL

IV) particle size range comprise a heterogeneous mixture of species overlapping very large HDL and containing APOE-only lipoproteins [27], so we focused our analyses on small (LDL IIIb [20.49–20.82 nm] and LDL IIIa [20.82–21.41 nm]), medium (LDL IIB [21.41–22 nm] and LDL IIA [22–22.46 nm]), and large (LDL I [22.46–23.33 nm]) LDL size ranges. Wild-type males were the only mice that exhibited robust statin-induced decreases in these LDL particles (Figs. 5A, 6, S3, and S4). For comparison, we also included pre-statin and on-statin human lipoprotein profiles from CAP clinical trial participants (Fig. S5). Similar to male mice, men experienced a significant reduction in lipoprotein particle concentrations in the medium and large LDL to small IDL size range after statin treatment, but unlike the sex difference in statin response in the mice, there were similar reductions in levels of these particles in women.

Male mice carrying one or two copies of the *Zfp335*<sup>R1092W</sup> mutation exhibited significantly blunted LDL statin responses compared to wild-type (Fig. 6,  $p = 0.0011$  and  $p = 0.0010$ , respectively). Female mice of any genotype experienced minimal changes in LDL particle concentrations with statin treatment (Figs. 5B, S3, and S4B); the statin-induced reductions in non-HDL cholesterol in female wild-type and heterozygous mice (Fig. 4C) seemed largely driven by particles in the IDL size range instead (Figs. 5B, S3, and S4B). We also went back to investigate whether there was a significant sex interaction of the original association of LCL *ZNF335* statin response with plasma LDL cholesterol statin response in





**Fig. 5** Statin-induced changes in wild-type **A**  $N=5$  male and **B**  $N=11$  female mouse lipoprotein profiles measured by ion mobility. Mouse profiles were measured before and after 4 weeks of simvastatin-containing diet. The size intervals designating the major lipoprotein subclasses are based on those defined in humans [29]


CAP. This association was also stronger in men than women, though the interaction of sex with gene expression change was not significant ( $p=0.28$ , Fig. S6).

### Discussion

In this study, we show evidence that zinc finger protein 335 (*ZNF335/Zfp335*) is a novel modulator of LDLC statin response. In our *in vitro* experiments, we identify 147 human genes whose *in vitro* statin responses correlate significantly with the *in vivo* plasma LDLC statin responses of the corresponding donors, with *ZNF335* and *CNOT3* topping the list. In our subsequent *in vivo* experiments in mice using a *Zfp335* hypomorphic missense (*Zfp335<sup>R1092W</sup>*) genetic model, we show for the first time that reduced ZFP335 function leads to reduced

plasma cholesterol levels and a blunted LDL response to statin, with significant sex differences.

*ZNF335* resides in the nucleus and regulates transcription, including enhancing transcriptional activation by nuclear hormone receptors [35, 36]. It can activate transcription by recruiting histone methyltransferase complexes to the promoters of genes [32]. *ZNF335* is widely expressed across human tissues [37] and has diverse functions, including roles in neuronal development [32] and T-cell maturation [23, 38, 39]. Variants associated with blood HDL cholesterol, triglyceride, non-HDL cholesterol, and LDL cholesterol levels in GWAS are located in and near the *ZNF335* gene [40]. Though the dominant contribution to this association signal is likely the nearby *PLTP* (phospholipid transfer protein) gene, which has



**Fig. 6** Statin-induced changes in male mouse LDL particle concentrations as measured by ion mobility. Sample sizes were  $N=5, 12,$  and  $5$  for wild-type, heterozygotes, and homozygous *Zfp335*<sup>R1092W</sup>, respectively. Values are mean  $\pm$  SD. Genotypes were compared using two-way ANOVA with Tukey's multiple comparison test.  $**p < 0.01$

well-established roles in lipoprotein metabolism [41], it is possible that *ZNF335* also contributes to this signal. Here, we demonstrate for the first time that *ZNF335* influences plasma cholesterol levels and statin response. Though the correlation between statin-induced changes in LCL *ZNF335* gene expression with donor plasma LDL-C statin response is quite modest, it is still a significant contribution given that only about 5% of the variation in statin response has been explained by genetic factors to date [5]. Further investigation is needed to determine which, if any, genes affecting cholesterol metabolism may be regulated transcriptionally by *ZNF335* and to determine the mechanism by which *ZNF335* influences plasma cholesterol phenotypes.

*ZNF335* physically interacts with members of the CCR4-NOT (carbon catabolite repression 4–negative on TATA-less) complex, including *CNOT6* (aka CCR4) and *CNOT9* (aka RCD1) [30]. The CCR4-NOT complex can regulate gene expression via several mechanisms, including transcription and mRNA deadenylation [42]. Though a physical interaction has not been reported between *ZNF335* and the CCR4-NOT complex regulatory subunit *CNOT3*, here we observe strong correlations between

their statin-induced gene expression changes, indicating they may be co-regulated by some of the same statin-responsive factors. *ZNF335* and *CNOT3* expression changes in LCLs also show similar strength and directionality of correlation with plasma LDL-C statin responses of the corresponding statin clinical trial participants, so they may be functionally related, and it is plausible that they could be acting as components of the same complex. *CNOT3* has been previously reported to have roles in regulation of lipid and glucose metabolism and to be differentially regulated in fasted and fed states based on studies in *CNOT3* haplodeficient mice [33]. A functional link between *ZNF335* and *CNOT3* could be further explored in future studies, as could effects of conditional knockdowns of both genes on lipids and statin response.

As with *Cnot3* [31], knockout of *Zfp335* in mice is embryonic lethal [32]. Thus, to study the potential role of *Zfp335* in cholesterol metabolism and statin response, here we use mice carrying a hypomorphic mutation (R1092W aka bloto) in *Zfp335*. Previous studies using *Zfp335*<sup>R1092W</sup> mice have shown that the R1092W mutation in the twelfth zinc finger of ZFP335 impairs its ability to bind a subset of its DNA targets [23] by modifying one of its two known DNA binding domains [43]. Previous studies of this genetic mouse model have focused on the role of ZFP335 in T-cell development, but here we also show that the mutation has an impact on body weight, blood cholesterol levels, and blood cholesterol statin response. It is interesting to note that for some of these phenotypes (e.g., body weight and HDL-cholesterol levels in male mice), heterozygotes display intermediate phenotypes typical of an additive relationship, while for other phenotypes, heterozygotes are phenotypically more similar to wild-type animals (e.g., non-HDL cholesterol statin response in female animals) or to *Zfp335*<sup>R1092W/R1092W</sup> homozygous mutant animals (e.g., LDL particle statin response in male animals). Perhaps ZFP335 acts as a component of distinct complexes to elicit some of these phenotypes, which could explain why the same genetic model is not common to them all, or perhaps the fact that the mutation is only a partial loss of function mutation plays a role.

We observe striking sex differences when studying the impact of the hypomorphic missense *Zfp335* mutation on mouse HDL cholesterol levels and LDL statin response. Similar, though less significant, sex differences are also observed for the human LCL *ZNF335* statin response correlation with plasma LDL-C statin response and the correlation of LCL *ZNF335* levels with plasma HDL-cholesterol levels, showing inter-species replication despite known differences in lipoprotein metabolism and profiles between the species. The differences in the

impact of the *Zfp335* mutation between sexes could be related to sex-specific differences in the hormonal milieu, given *ZNF335*'s role in nuclear hormone receptor signaling, but the hormone(s) involved and which of their downstream targets regulate cholesterol levels and statin response remain to be elucidated.

Sex differences are also observed in the lipoprotein profiles and lipoprotein response to statin in wild-type mice. Wild-type female mice have less plasma HDL cholesterol and total cholesterol than males (in agreement with other studies in the literature which have studied sexual dimorphism in mouse lipoprotein metabolism [44, 45]) and exhibit a blunted non-HDL cholesterol statin response compared to males. In this study, though male mice experience significant reductions in LDL with statin treatment, the statin-induced reduction in non-HDL cholesterol in female mice appears to be driven mainly by larger particles (including IDL) rather than LDL. In contrast, in humans, women experience LDL reductions quite similar to men.

This study illustrates the utility of clinical trial participant-derived cell lines in identifying novel genetic factors involved in drug response phenotypes. The correlation of in vitro drug-induced transcriptomic changes in cell lines to in vivo drug response phenotypes from the corresponding participants during the clinical trial can be a powerful approach, even in the transformed cell lines used here. Though LCLs may not always mirror what might be happening in hepatocytes, they express cholesterol biosynthesis enzymes (including *HMGCR*), *LDLR*, and many other key cholesterol metabolism genes and have been previously used to identify novel modulators of statin response that have been validated in human hepatocyte cell lines and/or in vivo [8, 10, 11]. The emergence of induced pluripotent stem cells as a model system holds additional promise in investigating genes that are not as universally expressed across cell types as the genes we identify here, since they can be differentiated into a variety of specialized cell types.

## Conclusions

Here, we identify a novel role for *ZNF335* in modifying circulating levels of LDL-cholesterol and LDL-cholesterol statin response by leveraging naturally occurring variation in *ZNF335* expression in statin clinical trial participant-derived cell lines and a hypomorphic missense mouse model for *ZNF335*. Thus, it is possible that inter-individual variation in *ZNF335* expression contributes to differences in cholesterol levels and statin response in humans.

## Abbreviations

LDLC	Low-density lipoprotein cholesterol
CVD	Cardiovascular disease

GWAS	Genome-wide association study
SREBF2	Sterol regulatory element binding transcription factor 2
ZNF335	Zinc finger protein 335 (human)
ZFP335	Zinc finger protein 335 (murine)
LCL	Lymphoblastoid cell line
CAP	Cholesterol and Pharmacogenetics
EBV	Epstein-Barr virus
AfrAm	African American
BMI	Body mass index
RNA-seq	Ribonucleic acid sequencing
ENU	N-ethyl-N-nitrosourea
TC	Total cholesterol
HDLC	High-density lipoprotein cholesterol
IM	Ion mobility
CCR4-NOT	Carbon catabolite repression-negative on TATA-less
FDR	False discovery rate
ANOVA	Analysis of variance
SD	Standard deviation
Q	False discovery rate adjusted <i>p</i> value
F	Female
M	Male
IDL	Intermediate density lipoprotein
VLDL	Very low density lipoprotein
PLTP	Phospholipid transfer protein

## Supplementary Information

The online version contains supplementary material available at <https://doi.org/10.1186/s13073-024-01366-9>.

Additional file 1: Supplementary tables. Table S1 CAP LCL gene expression statin responses. Table S2 Spearman correlations of statin-induced gene expression changes in LCLs with plasma LDL-cholesterol statin response in the corresponding CAP participants

Additional file 2: Supplementary figures. Fig. S1 Correlation of pre-statin plasma HDL-cholesterol with *ZNF335* LCL gene expression levels from the corresponding donors split by sex ( $N = 211$  men,  $N = 216$  women). Fig. S2 Mouse growth curves split by *Zfp335* genotype and sex. Adjusted *p* values were calculated using the Compare Groups of Growth Curves (CGGC) method with 10,000 permutation tests. Male sample sizes were  $N = 7, 19,$  and  $7$  and female sample sizes were  $N = 7, 8,$  and  $4$  for wild-type, heterozygotes, and homozygous *Zfp335*<sup>R1092W</sup>, respectively. Values are mean  $\pm$  SD. \* $p < 0.05$ , \*\* $p < 0.01$ . Fig. S3 Statin-induced changes in wild-type  $N = 5$  male and  $N = 11$  female mouse lipoprotein profiles measured by ion mobility. Mouse profiles were measured before and after 4 weeks of simvastatin-containing diet. Fig. S4 Statin-induced changes in (A) male and (B) female mouse plasma lipoprotein composition split by *Zfp335* genotype as measured by ion mobility. Male sample sizes were  $N = 5, 12,$  and  $5$  and female sample sizes were  $N = 11, 11,$  and  $5$  for wild-type, heterozygotes, and homozygous *Zfp335*<sup>R1092W</sup>, respectively. The size intervals designating the major lipoprotein subclasses are based on those defined in humans [29]. Values are mean  $\pm$  SD. Fig. S5 Ion mobility lipoprotein profiles for  $N = 803$  CAP participants before and after 6 weeks of 40 mg/day simvastatin treatment. Profiles shown are (A) pre-statin split by sex and race/ethnicity, (B) on-statin split by sex and race/ethnicity, (C) pre- and on-statin for all, and (D) pre-statin and on-statin split by sex.  $N = 142$  for female African American,  $N = 142$  for male African American,  $N = 243$  for female white, and  $N = 276$  for male white participants. Fig. S6 Correlation of plasma LDL-cholesterol statin response with *ZNF335* LCL gene expression statin response from the corresponding donors split by sex ( $N = 211$  men,  $N = 216$  women)

## Acknowledgements

We thank the CAP participants for their contributions to this study. We thank Arnie Acosta for culturing LCLs, Bahareh Sahami for lipid measurements, Jennifer Beckstead for mouse ion mobility measurements, Jinping An for conducting a pilot mouse study, Quest Diagnostics for providing the CAP IM data, and numerous UC Berkeley undergraduate volunteers for their assistance with this project. The mice used for this work were housed in a UCSF MLK Cores

Research Facility. We fondly remember the late Dr. Debbie Nickerson and her contributions to this work while leading the Northwest Genomics Center.

#### Authors' contributions

ET helped to design the experiments, analyzed the data, and drafted the manuscript. FYT conducted the mouse experiments, maintained the mouse line, and organized mouse data. YQ assisted with mouse experiments. KS and DN cultured LCLs and extracted RNA. SMK supervised the mouse Liasys and ion mobility analysis. NY assisted with mouse experiments. JO performed mouse ion mobility analysis. BYH and JGC generated the *Zfp335*<sup>R1092W</sup> (bloto) mutant mouse line. YIC and JIR contributed to the creation of the CAP LCLs. RMK conducted the CAP clinical trial, helped interpret the ion mobility results, and revised the manuscript. MWM supervised the work and revised the manuscript. All authors read and approved the final manuscript.

#### Funding

This project was supported by NIH U19 HL069757, NIH P50 GM115318, AHA 15POST21880006, and AHA 12POST10430005. This study was also supported in part by the NIH Pharmacogenomics Research Network (PGRN) RNA Sequencing Project. Establishment and storage of the CAP LCLs was partially supported by NIH P30 DK063491. The funders had no role in the design of the study, the collection, analysis, or interpretation of the data, or in writing the manuscript.

#### Availability of data and materials

The CAP (Cholesterol and Pharmacogenetics) RNA-seq and phenotype data used for these analyses are available in dbGaP (database of Genotypes and Phenotypes) under accession phs000481.v3.p2, [https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study\\_id=phs000481.v3.p2](https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000481.v3.p2) [46].

#### Declarations

##### Ethics approval and consent to participate

Written informed consent was obtained from all human participants, and the study was approved by the institutional review boards of the University of California, San Francisco and the University of California, Los Angeles. Mouse studies were approved by the Children's Hospital & Research Center Oakland and the University of California, San Francisco Institutional Animal Care and Use Committees. This study was conducted in compliance with the Declaration of Helsinki.

##### Consent for publication

Not applicable.

##### Competing interests

The authors declare that they have no competing interests.

##### Author details

<sup>1</sup>Department of Pediatrics, University of California San Francisco, Oakland, CA, USA. <sup>2</sup>Howard Hughes Medical Institute, Department of Microbiology and Immunology, University of California San Francisco, San Francisco, CA, USA. <sup>3</sup>The Institute for Translational Genomics and Population Sciences, Department of Pediatrics, The Lundquist Institute for Biomedical Innovation at Harbor-UCLA Medical Center, Torrance, CA, USA. <sup>4</sup>Department of Medicine, University of California San Francisco, Oakland, CA, USA.

Received: 27 June 2023 Accepted: 18 July 2024

Published online: 26 July 2024

#### References

- Arnett DK, Blumenthal RS, Albert MA, Buroker AB, Goldberger ZD, Hahn EJ, Himmelfarb CD, Khera A, Lloyd-Jones D, McEvoy JW, et al. 2019 ACC/AHA guideline on the primary prevention of cardiovascular disease: a report of the American College of Cardiology/American Heart Association Task Force on Clinical Practice Guidelines. *Circulation*. 2019;140:e596–646.
- Simon JA, Lin F, Hulley SB, Blanche PJ, Waters D, Shiboski S, Rotter JI, Nickerson DA, Yang H, Saad M, Krauss RM. Phenotypic predictors of response to simvastatin therapy among African-Americans and Caucasians: the Cholesterol and Pharmacogenetics (CAP) Study. *Am J Cardiol*. 2006;97:843–50.
- Chasman DI, Giulianini F, Macfadyen J, Barratt BJ, Nyberg F, Ridker PM. Genetic determinants of statin-induced low-density lipoprotein cholesterol reduction: the justification for the use of statins in prevention: an intervention trial evaluating rosuvastatin (JUPITER) trial. *Circ Cardiovasc Genet*. 2012;5:257–64.
- Deshmukh HA, Colhoun HM, Johnson T, McKeigue PM, Betteridge DJ, Durrington PN, Fuller JH, Livingstone S, Charlton-Menys V, Neil A, et al. Genome-wide association study of genetic determinants of LDL-c response to atorvastatin therapy: importance of Lp(a). *J Lipid Res*. 2012;53:1000–11.
- Postmus I, Trompet S, Deshmukh HA, Barnes MR, Li X, Warren HR, Chasman DI, Zhou K, Arseneault BJ, Donnelly LA, et al. Pharmacogenetic meta-analysis of genome-wide association studies of LDL cholesterol response to statins. *Nat Commun*. 2014;5:5068.
- Oni-Orisan A, Haldar T, Ranatunga DK, Medina MW, Schaefer C, Krauss RM, Iribarren C, Risch N, Hoffmann TJ. The impact of adjusting for baseline in pharmacogenomic genome-wide association studies of quantitative change. *NPJ Genom Med*. 2020;5:1.
- Brown MS, Goldstein JL. The SREBP pathway: regulation of cholesterol metabolism by proteolysis of a membrane-bound transcription factor. *Cell*. 1997;89:331–40.
- Medina MW, Theusch E, Naidoo D, Bauzon F, Stevens K, Mangravite LM, Kuang YL, Krauss RM. RHOA is a modulator of the cholesterol-lowering effects of statin. *PLoS Genet*. 2012;8:e1003058.
- Mangravite LM, Engelhardt BE, Medina MW, Smith JD, Brown CD, Chasman DI, Mecham BH, Howie B, Shim H, Naidoo D, et al. A statin-dependent QTL for GATM expression is associated with statin-induced myopathy. *Nature*. 2013;502:377–80.
- Kim K, Theusch E, Kuang YL, Dose A, Mitchel K, Cubitt C, Chen YI, Krauss RM, Medina MW. ZNF542P is a pseudogene associated with LDL response to simvastatin treatment. *Sci Rep*. 2018;8:12443.
- Mitchel K, Theusch E, Cubitt C, Dosé AC, Stevens K, Naidoo D, Medina MW. RP1–13D10.2 is a novel modulator of statin-induced changes in cholesterol. *Circ Cardiovasc Genet*. 2016;9:223–30.
- Friedewald WT, Levy RI, Fredrickson DS. Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge. *Clin Chem*. 1972;18:499–502.
- Medina MW, Gao F, Ruan W, Rotter JI, Krauss RM. Alternative splicing of 3-hydroxy-3-methylglutaryl coenzyme A reductase is associated with plasma low-density lipoprotein cholesterol response to simvastatin. *Circulation*. 2008;118:355–62.
- Theusch E, Kim K, Stevens K, Smith JD, Chen YI, Rotter JI, Nickerson DA, Medina MW. Statin-induced expression change of INSIG1 in lymphoblastoid cell lines correlates with plasma triglyceride statin response in a sex-specific manner. *Pharmacogenomics J*. 2017;17:222–9.
- Parkhomchuk D, Borodina T, Amstislavskiy V, Banaru M, Hallen L, Krobitsch S, Lehrach H, Soldatov A. Transcriptome analysis by strand-specific sequencing of complementary DNA. *Nucleic Acids Res*. 2009;37:e123.
- Theusch E, Chen YI, Rotter JI, Krauss RM, Medina MW. Genetic variants modulate gene expression statin response in human lymphoblastoid cell lines. *BMC Genomics*. 2020;21:555.
- Kim D, Pertea G, Trapnell C, Pimentel H, Kelley R, Salzberg SL. TopHat2: accurate alignment of transcriptomes in the presence of insertions, deletions and gene fusions. *Genome Biol*. 2013;14:R36.
- Zerbino DR, Achuthan P, Akanni W, Amode MR, Barrell D, Bhai J, Billis K, Cummins C, Gall A, Giron CG, et al. Ensembl 2018. *Nucleic Acids Res*. 2018;46:D754–61.
- Arvey A, Tempera I, Tsai K, Chen HS, Tikhmyanova N, Klichinsky M, Leslie C, Lieberman PM. An atlas of the Epstein-Barr virus transcriptome and epigenome reveals host-virus regulatory interactions. *Cell Host Microbe*. 2012;12:233–45.
- Anders S, Pyl PT, Huber W. HTSeq—a Python framework to work with high-throughput sequencing data. *Bioinformatics*. 2015;31:166–9.
- Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-Seq data with DESeq2. *Genome Biology*. 2014;15:550.

22. Benjamini Y, Hochberg Y. Controlling the false discovery rate: a practical and powerful approach to multiple testing. *J Royal Stat Soc Ser B*. 1995;57:289–300.
23. Han BY, Wu S, Foo CS, Horton RM, Jenne CN, Watson SR, Whittle B, Goodnow CC, Cyster JG. Zinc finger protein Zfp335 is required for the formation of the naive T cell compartment. *Elife*. 2014;3:e03549.
24. Elso CM, Roberts LJ, Smyth GK, Thomson RJ, Baldwin TM, Foote SJ, Handman E. Leishmaniasis host response loci (Imr1-3) modify disease severity through a Th1/Th2-independent pathway. *Genes Immun*. 2004;5:93–100.
25. Baldwin T, Sakthianandeswaren A, Curtis JM, Kumar B, Smyth GK, Foote SJ, Handman E. Wound healing response is a major contributor to the severity of cutaneous leishmaniasis in the ear model of infection. *Parasite Immunol*. 2007;29:501–13.
26. Chiu S, Bergeron N, Williams PT, Bray GA, Sutherland B, Krauss RM. Comparison of the DASH (Dietary Approaches to Stop Hypertension) diet and a higher-fat DASH diet on blood pressure and lipids and lipoproteins: a randomized controlled trial. *Am J Clin Nutr*. 2016;103:341–7.
27. Qin Y, Ting F, Kim MJ, Strelnikov J, Harmon J, Gao F, Dose A, Teng BB, Alipour MA, Yao Z, et al. Phosphatidylinositol-(4,5)-bisphosphate regulates plasma cholesterol through LDL (low-density lipoprotein) receptor lysosomal degradation. *Arterioscler Thromb Vasc Biol*. 2020;40:1311–24.
28. Musunuru K, Orho-Melander M, Caulfield MP, Li S, Salameh WA, Reitz RE, Berglund G, Hedblad B, Engström G, Williams PT, et al. Ion mobility analysis of lipoprotein subfractions identifies three independent axes of cardiovascular risk. *Arterioscler Thromb Vasc Biol*. 2009;29:1975–80.
29. Caulfield MP, Li S, Lee G, Blanche PJ, Salameh WA, Benner WH, Reitz RE, Krauss RM. Direct determination of lipoprotein particle sizes and concentrations by ion mobility analysis. *Clin Chem*. 2008;54:1307–16.
30. Garapaty S, Mahajan MA, Samuels HH. Components of the CCR4-NOT complex function as nuclear hormone receptor coactivators via association with the NRC-interacting Factor NIF-1. *J Biol Chem*. 2008;283:6806–16.
31. Neely GG, Kuba K, Cammarato A, Isobe K, Amann S, Zhang L, Murata M, Elmén L, Gupta V, Arora S, et al. A global in vivo *Drosophila* RNAi screen identifies NOT3 as a conserved regulator of heart function. *Cell*. 2010;141:142–53.
32. Yang YJ, Baltus AE, Mathew RS, Murphy EA, Evrony GD, Gonzalez DM, Wang EP, Marshall-Walker CA, Barry BJ, Murn J, et al. Microcephaly gene links trithorax and REST/NRSF to control neural stem cell proliferation and differentiation. *Cell*. 2012;151:1097–112.
33. Morita M, Oike Y, Nagashima T, Kadomatsu T, Tabata M, Suzuki T, Nakamura T, Yoshida N, Okada M, Yamamoto T. Obesity resistance and increased hepatic expression of catabolism-related mRNAs in *Cnot3*<sup>+/-</sup> mice. *EMBO J*. 2011;30:4678–91.
34. Suzuki T, Kikuguchi C, Nishijima S, Nagashima T, Takahashi A, Okada M, Yamamoto T. Postnatal liver functional maturation requires *Cnot* complex-mediated decay of mRNAs encoding cell cycle and immature liver genes. *Development*. 2019;146:dev168146.
35. Garapaty S, Xu CF, Trojer P, Mahajan MA, Neubert TA, Samuels HH. Identification and characterization of a novel nuclear protein complex involved in nuclear hormone receptor-mediated gene regulation. *J Biol Chem*. 2009;284:7542–52.
36. Mahajan MA, Murray A, Samuels HH. NRC-interacting factor 1 is a novel cotransducer that interacts with and regulates the activity of the nuclear hormone receptor coactivator NRC. *Mol Cell Biol*. 2002;22:6883–94.
37. GTEx Consortium, The GTEx Consortium atlas of genetic regulatory effects across human tissues. *Science*. 2020;369:1318–1330.
38. Ratiu JJ, Barclay WE, Lin E, Wang Q, Wellford S, Mehta N, Harnois MJ, DiPalma D, Roy S, Contreras AV, et al. Loss of Zfp335 triggers cGAS/STING-dependent apoptosis of post- $\beta$  selection thymocytes. *Nat Commun*. 2022;13:5901.
39. Liu H, Wang X, Ding R, Jiao A, Zheng H, Zhang C, Feng Z, Su Y, Yang X, Lei L, et al. The transcription factor Zfp335 promotes differentiation and persistence of memory CD8(+) T cells by regulating TCF-1. *J Immunol*. 2022;209:886–95.
40. Graham SE, Clarke SL, Wu KH, Kanoni S, Zajac GJM, Ramdas S, Surakka I, Ntalla I, Vedantam S, Winkler TW, et al. The power of genetic diversity in genome-wide association studies of lipids. *Nature*. 2021;600:675–9.
41. Jiang XC, Jin W, Hussain MM. The impact of phospholipid transfer protein (PLTP) on lipoprotein metabolism. *Nutr Metab (Lond)*. 2012;9:75.
42. Shirai YT, Suzuki T, Morita M, Takahashi A, Yamamoto T. Multifunctional roles of the mammalian CCR4-NOT complex in physiological phenomena. *Front Genet*. 2014;5:286.
43. Han BY, Foo CS, Wu S, Cyster JG. The C2H2-ZF transcription factor Zfp335 recognizes two consensus motifs using separate zinc finger arrays. *Genes Dev*. 2016;30:1509–14.
44. Norheim F, Hasin-Brumshtein Y, Vergnes L, Chella Krishnan K, Pan C, Seldin MM, Hui ST, Mehrabian M, Zhou Z, Gupta S, et al. Gene-by-sex interactions in mitochondrial functions and cardio-metabolic traits. *Cell Metab*. 2019;29:932–949.e934.
45. Bruell JH, Daroczy AF, Hellerstein HK. Strain and sex differences in serum cholesterol levels of mice. *Science*. 1962;135:1071–2.
46. CAP Investigators. Cholesterol and Pharmacogenetics (CAP) Study. The database of Genotypes and Phenotypes (dbGaP), [https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study\\_id=phs000496.v1.p1](https://www.ncbi.nlm.nih.gov/projects/gap/cgi-bin/study.cgi?study_id=phs000496.v1.p1).

## Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.