

UC Davis

UC Davis Previously Published Works

Title

Corrigendum: Characterization of the Spatial and Temporal Expression of Two Soybean miRNAs Identifies SCL6 as a Novel Regulator of Soybean Nodulation.

Permalink

<https://escholarship.org/uc/item/1sb397th>

Authors

Hossain, Md
Hoang, Nhung
Yan, Zhe
et al.

Publication Date

2019

DOI

10.3389/fpls.2019.01692

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



Corrigendum: Characterization of the Spatial and Temporal Expression of Two Soybean miRNAs Identifies SCL6 as a Novel Regulator of Soybean Nodulation

Md Shakhawat Hossain^{1†}, Nhung T. Hoang^{1†}, Zhe Yan¹, Katalin Tóth¹, Blake C. Meyers² and Gary Stacey^{1*}

¹ C.S. Bond Life Science Center, Divisions of Plant Sciences and Biochemistry, University of Missouri, Columbia, MO, United States, ² Donald Danforth Plant Science Center, St. Louis, MO, United States

OPEN ACCESS

Edited and reviewed by:

Aardra Kachroo,
University of Kentucky, United States

*Correspondence:

Gary Stacey
staceyg@missouri.edu

[†]These authors have contributed
equally to this work

Specialty section:

This article was submitted to
Plant Microbe Interactions,
a section of the journal
Frontiers in Plant Science

Received: 24 October 2019

Accepted: 29 November 2019

Published: 12 February 2020

Citation:

Hossain MS, Hoang NT, Yan Z,
Tóth K, Meyers BC and Stacey G
(2020) Corrigendum: Characterization
of the Spatial and Temporal
Expression of Two Soybean miRNAs
Identifies SCL6 as a Novel Regulator of
Soybean Nodulation.
Front. Plant Sci. 10:1692.
doi: 10.3389/fpls.2019.01692

Keywords: miRNA, miR171, GRAS TF, Scarecrow like-6, NSP2, nodulation, symbiosis, soybean

A Corrigendum on

Characterization of the Spatial and Temporal Expression of Two Soybean miRNAs Identifies SCL6 as a Novel Regulator of Soybean Nodulation.

By Hossain MdS, Hoang NT, Yan Z, Tóth K, Meyers BC and Stacey G (2019). *Front. Plant Sci.* 10:475. doi: 10.3389/fpls.2019.00475

It was brought to our attention that, in the original article, inappropriate manipulation of **Figures 1** and **5** took place, specifically, **Figures 1B, 1C, 1D, 1E, 1F, 1G** and **5B, 5C, 5D, 5E, 5F, 5G**. Upon review, we have agreed that these figures be removed from the article. The manipulation of the figures involved an effort to increase the contrast of the images to make them more presentable.

Although this did not change the interpretation of the data, the manipulation was clearly against our lab policy and the policies of the journal. Given that other data in the paper (e.g., qPCR and other experiments) confirm the data presented in these figures, we feel that the conclusions of the paper remain valid even with the figures removed.

To avoid any additional questions or concerns, we have also corrected **Figures 2D, 2E** and **2F**, as well as **Figures 4D, 4E, 4F**, and **4G**, and replaced these figures with original images. The corrected figures appear below. Due to the changes mentioned above, a correction has been made to the **Results**, subsection **Gma-miR171o and Gma-miR171q Exhibit Distinct Expression Patterns in Response to Bacterial Infection:**

“To understand the symbiotic role of gma-miR171o and gma-miR171q in soybean, we measured the relative expression level of these two miRNAs in nodules 3 weeks post-

inoculation with *B. japonicum*, as compared to uninfected root tissues (**Figure 1A**). Interestingly, these two miRNAs showed opposite expression patterns in response to *B. japonicum* infection (**Figure 1A**); *gma-miR171o* expression was suppressed upon bacterial infection, while *gma-miR171q* was induced.”

A correction has also been made to the **Results**, subsection **Gene Expression and Promoter Localization of *GmSCL6-1* and *GmNSP2.1* Are Inversely Correlated With *Gma-miR171o* and *Gma-miR171q***. Paragraph two has been removed entirely.

The authors apologize for these errors and state that they do not change the scientific conclusions of the article in any way. The original article has been updated.

Copyright © 2020 Hossain, Hoang, Yan, Tóth, Meyers and Stacey. This is an open-access article distributed under the terms of the Creative Commons Attribution License (CC BY). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.




FIGURE 1 | Expression patterns of *gma-miR171o* and *gma-miR171q* during symbiosis with *B. japonicum*.




FIGURE 2 | Overexpression of *gma-miR171o* and *gma-miR171q* inhibits soybean nodulation.




FIGURE 4 | *GmSCL6-1* and *GmNSP2.1* are required for soybean nodulation.




FIGURE 5 | Expression analysis of target genes, *GmSCL6-1* and *GmNSP2.1* in soybean hairy root transgenic tissues.