

Lawrence Berkeley National Laboratory

LBL Publications

Title

PRODUCTION OF PI-MESONS IN NUCLEON-NUCLEON COLLISIONS

Permalink

<https://escholarship.org/uc/item/1sc4555j>

Author

Brueckner, Keith Allan

Publication Date

1950-07-20

UNIVERSITY OF CALIFORNIA - BERKELEY

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

RADIATION LABORATORY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

Production of π -mesons in Nucleon-Nucleon Collisions

Keith Allan Brueckner

July 20, 1950

Berkeley, California

-2-

INSTALLATION:	No. of Copies
Argonne National Laboratory	8
Armed Forces Special Weapons Project	1
Atomic Energy Commission - Washington	2
Battelle Memorial Institute	1
Brush Beryllium Company	1
Brookhaven National Laboratory	4
Bureau of Medicine and Surgery	1
Bureau of Ships	1
Carbide and Carbon Chemicals Division (K-25 Plant)	4
Carbide and Carbon Chemicals Division (Y-12 Plant)	4
Chicago Operations Office	1
Columbia University (J. R. Dunning)	1
Columbia University (G. Failla)	1
Dow Chemical Company	1
H. K. Ferguson Company	1
General Electric, Richland	3
Harshaw Chemical Corporation	1
Idaho Operations Office	1
Iowa State College	2
Kansas City Operations Branch	1
Kellex Corporation	2
Knolls Atomic Power Laboratory	4
Los Alamos Scientific Laboratory	3
Mallinckrodt Chemical Works	1
Massachusetts Institute of Technology (A. Gaudin)	1
Massachusetts Institute of Technology (A. R. Kaufmann)	1
Mound Laboratory	3
National Advisory Committee for Aeronautics	1
National Bureau of Standards	3
Naval Radiological Defense Laboratory	2
New Brunswick Laboratory	1
New York Operations Office	3
North American Aviation, Inc.	1
Oak Ridge National Laboratory	8
Patent Branch (Washington)	1
Rand Corporation	1
Sandia Corporation	1
Santa Fe Operations Office	2
Sylvania Electric Products, Inc.	1
Technical Information Division (Oak Ridge)	15
USAF, Air Surgeon (Lt. Col. R. H. Blount)	1
USAF, Director of Armament (Captain C. I. Browne)	1
USAF, Director of Research and Development (Col. R. J. Mason, Fred W. Bruner)	2
USAF, Eglin Air Force Base (Major A. C. Field)	1
USAF, Kirtland Air Force Base (Col. Marcus F. Cooper)	1
USAF, Maxwell Air Force Base (Col. F. N. Moyers)	1
USAF, NEPA Office	2
USAF, Office of Atomic Energy (Col. H. C. Donnelly, A. A. Fickel)	2
USAF, Offutt Air Force Base (Col. H. R. Sullivan, Jr.)	1
USAF, Wright-Patterson Air Force Base (Rodney Nudenberg)	1

INSTALLATION	No. of Copies
U. S. Army, Atomic Energy Branch (Lt. Col. A. W. Betts)	1
U. S. Army, Army Field Forces (Captain James Kerr)	1
U. S. Army, Commanding General, Chemical Corps Technical Command (Col. John A. MacLaughlin thru Mrs. Georgia S. Benjamin)	1
U. S. Army, Chief of Ordnance (Lt. Col. A. R. Del Campo)	1
U. S. Army, Commanding Officer, Watertown Arsenal (Col. Carroll H. Deitrick)	1
U. S. Army, Director of Operations Research (Dr. Ellis Johnson)	1
U. S. Army, Office of Engineers (Allen O'Leary)	1
U. S. Army, Office of the Chief Signal Officer (Curtis T. Clayton thru Maj. George C. Hunt)	1
U. S. Army, Office of the Surgeon General (Col. W. S. Stone)	1
U. S. Geological Survey (T. B. Nolan)	2
USAF, Director of Plans and Operations (Col. R.L. Applegate)	1
U. S. Public Health Service	1
University of California at Los Angeles	1
University of California Radiation Laboratory	5
University of Rochester	2
University of Washington	1
Western Reserve University	2
Westinghouse Electric Company	4
Naval Medical Research Institute	1
University of Rochester	1
California Institute of Technology (R. F. Bacher)	1
TOTAL	139

Information Division
Radiation Laboratory
University of California
Berkeley, California

Production of π -mesons in Nucleon-Nucleon Collisions

Keith Allan Brueckner

July 20, 1950

ABSTRACT

The production of π -mesons in nucleon-nucleon collisions as predicted by scalar meson theory, pseudoscalar meson theory with pseudoscalar and pseudovector coupling, and vector theory with vector coupling is compared with the experimental results of the workers at Berkeley. The calculations are made on the basis of 3rd order perturbation theory using the methods of Feynman and Dyson and also using a phenomenological treatment of the nucleon-nucleon interaction. Account is taken of the fact that the final nucleons are not in plane wave states. It is shown that pseudoscalar theory with pseudovector coupling gives qualitative agreement with experiment.

I Introduction

A problem of considerable interest in meson theory is the production of mesons in nucleon-nucleon collisions. The intimate connection predicted by meson theory between this process and ordinary nucleon-nucleon scattering provided an excellent and basic opportunity for testing the fundamental assumptions of meson theory. The meson theory of nuclear forces assumes that π -mesons, found to interact strongly with nuclei, are responsible for the coupling between nucleons. Since this coupling via the meson field implies the existence of virtual mesons in the mutual field of two nucleons, it should be possible, if sufficient energy is available, for virtual mesons to be materialized as free and observable particles. A comparison of the predictions of the theory with the experimental measurements of nucleon-nucleon scattering and of meson production should indicate whether this basic assumption is quantitatively correct.

Experiments are now being carried out at Berkeley¹ which give information about the production of charged and neutral mesons in neutron-proton and proton-proton collisions. It is of interest to consider in a systematic manner the theory of the production in order to understand what can be learned from these experiments.

The production of mesons in nucleon-nucleon collisions has been studied in considerable detail by a number of theoretical workers.² The most thorough theoretical analyses which have been made can be divided into two types: 1) an application of the meson field theory of nuclear forces to the problem considered as a third order process, and 2) an attempt to describe phenomenologically the scattering of the nucleons associated with the meson emission and only the meson emission itself by meson field theory. The first method suffers from the well-known failure of meson theory to

describe nuclear forces in more than a qualitative way and from the more general failure of perturbation theory in the weak coupling approximation applied to problems in which the coupling is not weak. This method can, however, be applied rigorously in the lowest order and with this limitation gives a logically complete description of the process. Also, because of the close relation between high energy nucleon-nucleon scattering (virtual meson exchanges) and meson production (virtual meson exchanges together with a real meson emission), a theory which gives qualitatively correct results for the former process might be expected to be correct to a similar approximation for the latter.

The second method based on a more phenomenological approach describes the meson emission on the basis of field theory but separates the nucleon-nucleon scattering, which gives the momentum transfer necessary for over-all energy and momentum conservation, and attempts to describe this in terms of the experimentally measured potentials. This method is inadequate in as far as processes can occur in the meson production which cannot be described in terms of the scattering process preceded or followed by meson emission. Such processes are for example the interruption of a virtual meson exchange by a real meson emission. These processes are equivalent to scatterings which take place far off the energy shell, i.e., where energy and momentum are related very differently from the relationship for free particles. Only if such processes give unimportant contributions can the phenomenological approach be approximately correct. However, if this assumption is made, the calculation can be made in a manner much more independent of meson field theory than is the third order perturbation calculation.

The rigorous treatment of meson production as a third order process

-6-

is complicated by mathematical difficulties in integrating the cross-sections over the momenta of the final nucleons to give a result which depends only on the meson momentum. This problem has been partially solved by Cecille Morette² for pseudoscalar mesons with pseudoscalar coupling who, however, ignored the effects of the Pauli principle, and so introduced a rather large error (about 50%) in the cross-section near threshold. Her expressions for the cross-sections also have been averaged over meson and nucleon charges and so do not separately give the cross-sections for charged and neutral mesons for neutron-proton and proton-proton collisions. Since these separate quantities are those which can be determined experimentally, it is of interest to calculate them. We therefore have in Section II considered the calculation of the transition matrix elements on the basis of rigorous third order perturbation theory, and obtained the expressions for scalar theory, pseudoscalar theory with pseudoscalar and pseudovector coupling, and vector theory with vector coupling. The calculations are made in the center-of-mass system for energies near threshold where the velocities of the final particles are small. Corrections of the order of v^2/c^2 for the final nucleons and meson are neglected, so the results are only applicable for incident nucleon energies of 350 to 400 Mev corresponding to maximum meson energies in the center-of-mass system of 23 to 44 Mev.

The second method of calculation, treating the nucleon-nucleon interaction phenomenologically, has been carried out by Marshak and Foldy³ for scalar mesons and for pseudoscalar mesons with pseudovector coupling. They found a zero cross-section for scalar mesons and a very small cross-section (about 10^{-31} at 350 Mev) for pseudoscalar mesons. Both of these results disagree with the experimentally observed¹ large cross section of the order of 10^{-28} cm², however, their treatment suffers from an unrealistic choice

of the nuclear potentials since they assumed charge independence of the forces at the large momentum transfers necessary for meson production. Experimentally, high energy neutron-proton and proton-proton scattering are qualitatively different. Approximate agreement with the experimental results at high energy is given by the potentials⁴

$$\begin{aligned}
 & -g_{NP}(1 + P_x)/2 \exp(-\mu r)/4\pi r \text{ for N-P scattering, } g_{NP}^2/4\pi = .479 \quad (1) \\
 & -g_{PP}^2(\sigma_1 \cdot \nabla/\mu)(\sigma_2 \cdot \nabla/\mu) \exp(-\mu r)/4\pi r \text{ for P-P scattering,} \\
 & g_{PP}^2/4\pi = .0418
 \end{aligned}$$

where P_x is the space exchange operator. The choice of the P-P potential is not unique; any potential which predicts a very singular and strong interaction in P states would give approximate agreement with the high energy scattering. This potential is chosen since it corresponds to pseudoscalar theory with pseudovector coupling. Using these potentials it is of interest to carry out calculations similar to those done by Marshak and Foldy to see if sufficiently large cross sections can be obtained to explain the experimental results.

This method is applied in Section III to the calculation of the transition matrix elements for scalar theory, vector theory with vector coupling, and pseudoscalar theory with pseudovector coupling. For pseudoscalar coupling, it is possible to generalize the P-P potential to its relativistic form using the equivalence between pseudovector and pseudoscalar coupling pointed out by Nelson⁵. The equivalence theorem gives for the potential the result

$$g_{PP}^2(2M/\mu)^2 (\gamma_5)_1(\gamma_5)_2 \exp(-\mu r)/4\pi r \quad (2)$$

The calculation can then be carried out using this type of interaction.

However, it will be shown that for pseudoscalar coupling, processes occur

in the production of mesons which cannot be described in terms of a potential interaction and which give the largest contribution to the matrix element. A treatment on the basis of a potential model therefore is not justified. For such a theory the methods of third order perturbation theory should give more reliable results. These phenomenological calculations are made in the center-of mass system and are restricted to energies near threshold, i.e., less than 400 Mev for the incident nucleon in the laboratory system.

An additional important effect which has been ignored in these calculations must be considered before comparison with experiment can be made. In the ordinary approach to the problem of meson production, the approximation is made of replacing the wave functions of the initial and final nucleons by plane waves. This approximation is fairly good for the initial nucleons since it is equivalent to the use of Born approximation in high energy scattering. However, because of the large amount of energy carried off by the meson in its rest mass, the final nucleons are moving slowly, particularly near threshold. It was pointed out by E.W. Hart and Dr. Geoffrey Chew that as a consequence of this it is possible, if the final nucleons are a neutron and a proton, deuteron may be formed. In addition, even if the nucleons do not form a deuteron, use of the plane wave approximation for the final nucleons gives very inaccurate results. In the Appendix these effects are considered and shown to be important.

II Third Order Field-Theoretic Calculation of Transition Matrix Elements

The third order matrix element may be written down directly using the methods of Feynman and Dyson.⁶ A Feynman-Dyson diagram for meson production is given in Figure 1. Seven additional diagrams may be obtained for emission of the meson by the three other nucleons and for corresponding diagrams in which the two initial or final nucleons are

-9 -

interchanged. The matrix element for the diagram of Fig. 1 is

$$f^\lambda (f^\rho)^2 \bar{\Psi}(3) \left[\frac{U_\mu^2 \tau^\lambda \mathbb{1} \tau^\rho}{(P_1 - q)_\mu \gamma_\mu - iM} U_\nu^1 \right] \Psi(1) \bar{\Psi}(4) \frac{U^1 \tau^\rho \Psi(2)}{(P_4 - P_2)^2 + \mu^2} \phi_\mu^\lambda(q) \quad (3)$$

where $U_\mu^1 = U_\mu^2 = \delta_{\mu 0}$ for S(III)* and $= \gamma_5 \delta_{\mu 0}$ for PsPs(III);

$U_\mu^1 = \delta_{\mu 0} \gamma_5 / \mu (P_4 - P_2)_\nu \gamma_\nu$ and $U_\mu^2 = \delta_{\mu 0} \gamma_5 / \mu q_\nu \gamma_\nu$ for Ps.Pv(III);

$U_\mu^1 = U_\mu^2 = \gamma_\mu$ for V.V(III); $\phi_\mu^\lambda = \phi^\lambda \delta_{\mu 0}$ for S(III), Ps.Ps(III), and

Ps.Pv(III); and $\phi_\mu^\lambda = \phi^\lambda$ for V.V(III). We use the symmetrical isotopic

spin notation⁷ $f^\lambda \tau^\lambda = f_1 \tau_1 + f_2 \tau_2 + f_3 \tau_3 + f_4 \tau_4$. Charge symmetry requires

that $f_1 = f_2 = \sqrt{2}/2f$; the use of both f_3 and f_4 corresponds to two choices

of sign for the coupling of neutral mesons to nucleons, since the expectation value of τ_3 is positive for a neutron and negative for a proton,

while the expectation value of τ_4 is positive for both nucleon states.

The seven additional matrix elements can be obtained by various permu-

tations of P_1, P_2, P_3 , and P_4 and changes of the sign of q_μ . These matrix elements are approximately in the center-of-mass system

$$f^\lambda (f^\rho)^2 [\Psi^*(3) \tau^\lambda \tau^\rho U_1 \Psi(1)] [\Psi^*(4) \tau^\rho U_2 \Psi(2)] / (2M^2 \mu^5) \quad (4)$$

where $U_1 = U_2 = 1$, S(III); $U_1 = 1, U_2 = \sigma \cdot P_2 / \mu / 4M^2$, Ps.Ps(III);

$U_1 = \sigma \cdot \underline{q} \cdot P_1 / \mu M, U_2 = \sigma \cdot P_2 / M$, Ps-Pv(III); $U_1 = |\vec{q}| / \mu, U_2 = 1$, V.V(III).

The total transition matrix elements given by these expressions, squared and averaged over the spins of the nucleons are given by the expressions

$$\begin{aligned} |H_{if}|^2 &= G_{\text{III}}^6 / (2M^2 \mu^5) && \text{S(III), Ps.Ps(III)} \\ &= G_{\text{III}}^6 / (2M^2 \mu^5) (|\vec{q}|^2 / \mu^2) && \text{Ps.Pv(III), V.V(III)} \end{aligned} \quad (5)$$

where the values of G_{III}^6 for the various theories and processes are given in Table I. It is apparent that the results are sensitive to the relative

* We shall in what follows refer to the 3rd order computations for scalar, pseudoscalar with pseudoscalar or pseudovector coupling, and vector theory with vector coupling as S(III), Ps.Ps(III), Ps.Pv(III), and V.V(III) respectively.

choice of sign for the coupling of neutral mesons to neutrons and protons, since the use of f_3 with f_4 set equal to zero corresponds to the opposite choice of sign for the couplings, while the use of f_4 with $f_3 = 0$ corresponds to the same choice of sign of the couplings. It is interesting to observe that the only theory which predicts production of neutral mesons in P-P collisions with a cross section comparable with that for charged mesons is Ps.Ps(III). For the other theories cancellations occur between matrix elements corresponding to emission of the final meson by the initial or final nucleons which make the cross section the order of $(M/M)^2$ smaller than for charged production.

-11-

III Phenomenological Calculation of Production

In this type of calculation, we assume that the process of production can be described by meson emission preceded or followed by the nucleon-nucleon scattering. This, however, is not quite true since virtual processes occur which cannot be described in terms of such a separation. An analysis of the process considered as taking place in third order indicates three basic ways in which a typical process can take place. These are

$$\begin{array}{l}
 (1) \\
 (2) \\
 (3)
 \end{array}
 \begin{array}{l}
 P_1 + P^0 + \omega^0 \rightarrow P^0 + P_3 \\
 P_1 + P_2 \rightarrow \left\{ \begin{array}{l} P_2 + P^0 + \omega^0 \rightarrow P^0 + P_3 \\ P_2 + P^0 + \omega^0 \rightarrow P_2 + P_4 + \omega + \omega^0 \end{array} \right. \\
 P_2 + P^0 + \omega^0 \rightarrow P_2 + P_4 + \omega + \omega^0
 \end{array}
 \rightarrow P_3 + P_4 + \omega \quad (6)$$

It is clear that the virtual meson (ω^0) exchange in processes (1) and (2) is analogous to that which occurs in scattering; therefore the exchange can be replaced by the effects of the potential which predicts the scattering. However, in process (3) the real meson (ω) is emitted between the emission and reabsorption of the virtual meson. Such a process cannot occur in the scattering of real nucleons; therefore its effect cannot be given in terms of the potential model. However, it can be shown rather easily that such processes will give a rather small contribution, at least for theories in which negative energy states are not important for the virtual nucleons. The energy denominator for these matrix elements is given by

$$1 / (E_0 - E^0_1) (E_0 - E^0_2) \quad (7)$$

where E_0 is the total energy and E^0_1 and E^0_2 are the energies of the two intermediate states. These denominators are (ignoring negative energy states) for processes (1) and (2)

$$1 / (E_0/2 - E_4 - \omega^0) (E_0 - 2E_4) \cong 1 / (\omega/2 - \omega^0) \omega \quad (8)$$

and for process (3)

$$1 / (E_0/2 - E_4 - \omega^0) (E_0/2 - \omega - \omega^0 - E_3) \cong 1 / (\omega/2 - \omega^0) (-\omega/2 - \omega^0) \quad (9)$$

Now since near threshold the energy ω' of the virtual mesons is much larger than the energy ω of the real meson, these are approximately $-1/\omega\omega'$ for (1) and (2), and $1/\omega'^2$ for (3). Therefore the contribution from process (3) is smaller than the contribution from processes (1) and (2) in the ratio $\omega/2\omega'$. Since the momentum of the virtual meson is equal to the difference of the momentum of the initial and final nucleons which at threshold is about $(\mu M)^{1/2}$, the ratio $\omega/2\omega'$ is about $1/2(\mu/M)^{1/2}$ which is about 20 percent. This contribution is negligible only if the ratio of the masses of the meson and nucleon is small; actually an error of the order of 20 percent in the matrix element can be expected if this term is ignored. A further error arises from the neglect of negative energy states for the virtual nucleons; however, because of the largeness of the energy denominators for such processes, the contribution is negligible for all theories except for pseudoscalar coupling. This case will be discussed in detail below.

We now consider processes leading from the initial state of two nucleons to the final state of two nucleons and a meson. This can take place in two ways, either through a scattering of the two initial nucleons followed by the meson emission, or with the order of these events reversed. We therefore have for the matrix element

$$\int d\vec{r} d\vec{r}' [\Psi_F^*(\vec{r}, \vec{r}') H(\vec{r}, \vec{r}') \Psi_I(\vec{r}, \vec{r}')] \int d\vec{r} d\vec{r}' [\Psi_I^*(\vec{r}, \vec{r}') S_I(\vec{r}, \vec{r}') \Psi_I(\vec{r}, \vec{r}')] / E_0 - E' + \int d\vec{r} d\vec{r}' [\Psi_F^*(\vec{r}, \vec{r}') S_{II}(\vec{r}, \vec{r}') \Psi''(\vec{r}, \vec{r}')] \int d\vec{r} d\vec{r}' [\Psi^{*''}(\vec{r}, \vec{r}') H(\vec{r}, \vec{r}') \Psi_I(\vec{r}, \vec{r}')] / E_0 - E'' \quad (10)$$

In this expression S_I and S_{II} are the potentials describing the interaction of the initial and final nucleons, H is the operator for meson emission, and E' and E'' are the energies of the two intermediate states. The energy denominators are

$$E_0 - E' = P_1^2/2M - P_2^2/2M - P_1'^2/2M = (P_0^2 - P_1'^2)/M \quad (11)$$

$$E_0 - E'' = P_0^2/M - P''^2/M - \omega$$

A. Scalar meson theory, vector meson theory

We shall now consider the case of two initial protons P_1, P_2 going into a final neutron N_3 , proton P_4 and positive meson q . For this case we have the scattering before meson emission in the P-P potential and the scattering after emission in the N-P potential. If we take scalar coupling for the meson, then

$$H = f^\lambda \tau^\lambda \exp(-iq \cdot r) / (2\omega)^{1/2} \quad (12)$$

The case of vector coupling can be considered simultaneously since the coupling is

$$H = f^\lambda \tau^\lambda \gamma_\mu \phi_\mu(\vec{r}) \quad (13)$$

where γ_μ is the 4-vector formed from the Dirac matrices. This coupling is approximately for longitudinally polarized mesons

$$f^\lambda \tau^\lambda |\vec{q}|/\mu \exp(-i\vec{q} \cdot \vec{r}) / (2\omega)^{1/2} \quad (14)$$

if corrections of the order of v/c for the nucleons are ignored. Therefore we can obtain this result from that for scalar theory by multiplying the matrix element by $|\vec{q}|/\mu$.

We find for the matrix element for $S(\text{phen})^*$

$$\begin{aligned} & -(2)^{1/2} f g_{PP}^2 M(1 - P_{12}) \frac{(\chi_3^* \vec{\sigma} \cdot (\vec{P}_1 + \vec{P}_4) \chi_1) (\chi_4^* \vec{\sigma} \cdot (\vec{P}_4 + \vec{P}_1) \chi_2)}{(2\omega)^{1/2} (P_1^2 - P_4^2) (\mu^2 + (\vec{P}_1 + \vec{P}_4)^2)} \quad (15) \\ & + (2)^{1/2} f g_{NP}^2 (1 - P_{12}) \frac{(1 + P_{34}) (\chi_3^* \chi_1) (\chi_4^* \chi_2)}{2 (2\omega)^{1/2} \omega (\mu^2 + (\vec{P}_1 + \vec{P}_4)^2)} \end{aligned}$$

Near threshold, $P_1 \gg P_3$ or P_4 . We can also disregard μ^2 relative to P_1^2 .

We then note that $(1 - P_{12})(1 + P_{34})(\chi_3^* \chi_2)(\chi_4^* \chi_1) \equiv 0$. Therefore the second term of the matrix element vanishes, and the expression simplifies to

$$- f g_{PP}^2 (1 - P_{12}) (\chi_3^* \sigma \cdot P_1 \chi_1) (\chi_4^* \sigma \cdot P_1 \chi_2) / M \mu^4 (\mu)^{1/2} \quad (16)$$

where we have set $P_1^2/2M = \mu/2$ and $\omega = \mu$, which are their values at threshold.

* We shall refer to the phenomenological treatment of scalar, vector, and pseudo-scalar theory with pseudovector coupling as $S(\text{phen})$, $V \cdot V(\text{phen})$, and $P_s \cdot P_v(\text{phen})$, respectively.

-14-

A similar expression is obtained for an initial neutron and proton going into two final protons and a meson. For processes involving the scattering of two neutrons, the result depends on the choice of the N-N interaction. However, in the absence of any direct information, we shall assume that this is the same as the P-P interaction. The results for these processes are then identical with those given above.

For the production of neutral mesons, the analysis is similar to that given here. In this case, however, the nucleon charge is unchanged by the meson emission, and the scattering takes place before and after emission in the same potential. We find that cancellation occurs between the terms representing scattering before or after emission so that a zero cross-section is predicted for neutral mesons in either N-P or P-P collisions.

B. Pseudoscalar meson with pseudovector coupling

The analysis for pseudovector coupling can be carried out in an exactly similar way. Here we have

$$H = f^\lambda \tau^\lambda \sigma \cdot q / \mu \exp(-i\vec{q} \cdot \vec{r}) / (2\omega)^{1/2} \quad (17)$$

This gives for the production of charged mesons:

$$-f(1-P_{12})(g_{PP})^2 (\chi_3^* \sigma \cdot q \sigma \cdot P_1 \chi_1) (\chi_4^* \sigma \cdot P_1 \chi_2) / \mu^2 - g_{NP}^2 \frac{(1+P_{34})}{2} (\chi_3^* \sigma \cdot q \chi_1) (\chi_4^* \chi_2) / M_\mu^3 (\mu)^{1/2} \quad (18)$$

For neutral mesons from N-P collisions, we have the matrix element

$$g_{NP}^2 f_3 (1-P_{12}P_{34}) (\chi_4^* \sigma \cdot q \chi_2) (\chi_3^* \chi_1) / M_\mu^4 (2\mu)^{1/2} \quad (19)$$

and for P-P collisions the result is again zero.

C. Pseudoscalar meson with pseudoscalar coupling

For the case of pseudoscalar coupling, we have

$$H = f^\lambda \tau^\lambda \gamma_5 \exp(-i\vec{q} \cdot \vec{r}) / (2\omega)^{1/2} \quad (20)$$

We shall now use the relativistic generalization of the P-P potential, which is

$$g_{PP}^2 (2M/\mu)^2 (\gamma_5)_1 (\gamma_5)_2 \exp(-\mu r) / 4\pi r \quad (21)$$

Since negative energy states are important with this form of interaction, we must reconsider the energy denominators in their relativistic form. If the intermediate nucleon is in a negative energy state, its energy is approximately equal to the negative of its rest mass, and we have for the energy denominators as given in Eq. 6, for processes (1) and (2) approximately $1/2M^2$ and for process (3) $-1/2M\omega$. We also must consider the behaviour of the matrix elements describing the emission and reabsorption of the mesons. For transitions between positive energy states, $(\gamma_5) (\gamma_5) \sim (v/c)^2$ where v is the velocity of the nucleons. For transitions to and from negative energy states, $(\gamma_5) (\gamma_5)$ is about 1. Combining these results we find for the approximate magnitude of the matrix elements

$$\text{Processes (1) and (2)} \quad 2(v/c)^2/\omega^2 \sim 1/2M\omega^2 (\mu/M)^{1/2} \cong .40/2M\omega^2 \quad (\text{transitions to positive energy states}) \quad (22)$$

$$\text{Process (3)} \quad 1/2M\omega^2 \quad (\text{transition to negative energy state})$$

We see therefore that near threshold we cannot neglect the non-potential like terms corresponding to process (3) with the virtual nucleon undergoing transitions to negative energy states, since they give the largest contribution to the matrix element. We must refer to the third order perturbation theory results for a more adequate description of the process for the anomalous case of pseudoscalar coupling.

D. Summary of results for phenomenological calculation

The squares of the magnitudes of the matrix elements given by the application of the phenomenological method to the meson production problem are given by

$$\begin{aligned} |H_{if}|^2 &= G_{\text{phen}}^6 / 2M^2 \mu^5 && S(\text{phen}) \\ &= (G_{\text{phen}}^6 / 2M^2 \mu^5) q^2 / \mu^2 && P_s \cdot P_v(\text{phen}), V \cdot V^0(\text{phen}) \end{aligned} \quad (23)$$

where the values of G_{phen}^6 are listed in Table II. A comparison of these results with those obtained in Section II by the third order calculation for the production of charged mesons is given in Table III. It is apparent that the results

obtained by these two methods are approximately equal for not unreasonable choices of the coupling constants. The use of symmetric theory in which $f = f_3$ and f_4 is set equal to zero, however, would predict a zero cross section for S(III) and V·V(III). It is certainly not necessary, however, that such a choice of the coupling constants be made.

IV Calculation of Differential Cross-Section

We shall consider the specific case of production of positive mesons in P-P collisions. The generalization to other cases can easily be made. To the approximation used in these calculations, the differential cross section in the center-of-mass coordinate system for the production of two nucleons and a meson in a nucleon-nucleon collision is given by the expression

$$d\sigma/d\Omega dT = 2(2)^{1/2} M \mu (T(T_m - T))^{1/2} |H_{if}|^2 / (4\pi)^3 \pi \quad (24)$$

where T is the meson kinetic energy, T_m is the maximum meson kinetic energy, and H_{if} is the transition matrix element including the effects discussed in the Appendix of the interaction of the final nucleons. Using the results of that section, we can write

$$|H_{if}|^2 = \left| \psi_t(\vec{r}=0) M_{if}(0)(\text{triplet}) \right|^2 + \left| \psi_s(\vec{r}=0) M_{if}(0)(\text{singlet}) \right|^2 \quad (25)$$

where $M_{if}(0)(\text{triplet})$ is the transition matrix element to a triplet state calculated in Sections II and III, evaluated at zero relative momentum for the final nucleons, similarly for $M_{if}(0)(\text{singlet})$. Substituting the values for the wave functions, this becomes (setting $E_f = T_m - T$)

$$d\sigma/d\Omega dT = 2(2)^{1/2} M \mu (T(T_m - T))^{1/2} \left(\left| M_{if}(0)(\text{triplet}) \right|^2 (v_t + T_m - T) / (\mathcal{E}_t + T_m - T) \right. \\ \left. + \left| M_{if}(0)(\text{singlet}) \right|^2 (v_s + T_m - T) / (\mathcal{E}_s + T_m - T) \right) \quad (26)$$

The values of $M_{if}(0)(\text{triplet})$ and $M_{if}(0)(\text{singlet})$ are given in Table IV. The relation between the constants G_{III}^6 and G_{phen}^6 and the coupling constants f , f_3 , and f_4 is given in Tables I and II.

The total cross section then is given by the expressions

$$\frac{4\pi(2)^{\frac{1}{2}}M^2}{(4\pi)^3} \mu T_m^2 \left| M_{if}(0)(\text{triplet}) \right|^2 \left(1/4 + (V_t - \mathcal{E}_t)/T_m (1 + 2\mathcal{E}_t/T_m - 2(\mathcal{E}_t/T_m - (\mathcal{E}_t/T_m)^2)^{\frac{1}{2}}) \right)^{\frac{1}{2}} + \quad (27)$$

+ (corresponding term for singlet) for S and Ps.Ps

and

$$\frac{8\pi(2)^{\frac{1}{2}}M^2}{(4\pi)^3} \mu T_m^3 \left| \frac{M_{if}(0)(\text{triplet})}{T} \right|^2 \left(1/16 + (V_t - \mathcal{E}_t)/T_m (3/8 + 3/2 \mathcal{E}_t/T_m + (\mathcal{E}_t/T_m)^2 - \right. \quad (28)$$

- (1 + \mathcal{E}_t/T_m)^{3/2} (\mathcal{E}_t/T_m)^{\frac{1}{2}}) + (corresponding term for singlet)) for V.V. and Ps.Pv.

We can now use this cross section ignoring the possibility of deuteron formation to estimate the values of the coupling constants. If we assume a cross section at 350 Mev of $2 \times 10^{-28} \text{ cm}^2$ with the corresponding maximum value of the meson kinetic energy in the center-of-mass system of 24.8 Mev, the corresponding values of the constants $G_{\text{III}}^6/(4\pi)^3$ and $G_{\text{phen}}^6/(4\pi)^3$ are given in Table V. The values of the coupling constants f , f_3 , and f_4 calculated from these values of $G^6/(4\pi)^3$, by the relation given in Tables I and II are given in Table VI. For comparison the values of the coupling constants adjusted to agree with the magnitude of P-P scattering at 350 Mev⁴ (about 25 millibarns) are also given. It is apparent that the coupling constants predicted by these two processes, meson production and high energy scattering, are well within the same order of magnitude. Conversely we can conclude that any of the 3rd order or phenomenological results can correctly predict the order of magnitude of the meson production cross section, for not unreasonable choices of the coupling constants.

Finally we must consider the additional process which can contribute to meson production, the formation of a deuteron and a meson in a proton-proton collision. The cross section can be calculated easily; we have

$$\begin{aligned} d\sigma/d\Omega &= \left| M_{if}(0)(\text{triplet}) \psi_D(r=0) \right|^2 M\mu/(8\pi^2 P_0) \\ &= \left| M_{if}(0)(\text{triplet}) \right|^2 4(2)^{1/2} (T_m \mathcal{E}_t)^{1/2} V_t M^2 \mu / (4\pi)^3 \end{aligned} \quad (29)$$

Using the values of the matrix elements given in Table IV the cross sections are

$$\begin{aligned} d\sigma/d\Omega = & (G_{\text{III,phen}}^2/4\pi)^3 (T_{\text{M}}/\mu)^{3/2} 6.66 \times 10^{-26} \text{ cm}^2 \quad \text{Ps}\cdot\text{Pv(III)}, \text{Ps}\cdot\text{Pv(phen)} \\ & (G_{\text{III}}^2/4\pi)^3 (T_{\text{M}}/\mu)^{1/2} 2.22 \times 10^{-26} \text{ cm}^2 \quad \text{Ps}\cdot\text{Ps(III)} \end{aligned} \quad (30)$$

At 350 Mev, using the values of the constants $G^2/4\pi$ given in Table V, the total cross sections for formation of a deuteron and a meson are

$$\begin{aligned} \sigma = 7.63 \times 10^{-28} \text{ cm}^2 & \quad \text{Ps}\cdot\text{Pv(III)}, \text{Ps}\cdot\text{Pv(phen)} \\ 2.57 \times 10^{-28} \text{ cm}^2 & \quad \text{Ps}\cdot\text{Ps(III)} \end{aligned} \quad (31)$$

These cross sections are larger than those in which the final nucleons are not bound (assumed to be $2 \times 10^{-28} \text{ cm}^2$).

The differential cross sections at 350 Mev for proton-proton production of positive mesons are given in Figs. 2 and 3 for scalar theory and pseudoscalar theory with pseudovector coupling. For comparison the cross sections obtained when the Born approximation was made of treating the final nucleon wave functions as plane waves are also given (dashed curves). The very striking effects of the interactions of the final nucleons are obvious. The variation of the total cross sections with energy is given in Table VII, including the contribution to the cross section when a deuteron is formed. The normalization is again to a total cross section of $2 \times 10^{-28} \text{ cm}^2$ at 350 Mev for the production in which the two final nucleons are unbound.

V Conclusions

The experimental results of the Berkeley workers¹ indicate that the cross section per nucleon for protons bombarding carbon is about $2 \times 10^{-28} \text{ cm}^2$ for both charged and neutral mesons. For protons bombarding free protons, the cross section is about the same for production of charged mesons, but appears to be perhaps an order of magnitude smaller for neutral meson production. It is apparent from the results of Sections II, III, and IV that the relative size

of these cross sections is predicted successfully only by the third order and the phenomenological results for pseudoscalar mesons with pseudovector coupling. The phenomenological result for scalar and vector theory fails in that a zero cross section is predicted for production of neutral mesons in N-P collisions. The third order result for pseudoscalar theory with pseudoscalar coupling is of the right order of magnitude except for neutral mesons produced in proton-proton collisions, where a cross section is predicted comparable with that for charged mesons, in contradiction with the experimental result. The 3rd order result for scalar and vector theories is peculiar in that the cross section for neutral mesons vanishes for neutron-proton collisions and also vanishes for charged mesons if f^2 is taken equal to $1/2 f_3^2$ (see Table I) corresponding to the use of symmetrical theory. It is interesting to observe that a small cross section is predicted for neutral meson production in proton-proton collisions by all of the theories except pseudoscalar theory with pseudoscalar coupling. It is somewhat questionable, however, that such cancellations as those which appear in these calculations are to be quantitatively believed since it is possible that higher order virtual effects would remove the cancellation which appears in lowest order⁸.

The phenomenological calculations for pseudovector coupling are successful in predicting a sufficiently large cross section for agreement with experiment because an unsymmetrical choice of the neutron-proton and proton-proton potentials was made. The calculation made by Marshak and Foldy used a symmetrical interaction and cancellations which occurred reduced the cross section by 2 or 3 orders of magnitude. It is not necessarily true, however, that all symmetrical theories would give a similar result. A theory which while symmetrical could also predict the high energy nucleon-nucleon scattering would presumably give the same general features as the potential models used

in these phenomenological calculations. It is interesting to note that the use of symmetrical theory in the third order calculation does predict correctly the general magnitudes of the cross sections for pseudovector coupling but gives zero cross sections for scalar and vector theory.

The experiments of meson production by protons bombarding free protons¹ provide a good opportunity for verifying the detailed predictions of the differential energy spectra. The experimental results of Cartwright, et al, are shown in Fig. 4 in comparison with the predictions of pseudoscalar theory with pseudovector coupling. It is apparent that agreement with experiment can be obtained only if the effects of the interactions of the final nucleons are taken into account. The predicted fine structure of the high energy peak resulting from the deuteron formation cannot be resolved with the present experimental data; presumably an improvement of experimental techniques will make it possible to test this prediction of the theory.

The author wishes to thank Professor Robert Serber and Dr. K. M. Watson for many interesting discussions of the theoretical results derived in this paper. He also wishes to thank the experimental workers at Berkeley, particularly Drs. Chaim Richman, Herb York, and Vince Peterson for continuous information about the preliminary results of their work, and for aid in interpreting the experiments. In particular he wishes to thank William Cartwright and Marian Whitehead for permission to quote the results of their experiment in advance of publication.

APPENDIX

Keith Brueckner, Geoffrey Chew, and Edward Hart

Effects of interaction of Final Nucleons

For simplicity we restrict ourselves to the case of two initial protons leading to a final neutron, proton, and positive meson. In the calculations described in Sections II and III, we have made the approximation of representing the wave function of the final nucleons by plane waves. This is equivalent to using the Born approximation to describe the nucleon-nucleon scattering. However, near threshold where the final nucleons have low energies, the scattering into these final states is poorly represented by the Born approximation applied to the potentials. The calculations can be done in a more satisfactory way if the actual wave function of the final nucleons is used. We can then resolve this into plane waves by the relation

$$\Psi_F(\mathbf{r}) = \int \alpha_k \exp(i\mathbf{k}\cdot\mathbf{r}) d\vec{k} \quad (1)$$

where

$$\alpha_k = 1/(2\pi)^3 \int \Psi_F(\vec{r}) \exp(-i\vec{k}\cdot\vec{r}) d\vec{r} \quad (2)$$

The calculations which we have made can then be considered to represent one of the Fourier components of this momentum distribution. We can represent the transition matrix element which leads from the initial state to the final state in which we have plane outgoing waves of relative momentum \vec{k} by $M_{if}(\vec{k})$. The transition matrix element to a state $\Psi_F(\vec{r})$ then will be given by the expression

$$H_{if} = \int d\vec{k} \alpha_k M_{if}(\vec{k}) \quad (3)$$

If we wish further to separate the final state into singlet and triplet spin states, we must consider separately the matrix elements of $M_{if}(\vec{k})$ leading to these spin states.

If we insert the definition of α_k , we have

$$H_{if} = 1/(2\pi)^3 \iint d\vec{r} d\vec{k} \exp(-i\vec{k}\cdot\vec{r}) \Psi_F(\vec{r}) M_{if}(\vec{k}) \quad (4)$$

Now if we define

$$1/(2\pi)^3 \int d\vec{k} \exp(-i\vec{k}\cdot\vec{r}) M_{if}(\mathbf{k}) = M'_{if}(\vec{r}) \quad (5)$$

then we have for the matrix element

$$\begin{aligned}
 H_{if} &= \int \psi_F(\vec{r}) M'_{if}(\vec{r}) d\vec{r} \\
 &= \psi_F(\vec{r}_{\text{average}}) \int M'_{if}(\vec{r}) d\vec{r} \\
 &= \psi_F(\vec{r}_{\text{average}}) M_{if}(0)
 \end{aligned}
 \tag{6}$$

where $\psi_F(\vec{r}_{\text{av}})$ is the value of the final wave function at an average value of \vec{r} . This separation can be made more acceptable if we note that $M'_{if}(\vec{r})$ must be large only for \vec{r} considerably less than the range of the forces since the large momentum transfers necessary for meson production lead to a rather singular form for $M'_{if}(\vec{r})$. Then, since for low energy nucleons, $\psi_F(\vec{r})$ is slowly varying over a region of the size of the meson Compton wave length, the result will not be sensitive to the particular value of \vec{r}_{av} , as long as \vec{r}_{av} is considerably less than $\hbar/\mu c$.

This simple result can be applied to the problem of interest. If we wish to calculate the probability that a deuteron is formed, we can take the expectation value of the transition matrix element $M_{if}(0)$ between the initial state of arbitrary spin and the final triplet state, and multiply this by the deuteron wave function evaluated at \vec{r}_{av} . Similarly, if we wish to include the effects of the interaction of the slowly moving final nucleons in a singlet or triplet state, we multiply the appropriate values of the matrix elements by the singlet or triplet wave functions evaluated at \vec{r}_{av} . We shall use the approximate wave functions for a square well⁴ of range 1.53×10^{-13} cm, triplet depth V_t of 52.9 Mev, singlet depth V_s of 41.1 Mev. This gives simple analytic expressions for the wave functions of the deuteron and of the unbound system of neutron and proton of low relative momentum. The use of this non-singular potential may underestimate the magnitude of the wave functions for small separations; however, the results will not be qualitatively incorrect. The approximate wave functions are

$$\begin{aligned}\Psi_D(r) &= \sin((MV_t)^{1/2}r)/r \ ((M\varepsilon_t)^{1/2}/2\pi)^{1/2} \\ \Psi_{t,s}(r) &= \sin((M(V_{t,s} - E_f))^{1/2}r)/r \ (M(\varepsilon_{t,s} + E_f))^{-1/2}\end{aligned}\quad (7)$$

where ε_s , ε_t are the singlet or triplet binding energy (in magnitude) and E_f is the energy of the final nucleons. The magnitude of these wave functions is quite insensitive to the choice of \vec{r}_{av} for \vec{r}_{av} less than $\hbar/\mu c$; for simplicity we shall evaluate them at \vec{r}_{av} equal to zero. We then have

$$\begin{aligned}\Psi_D(0) &= (MV_t)^{1/2} \ ((M\varepsilon_t)^{1/2}/2\pi)^{1/2} \\ \Psi_{t,s}(0) &= ((V_{t,s} + E_f)/(\varepsilon_{t,s} + E_f))^{1/2}\end{aligned}\quad (8)$$

It is apparent from these results that the value of the matrix element is considerably increased by the factor $\Psi_F(0)$ near threshold, where the final nucleon energy E_f is considerably less than the well depth of about 50 Mev.

The factor

$$(V_{s,t} + E_f)/(\varepsilon_{s,t} + E_f) \quad (9)$$

only approaches one for $E_f \gg V$; this condition is not satisfied until the incident nucleon energies are the order of a Bev. At 350 Mev where the final nucleon energy varies from 0 to about 25 Mev, this factor varies from about 2 to 25 for the triplet state and from 2 to about 600 for the singlet state. This has the effect of raising the cross section by a factor of about 3 or 4. The effects of the interaction of the final nucleons therefore clearly are large and cannot be ignored.

-24-

Table I

Values of G_{III}^6 for production of charged mesons and neutral mesons of type 3 (coupled through γ_3) or of type 4 (coupled through γ_4).

		S(III), V.V.(III)	Ps.Ps(III) $\times \mu^3/16M^3$	Ps.Pv(III) $\times(M/\mu)^2$
N=P	π^+, π^-	$f^2(f^2 - 1/2 f_3^2)^2$	$f^2(f^2 + 1/2 f_4^2)^2$	$f^2(f^2 + 1/2 f_3^2)^2$
	$\pi^0(3)$	0	$8f_3^2(f_3^2 - f_4^2)^2$	$8f_4^2 f_3^2$
	$\pi^0(4)$	0	$8f_4^2((f_3^2 - f_4^2)^2 + f_4^4)$	0
P=P	π^+	$f^2(f^2 - 1/2 f_3^2)^2$	$2f^2(3f_4^4 - f_3^2 f_4^2 + 3/4 f_4^4)$	$f^2(f^2 + 1/2 f_3^2)^2$
	$\pi^0(3)$	0	$16f_3^2(f_3^2 + f_4^2)^2$	0
	$\pi^0(4)$	0	$16f_4^2(f_3^2 + f_4^2)^2$	0

Table II

Values of G_{phen}^6 for production of charged mesons and neutral mesons of type 3 and type 4.

		S(phen), V.V.(phen)	Ps.Pv(phen)
Charged		$f^2(M/\mu g_{\text{PP}}^2)^2$	$f^2(M/\mu g_{\text{PP}}^2 + g_{\text{NP}}^2)^2$
N=P	$\pi^0(3)$	0	$2f_3^2(g_{\text{NP}}^2)^2$
	$\pi^0(4)$	0	0
P=P	$\pi^0(3)$	0	0
	$\pi^0(4)$	0	0

-25-

Table III

Ratio of 3rd order to phenomenological matrix elements for production of charged mesons.

Scalar, vector	Pseudovector coupling
$((f^2 - 1/2 f_3^2)/4\pi)/.266)^2$	$((f^2 + 1/2 f_4^2)/4\pi)/.114)^2$

Table IV

Values of square of magnitude of transition matrix elements, for P-P production of positive mesons, evaluated at zero relative momentum for the final nucleons, leading to singlet and triplet spin state. All are to be multiplied by $1/2 M^2 \mu^5$.

	S(III)	S(phen)	V.V(III)	V.V(phen)	Ps.Ps(III)	Ps.Pv(III)	Ps.Pv(phen)
$ M_{if}(0)(\text{triplet}) ^2$	0	0	0	0	$2/3 G_{III}^6$	$G_{III}^6 2Tm/\mu$	$G_{phen}^6 2Tm/\mu$
$ M_{if}(0)(\text{singlet}) ^2$	G_{III}^6	G_{phen}^6	$G_{III}^6 2T/\mu$	$G_{phen}^6 2T/\mu$	$1/3 G_{III}^6$	0	0

Table V

Values of the constants $G_{III}^2/4\pi$ and $G_{phen}^2/4\pi$ to give a total cross section at 350 Mev of $2 \times 10^{-28} \text{ cm}^2$ for production of a positive meson, neutron, and proton in a proton-proton collision.

	S(III), S(phen)	V.V.(III), V.V.(phen)	Ps.Ps(III)	Ps.Pv(III), Ps.Pv(phen)
$G^2/4\pi$.268	.423	.298	.513

Table VI

Values of the coupling constants $f, f_3,$ and f_4 given by Tables I, II, and V, and also as predicted by P-P scattering at 350 Mev.

	Meson Production	P-P Scattering
S(III)	$(f^2 (f^2 - 1/2 f_3^2)^2)^{1/3} / 4\pi = .268$	$(f_3^2 + f_4^2) / 4\pi = .283$
S(phen)	$f^2 / 4\pi = .273$	
V.V.(III)	$(f^2 (f^2 - 1/2 f_3^2)^2)^{1/3} / 4\pi = .223$	$(f_3^2 + f_4^2) / 4\pi = .283$
V.V.(phen)	$f^2 / 4\pi = .613$	
Ps.Ps(III)	$(f^2 (3f^4 - f^2 f_4^2 + 3/4 f_4^4)^2)^{1/3} / 4\pi = 3.79$	$f^2 / 4\pi \sim 6$
Ps.Pv(III)	$(f^2 (f^2 + 1/2 f_3^2)^2)^{1/3} / 4\pi = .145$	$(f_3^2 + f_4^2) / 4\pi = .0418$
Ps.Pv(phen)	$f^2 / 4\pi = .189$	

Table VII

Variation with energy of total cross section for positive meson production in P-P collisions, in units of 10^{-28} cm². The columns headed Unbound are for production leading to a neutron, proton, and meson; those headed Deuteron are for production leading to a deuteron and a meson.

Mev	Scalar	Vector	Ps.Ps.		Ps.Pv.	
	Unbound	Unbound	Unbound	Deuteron	Unbound	Deuteron
290	0	0	0	0	0	0
325	1.10	0.64	0.83	2.05	0.53	3.85
350	2.00	2.00	2.00	2.57	2.00	7.63
375	3.08	4.51	3.48	3.07	5.01	13.0

Figure Captions

- Fig. 1 Feynman-Dyson diagram for meson production by nucleon-nucleon collisions in lowest order. The solid lines represent the nucleons, the dashed lines represent mesons.
- Fig. 2 Differential cross section at 350 Mev in the laboratory system for production of positive scalar mesons in proton-proton collisions. The solid curve includes the effects of the interactions of the final particles; the dashed curve is the result of the calculations using Born approximation throughout.
- Fig. 3 Differential cross section at 350 Mev in the laboratory system for production of positive pseudoscalar mesons with pseudovector coupling in proton-proton collisions. The delta function representing deuteron formation is averaged over a 5 Mev energy interval.
- Fig. 4 Comparison of the experimental results of Cartwright, et. al., for production of positive mesons by 340 Mev protons bombarding free protons. The curve is in the laboratory system for mesons produced in the direction of the proton beam.

REFERENCES

1. C. Richman and H. Wilcox, Am. Phys. Soc 24, No. 8, (1949)
B. Bjorkland, W.E. Crandall, B.J. Moyer, H.F. York, Phys. Rev. 77,
213 (1950)
W. Cartwright, C. Richman, M. Whitehead, H. Wilcox, Phys.Rev.(in press)
V.Z. Peterson, Ph.D. thesis, U. of Calif., 1950.
W. Cartwright and M. Whitehead, private communication
2. For references, see Cecille Morette, Phys. Rev. 76, 1432 (1949)
3. R. Marshak and L. Foldy, Phys. Rev. 75, 1493 (1949)
4. R. Christian and E. Hart, Phys. Rev. 77, 441 (1950)
R. Christian and H.P. Noyes, Phys. Rev. (in press)
5. E.C. Nelson, Phys. Rev. 60, 830 (1941)
6. R.P. Feynman, Phys. Rev. 76, 769 (1949)
F.J. Dyson, Phys. Rev. 75, 1736 (1949)
7. K.M. Case, Phys. Rev. 76, 1 (1949)
8. K.A. Brueckner and K. Watson found for neutral meson production by photons that a similar cancellation in pseudoscalar theory was removed in higher order (Phys Rev. in press).







FIG 1

MU 633





162061



MU 563

FIG. 4

163381