

UCLA

UCLA Previously Published Works

Title

Meeting report on the NIDDK/AUA Workshop on Congenital Anomalies of External Genitalia: challenges and opportunities for translational research

Permalink

<https://escholarship.org/uc/item/1vk2c98g>

Journal

Journal of Pediatric Urology, 16(6)

ISSN

1477-5131

Authors

Stadler, H Scott
Peters, Craig A
Sturm, Renea M
et al.

Publication Date

2020-12-01

DOI

10.1016/j.jpurol.2020.09.012

Peer reviewed



Review Article

Meeting report on the NIDDK/AUA Workshop on Congenital Anomalies of External Genitalia: challenges and opportunities for translational research[☆]



^aDepartment of Skeletal Biology, Shriners Hospital for Children, 3101 SW Sam Jackson Park Road, Portland, OR, Oregon Health & Science University, Department of Orthopaedics and Rehabilitation, Portland, 97239, OR, USA

H. Scott Stadler ^{a,***,1}, Craig A. Peters ^{b,c,**,1}, Renea M. Sturm ^{d,1}, Linda A. Baker ^b, Carolyn J.M. Best ^e, Victoria Y. Bird ^{f,g}, Frank Geller ^h, Deborah K. Hoshizaki ⁱ, Thomas B. Knudsen ^j, Jenna M. Norton ⁱ, Rodrigo L.P. Romao ^k, Martin J. Cohn ^{l,*,1}

^bDepartment of Urology, University of Texas Southwestern, 5323 Harry Hines Blvd., Dallas, 75390-9110, TX, USA

^cPediatric Urology, Children's Health System Texas, University of Texas Southwestern, Dallas, 75390, TX, USA

^dDepartment of Urology, Division of Pediatric Urology, University of California Los Angeles, 200 Medical Plaza #170, Los Angeles, 90095, CA, USA

^eAmerican Urological Association, 1000 Corporate Boulevard, Linthicum, 21090, MD, USA

^fDepartment of Urology, University of Florida, Gainesville, 32610, FL, USA

^gNational Medical Association and Research Group, 5745 SW 75th Street, #507, Gainesville, 32608, FL, USA

^hDepartment of Epidemiology Research, Statens Serum Institut, 5 Artillerivej, Copenhagen S, DK-2300, Denmark

ⁱNational Institute of Diabetes and Digestive and Kidney Diseases, National Institutes of Health, 6707 Democracy Boulevard, Bethesda, 20892, MD, USA

^jUS Environmental Protection Agency, Office of Research and Development, Center for Computational Toxicology and Exposure, Research Triangle Park, 27711, NC, USA

Summary

Congenital anomalies of the external genitalia (CAEG) are a prevalent and serious public health concern with lifelong impacts on the urinary function, sexual health, fertility, tumor development, and psychosocial wellbeing of affected individuals. Complications of treatment are frequent, and data reflecting long-term outcomes in adulthood are limited. To identify a path forward to improve treatments and realize the possibility of preventing CAEG, the National Institute of Diabetes and Digestive and Kidney Diseases and the American Urological Association convened researchers from a range of disciplines to coordinate research efforts to fully understand the different etiologies of these common conditions, subsequent variation in clinical phenotypes, and best practices for long term surgical success. Meeting participants concluded that a central data hub for clinical evaluations, including collection of DNA samples from patients and their

parents, and short interviews to determine familial penetrance (small pedigrees), would accelerate research in this field. Such a centralized datahub will advance efforts to develop detailed multi-dimensional phenotyping and will enable access to genome sequence analyses and associated metadata to define the genetic bases for these conditions. Inclusion of tissue samples and integration of clinical studies with basic research using human cells and animal models will advance efforts to identify the developmental mechanisms that are disrupted during development and will add cellular and molecular granularity to phenotyping CAEG. While the discussion focuses heavily on hypospadias, this can be seen as a potential template for other conditions in the realm of CAEG, including cryptorchidism or the exstrophy–epispadias complex. Taken together with long-term clinical follow-up, these data could inform surgical choices and improve likelihood for long-term success.

Introduction

Congenital anomalies of the external genitalia (CAEG) are a prevalent and serious public health concern with lifelong impacts on the urinary function, sexual health, fertility, tumor development, and psychosocial wellbeing of affected individuals. CAEG include a broad set of conditions that present in a wide spectrum, ranging from mild variants of normal to extreme anomalies that interfere with function (Table 1). Although many conditions require surgical management, complications are frequent and high-quality epidemiological data reflecting long-term

outcomes in adulthood are limited [1–8]. Improving treatments and realizing the possibility of preventing CAEG will require the coordinated efforts of diverse research disciplines to fully understand the different etiologies of these common conditions, subsequent variation in clinical phenotypes, and best practices for long term surgical success. The National Institute of Diabetes and Digestive and Kidney Diseases (NIDDK) and the American Urological Association (AUA) convened researchers from a range of disciplines, including pediatric urology, developmental biology, genetics & genomics, computational systems biology, toxicology, endocrinology, and epidemiology at AUA

[☆] The authors served as the Writing Group of the Workshop on Congenital Anomalies of the External Genitalia.

^kDepartments of Surgery and Urology, IWK Health Centre, Dalhousie University, Halifax, Nova Scotia, Canada

^lDepartment of Molecular Genetics and Microbiology, Department of Biology, And UF Genetics Institute, University of Florida, PO Box 103610, Gainesville, 32610, FL, USA

* Corresponding author.

** Corresponding author. Department of Urology, University of Texas Southwestern, 5323 Harry Hines Blvd., Dallas, 75390-9110, TX, USA.

*** Corresponding author. stadlers@ohsu.edu (H.S. Stadler) craig.peters@utsouthwestern.edu (C.A. Peters) mjcohn@ufl.edu (M.J. Cohn)

Keywords

Hypospadias; Genitourinary development; Chordee; Epispadias; Genetics; Endocrine disruption; Penis; Urethra

Abbreviations

ABM, agent-based models; AOP, adverse outcome pathway; AUA, American Urological Association; BMP, bone morphogenetic protein; CAEG, Congenital anomalies of the external genitalia; DNA, deoxyribonucleic acid; EDC, endocrine disrupting chemicals; EGF, epidermal growth factor; EFNB, ephrin B; EHR, electronic health record; EYA1, EYA transcriptional coactivator and phosphatase 1; FGF, fibroblast growth factor; GWAS, genome-wide association study; HGF, Hepatocyte growth factor; ICD-10, International Classification of Diseases, 10th Revision; IGF, Insulin-like growth factor; IRX, Iroquois-class homeodomain; RNA, ribonucleic acid; SHH, Sonic hedgehog; SNP, single nucleotide polymorphism; VP, phenotypic variance; VG, genomic variance; VE, environmental variance; VGE, genome–environmental interaction variance; WNT, Wingless-INT

Available online 16 September 2020

Headquarters in Linthicum, Maryland in May 2018 to discuss the best way forward to address this public health concern.

Meeting participants discussed research needs relating to: 1) defining the molecular determinants of normal external genital development to improve understanding of CAEG etiologies, 2) mapping molecular pathways to the cellular behaviors that drive genitourinary development, 3) systems modeling to unravel complex tissue interactions that coordinate morphogenesis 4), improving the quality of epidemiological reporting in the era of “Big Data,” and 5) new approaches to clinical phenotyping. The workshop resulted in development of a general framework for advancing research and clinical care to improve outcomes for people with CAEG, which we describe in this article (See [Box 1](#)). The goal was to provide a broad and inclusive framework from varied perspectives, including basic scientists, epidemiologist, clinicians, environmental scientists and others. This framework is intended to serve as a template for organizing current and future knowledge to better develop understanding of these conditions in order to facilitate possible prevention and clinical care. There was broad support for future workshops that target specific needs, such as development of more granular clinical phenotyping protocols that incorporate anatomic, molecular, and genetic data.

Defining the molecular determinants of normal external genital development to better understand the causes of CAEG

During embryogenesis, external genital development involves a complex and dynamic series of cellular events, including differential cell proliferation, apoptosis, migration, and sorting [9–14]. Mechanistically, these cellular processes are controlled by tissue-specific growth factor signaling and sex hormones, which regulate the activity of target genes that orchestrate development of the genitalia in a sex-specific manner [14–18]. At present, our understanding of how these molecular signals are integrated to control the growth and differentiation of the primitive genital anlagen is incomplete, as the full complement of the signaling factors necessary for genital development remains to be elucidated. Equally important is the lack of information regarding how these molecular signals are regulated (or perturbed) by genomic or environmental factors, which, when identified, will provide mechanistic insights into the molecular pathology of CAEG.

The external genitalia arise as paired genital swellings located on each side of the embryonic cloaca [19]. During development (around embryonic day 28 in human and 10.5 in mice) these genital swellings fuse to form the genital tubercle, the structure from which the external genitalia is derived. Genital tubercle development involves some of the same processes as embryonic limb and digit development, including the specification and patterning of tissues along three spatial planes or axes, namely proximodistal (base-to-tip), anteroposterior (cranial side, which will become the dorsal aspect of the phallus, to caudal side, which will become ventral), and mediolateral (midline to left and right sides) [20,21]. To achieve this differential patterning, the endodermal urethral plate epithelium grows out from the cloaca with the genital swellings and functions as the critical organizing center of the genital tubercle. Signaling interactions between the urethral plate and adjacent mesenchyme are mediated by secreted proteins - including SHH, BMP4, BMP7, WNT5a, and FGF10 - that control many of the cellular processes required for genital development [10,13,14,22,23].

For the most part, the developmental processes regulated by these growth factors during genital development have been investigated through the use of genetically engineered mice, which determined that the formation of the urethral tube proceeds by central canalization of the urethral plate [9,24]. While anatomical differences clearly exist between the human and mouse penis, including the presence of bone (os penis or baculum) and distal fibrocartilage (occasionally referred to as the male urogenital mating protuberance) in mice, it is likely that the fundamental developmental processes and molecular signals mediating genital outgrowth and urethral tube formation, two sites commonly affected in CAEG, are generally conserved. However, comparative studies of urethral tube formation in mouse and human have identified potential differences in the modes of lumen formation; human urethral tube development appears to involve a combination of fusion and canalization processes [25,26], whereas in mice the urethral lumen seems to arise by cavitation of the bilaminar urethral plate [9,24,27]. The extent to which the underlying molecular mechanisms differ in mouse and human urethrae will require further investigation. Genomic analyses of individuals affected by CAEG have identified mutations and copy number variants in several of the developmental genes identified as essential for mouse genital development, including EPHB2, EFNB2, EYA1, FGFR2, and HOXA13, highlighting the translational

Table 1 Congenital anomalies of external genitalia.

Virilized or Masculinized Phenotype			Nonvirilized or Feminized Phenotype		
Prepuce	Structure	Hypospadias-associated vs isolated Phimosis	Clitoris	Structure / size	Enlarged (CAH, androgen exposure) Diminutive (CAIS)
Penis	Structure	Congenital megaprepuce Penoscrotal webbing Micropenis	Vestibule/ Urethra	Structure	Labial fusion Urethral prolapse Urethral polyps (tumor excluded) Paraurethral or Gartner’s cysts Ectopic ureterocele, ectopic ureteral insertion
	Alignment	Torsion Chordee with hypospadias Congenital chordee		Meatal location	See DSD: Ambiguous genitalia Pure urogenital sinus anomaly Persistent cloacal anomalies
	Number	Aphallia Diphallia			
External urethra	Structure	Megalourethra (prune belly syndrome) Anterior urethral valves/diverticula	Vagina	Structure	Female epispadias Hymenal anomaly (skin tag, imperforate) Transverse or longitudinal vaginal septum Vaginal atresia or agenesis (MRKH) Urogenital sinus/cloacal anomalies, aplasia, see DSD
	Meatal location	Hypospadias Epispadias (with vs without exstrophy) Fistula (with vs without anorectal malformation)		Meatal location /number	
Scrotal	Number	Urethral duplication (with vs without bladder duplication)	Cervix/uterus	Structure	Uterine anomalies of lateral or vertical fusion Cervical stenosis, absence or lack of patency Embryonic cyst Congenital fistula Uterine or cervical agenesis/aplasia/hypoplasia
	Location/structure	Penoscrotal transposition Bifid scrotum Ectopic scrotum Scrotal hypoplasia		Number	
External lesions	Structure	Vascular malformations Penile or urethral mass/cyst	Gonadal / Fallopiian tubes	Structure	Duplication with or without insertion anomaly See DSD: includes streak gonad
Testis	Location	Undescended testis Retractile testis Splenogonadal fusion Transverse testicular ectopia		Number	Absence/torsion Accessory ovary
	Number	Vanished’ testis or nubbin	Duplication anomalies of	Miscellaneous	Fallopiian tube cyst Embryonic cyst of the broad ligament Urogenital sinus/cloacal anomalies,

(continued on next page)

Table 1 (continued)

Virilized or Masculinized Phenotype		Nonvirilized or Feminized Phenotype	
		Perinatal torsion Polyorchidism See DSD: Disorders of gonadal development	<i>clitoris, urethra, vagina, uterus (with vs without bladder duplication)</i> Hernia/hydrocele Epididymal cyst Congenital absence of the vas deferens Anomalous vasal insertion
Anomalies of the vas deferens, epididymis, processus vaginalis	Gonadal development Structure		aplasia, see DSD
Differences of Sexual Development			
46,XX DSD		46,XY DSD	
Disorders of Gonadal Development	Gonadal dysgenesis Testicular DSD (e.g. SRY+, dup SOX9) Ovotesticular DSD	Disorders of Gonadal Development	Complete gonadal dysgenesis Partial gonadal dysgenesis Bilateral vanishing testis/ testicular regression syndrome Ovotesticular DSD
Androgen Excess	Fetal (congenital adrenal hyperplasia) Fetoplacental (aromatase deficiency) Maternal (ovarian, adrenal tumors, exogenous)	Disorders of Androgen Synthesis or Action	Androgen biosynthesis defect (17, 20-Lyase deficiency) Defect in androgen activity (complete or partial androgen insensitivity syndrome) LH receptor defects (Leydig cell agenesis/ unresponsiveness)
Other 46,XX DSD	Cloacal exstrophy Mayer–Rokitansky–Küster–Hauser syndrome		Disorders of AMH and AMH receptor (persistent Müllerian duct syndrome)
47,XXY DSD	Klinefelter syndrome		Disorders of testosterone metabolism by peripheral tissues (5 α -reductase deficiency)
Disorders of Gonadal Differentiation	Seminiferous tubule dysgenesis Klinefelter Syndrome 46, XX Male	45,X DSD 45,X/46,XY, 46,XX/46,XY DSD Unclassified	Turner syndrome Ovotesticular DSD/mixed gonadal dysgenesis
Syndromes of gonadal dysgenesis	Turner syndrome Pure gonadal dysgenesis Mixed gonadal dysgenesis Partial gonadal dysgenesis		

Wein AJ et al. Campbell-Walsh Urology. 10th ed. Philadelphia PA: Elsevier 2012.

World Health Organization. International statistical classification of diseases and related health problems. 10th revision, 5th ed. Sweden: WHO 2016.

Levin TL, Han B, Little BP. Congenital Anomalies of the Male Urethra. 2007; 37(9):851-62.

Grimbizis GF, Gordts S, Sardo ADS, Brucker S et al. The ESHRE/ESGE Consensus on the Classification of Female Genital Tract Congenital Anomalies. 2013; 28(8):2032-44.

relevance of developmental genetic analysis of CAEG in mice [11,28–31].

What is lacking is a molecular understanding of how these signals are regulated and integrated to facilitate the growth and closure of the external genitalia. Indeed, most CAEG have unknown etiologies with a presumed mix of monogenic and/or multifactorial processes involving both genetic and environmental factors (Table 2). Recent advances in

genomic sequencing, such as whole exome sequencing, RNA-seq, single cell RNA-seq, ATAC-seq, ChIP-seq, and ChIRP-seq, will be instrumental in interrogating patient samples for the genetic and epigenetic causes underlying CAEG. Fundamental research is required to address knowledge gaps in our understanding of the earliest stages of genital development, including the mechanisms that determine the positioning, outgrowth, and sexual differentiation of the external genital

anlagen as well as the mechanisms by which environmental factors disrupt these processes. Critical morphogenetic processes needing further study include epithelial–mesenchymal transition, cell polarity, cell adhesive versus repulsive cues, cell death, mechano-transduction, and the roles of extracellular matrix, vascular endothelial cells, neuronal cells, and neural crest cells. The roles of epigenetic factors, long noncoding RNAs, and miRNAs have just begun to be interrogated in genital tubercle development [32,33], but are generally important in embryogenesis [34,35]. Thus, the quest to identify the pivotal molecular mediators needed for human embryonic external genital morphogenesis should continue, as understanding their identities and functions will be required for insight into the epigenetic, genetic, and environmental causes of human CAEG.

Mapping molecular pathways to the cellular behaviors that drive genitourinary development

Only the minority of hypospadias cases have an identifiable monogenic or chromosomal abnormality, although recent discoveries of variation in gene copy number [28,36] highlight the importance of distinguishing normal genetic variation from variants associated with phenotypic disorders. Challenges in identifying causes of external genital anomalies might indicate either non-genetic underpinnings, such as environmental factors, or technical limitations of screening methods. Whether primary causes are genetic, environmental, or a combination of the two, abnormal external genital development results from disruptions in the cellular and molecular processes that regulate genital morphogenesis. Indeed, animal model studies are beginning to show that mutations in developmental control genes and exposures to endocrine disrupting chemicals (EDC) during critical developmental periods can result in mis-regulation of the same genetic pathways and cellular processes [37–39]. Thus, two major challenges are (a) identifying the molecular and cellular mechanisms that control normal outgrowth, tissue patterning, and urethral tube formation, and (b) understanding how these processes are disrupted in individuals with CAEG. Knowing which genes orchestrate these processes during normal development will allow investigators to determine the range and scope of mutations and allelic differences that contribute to the commonality of these defects (see [Box 1](#)).

Potential antiandrogenic and estrogenic actions of endocrine disruptors

Experimental models demonstrate that several classes of chemicals ubiquitously present in the environment possess anti-androgenic and estrogenic activity in a dose-dependent, additive manner [40]. These environmental EDCs include industrial products like phthalates, bisphenol A, and PCBs, agricultural biocides such as DDT, hexachlorobenzene, and vinclozolin, as well as pharmaceutical and personal care products. Environmental EDCs have been suggested as a potential cause for an increase in the

prevalence of male reproductive disorders, such as hypospadias, cryptorchidism, infertility and testicular cancer, referred to as the testicular dysgenesis syndrome hypothesis [41].

Evidence in humans is limited to pregnancy cohort studies focused on anthropometric measurements of the external genitalia (e.g., anogenital distance, penile length/width) [42–44], and small case–control studies concentrated on specific classes of compounds and occurrence of hypospadias and cryptorchidism in the offspring [45]. A recent systematic review concludes that limited epidemiological evidence exists for a small increase in the risk of male reproductive disorders after exposure to persistent environmental EDCs (e.g. pesticides) but not for rapidly metabolized chemicals (e.g. phthalates). However, the authors also conclude that future, higher-quality epidemiological studies could change the weight of the evidence in either direction [46]. Potential antiandrogenic and estrogenic action of EDCs deserves continued attention from the research community.

Human genetic studies on hypospadias

Epidemiological evidence for a genetic component in hypospadias was observed in a familial aggregation study of isolated hypospadias, with recurrence risk ratios of 50.8, 11.6 and 3.27 for male twin pairs and the first- and second-degree relatives of a case, respectively [47]; in additional heritability analyses on twins of known zygosity, the model with additive genetics and unique environment provided the best fit with a heritability estimate of 77% (95% CI: 57%–90%).

Human genetic studies long focused on rare mutations in candidate genes. These candidate genes were selected from multiple pathways, including SHH, WNT, BMP, FGF, androgen and estrogen signaling, and ephrin receptor and homeobox genes [48]. Identified variants are usually rare, and moderate sample sizes and sparse follow-up in independent cohorts made verifying specific hypospadias risk variants difficult.

Improvements in microarray technologies enabled more comprehensive approaches investigating common variation over the whole genome. A genome-wide association study (GWAS) with 1000 surgery-confirmed isolated cases of hypospadias and more than 5000 controls identified and replicated 18 loci at genome-wide significance, explaining 9% of hypospadias risk [29]. The identified regions were enriched for genes with key roles in embryonic development (e.g., *HOXA4*, *IRX5*, *IRX6* and *EYA1*). Skin and the urogenital system—two physiological systems with an obvious link to hypospadias—yielded significant results in pathway analysis. Findings for the musculoskeletal system require further investigation. Analyzing the heritability explained by all the genetic information captured in the genome-wide SNP data yielded a 57% estimate. Thus, extending the GWAS by SNP-array genotyping of an additional large group of hypospadias cases has potential to identify more associated loci and further elucidate underlying mechanisms. Because the projects generating these data did not specifically target hypospadias, implementing functional data such as DNA methylation in preputial tissue

Table 2 Diagnosis, description, incidence, and developmental/genetic causes of hypospadias.

Diagnosis	Genital organ	Anatomic Variance	Epidemiology	Developmental cause(s)	Genetic Associations/Molecular Mediators
Hypospadias	<ul style="list-style-type: none"> Preputial structure (deficient, hooded) Chordee (ventral penile curvature) Urethral structure and meatal location (ectopic, proximal) 	<p>Distal (85%) vs mid to proximal forms based on location of meatus</p> <p>Varied classification schemes</p> <ul style="list-style-type: none"> Forme fruste Standard hypospadias Severe hypospadias Hypospadias variations <p>Potential variants:</p> <ul style="list-style-type: none"> Megameatus intact prepuce Chordee DSD associated defects (e.g. micropenis, ambiguous genitalia, cryptorchidism or testicular dysgenesis) Penoscrotal webbing or transposition, bifid scrotum 	<p>0.8% of male births</p> <ul style="list-style-type: none"> Most common male external genital birth defect 10–15% familial Recurrence risk ratios for male twin pairs, first- and second-degree relatives of a case: 50.8, 11.6 and 3.3 respectively ~10% of cases explained by identified genetic variants <p>>280 OMIM citations (www.omim.gov)</p> <p>Environmental/ Parental Factors:</p> <ul style="list-style-type: none"> Chemical: pharmaceutical and environmental endocrine-disrupting chemicals, biocides Parental: <p>Placental insufficiency, maternal age, valproic acid use</p> <p>- examples of factors that may increase risk</p>	<p>Abnormal urethrogenesis with complex etiology involving both environmental and genetic factors</p> <p>Computational modeling of adverse outcome pathways: urethral tube closure dependent upon molecular induction of mesenchymal proliferation and epithelial morphogenesis, linked to androgen signaling</p> <p>Data needed at cellular level regarding how molecular cues are translated into morphological anomalies</p> <ul style="list-style-type: none"> Temporal Tissue-specific (cell–cell, cell–matrix interactions) Biomechanical forces 	<p>A minority of cases are secondary to an identified monogenic or chromosomal abnormality.</p> <p>1) Penile development/midline body patterning</p> <ul style="list-style-type: none"> MID1; Xq22: Midline 1; X-linked Opitz Syndrome 22q11.2: Opitz G/BBB Autosomal Dominant Syndrome HOXA13; 7p15-p14.2: Homeobox gene A13 HOXD13; 2q31-q32: Homeobox gene D13 SHH; 7q36: Sonic hedgehog BMP4; 14q22-q23: Bone morphogenetic factor 4 BMP7; 20q13: Bone morphogenetic factor 7 FGF8; 10q24: Fibroblast growth factor 8 FGFR2; 10q26: Fibroblast growth factor receptor 2 FOXA1/A2; 14q21.1/20p11.21 Forkhead Box A1/A2 GLI3; 7p14.1: GLI family zinc finger 3 <p>2) Testis determining</p> <ul style="list-style-type: none"> NR5A1; 9q33.3: Steroidogenic factor-1 SRY; Yp11.3: Sex-determining region of Y chromosome WT-1; 11p13: Wilms' tumor –1 NR0B1; Xp21.3-p21.2: DAX-deleted in azoospermia LHCGR; 2p21: Luteinizing hormone receptor <p>3) Androgen production/signaling</p> <ul style="list-style-type: none"> AR; Xq11.2-q12: Androgen receptor

- SRD5A2; 2p23: 5 α -reductase, type II
 - HSD17B3; 9q22: 17 β -hydroxysteroid dehydrogenase type 3
 - HSD3B2; 1p13.1: 3 β -hydroxysteroid dehydrogenase type 2
 - ESR1; 6q25.1: Estrogen receptor 1
 - ESR2; 14q23.2: Estrogen receptor 2
 - MAML1; Xq28: Mastermind-like domain containing 1
 - ATF3; 1q32.3: Activating transcription factor 3
- 4) Potential GWAS-identified:
- DGKK; Xp11.22: diacylglycerol kinase kappa
 - EYA1; 8q13.3: EYA transcriptional coactivator and phosphatase 1
 - HOXA4; 7p15.2: Homeobox gene A4
 - IRX5; 16q12.2/IRX6; 16q12.2: Iroquois homeobox genes
 - SP1/SP7; 12q13.13: Sp1/7 transcription factor
- 5) **Key susceptibility loci to environmental exposures:** the interaction between genetic and endocrine signals
- 6) Potential role of **epigenetic mechanisms, long noncoding RNAs, miRNAs, unfolded protein response (autophagy-apoptosis), metabolome**

Table 3 Elements of clinical phenotyping of CAEG.

Element	Comment
Physical - structural	Templated descriptions Standardized measurements 3D imaging with quantitative analysis/ AI
Functional	Urinary Sexual
Genetic	Genotyping Family pedigree
Tissue and cellular	Tissue typing Cell lineage Cellular fingerprinting
Molecular	Gene/RNA expression Protein expression Molecular fingerprinting
Associated conditions	Other potentially involved and related systems – e.g. cardiac, neurological

or fetal murine gene expression during urethral development could further improve pathway analyses.

Re-thinking genetics

Beyond searching for mutations in coding and regulatory regions of developmental control genes, we must define key susceptibility loci that influence individual environmental exposure response. Differential sensitivity to anti-androgenic or estrogenic chemicals, for example, may have a genetic basis. Much of external genital development occurs before sexual differentiation of the gonads and is virtually identical in males and females. After gonadal differentiation and onset of steroidogenesis, systemically circulating sex hormones direct the genital tubercle down masculinizing or feminizing pathways, which involve strikingly different morphogenetic processes. Understanding how these pathways affect expression of genes that govern genital morphogenesis in target tissues will be essential in determining how chemicals may disrupt or mimic sex hormones, leading to CAEG. Interactions between genetic and endocrine signals in the external genitalia are largely unknown at present. Because the incidence of hypospadias appears to have outpaced the mutation rates for most protein coding genes, characterizing epigenetic mechanisms capable of influencing the expression/function of loci involved in penile development and urethral closure will be critical. Candidate epigenetic mechanisms to consider for potential roles in penile development include DNA methylation, modifications affecting chromatin condensation (such as histone acetylation/methylation), and the expression of non-coding RNAs (micro RNAs and long non-coding RNAs) that can regulate the expression of multiple protein-coding genes. Traditionally, perturbations in these epigenetic mechanisms have not been examined in clinical genetic evaluations and are likely to be identified as contributing factors in the pathology of CAEG due to their capacity to regulate the expression of susceptibility loci. Establishing the relevant linkages between genetic variation, epigenetic influences, and environmental factors will

Box 1. Key points: Essential elements for advancing research and treatment of CAEG.

- Clinical phenotyping consensus
- Novel phenotyping Resources and Technologies
- Tissue banking and bio-repositories
- Standardization Workshops
- Surgical follow-up standardization
- Clinical Pedigree resources
- Collaborative Networks
- Data analysis resources
- Data integration centers

be essential for a true understanding of many of the CAEG conditions. Moreover, identification of mechanisms of these interactions will permit a functional understanding of the causes of CAEG and has the potential to generate markers of specific processes and lead to therapeutic interventions.

Focus on the cell: connecting molecules to morphological anomalies

The paucity of data at the cellular level has hampered efforts to connect genotype to phenotype in the external genitalia. Determining CAEG etiologies will require understanding of how molecular cues, whether intrinsic or environmental, are translated into morphological anomalies. Areas of exploration include the relationship of gene expression to cellular behavior in a temporal and tissue-specific manner during external genital development; the roles of cell–cell and cell–matrix communication in urethral tube development and how these processes influence cell and tissue mechanics, integrity, and morphogenesis. Understanding the causes of failed urethral tube development will require detailed investigations into the consequences of gene expression patterns on cell polarity, shape, migration, apoptosis, and proliferation. These questions present an opportunity to develop experimental tests that will define the actual mechanisms that regulate cellular processes during human genital development.

Experimental models and approaches

Animal models

Some of these challenges cannot be addressed directly in human cells or tissues and require animal models. The mouse remains a system of choice for numerous reasons, including its well-developed genetic and genomic infrastructure, advantages as a developmental system, and amenability to genetic manipulation. External genitalia of mice and humans have considerable differences in their anatomy and development, including urethral tube formation, which must be considered when extrapolating between species. Nonetheless, molecular genetic studies suggest that the developmental mechanisms are largely conserved between mice and humans.

Although rodent models will remain essential tools in studies of external genital development, additional models are needed. Historical thinking suggested that phalluses of different vertebrates were related only by common functions with no shared evolutionary history, raising questions about the wisdom of studying genital development in an organism in which the penis could be wholly unrelated to that of humans. However, recent work indicates the amniote penis evolved once, and the external genitalia of mammals, birds, and reptiles are homologous, with similarities due to common ancestry [19,49,50]. This discovery opens possibilities for use of novel, comparative models of genital development with potential advantages over current models, exemplified in recent use of bird and reptile embryos to fate map the progenitor cells of the genital tubercle [51,52], discovery of similarities between guinea pig and human urethral tube formation [53], and use of marsupials to study genital morphogenesis during their extended period of development in the pouch [54,55]. The “optimal” animal model depends on the question under study; therefore, the field must be open to use of non-traditional models that could enable new experimental approaches and insights applicable to humans.

In vitro models

In vitro models using human induced pluripotent stem cells or genital tissues will enable identification, confirmation, and integration of molecular and cellular processes required for genital development. These investigations may be enhanced by the use of three-dimensional organoids, which have revolutionized studies of brain, kidney, and heart development [56–58]. Research in external genital development has not leveraged organoid systems, which represent an untapped resource for modeling, testing, and perturbing the processes hypothesized to underlie CAEG. Additionally, by procuring human tissues from clinical settings, organ culture may enable identification of cellular processes mediating human genital development and closure (i.e., matrix deposition, cell migration, epithelial–mesenchymal transition, apoptosis). Clinical samples will also facilitate development of cell lines to establish the genetic and epigenetic hierarchies functioning in the external genitalia and provide high-throughput avenues to assess susceptibility of these hierarchies to various environmental factors. Finally, such *in vitro* approaches may allow determination of the genetic and environmental inputs necessary to engineer new tissues, providing clinically relevant alternatives to repair or replace tissues affected by congenital malformation, injury, or disease.

Systems modeling of complex tissue interactions in silico

Understanding how tissues and organs arise during embryogenesis is a central question in developmental biology and a challenge for predicting developmental toxicity. While alterations in key signaling pathways (e.g., SHH, WNT, TGF β , EPH receptor tyrosine kinases, retinoic acid) can disrupt genitourinary development, the critical developmental processes causal in CAEG are poorly understood. To address this knowledge-gap, it is essential that approaches complementary to genetic and toxicological screens be used. At the forefront of these approaches is systems modeling, which allows complex defects such as

CAEG to be computationally interrogated. This approach provides a mechanism to assess adverse outcomes pathways in a high throughput manner, identifying the roles of genetic mutations or putative teratogens in the pathology of CAEG [59]. Indeed, recent applications of systems modeling to kidney development identified a Turing mechanism by which GDNF, RET, and WNT11 mediate the complex tissue dynamics controlling branching morphogenesis [60,61]. Systems modeling will also facilitate analysis of how disruption of signaling events functioning simultaneously during the genital development causes CAEG, as many of these signaling events are often integrated during formation of the external genitalia [59,62,63].

For hypospadias, computational models successfully recapitulated sexually dimorphic development of the genital tubercle controlled by SHH, FGF10, and androgen pathways through modulation of stochastic cell behaviors, including differential adhesion, motility, proliferation, and apoptosis. Proper urethral tube closure in this model was shown to depend quantitatively on SHH- and FGF10-induced effects on mesenchymal proliferation and epithelial apoptosis. Moreover, the model also linked these cellular processes to androgen signaling. In the absence of androgen, genital tubercle development was feminized and with partial androgen deficiency, the model resolved with incomplete urethral tube closure, providing an *in-silico* platform for probabilistic prediction of hypospadias risk across combinations of minor perturbations to the system [64,65].

Finally, in a human cell-based ‘biomimetic system’ designed to assess the influence of environmental factors on controlled tissue events [66], organoid survival was dependent on signaling through EGF, IGF, HGF, and FGF pathways, and organoid fusion was disrupted by inhibition of BMP signaling. Concordance between the effects of EGF, FGF, and BMP inhibitors on palatal organoid fusion and epithelial cell migration *in vitro* suggested critical dependence on epithelial morphogenesis and, potentially, an innate feedback mechanism. These findings demonstrate promising utility of integrative modeling and adverse outcome pathways to decode the “toxicological blueprint of active substances” that interact with the developing embryo [67].

Distinguishing pattern from process: imaging developmental dynamics

Understanding the morphogenetic processes and tissue movements involved in human and mouse urethral development has been confounded by descriptions that conflate anatomical patterns with developmental processes [26,68]. Recent advances in imaging technology enable real-time, three-dimensional imaging of genital morphogenesis [69], providing opportunities to directly observe processes previously inferred from static, two-dimensional images (e.g., histological sections) [25,26]. When applied to mouse models that allow single cell labeling and lineage tracing, such live imaging approaches can reveal the spatiotemporal processes involved in urethral tube formation.

Developmental biomechanics

Direct investigation of biomechanical forces speculated to act during urethral tube development are needed. For example, the ventral aspect of the urethral plate is

believed to be under tension during lumen formation and may be susceptible to rupture when epithelial mechanical integrity is compromised [37]. Such investigation may reveal the relationship between epithelial maturation and lumen formation and commonalities in tubulogenesis across organ systems. Application of engineering principles to genital developmental systems research will enhance understanding of genetic and environmental effects on the cellular processes that drive morphogenesis.

Improving the quality of epidemiological reporting in the era of “Big Data”

Recent epidemiological studies have yielded conflicting results regarding prevalence of CAEG. Conflicts may result from inconsistent recognition and classification of CAEG, due to the lack of a standardized classification system. A 30-year, population-based Danish study found hypospadias rates doubled [70], with the final prevalence similar to the rate seen in Nova Scotia, Canada around the same time [71]. However, prevalence in Nova Scotia remained stable over the same time period [71]. A recent meta-analysis of studies around the globe, covering over 90 million births, reported the mean prevalence of hypospadias (per 10,000 births) as 19.9 in Europe, 34.2 in North America, 5.2 in South America, 0.6–69 in Asia, 5.9 in Africa, and 17.1–34.8 in Australia [72]. The authors noted the extreme heterogeneity of published datasets, confounding factors, such as the absence of standardized definitions of hypospadias, conflicting results of studies from the same regions, and a complete absence of data from some regions [72]. Multiple reports suggest an increasing prevalence, although we lack clarity as to how increased numbers of registries, better data capture and reporting, or over reporting of mild cases may affect prevalence estimates. Inconsistencies across global hypospadias prevalence data suggest misclassification might be an issue. Hence, accurate epidemiology will require improved quality of CAEG recognition and reporting, including diagnosis,

comorbidities and systematic phenotypic descriptions (see section entitled “New approaches to clinical phenotyping” below for further details).

Data sources analyzed for reporting CAEG in recent decades have been inconsistent, challenging determination of the true CAEG prevalence. Needs include: (i) registry standardization, including consistent reporting of long-term data, reduced site variation, and multi- (versus single-) institution reporting; (ii) persistent data collection over time and across geography; (iii) consistent case discovery efforts; and (iv) a global standard that aligns the full range of phenotypes with classification codes (e.g., International Classification of Diseases, 10th Revision [ICD-10]).

Strategies are in place to systematically mine existing electronic health record (EHR) databases through graphical interfaces of integrated data repository systems (e.g., [Informatics for Integrating Biology and the Bedside \[I2B2\]](#)). Such state and national databases will aid in identifying regional prevalence patterns and variables correlated to specific CAEG using existing cohorts identified by ICD-10 or other coding systems. Zip codes available in EHRs can facilitate geospatial analysis and mapping of CAEG to identify potential hot spots and target interventions. Machine learning algorithms [73] can leverage EHR data for prediction models, and may integrate genetic variables from databases with curated clinical and genetic information (e.g., [BioVU](#)). Such models may predict surgical outcomes (e.g., complication or repeat surgery risk after correction of hypospadias) based on phenotypic, genetic and epigenetic relationships. Large databases may facilitate identification of maternal exposures before and during pregnancy associated with CAEG based on demographic, geographic and temporal information. Thus, by using high performance statistical software to cluster large EHR datasets, phenotypic variations (VP) associated with CAEG could be defined as a summation of variances, including genomic (VG), environmental (VE) and genome–environmental interaction (VGE), such that $VP = VG + VE + VGE$.




Figure 1 A model for the structure of a clinical registry that will utilize standardized criteria for phenotyping, tissue banking, analysis of tissues, integration of clinical outcomes, and data synthesis to determine risk and improve patient outcomes.

New approaches to clinical phenotyping

Enhancing understanding of CAEG will rely on robust phenotyping with precise and objective characterization of individual genital patterns, their developmental phenotype, that of their family, and genetic subtyping through individual and family genotyping and pedigree analysis. Robust description may include tissue, cellular and molecular phenotyping as well (See Table 3). Phenotypes must be based on reproducible categories that consistently apply, thus allowing for the transfer of knowledge over time. Description of genital anatomy requires objective measurable parameters that do not presuppose any particular pattern or relationship. CAEG are characterized by a wide variation in severity, yet phenotypic overlap complicates pattern recognition, as distinct characteristics seem associated with various patterns. Viewing CAEG as a mosaic of elements that contribute to the ultimate individual phenotype may improve categorization and better reflect developmental patterns.

Multiple potential methods may enable accurate and objective genital phenotyping, including standardized photographs, templated measurements, or ultrasonography. More complex methods, including physical molding and 3-D surface imaging, may prove valuable. Hypospadias description, for example, begins with meatal location, degree of ventral curvature, penoscrotal transposition, and bifid scrotum, but will need to include further parameters. Descriptive templates exist [74], but some may inappropriately categorize presumed yet unproven relationships. A single, consensus-based descriptive template—enhanced by visual imaging and recording—may enable objective comparison and categorization of patterns.

Considerations for clinical implementation

Optimal phenotypic description must be both practical and efficient to be integrated into clinical care and must be implemented in a systematic manner to enable consistent assessment of individuals as they grow, develop or receive surgical intervention. Yet, an optimal tool must balance this consistency over time with the flexibility to adapt to evolving knowledge in the field. The method must account for the sensitive nature of CAEG phenotyping assessments, including risks and demands of patient confidentiality. Optimal patient phenotyping must also enable identification of associated concurrent conditions, facilitating identification of associated developmental patterns as clues to etiology.

Development of such a rigorous system requires multi-center participation. Phenotyping in CAEG may be informed by efforts of interdisciplinary multi-institutional consortia to standardize terminology, categorize and quantitate phenotypic features and catalog clinical phenotypes outside of CAEG via EHR-based data templates that fulfill clinical and research needs. This approach permits longitudinal follow up, access to clinical records, recording of biobanking, and potential for interoperability with other health IT systems, enabling cross-site collaboration. Resulting data must be shared and accessible while remaining secure. Such database systems will require broad coordination, oversight, and maintenance.

Standardizing surgical interventions and follow-up

It is premature to consider standardized surgical interventions because optimal interventions remain incompletely defined. A future definition of best practices may be developed based on robust outcomes analysis and linkage with mechanistic understanding of clinical patterns. Templated follow-up patterns will likely be most useful to evaluate interventions, and will include structural, functional and psychological parameters.

Inherent to defining outcomes is the challenge of maintaining follow-up in a highly mobile population. Providing value to patients and families in terms of knowledge, counseling and engagement in the development of new knowledge may enable improved monitoring, and in turn, better understanding of the underlying developmental pattern and clinical outcomes. Some hospital systems have research opt-out policies wherein patients may be contacted for research unless they specifically request no contact, which has increased research participation.

Potential to glean information from tissue sampling

Despite the delay between initiation of the condition during fetal development and tissue sampling long after development is completed, sampling of genital and somatic tissue may yield clues regarding underlying pathophysiology. Tissue sampling also may provide information about specific properties, such as wound healing, hormonal responsiveness, growth regulation, and fibrosis, which might influence surgical outcomes. Developing standardized protocols for tissue sampling during genital reconstruction may be of value. Identifying flexible preservation techniques that allow for current and future methodologies is essential. Since we cannot accurately predict all future methods, optimal methods might be those that minimize tissue effects.

Effective data-gathering, recording and assessment structure within a longitudinal timeframe is essential and will be most productive within the context of a collaborative network of investigators. A key task of this collaboration would be the development and implementation of standards of participation for all investigators. The broad view of this collaboration (Fig. 1) should involve iterative processes of change based on new knowledge. Machine learning strategies may facilitate integration of clinical, genetic, and biological parameters and provide relevant translational insights. These insights may feed back into the informational loop, furthering our understanding of the developmental, biological and clinical processes underlying each phenotype and associated outcomes, thereby enabling optimized selection of treatments to unique phenotypes.

Conclusions

At the conclusion of the 2018 CAEG Workshop held at the AUA Headquarters, the authors of this report were charged with writing a synthesis of the current state of research (clinical and basic science) and the key knowledge gaps in

CAEG (See [Box 1](#)). As indicated by this report, hypospadias was a major focus of the presentations and discussions at the workshop, and we propose that this roadmap for hypospadias research also serve as a model for investigations of other genitourinary anomalies. At the clinical level, a major challenge identified by workshop participants is the need for a standardized method to phenotype CAEG patients. Standardization of clinical phenotyping should also include the creation of a phenotype-associated biobank of DNA and possibly other samples (e.g., blood) from patients and both parents. This resource should be in a searchable format, allowing for future studies aimed at identifying key loci involved in discrete malformations as well as environmental factors that may contribute to defect etiology.

At the basic science level, a need was identified for additional studies aimed at defining the gene regulatory networks that control the developmental processes required for growth, generation of specific cell and tissue types, formation of a functional urethral tube, and sexual differentiation of the external genitalia. As the repertoire of single cell genomic tools expands, it is anticipated that new insights into molecular processes that regulate the growth and differentiation of the external genitalia will be gained. Included in these novel approaches will be studies aimed at defining the role of epigenetic factors—such as non-coding RNAs, DNA and/or RNA methylation, and exogenous (e.g., environmental) regulators of gene expression—in mediating the development of the external genitalia. Finally, at the cellular level, recent advances in live cell imaging should provide the necessary approaches to determine common and species-specific cellular and molecular processes necessary and/or sufficient for the growth and differentiation of the external genitalia.

By addressing these challenges in future clinical and basic science studies, more streamlined and effective mechanisms to define the molecular basis for CAEG should be realized. More importantly, the integration of clinical and basic science approaches to studying CAEG should provide the best roadmap towards reducing and eventually preventing CAEG in the global population, affording directed therapeutic approaches to patients and families, as well as providing new insights into mechanisms underlying patient susceptibility to these defects.

Disclaimers

The views expressed in this article are those of the authors and do not necessarily reflect the views or policies of their employers or funding agencies. Mention of trade names or commercial products does not constitute endorsement or recommendation for use.

Conflicts of interest

The authors declare no conflicts of interest.

Acknowledgments

The authors served as the writing group for the NIDDK/AUA Workshop on Congenital Anomalies of the External

Genitalia and gratefully acknowledge the contributions of the following workshop participants: Andrew Adey (Oregon Health and Science University), Darius Bagli (University of Toronto), Larry Baskin (University of California - San Francisco), Spyros Darmanis (Chan Zuckerberg Biohub), L. Earl Gray (U.S. Environmental Protection Agency), Dorrie Lamb (Weill Cornell Medicine), Sean Li (Boston Children's Hospital), Liang Ma (Washington University), Andrew Pask (University of Melbourne), Eric Vilain (Children's National Medical Center and George Washington University), and Duncan Wilcox (University of Texas Southwestern Medical Center).

References

- [1] Chertin B, Natsheh A, Ben-Zion I, Prat D, Kocherov S, Farkas A, et al. Objective and subjective sexual outcomes in adult patients after hypospadias repair performed in childhood. *J Urol* 2013;190:1556–60.
- [2] Kiss A, Sulya B, Szasz AM, Romics I, Kelemen Z, Toth J, et al. Long-term psychological and sexual outcomes of severe penile hypospadias repair. *J Sex Med* 2011;8:1529–39.
- [3] Ortqvist L, Fossum M, Andersson M, Nordenstrom A, Frisen L, Holmdahl G, et al. Long-term followup of men born with hypospadias: urological and cosmetic results. *J Urol* 2015;193:975–81.
- [4] Ruppen-Greeff NK, Weber DM, Gobet R, Landolt MA. Health-related quality of life in men with corrected hypospadias: an explorative study. *J Pediatr Urol* 2013;9:551–8.
- [5] Ruppen-Greeff NK, Landolt MA, Gobet R, Weber DM. Appraisal of adult genitalia after hypospadias repair: do laypersons mind the difference? *J Pediatr Urol* 2016;12:32.e1–8.
- [6] Rynja SP, Wouters GA, Van Schaijk M, Kok ET, De Jong TP, De Kort LM. Long-term followup of hypospadias: functional and cosmetic results. *J Urol* 2009;182:1736–43.
- [7] Rynja SP, de Jong TP, Bosch JL, de Kort LM. Functional, cosmetic and psychosexual results in adult men who underwent hypospadias correction in childhood. *J Pediatr Urol* 2011;7:504–15.
- [8] Springer A. Assessment of outcome in hypospadias surgery - a review. *Front Pediatr* 2014;2.
- [9] Seifert AW, Harfe BD, Cohn MJ. Cell lineage analysis demonstrates an endodermal origin of the distal urethra and perineum. *Dev Biol* 2008;318:143–52.
- [10] Morgan EA, Nguyen SB, Scott V, Stadler HS. Loss of Bmp7 and Fgf8 signaling in Hoxa13-mutant mice causes hypospadias. *Development* 2003;130:3095–109.
- [11] Yucel S, Liu W, Cordero D, Donjacour A, Cunha G, Baskin LS. Anatomical studies of the fibroblast growth factor-10 mutant, Sonic Hedge Hog mutant and androgen receptor mutant mouse genital tubercle. *Adv Exp Med Biol* 2004;545:123–48.
- [12] Dravis C, Yokoyama N, Chumley MJ, Cowan CA, Silvano RE, Shay J, et al. Bidirectional signaling mediated by ephrin-B2 and EphB2 controls urorectal development. *Dev Biol* 2004;271:272–90.
- [13] Lin C, Yin Y, Bell SM, Veith GM, Chen H, Huh SH, et al. Delineating a conserved genetic cassette promoting outgrowth of body appendages. *PLoS Genet* 2013;9:e1003231.
- [14] Seifert AW, Zheng Z, Ormerod BK, Cohn MJ. Sonic hedgehog controls growth of external genitalia by regulating cell cycle kinetics. *Nat Commun* 2010;1:23.
- [15] Suzuki H, Matsushita S, Suzuki K, Yamada G. 5 α -Dihydrotestosterone negatively regulates cell proliferation of the periurethral ventral mesenchyme during urethral tube formation in the murine male genital tubercle. *Andrology* 2017;5:146–52.

- [16] Larkins CE, Enriquez AB, Cohn MJ. Spatiotemporal dynamics of androgen signaling underlie sexual differentiation and congenital malformations of the urethra and vagina. In: *Proc Natl Acad Sci U S A*, 113; 2016. E7510–7.
- [17] Matsushita S, Suzuki K, Murashima A, Kajioaka D, Acebedo AR, Miyagawa S, et al. Regulation of masculinization: androgen signalling for external genitalia development. *Nat Rev Urol* 2018;15:358–68.
- [18] Zheng Z, Armfield BA, Cohn MJ. Timing of androgen receptor disruption and estrogen exposure underlies a spectrum of congenital penile anomalies. *Proc Natl Acad Sci U S A* 2015; 112:E7194–203.
- [19] Gredler ML, Larkins CE, Leal F, Lewis AK, Herrera AM, Perriton CL, et al. Evolution of external genitalia: insights from reptilian development. *Sex Dev* 2014;8:311–26.
- [20] Perriton CL, Powles N, Chiang C, Maconochie MK, Cohn MJ. Sonic hedgehog signaling from the urethral epithelium controls external genital development. *Dev Biol* 2002;247: 26–46.
- [21] Ipulana LA, Raga D, Suzuki K, Murashima A, Matsumaru D, Cunha G, et al. Investigation of sexual dimorphisms through mouse models and hormone/hormone-disruptor treatments. *Differentiation; research in biological diversity*. 2015.
- [22] Ching ST, Infante CR, Du W, Sharir A, Park S, Menke DB, et al. *Isl1* mediates mesenchymal expansion in the developing external genitalia via regulation of *Bmp4*, *Fgf10* and *Wnt5a*. *Hum Mol Genet* 2018;27:107–19.
- [23] Kajioaka D, Suzuki K, Nakada S, Matsushita S, Miyagawa S, Takeo T, et al. *Bmp4* is an essential growth factor for the initiation of genital tubercle (GT) outgrowth. *Congenital anomalies*; 2019.
- [24] Hynes PJ, Fraher JP. The development of the male genitourinary system: III. The formation of the spongiosae and glandular urethra. *Br J Plast Surg* 2004;57:203–14.
- [25] Shen J, Overland M, Sinclair A, Cao M, Yue X, Cunha G, et al. Complex epithelial remodeling underlie the fusion event in early fetal development of the human penile urethra. *Differentiation; research in biological diversity* 2016;92:169–82.
- [26] Li Y, Sinclair A, Cao M, Shen J, Choudhry S, Botta S, et al. Canalization of the urethral plate precedes fusion of the urethral folds during male penile urethral development: the double zipper hypothesis. *J Urol* 2015;193:1353–9.
- [27] Hynes PJ, Fraher JP. The development of the male genitourinary system: II. The origin and formation of the urethral plate. *Br J Plast Surg* 2004;57:112–21.
- [28] Tannour-Louet M, Han S, Corbett ST, Louet JF, Yatsenko S, Meyers L, et al. Identification of de novo copy number variants associated with human disorders of sexual development. *PLoS One* 2010;5:e15392.
- [29] Geller F, Feenstra B, Carstensen L, Pers TH, van Rooij IA, Korberg IB, et al. Genome-wide association analyses identify variants in developmental genes associated with hypospadias. *Nat Genet* 2014;46:957–63.
- [30] Wang C, Gargollo P, Guo C, Tang T, Mingin G, Sun Y, et al. *Six1* and *Eya1* are critical regulators of peri-cloacal mesenchymal progenitors during genitourinary tract development. *Dev Biol* 2011;360:186–94.
- [31] Goodman FR, Bacchelli C, Brady AF, Brueton LA, Fryns JP, Mortlock DP, et al. Novel *HOXA13* mutations and the phenotypic spectrum of hand-foot-genital syndrome. *Am J Hum Genet* 2000;67:197–202.
- [32] Chen Y, Kuroki Y, Shaw G, Pask AJ, Yu H, Toyoda A, et al. Androgen and oestrogen affect the expression of long non-coding RNAs during phallus development in a marsupial. *Noncoding RNA* 2018;5.
- [33] Carlson HL, Quinn JJ, Yang YW, Thornburg CK, Chang HY, Stadler HS. LncRNA-HIT functions as an epigenetic regulator of chondrogenesis through its recruitment of p100/CBP complexes. *PLoS Genet* 2015;11:e1005680.
- [34] Schmitz SU, Grote P, Herrmann BG. Mechanisms of long non-coding RNA function in development and disease. *Cell Mol Life Sci* : CM 2016;73:2491–509.
- [35] Ivey KN, Srivastava D. microRNAs as developmental regulators. *Cold Spring Harbor perspectives in biology* 2015;7: a008144.
- [36] Harrison SM, Seideman C, Baker LA. DNA copy number variations in patients with persistent cloaca. *J Urol* 2014;191: 1543–6.
- [37] Gredler ML, Seifert AW, Cohn MJ. Tissue-specific roles of *Fgfr2* in development of the external genitalia. *Development* 2015; 142:2203–12.
- [38] Liu D, Shen L, Tao Y, Kuang Y, Cai L, Wang D, et al. Alterations in gene expression during sexual differentiation in androgen receptor knockout mice induced by environmental endocrine disruptors. *Int J Mol Med* 2015;35:399–404.
- [39] Ross AE, Marchionni L, Phillips TM, Miller RM, Hurley PJ, Simons BW, et al. Molecular effects of genistein on male urethral development. *J Urol* 2011;185:1894–8.
- [40] Howdeshell KL, Wilson VS, Furr J, Lambright CR, Rider CV, Blystone CR, et al. A mixture of five phthalate esters inhibits fetal testicular testosterone production in the sprague-dawley rat in a cumulative, dose-additive manner. *Toxicol Sci* : an official journal of the Society of Toxicology 2008;105: 153–65.
- [41] Skakkebaek NE, Rajpert-De Meyts E, Buck Louis GM, Toppari J, Andersson AM, Eisenberg ML, et al. Male reproductive disorders and fertility trends: influences of environment and genetic susceptibility. *Physiol Rev* 2016;96:55–97.
- [42] Arbuckle TE, Agarwal A, MacPherson SH, Fraser WD, Sathyanarayana S, Ramsay T, et al. Prenatal exposure to phthalates and phenols and infant endocrine-sensitive outcomes: the MIREC study. *Environ Int* 2018;120:572–83.
- [43] Bornehag CG, Carlstedt F, Jonsson BA, Lindh CH, Jensen TK, Bodin A, et al. Prenatal phthalate exposures and anogenital distance in Swedish boys. *Environ Health Perspect* 2015;123: 101–7.
- [44] Swan SH, Sathyanarayana S, Barrett ES, Janssen S, Liu F, Nguyen RH, et al. First trimester phthalate exposure and anogenital distance in newborns. *Human reprod (Oxford, England)* 2015;30:963–72.
- [45] Poon S, Koren G, Carnevale A, Aleksa K, Ling J, Ozsarfaty J, et al. Association of in utero exposure to polybrominated diphenyl ethers with the risk of hypospadias. *JAMA pediatrics* 2018;172:851–6.
- [46] Bonde JP, Flachs EM, Rimborg S, Glazer CH, Giwercman A, Ramlau-Hansen CH, et al. The epidemiologic evidence linking prenatal and postnatal exposure to endocrine disrupting chemicals with male reproductive disorders: a systematic review and meta-analysis. *Hum Reprod Update* 2016;23:104–25.
- [47] Schnack TH, Zdravkovic S, Myrup C, Westergaard T, Christensen K, Wohlfahrt J, et al. Familial aggregation of hypospadias: a cohort study. *Am J Epidemiol* 2008;167:251–6.
- [48] Bouty A, Ayers KL, Pask A, Helouy Y, Sinclair AH. The genetic and environmental factors underlying hypospadias. *Sex Dev* 2015;9:239–59.
- [49] Sanger TJ, Gredler ML, Cohn MJ. Resurrecting embryos of the tuatara, *Sphenodon punctatus*, to resolve vertebrate phallus evolution. *Biol Lett* 2015;11.
- [50] Brennan PL. Evolution: one penis after all. *Curr Biol* 2016;26: R29–31.
- [51] Herrera AM, Cohn MJ. Embryonic origin and compartmental organization of the external genitalia. *Sci Rep* 2014;4:6896.
- [52] Tschopp P, Sherratt E, Sanger TJ, Groner AC, Aspiras AC, Hu JK, et al. A relative shift in cloacal location repositions

- external genitalia in amniote evolution. *Nature* 2014;516:391–4.
- [53] Wang S, Shi M, Zhu D, Mathews R, Zheng Z. External genital development, urethra formation, and hypospadias induction in Guinea pig: a double zipper model for human urethral development. *Urology* 2018;113:179–86.
- [54] Chew KY, Pask AJ, Hickford D, Shaw G, Renfree MB. A dual role for SHH during phallus development in a marsupial. *Sex Dev* 2014;8:166–77.
- [55] Chew KY, Renfree MB. Inducing sex reversal in marsupial mammals. *Sex Dev* 2016;10:301–12.
- [56] Lancaster MA, Renner M, Martin CA, Wenzel D, Bicknell LS, Hurler ME, et al. Cerebral organoids model human brain development and microcephaly. *Nature* 2013;501:373–9.
- [57] Lee EJ, Kim DE, Azeloglu EU, Costa KD. Engineered cardiac organoid chambers: toward a functional biological model ventricle. *Tissue Eng* 2008;14:215–25.
- [58] Takasato M, Er PX, Chiu HS, Maier B, Baillie GJ, Ferguson C, et al. Kidney organoids from human iPS cells contain multiple lineages and model human nephrogenesis. *Nature* 2015;526:564–8.
- [59] Kleinstreuer N, Dix D, Rountree M, Baker N, Sipes N, Reif D, et al. A computational model predicting disruption of blood vessel development. *PLoS Comput Biol* 2013;9:e1002996.
- [60] Menshykau D, Michos O, Lang C, Conrad L, McMahon AP, Iber D. Image-based modeling of kidney branching morphogenesis reveals GDNF-RET based Turing-type mechanism and pattern-modulating WNT11 feedback. *Nat Commun* 2019;10:239.
- [61] Multerer MD, Wittwer LD, Stopka A, Barac D, Lang C, Iber D. Simulation of morphogen and tissue dynamics. *Methods Mol Biol* 2018;1863:223–50.
- [62] Hutson MS, Leung MCK, Baker NC, Spencer RM, Knudsen TB. Computational model of secondary palate fusion and disruption. *Chem Res Toxicol* 2017;30:965–79.
- [63] Leung MC, Phuong J, Baker NC, Sipes NS, Klinefelter GR, Martin MT, et al. Systems toxicology of male reproductive development: profiling 774 chemicals for molecular targets and adverse outcomes. *Environ Health Perspect* 2016;124:1050–61.
- [64] Knudsen TB, Klieforth B, Slikker Jr W. Programming microphysiological systems for children's health protection. *Exp Biol Med* 2017;242:1586–92.
- [65] Leung MC, Hutson MS, Seifert AW, Spencer RM, Knudsen TB. Computational modeling and simulation of genital tubercle development. *Reprod Toxicol* 2016;64:151–61.
- [66] Belair DG, Wolf CJ, Moorefield SD, Wood C, Becker C, Abbott BD. A three-dimensional organoid culture model to assess the influence of chemicals on morphogenetic fusion. *Toxicol Sci* 2018;166(2):394–408.
- [67] Sturla SJ, Boobis AR, FitzGerald RE, Hoeng J, Kavlock RJ, Schirmer K, et al. Systems toxicology: from basic research to risk assessment. *Chem Res Toxicol* 2014;27:314–29.
- [68] Liu G, Liu X, Shen J, Sinclair A, Baskin L, Cunha GR. Contrasting mechanisms of penile urethral formation in mouse and human. *Differentiation; research in biological diversity* 2018;101:46–64.
- [69] Hashimoto D, Hyuga T, Acebedo AR, Alcantara MC, Suzuki K, Yamada G. Developmental mutant mouse models for external genitalia formation. *Congenital Anom* 2019;59:74–80.
- [70] Lund L, Engebjerg MC, Pedersen L, Ehrenstein V, Norgaard M, Sorensen HT. Prevalence of hypospadias in Danish boys: a longitudinal study, 1977-2005. *Eur Urol* 2009;55:1022–6.
- [71] Lane C, Boxall J, MacLellan D, Anderson PA, Dodds L, Romao RLP. A population-based study of prevalence trends and geospatial analysis of hypospadias and cryptorchidism compared with non-endocrine mediated congenital anomalies. *J Pediatr Urol* 2017;13:284.e1–7.
- [72] Springer A, van den Heijkant M, Baumann S. Worldwide prevalence of hypospadias. *J Pediatr Urol* 2016;12:152.e1–7.
- [73] Akbulut A, Ertugrul E, Topcu V. Fetal health status prediction based on maternal clinical history using machine learning techniques. *Comput Methods Progr Biomed* 2018;163:87–100.
- [74] Ahmed SF, Khwaja O, Hughes IA. The role of a clinical score in the assessment of ambiguous genitalia. *BJU Int* 2000;85:120–4.