

UC Irvine

UC Irvine Previously Published Works

Title

Inclusive Lepton Production in e^+e^- Annihilation at 29 GeV

Permalink

<https://escholarship.org/uc/item/1x37s684>

Journal

Physical Review Letters, 60(25)

ISSN

0031-9007

Authors

Ong, RA
Weir, AJ
Abrams, GS
et al.

Publication Date

1988-06-20

DOI

10.1103/physrevlett.60.2587

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Inclusive Lepton Production in e^+e^- Annihilation at 29 GeV

R. A. Ong,^(a) A. J. Weir,^(b) G. S. Abrams, D. Amidei,^(a) A. R. Baden, T. Barklow, A. M. Boyarski, J. Boyer, P. R. Burchat,^(c) D. L. Burke, F. Butler, J. M. Dorfan, G. J. Feldman, G. Gidal, L. Gladney,^(d) M. S. Gold, G. Goldhaber, L. Golding,^(e) J. Haggerty,^(f) G. Hanson, K. Hayes, D. Herrup, R. J. Hollebeek,^(d) W. R. Innes, J. A. Jaros, I. Juricic, J. A. Kadyk, D. Karlen, S. R. Klein, A. J. Lankford, R. R. Larsen, B. W. LeClaire,^(g) M. Levi, N. S. Lockyer,^(d) V. Lüth, M. E. Nelson,^(h) M. L. Perl, A. Petersen,⁽ⁱ⁾ B. Richter, K. Riles, P. C. Rowson,^(j) T. Schaad,^(k) H. Schellman,^(a) W. B. Schmidke, P. D. Sheldon,^(l) G. H. Trilling, D. R. Wood,^(m) and J. M. Yelton⁽ⁿ⁾

*Stanford Linear Accelerator Center, Stanford University, Stanford, California 94309, and
Lawrence Berkeley Laboratory and Department of Physics, University of California,
Berkeley, California 94720, and*

Harvard University, Cambridge, Massachusetts 02138

(Received 29 April 1988)

We use data taken by the Mark II detector at the SLAC storage ring PEP to study inclusive lepton production in e^+e^- annihilation at 29 GeV. From fits to the inclusive lepton momentum and transverse-momentum spectra we measure the semileptonic branching ratios for charm and bottom hadrons. We also determine with good precision the mean value of the bottom fragmentation function using both electron and muon samples.

PACS numbers: 13.20.Fc, 13.20.Jf, 13.65.+i, 13.87.Fh

Semileptonic decays of charm and bottom hadrons are the primary sources of leptons in hadronic events from high-energy e^+e^- annihilation. The momentum and transverse-momentum distributions of these leptons depend on the parent quark masses and fragmentation functions; the rates of production depend on the weak-decay semileptonic branching ratios. We separate the charm and bottom contributions to the prompt-lepton signal by using the harder transverse-momentum spectra of leptons from B decay. After making this separation, we use the lepton momentum spectra to study the fragmentation properties of charm and bottom hadrons. We have previously reported an analysis of inclusive electrons for a data sample with an integrated luminosity of 35 pb^{-1} .¹ In this Letter, we present new results for both inclusive muon and electron production from our full PEP data sample of 204 pb^{-1} .² The increased amount of data represents a significant statistical improvement over previous results and allows us to perform extensive crosschecks on our analysis procedure.

The data were taken with the Mark II detector at the PEP e^+e^- storage ring at the Stanford Linear Accelerator Center. The Mark II detector has been described previously,³ as have the elements necessary for electron identification.¹ We describe here the components essential for the identification of muons. The momenta of charged particles are measured in two cylindrical drift chambers immersed in an axial magnetic field of 2.3 kG. The rms momentum resolution is given by $(\delta p/p)^2 \approx (0.025)^2 + (0.011 p_\perp)^2$, where $p_\perp$, the momentum perpendicular to the beam direction, is in units of GeV/c. The muon detectors consist of layers of hadron absorber and proportional chambers mounted on

four sides of the central detector. Each wall has four alternating layers of iron and chambers; the innermost layer measures the polar angle of the track while the outer three layers measure the azimuthal angle. The proportional chamber wires are separated by 2.5 cm. Including the electromagnetic calorimeter, there are 7.3 nuclear interaction lengths preceding the fourth muon layer. The solid-angle coverage of the outer layer of the muon system is 45% of 4π .

Muons with momenta greater than 1.8 GeV/c are expected to penetrate to the outermost muon layer, leaving proportional chamber hits in all four layers. Hadrons, however, are expected to be absorbed in the iron as a result of inelastic nuclear collisions. In the muon-hadron separation algorithm, drift-chamber tracks are extrapolated into the muon system. At each of the four muon layers a mean standard deviation due to Coulomb scattering, extrapolation error, and the proportional chamber intrinsic resolution is calculated based on the total amount of material traversed. A muon candidate is required to leave a hit within two standard deviations of the track extrapolation in all four layers. The muon identification efficiency has been measured with the use of $e^+e^- \rightarrow e^+e^-\mu^+\mu^-$ events and varies from 78% at $p = 2 \text{ GeV/c}$ to 91% at high momenta.⁴

The background to the lepton sample has two components, decay and misidentification of hadrons. What we call *decay* muons arise from charged-pion and -kaon decays, and *decay* electrons come from Dalitz π^0 decays and photon conversions in the drift chamber. *Misidentified* muons consist of hadronic primary or secondary punchthrough in the muon system and *misidentified* electrons arise from hadronic interaction or

photon overlap in the calorimeter. We describe here our estimation of these backgrounds to the muon signal; the backgrounds to the electron signal are discussed in detail in Ref. 1 (typical electron misidentification probabilities are 0.5% but are as high as 1.6% for 2-GeV/ c tracks in the core of a jet). The muon decay background is estimated with a detailed Monte Carlo simulation of the decays of pions and kaons in the drift chamber. The per-track probability for such a background decay varies from 0.4% at 2 GeV/ c to 0.25% at 8 GeV/ c with a systematic uncertainty of 20%. The muon misidentification background has been estimated in three ways: (1) with a source of pions from $\tau^\pm \rightarrow \pi^\pm \pi^\pm \pi^\mp \nu$ events, (2) by our performing a fit to the distribution of muon chamber hits among the four layers for tracks in hadronic events failing the muon identification criteria,⁵ and (3) with a detailed Monte Carlo simulation⁶ of hadronic interactions in the material preceding and included in the muon system. From these studies we estimate the per-track misidentification probability to vary from 0.28% at 2 GeV/ c to 0.45% at 8 GeV/ c with a systematic uncertainty of 40%.

The hadronic-event sample was selected by our requiring events to have at least six charged tracks. We required the total scalar momenta of the charged tracks to be greater than 7.25 GeV. The event vertex was required to lie within 4 cm of the beam position in the direction perpendicular to the beams and within 10 cm in the direction along the beams. In each event the thrust direction was calculated from charged tracks (including any lepton candidates). To ensure that the event was well contained in the detector, we required that the absolute value of the cosine of the angle between the thrust direction and the beam direction was less than 0.7. After these cuts there were 64 459 events.

We searched in these hadronic events for lepton candidates with momenta greater than 2 GeV/ c . Background electrons from Dalitz π^0 decays and photon conversions were removed by a pair-finding algorithm.⁷ We also required each lepton candidate to have at least two associated hits in the inner drift chamber to suppress conversions occurring between the inner and main drift chambers.

There is a background to the inclusive electron signal from the two-photon process $e^+e^- \rightarrow e^+e^- + \text{hadrons}$ where one of the electrons scatters into the central detector. Momentum conservation in the plane perpendicular to the beams requires that the hadronic shower recoils against the scattered electron; as a result, the two-photon events often have the topology of an isolated energetic electron opposite a low-energy hadron jet. We imposed several cuts taking advantage of this topology to suppress the two-photon background.⁸ These cuts removed approximately 5% of the electron sample. We measured the sensitivity of these cuts by using a Monte Carlo simulation that incorporates complete lowest calculations for the $e^+e^- \rightarrow q\bar{q}$ process,⁹ followed by fragmentation

of the quarks into hadrons using the LUND 6.3 shower code.¹⁰ From this study, we found good agreement between the number of events removed by the two-photon cuts and the number predicted by the simulation. We estimate the background from two-photon processes to contribute less than 2% contamination in the electron sample. The background in the lepton signals from τ pair production is negligible.

After all cuts, we were left with a sample of 2621 electron and 1230 muon candidates. We determined the transverse momentum of each lepton measured relative to the thrust axis. The lepton candidates were separated into bins of momentum (p) and transverse momentum (p_t), of width 0.5 and 0.25 GeV/ c , respectively.

We performed binned maximum-likelihood fits to the observed lepton p , p_t distributions. The sources of these observed leptons are (1) background from misidentified hadrons, (2) background from decay of hadrons, (3) primary bottom-quark decay, (4) primary charm-quark decay, and (5) decay of charm quarks in $b\bar{b}$ events (secondary b decay). To estimate the contribution from charm and bottom decay we used a Monte Carlo program to produce bottom and charm quarks and string fragmentation.¹⁰ We used the Peterson parametrization¹¹ of the fragmentation function

$$f(z) = \{z[1 - 1/z - \epsilon/(1 - z)]^2\}^{-1},$$

where $z \equiv (E + p_\parallel)_{\text{hadron}}/(E + p)_{\text{quark}}$ and ϵ is a free parameter. The variables in the fits were the average semileptonic branching ratios for B - and C -hadron decay and $\langle z_b \rangle$, the mean of the fragmentation function for b -quark hadronization. We also allowed the background levels to vary by including multiplicative scale factors for the amount of misidentification and decay backgrounds. The values of the scale factors obtained from the fit provided a check of our background estimates. We generated $b\bar{b}$ events at different values of $\langle z_b \rangle$; in the fit we interpolated between them to obtain the value which best fitted the data. The value of $\langle z_c \rangle$ was chosen to be 0.68 in agreement with exclusive D^* measurements¹² and the results of other inclusive lepton analyses.¹³

The results obtained from the fit are shown in Table I. The first error for each result is statistical and the second

TABLE I. Inclusive lepton results. The second column shows the results of this analysis. In the third column, the world average values are given. The measured quantities are described in the text.

Quantity	This analysis	World average
$B(C \rightarrow l)(\%): \mu$	$7.8 \pm 0.9 \pm 1.2$	$7.9 \pm 0.7 \pm 1.1$
e	$9.6 \pm 0.7 \pm 1.5$	$10.5 \pm 0.6 \pm 0.8$
$B(B \rightarrow l)(\%): \mu$	$11.8 \pm 1.2 \pm 1.0$	$12.6 \pm 0.9 \pm 0.8$
e	$11.2 \pm 0.9 \pm 1.1$	$12.5 \pm 1.0 \pm 1.0$
$\langle z_b \rangle: \mu$	$0.82 \pm 0.04 \pm 0.05$	$0.84 \pm 0.03 \pm 0.03$
e	$0.85 \pm 0.03 \pm 0.05$	$0.79 \pm 0.03 \pm 0.03$

is systematic. The lepton momentum and transverse-momentum spectra are shown in Fig. 1. The systematic errors arise from a number of sources of which the uncertainty in the background estimate is dominant.

To account for background uncertainties, the fits were repeated with the background level allowed to vary within the systematic errors assigned during its estimation. We also allowed the shape of the background contribution to vary (to account, for example, for a different K/π ratio from that assumed). The value of $\langle z_c \rangle$ was allowed to vary within the range of 0.68 ± 0.06 , accounting for the finite precision of the exclusive D^* measurements and the possible inclusion of charmed hadrons other than the D^* . Although we did not include a term to account for leptons from $b \rightarrow \tau \rightarrow l$ decays, we allowed this branching ratio to be as large as 5% for the estimate of the systematic errors. We accounted for uncertainties in the lepton-identification efficiency, the fiducial acceptance, and the number of nonhadronic events in the data. The systematic error resulting from uncertainty in the rest-frame momentum and p_t distributions of leptons from heavy-quark decays was found to be negligible. In the fits we assumed that b quarks decay 100% of the time to c quarks. The systematic effect of the possible inclusion of 5% fraction of $b \rightarrow u$ transitions had a negligible effect on the lepton momentum spectra.

The summed χ^2 for the fits are 51.7 and 85.8 for the muon and electron distributions, respectively, for 58 degrees of freedom. The electron fit has a small number of bins with large χ^2 values. These bins do not cluster at specific values of p or p_t , indicating that the large χ^2 for

the electron fit is likely due to statistical fluctuations. The values for the background scale factors obtained from the fit agree very well with our estimates of the decay and misidentification backgrounds. We have verified the analysis procedure by fitting a sample of 100 000 hadronic Monte Carlo events. The values obtained from the fit agree well with those used in the generation.

World average values for the branching ratios and fragmentation $\langle z_b \rangle$ are given in Table I. These values are calculated with use of previously published measurements from experiments at the SLAC and DESY storage rings PEP and PETRA;^{12,13} earlier results from the Mark II Collaboration are not included.¹⁴ The measurements presented in this analysis are in good agreement with the earlier results and have comparable precision to the world average values. The measurements also indicate that B fragmentation is peaked towards high values of $\langle z_b \rangle$, confirming previous results.

In conclusion, we have reported precise measurements of the momentum and transverse-momentum distributions for prompt muons and electrons in hadronic events in e^+e^- annihilation at 29 GeV. From a fit to these distributions we determine the semimuonic branching ratios to be $B(B \rightarrow \mu) = (11.8 \pm 1.2 \pm 1.0)\%$ and $B(C \rightarrow \mu) = (7.8 \pm 0.9 \pm 1.2)\%$, and the semielectronic branching ratios to be $B(B \rightarrow e) = (11.2 \pm 0.9 \pm 1.1)\%$ and $B(C \rightarrow e) = (9.6 \pm 0.7 \pm 1.5)\%$. The mean value of the fragmentation function for the b quark is found to be $\langle z_b \rangle(\mu) = 0.82 \pm 0.04 \pm 0.05$ and $\langle z_b \rangle(e) = 0.85 \pm 0.03 \pm 0.05$.¹⁵

This work is supported in part by the Department of




FIG. 1. Lepton momentum and transverse-momentum spectra for the muon and electron samples. The histograms represent the contributions from background, charm decay, and bottom decay (including secondary b decay). The lepton momentum spectra for (a),(d) $p_t < 1.0$ GeV/c; (b),(e) $p_t > 1.0$ GeV/c. (c),(f) The lepton transverse-momentum spectra. The highest-momentum bins in (a), (b), (d), and (e) contain all tracks with momenta greater than 6 GeV/c; the highest- p_t bins in (c) and (f) contain all tracks with p_t greater than 1.5 GeV/c. The data points are shown with statistical errors only.

Energy Contracts No. DE-AC03-76SF00515, No. DE-AC03-76SF00098, No. DE-AC02-76ER03064, and No. DE-AC03-81-ER40050.

^(a)Present address: University of Chicago, Chicago, IL 60637.

^(b)Present address: California Institute of Technology, Pasadena, CA 91125.

^(c)Present address: University of California, Santa Cruz, CA 95064.

^(d)Present address: University of Pennsylvania, Philadelphia, PA 19104.

^(e)Present address: Therma-Wave Corporation, Fremont, CA 94539.

^(f)Present address: Brookhaven National Laboratory, Upton, NY 11973.

^(g)Present address: University of Wisconsin, Madison, WI 53706.

^(h)Present address: California Institute of Technology, Pasadena, CA 91125.

⁽ⁱ⁾Present address: SCS, Hamburg, Federal Republic of Germany.

^(j)Present address: Columbia University, New York, NY 10027.

^(k)Present address: Université de Genève, CH-1211, Genève 4, Switzerland.

^(l)Present address: University of Illinois, Urbana, IL 61801.

^(m)Present address: CERN, CH-1211, Genève 23, Switzerland.

⁽ⁿ⁾Present address: University of Florida, Gainesville, FL 32611.

¹M. E. Nelson *et al.*, Phys. Rev. Lett. **50**, 1542 (1983).

²R. A. Ong, Ph.D. thesis, Stanford University, 1987, unpublished.

³R. H. Schindler *et al.*, Phys. Rev. D **24**, 78 (1981).

⁴The muon chamber hits in the outer three layers are somewhat correlated.

⁵The parameters of the fit are the punchthrough probabilities to the first three layers. The punchthrough to the fourth

layer is obtained by extrapolation.

⁶We used the hadronic-interaction simulation program FLUKA; P. A. Aarnio *et al.*, CERN Report No. CERN-TIS-RP/168, 1986 (unpublished).

⁷M. E. Nelson, Ph.D. thesis, University of California, Berkeley, Report No. LBL-16724, 1983 (unpublished).

⁸In particular, events are rejected if they contain an electron track satisfying any of the following cuts: (1) momentum greater than 9 GeV/c; (2) transverse momentum relative to the thrust axis greater than 2.5 GeV/c; or (3) the electron track isolated by at least 90° (in the plane perpendicular to the beams) from the nearest charged track.

⁹F. A. Berends, P. H. Daverveldt, and R. Kleiss, Nucl. Phys. **B253**, 441 (1985), and Comput. Phys. Commun. **40**, 271 (1986).

¹⁰T. Sjöstrand, Comput. Phys. Commun. **39**, 347 (1986).

¹¹C. Peterson *et al.*, Phys. Rev. D **27**, 105 (1983).

¹²W. Bartel *et al.*, Z. Phys. C **33**, 339 (1987).

¹³H. J. Behrend *et al.*, Z. Phys. C **19**, 291 (1983); B. Aveda *et al.*, Phys. Rev. Lett. **50**, 443 (1983); E. Fernandez *et al.*, Phys. Rev. Lett. **50**, 2054 (1983); M. Althoff *et al.*, Z. Phys. C **22**, 219 (1984); M. Althoff *et al.*, Phys. Lett. **146B**, 443 (1984); D. E. Koop *et al.*, Phys. Rev. Lett. **52**, 970 (1984); H. Aihara *et al.*, Phys. Rev. D **31**, 2719 (1985); H. Aihara *et al.*, Z. Phys. C **28**, 31 (1985); T. Pal *et al.*, Phys. Rev. D **33**, 2708 (1986). In addition, there are branching-ratio measurements from other e^+e^- experiments at the $\psi(3770)$ and $Y(4S)$ resonances; these measurements are summarized in Ref. 12.

¹⁴The world averages were computed by weighting of individual results by their statistical and systematic errors combined in quadrature. Fragmentation $\langle x \rangle$ values were scaled into the corresponding $\langle z \rangle$ values as discussed in Ref. 12. The systematic error for an average value was taken to be the smallest systematic error of the individual measurements. This conservative approach was adopted because the measurements largely have common systematic errors.

¹⁵The value of the parameter ϵ_b in the Peterson formula was determined to be $0.0069^{+0.0067+0.0089}_{-0.0038-0.0045}$ with muons, and $0.0038^{+0.0031+0.0039}_{-0.0019-0.0027}$ with electrons. These values were obtained from $\langle z_b \rangle$.




FIG. 1. Lepton momentum and transverse-momentum spectra for the muon and electron samples. The histograms represent the contributions from background, charm decay, and bottom decay (including secondary b decay). The lepton momentum spectra for (a),(d) $p_t < 1.0$ GeV/c; (b),(e) $p_t > 1.0$ GeV/c. (c),(f) The lepton transverse-momentum spectra. The highest-momentum bins in (a), (b), (d), and (e) contain all tracks with momenta greater than 6 GeV/c; the highest- p_t bins in (c) and (f) contain all tracks with p_t greater than 1.5 GeV/c. The data points are shown with statistical errors only.