

UC Irvine

Clinical Practice and Cases in Emergency Medicine

Title

Nystagmus Associated with Carbamazepine Toxicity

Permalink

<https://escholarship.org/uc/item/1xp1p2cb>

Journal

Clinical Practice and Cases in Emergency Medicine, 1(4)

Authors

Wirfs, Lauran
Whitworth, Kristen
Kellar, Jesse

Publication Date

2017

DOI

10.5811/cpcem.2017.6.34772

Supplemental Material

<https://escholarship.org/uc/item/1xp1p2cb#supplemental>

Copyright Information

Copyright 2017 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Nystagmus Associated with Carbamazepine Toxicity

Lauran Wirfs, BS*
Kristen Whitworth, DO†
Jesse Kellar, MD†

*Midwestern University, Chicago College of Osteopathic Medicine, Downers Grove, Illinois

†Lakeland Health, Department of Emergency Medicine, Saint Joseph, Michigan

Section Editor: Rick A. McPheeters, DO

Submission history: Submitted May 6, 2017; Revision received June 2, 2017; Accepted June 8, 2017

Electronically published October 18, 2017

Full text available through open access at http://escholarship.org/uc/uciem_cpce

DOI: 10.5811/cpcem.2017.6.34772

[Clin Pract Cases Emerg Med. 2017;1(4):441–442.]

CASE PRESENTATION

A 42-year-old male with past medical history significant for epilepsy presented to the emergency department (ED) complaining of dizziness, difficult ambulation, and blurred vision. Vitals were only significant for a blood pressure of 143/89 mm Hg. His neurogenic exam revealed gaze-evoked nystagmus (GEN) (Video) as well as subtle dysmetria and ataxia. The remainder of his physical exam was unremarkable. Computed tomography of the head without contrast showed no acute intracranial abnormality. Upon further questioning, it was discovered he had been unintentionally doubling his dose of carbamazepine 400mg BID because two physicians had prescribed him the medication at the same time. His laboratory studies showed an elevated carbamazepine level at 18.2 ug/mL (normal range 4-12 ug/mL), and he was admitted for observation. The next morning all his symptoms (including his nystagmus) had resolved.

DIAGNOSIS

Nystagmus Associated with Carbamazepine Toxicity

Mild GEN may be seen in the normal population with no underlying pathological cause.¹⁻² However, exaggerated GEN is pathologic and can be caused by drugs, structural brain abnormalities, or certain diseases (e.g. myasthenia gravis, or demyelinating diseases).²⁻³ Carbamazepine, whose structure is related to tricyclic antidepressants, is an anticonvulsant drug used in the treatment of epilepsy, bipolar affective disorders, and trigeminal neuralgia.⁴⁻⁶ The plasma-level therapeutic range of carbamazepine effective for seizure prophylaxis is 4-12 ug/mL. However, even at higher ends of the therapeutic range, patients can experience adverse effects such as diplopia, blurred vision, nystagmus, or ataxia.⁴ In severe toxicity, carbamazepine may induce seizures or altered level of consciousness, and progress to coma.⁴⁻⁷ Treatment of carbamazepine toxicity consists of trending serum levels of carbamazepine and

CPC-EM Capsule

What do we already know about this clinical entity?

Carbamazepine is an anticonvulsant used to treat epilepsy, bipolar disorder and trigeminal neuralgia. High doses can elicit diplopia, blurred vision, nystagmus, or ataxia. Severe toxicity may induce seizures and progress to coma.

What is the major impact of the image(s)?

This video exhibits exaggerated gaze-evoked nystagmus (GEN) secondary to carbamazepine toxicity.

How might this

improve emergency medicine practice?

Mild GEN can occur with no underlying pathology. Exaggerated GEN is pathologic and can be caused by drugs, structural brain abnormalities, or disease. The physician should consider these differentials in patients with exaggerated GEN.

supportive care with intravenous fluids as needed for hypotension and benzodiazepines to control seizures.⁷ Patients should be monitored until signs and symptoms of toxicity resolve, and they are deemed medically stable.⁵⁻⁷

Video. Gaze-evoked nystagmus on exam.

Address for Correspondence: Kristen Whitworth, DO, Lakeland Health, Department of Emergency Medicine, 1234 Napier Avenue, Saint Joseph, MI 49085. Email: Kwhitworth1@lakelandhealth.org.

Conflicts of Interest: By the *CPC-EM* article submission agreement, all authors are required to disclose all affiliations, funding sources and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

Copyright: © 2017 Wirfs et al. This is an open access article distributed in accordance with the terms of the Creative Commons Attribution (CC BY 4.0) License. See: <http://creativecommons.org/licenses/by/4.0/>

REFERENCES

1. Horton J, Kasper D, Fauci A, et al. Harrison's Principles of Internal Medicine. 19th ed. New York: McGraw-Hill; 2014:210-211.
2. Aminoff MJ, Greenberg DA, Simon RP. Clinical Neurology. 9th ed. New York: McGraw-Hill; 2015: Neuro-Ophthalmic Disorders.
3. Wong A. Eye Movement Disorders. 1st ed. New York: Oxford University Press; 2007: Nystagmus.
4. Bauer L. Applied Clinical Pharmacokinetics. 3rd ed. New York: McGraw-Hill; 2014: Carbamazepine.
5. Kearney T, Olson K. Poisoning & Drug Overdose. 6th ed. New York: McGraw-Hill; 2012: Carbamazepine and Oxcarbazepine.
6. Gaudins A, Peden G, Dowsett R. Massive overdose with controlled-release carbamazepine resulting in delayed peak serum concentrations and life-threatening toxicity. *Emerg Med (Fremantle)*. 2002;14(1):89-94.
7. Doyon S, Hoffman R, Howland M, et al. Goldfrank's Toxicologic Emergencies. 10th ed. New York: McGraw-Hill; 2015: Antiepileptics.