

# UC Irvine

## UC Irvine Previously Published Works

### Title

Cortical Network Dynamics in Acute Ischemic Stroke

### Permalink

<https://escholarship.org/uc/item/1zm0m5n5>

### Journal

STROKE, 44(2)

### ISSN

0039-2499

### Authors

Liu, Virginia  
Winter, William  
Xie, Kate  
et al.

### Publication Date

2013

### Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



# Stroke

 FREE ACCESS | **ARTICLE**

INTERNATIONAL STROKE CONFERENCE POSTER ABSTRACTS  
DIAGNOSIS POSTERS I

## Abstract WP216: Cortical Network Dynamics in Acute Ischemic Stroke

Virginia Liu, William Winter, Kate Xie, Shrey Kanjiya, George Tran, Ramesh Srinivasan, and Steven C Cramer Univ of California, Irvine, Orange, CA

Originally published 1 Feb 2013 | Stroke. 2013;44:AWP216

### Abstract


**INTRODUCTION:** Measures of brain function are a unique source of insight into stroke effects, but the most commonly used methods (fMRI, PET) are difficult to implement in acutely ill patients, and so little is known about changes in brain function in the days after stroke. Advances in dense-array electroencephalography (dEEG) make possible the study of cortical network dynamics at the bedside. The current study addressed the hypothesis that changes in the motor network are measurable in first days post-stroke, and that these network changes are related to behavioral deficits.


**METHODS:** Nine patients with acute (<120 hours) ischemic stroke underwent dEEG and behavioral testing that included Box & Blocks score (1 min). The dEEG signals from 256 scalp leads were recorded at 1000Hz, along with bilateral forearm EMG, during cued 0.14 Hz wrist flexions. Nine age-matched healthy controls underwent the same assessments. Cortical network dynamics were assessed in the motor system using EEG coherence between leads, focusing on  $\beta$  range (20-30 Hz) signals. Spectral power in this range was also obtained.


**RESULTS:** Patients were  $62 \pm 13$  years old, with median NIHSS score=3 (range 1-12), and could move  $26 \pm 18$  blocks (range 0-50) on the Box & Blocks test, fewer than controls ( $59 \pm 7$  blocks,  $p < 0.0001$ ). In patients, as compared to controls, the region overlying ipsilesional hand motor cortex showed increased coherence with [A] contralesional hand motor cortex ( $0.13 \pm 0.06$  vs  $0.07 \pm 0.05$ ,  $p = 0.03$ ) and [B] supplementary motor area ( $0.27 \pm 0.12$  vs  $0.15 \pm 0.07$ ,  $p = 0.02$ ) during flexion of the affected wrist. Spectral power did not differ between groups in these regions. Across all subjects, the magnitude of coherence between hand motor cortex and supplementary motor area was inversely related to Box & Blocks score ( $r = -0.61$ ,  $p = 0.008$ ).

**CONCLUSIONS:** Assessment of cortical dynamics in the motor system is feasible in the acute stroke setting, with hemiparetic patients demonstrating significantly increased  $\beta$  coherence. Increased EEG coherence between two motor cortex regions after stroke might reflect increased functional connectivity due to the reduced cortical inhibition that is seen early after stroke, effects of deafferentation, or increased input from a common area such as the thalamus.


## Footnotes



 Back to top



# Stroke

**AHA Journals**

Arteriosclerosis, Thrombosis, and Vascular Biology (ATVB)

Circulation

Circ: Arrhythmia and Electrophysiology

Circ: Genomic and Precision Medicine

Circ: Cardiovascular Imaging

Circ: Cardiovascular Interventions

Circ: Cardiovascular Quality & Outcomes

Circ: Heart Failure

Circulation Research

Hypertension

Stroke

Journal of the American Heart Association (JAHA)

**Journal Information**

About Stroke

Editorial Board

Reprints