

Lawrence Berkeley National Laboratory

Recent Work

Title

EXAFS and thermodynamic studies of uranyl complexes of HEDPA in neutral and alkaline solutions

Permalink

<https://escholarship.org/uc/item/2176v3b5>

Author

Nash, K.L.

Publication Date

2002-04-04

“EXAFS and thermodynamic studies of uranyl complexes of HEDPA in neutral and alkaline solutions”

W. A. Reed^a, A. Yu. Garnov^a, L. Rao^a and K. L. Nash^b

^a *Glenn T. Seaborg Center, Lawrence Berkeley National Laboratory, 1 Cyclotron Road, Berkeley, California 94720*

^b *Chemistry Division, Argonne National Laboratory, 9700 S. Cass Avenue, Argonne, Illinois 60439*

HEDPA (1-hydroxyethane-1,1-diphosphonic acid) and other diphosphonic acids are remarkably powerful chelating agents in moderately acidic solutions and have found applications in separations and waste treatments. Their coordination chemistry with lanthanides and actinides have been studied, but mostly in acidic solutions. There are structural data on the complexes of HEDPA with lanthanide elements, but not the complex with uranyl(VI). This study is focused on the coordination of HEDPA with uranyl(VI). Efforts are made to evaluate the complexes in neutral to basic solutions using EXAFS and stability constant data.

This work was supported by the US Department of Energy, under Contract No. DE-AC03-76SF00098.