

UC Davis

UC Davis Previously Published Works

Title

Associations among MHC genes, latitude, and avian malaria infections in the rufous-collared sparrow (*Zonotrichia capensis*).

Permalink

<https://escholarship.org/uc/item/21g3b352>

Journal

Ecology and Evolution, 14(7)

ISSN

2045-7758

Authors

Rivero de Aguilar, Juan

Barroso, Omar

Bonaccorso, Elisa

et al.

Publication Date

2024-07-01

DOI

10.1002/ece3.11634















Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

RESEARCH ARTICLE

Associations among MHC genes, latitude, and avian malaria infections in the rufous-collared sparrow (*Zonotrichia capensis*)

Juan Rivero de Aguilar^{1,2,3}  | Omar Barroso¹  | Elisa Bonaccorso^{4,5}  |
 Hector Cadena⁵  | Lucas Hussing² | Josefina Jorquera⁶ |
 Javier Martinez⁷  | Josué Martínez-de la Puente^{8,9}  | Alfonso Marzal^{10,11}  |
 Fabiola León Miranda⁶ | Santiago Merino¹²  | Nubia E. Matta¹³  |
 Marilyn Ramenofsky¹⁴  | Ricardo Rozzi¹  | Carlos E. Valeris-Chacín^{1,2}  |
 Rodrigo A. Vásquez^{1,2,3} | Juliana A. Vianna^{6,15}  | John C. Wingfield¹⁴ 

¹Centro Subantártico Cabo de Hornos (CHIC), Universidad de Magallanes, Puerto Williams, Chile

²Departamento de Ciencias Ecológicas, Facultad de Ciencias, Universidad de Chile, Santiago, Chile

³Instituto de Ecología y Biodiversidad, Facultad de Ciencias, Universidad de Chile, Santiago, Chile

⁴Laboratorio de Biología Evolutiva, and Instituto Biósfera, Colegio de Ciencias Biológicas y Ambientales, Universidad San Francisco de Quito, Quito, Ecuador

⁵Centro de la Biodiversidad y Cambio Climático, Universidad Tecnológica Indoamérica, Quito, Ecuador

⁶Departamento de Ecología, Instituto Para el Desarrollo Sustentable, Facultad de Ciencias Biológicas, Pontificia Universidad Católica de Chile, Santiago, Chile

⁷Departamento de Biomedicina y Biotecnología (Area de Parasitología), Universidad de Alcalá de Henares, Madrid, Spain

⁸Departamento de Parasitología, Universidad de Granada, Granada, Spain

⁹Ciber de Epidemiología y Salud Pública (CIBERESP), Madrid, Spain

¹⁰Departamento de Anatomía, Biología Celular y Zoología, Facultad de Biología, Universidad de Extremadura, Badajoz, Spain

¹¹Grupo de Investigaciones en Fauna Silvestre, Universidad Nacional de San Martín, Tarapoto, Peru

¹²Departamento de Ecología Evolutiva, Museo Nacional de Ciencias Naturales CSIC, Madrid, Spain

¹³Departamento de Biología, Facultad de Ciencias, Universidad Nacional de Colombia, Bogotá, Colombia

¹⁴Department of Neurobiology, Physiology and Behavior, University of California, Davis, California, USA

¹⁵Millennium Institute Center for Genome Regulation (CRG), Millennium Institute of Biodiversity of Antarctic and Subantarctic Ecosystems (BASE), Millennium Nucleus of Patagonian Limit of Life (LiLi), Santiago, Chile

Correspondence

Juan Rivero de Aguilar, Cape Horn
 International Center (CHIC), University of
 Magallanes, Puerto Williams, Cape Horn,
 Chile.
 Email: juan.rivero@umag.cl

Funding information

Universidad San Francisco de
 Quito, Grant/Award Number: HUBI
 12424; FONDECYT - CONICYT
 Postdoctorado Programa Iniciativa
 Milenio - ICN2021_044 (CGR), Grant/
 Award Number: 3170211; CONICYT PIA
 APOYO CCTE, Grant/Award Number:
 AFB170008 and IEB-FB210006; ANID
 - Convocatoria Nacional Subvención a la

Abstract

The major histocompatibility complex (MHC) is a genetic region in jawed vertebrates that contains key genes involved in the immune response. Associations between the MHC and avian malaria infections in wild birds have been observed and mainly explored in the Northern Hemisphere, while a general lack of information remains in the Southern Hemisphere. Here, we investigated the associations between the MHC genes and infections with *Plasmodium* and *Haemoproteus* blood parasites along a latitudinal gradient in South America. We sampled 93 rufous-collared sparrows (*Zonotrichia capensis*) individuals from four countries, Colombia, Ecuador, Peru, and Chile, and estimated MHC-I and MHC-II allele diversity. We detected between 1–4 (MHC-I) and 1–6 (MHC-II) amino acidic alleles per individual, with signs of positive

This is an open access article under the terms of the [Creative Commons Attribution](https://creativecommons.org/licenses/by/4.0/) License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2024 The Author(s). *Ecology and Evolution* published by John Wiley & Sons Ltd.

Instalación en la Academia, 2020, Grant/Award Number: PAI77200078; Cape Horn International Center (CHIC), Grant/Award Number: ANID/BASAL FB210018; ANID - Programa Iniciativa Milenio, Grant/Award Number: ICN2021_002, NCN2021-050 and ICN2021_044; Ministerio de Ciencia e Innovación, Grant/Award Number: PID2020-118205GB-I00; ERDF; FEDER Extremadura 2021-2027 Operational Program of the Recovery, Transformation and Resilience Plan; Consejería de Economía e Infraestructura of the Junta de Extremadura; European Regional Development Fund, Grant/Award Number: IB20089

selection. We obtained generalized additive mixed models to explore the associations between MHC-I and MHC-II diversity and latitude. We also explored the relationship between infection status and latitude/biome. We found a non-linear association between the MHC-II amino acid diversity and latitude. Individuals from north Chile presented a lower MHC genetic diversity than those from other locations. We also found an association between deserts and xeric shrublands and a lower prevalence of *Haemoproteus* parasites. Our results support a lower MHC genetic diversity in arid or semi-arid habitats in the region with the lower prevalence of *Haemoproteus* parasites.

KEYWORDS

haemosporidian parasites, major histocompatibility complex, parasite-mediated selection

TAXONOMY CLASSIFICATION

Disease ecology

1 | INTRODUCTION

The major histocompatibility complex (MHC) is a genetic region in jawed vertebrates harboring key immune genes involved in the immune response (Kaufman, 2018). Among them, the classical MHC-I genes and MHC-II genes code for proteins that conform to the so-called “MHC molecules” (Martin & Kaufman, 2022), which are involved in the presentation of parasite antigens (in the form of short peptides) to T lymphocytes (Rock et al., 2016). If peptides are recognized as non-self, it triggers the activation of cytotoxic T cells or switching on B cells to produce antibodies (Radwan et al., 2020). Because of this key role in immune defenses, MHC genes have been described as the “center of the immunological universe” (Trowsdale, 1995). MHC-I molecules mainly present peptides from intracellular antigens, in contrast, MHC-II present those from extracellular ones (Hess & Edwards, 2002).

Depending on the species, avian MHC genes have been found being both polygenic (one to several loci) and polymorphic (multiple alleles at each locus). Individual MHC is characterized by the number of gene copies (specific to each species) (Minias et al., 2019; Westerdahl et al., 2022) and by the heterozygosity at each locus (Alcaide et al., 2008). These will determine the number of MHC molecules (or MHC diversity) expressed by an individual, and collectively that of an entire population. MHC genes are codominantly expressed (both alleles at each locus) (Murphy & Weaver, 2017); thus, depending on the MHC genotype, each individual will express different MHC molecules, varying in their ability to present antigens (Nikolich-Zugich, 2004).

MHC are the most polymorphic genes found in vertebrates (Borghans et al., 2004), and this polymorphism results from the elevated variability observed in the peptide binding region (PBR), the cleft where peptides accommodate in the MHC molecule (Lenz, 2011). Different non-exclusive mechanisms have been proposed to explain MHC diversity (Edwards & Hedrick, 1998; Piertney & Oliver, 2006; Radwan et al., 2020; Spurgin & Richardson, 2010; van Oosterhout, 2009). Among them, the parasite-mediated

selection hypothesis states that MHC diversity is maintained by selective pressures caused by parasites (Piertney & Oliver, 2006). MHC should reflect the past and current selective pressures that a species (or population) has suffered over evolutionary time (Hasselquist, 2007; Levy et al., 2020; Minias et al., 2019; O'Connor et al., 2019). Within birds, passerines generally have more MHC gene copies than non-passerines, evidencing different evolutionary histories (Minias et al., 2019; O'Connor et al., 2019). Having an elevated MHC allele diversity (heterozygous advantage hypothesis) or rare MHC alleles (rare allele advantage hypothesis) would increase the recognition and presentation of peptides from parasites to the immune system cells (Spurgin & Richardson, 2010). In turn, spatially heterogeneous selective pressures from parasites would maintain different MHC alleles at a local scale (Loiseau et al., 2009).

Associations between MHC and the occurrence of diseases have been found in humans and other vertebrates (Lundie et al., 2008; Sanchez-Mazas, 2020), supporting the role of parasites in the evolution of MHC. Several resistance/susceptibility associations in birds have been observed in both experimental and field studies (reviewed in O'Connor et al., 2019), thus stimulating further research on the role of MHC in wild birds (Hasselquist, 2007; Minias et al., 2019; O'Connor et al., 2019). Due to its global distribution and relative ease of sampling, avian malaria parasites *Plasmodium* and *Haemoproteus* have become an excellent model for studying the ecology of host-parasite interactions in birds (Santiago-Alarcon & Marzal, 2020). Parasites of the *Plasmodium* and *Haemoproteus* genera have a cosmopolitan distribution and include diverse molecular and morphological species (Clark et al., 2014). These parasites are blood protozoan parasites that commonly infect birds in the wild (Valkiūnas, 2005) and are transmitted by dipteran vectors of the Family Culicidae (*Plasmodium*), Ceratopogonidae (Subgenus *Parahaemoproteus*), and Hippoboscidae (Subgenus *Haemoproteus*) (Santiago-Alarcon et al., 2012). Local extinctions of bird species have evidenced their impact on avian hosts (Atkinson et al., 2000), detrimental effects on individual fitness (Asghar et al., 2015; Merino

et al., 2000), survival (Martínez-de la Puente et al., 2010), and virulence (Videvall et al., 2020). However, infections are commonly detected as low parasitemia chronic infections with mild or not apparently detrimental effects on individuals (Asghar et al., 2011).

To test the parasite-mediated selection hypothesis and its role in MHC allele diversity, we investigated the associations between MHC-I and MHC-II allele diversity and the prevalence of *Plasmodium* and *Haemoproteus* parasites in the rufous-collared sparrow (*Zonotrichia capensis*) (P. L. Stadius Müller, 1776) across a latitudinal gradient along South America. The rufous-collared sparrow is a small passerine (16.8–31 g) found in open spaces from sea level to high elevation (~4600 m) (Rising & Jaramillo, 2023). Its distribution range spans Central and South America, from southern Mexico to Cape Horn in Chile. Numerous morphological subspecies have been proposed resulting from geographical differences in plumage, morphology and song (Chapman, 1940; Handford, 1985). All subspecies are mainly sedentary except for the southernmost species, the long-distance migratory *Z. capensis australis*. However, short-distance movements have been observed in resident subspecies related to altitudinal movements during winter/non-winter seasons (Poblete et al., 2023; Rising & Jaramillo, 2023). The rufous-collared sparrow is commonly infected by these parasites throughout its distribution range (Cadena-Ortiz et al., 2019; Doussang et al., 2019; Jones et al., 2015; Mantilla et al., 2016).

Based on the decreasing latitudinal gradient in biological diversity (LGD) from the Equator to high latitudes (Willig et al., 2003), we hypothesize that birds from locations close to the Equator would present higher MHC allele diversity and higher parasite prevalence compared with birds from the austral region of South America. By suffering a higher selective pressure from parasites, individuals would have evolved a vast array of immune defenses, that is, MHC allele diversity (Biedrzycka et al., 2018; Demas & Nelson, 2012; Hasselquist, 2007; Møller, 1998; Owen-Ashley et al., 2008). Previous studies have found evidence of a latitudinal gradient of *Plasmodium* and *Haemoproteus* infections in South America, with a general decreasing of both parasites to the south (Clark et al., 2014; Durrant et al., 2006; Fecchio et al., 2019; Merino et al., 2008; White et al., 1978). However, other studies have not observed this trend (Clark, 2018; Doussang et al., 2019). Thus, if avian malaria parasites do not follow a latitudinal gradient in prevalence, the MHC allele diversity would be lower in regions where parasite prevalence is low or absent (Hawley & Fleischer, 2012; Johnson & Haas, 2021). In this situation, biomes (Dinerstein et al., 2017; Olson et al., 2001) could represent a better approach to explain MHC diversity, since the biotic and abiotic factors of these biomes may affect the constitution of distinct assemblages of parasites, vectors, and hosts (Chapa-Vargas et al., 2020; Cuevas et al., 2020; Doussang et al., 2019; Garcia-Longoria et al., 2022; Hussing, 2020). In order to test this hypothesis, we specifically (i) estimated MHC-I and MHC-II allele diversity in rufous-collared sparrows in a latitudinal gradient in South America (from Colombia to Cape Horn in Chile), (ii) explored signals of positive selection on MHC alleles, (iii) investigated the associations among MHC allele diversity and latitude/infection, and

(iv) analyzed the associations among parasite infection status and latitude/biomes.

2 | MATERIALS AND METHODS

2.1 | Study area and bird sampling

We investigated MHC-I and MHC-II genes in 93 rufous-collared from Colombia, Ecuador, Peru and Chile, sampled in different studies from 2011 to 2018 (Basto et al., 2006; Cadena, 2015; Cadena-Ortiz et al., 2019; González et al., 2015; Hussing, 2020; Martínez et al., 2016; Marzal et al., 2015) (Figure 1, Table S1). Twenty-one individuals were sampled in Colombia, 22 in Ecuador, 11 in Peru, and 39 in Chile.

2.2 | MHC primers design, amplification, and sequencing

Genomic DNA was extracted by different methods depending on the laboratory of origin (Cadena, 2015; Cadena-Ortiz et al., 2019; Doussang et al., 2019; González et al., 2015; Martínez et al., 2016; Marzal et al., 2015). In Chile, and before molecular analyses, DNA samples were quantified in a Qubit fluorometer (Thermo Fisher). A Sequencing Library Preparation protocol (Illumina) was applied for the Illumina MiSeq System to obtain MHC-I and MHC-II sequences. The workflow consisted of an initial MHC primer design (Macrogen) with an overhang adapter attached to MHC-I and MHC-II forward and reverse primers. For MHC-I, we tested two different sets of primers, GCA21M (5'-CGTACAGCGGCTTGTGGCTGTGA-3') and fA23M (5'-GCGCTCCAGCTCCTTCTGCCATA-3') (Jones et al., 2014), and MhcPasCI-FW (5'-CSCSCAGGTCTSCACAC-3') and MhcPasCI-RV (5'-CWCARKAATTCTGYTCHCACC-3') (Alcaide et al., 2013). For MHC-II, we tested two sets of primers, HOPE1 (5'-GAAAGCTCGAGTGTCACTTACGAAACGGC-3') and HOPE 10 (5'-TCCACGCTGAACGGGCGGAACACCTC-3') (Sato et al., 2011), and 2zffw1 (5'-TGTCACCTTTCAYKAACGGCACGGAG-3') and 2zfrv1 (5'-GTAGTTGTCCGGCAGTACGTGTC-3') (Balakrishnan et al., 2010). We tested primers with the overhang adapters in five individuals through a standard PCR (Sallaberry-Pincheira et al., 2016). All primers worked successfully for the rufous-collared sparrow, however, we finally selected GCA21M/fA23M and HOPE1/HOPE10 because they were applied before in this species successfully (Jones et al., 2014; Sato et al., 2011). The GCA21M/fA23M primers amplified a region of 214 base pairs (bp) inside exon 3 of the alpha chain, a part of the PBR, whereas HOPE1/HOPE10 amplified 189 bp inside exon 2 of the beta chain, also part of PBR. PCR conditions for GCA21M/fA23M consisted of 94°C for 10 min, 35 cycles of 94°C for 30 s, 64°C for 30 s, and 72°C for 1 min, and at 72°C for 10 min. PCR conditions for HOPE1/HOPE10 consisted of a PCR of 94°C for 10 min, 35 cycles of 94°C for 30 s, 58°C for 30 s, 72°C for 1 min, and 72°C for 10 min (Jones



FIGURE 1 Sampling locations and biogeographical biomes of rufous-collared sparrows. Biomes and map where obtained with QGIS.

et al., 2014; Sato et al., 2011). Amplification success was confirmed by observing UV bands of appropriate size in an agar 1% gel stained with SYBR Green (Sigma Aldrich). The MHC-I and MHC-II amplicons were sequenced in MacroGen. We included a PCR mix reaction without a DNA template in every PCR run as the negative control.

2.3 | Library preparation and sequencing

Purified PCR products in a final 30- μ L volume with MHC-I and MHC-II amplicons were quantified in Nanodrop. After the purification Nextera XT v2 Index Kit A indices were added (Illumina). After a second purification with AMPure (Beckman Coulter), the libraries were diluted in 10 nM before making the equimolar pool, to which a 10% PhiX control library (Illumina) was added. The entire pool was then diluted to 10 pM, which was loaded onto the MiSeq sequencer (Illumina) with the Kit v3 (600Cycles) and mode of sequencing 2 \times 300pb (PE).

2.4 | Bioinformatics

Each MHC-I and MHC-II FASTQ sequences were merged with the AmpliMERGE tool in AmpliSAT, and then, the quality was checked with AmpliCHECK (Sebastian et al., 2016). Clustering and filtering sequences were performed in AmpliSAS by selecting the Illumina

platform. Elimination of chimeras and sequences with a length different from the expected sequence size was also chosen in this step (Rekdal et al., 2018). A total of 21,318,682 reads (59.5% quality score \geq Q30) were obtained from sequencing. From them, we obtained the total number of MHC-I and MHC-II nucleotide sequences for every individual. MHC-I and MHC-II sequences were aligned in Bioedit (Hall, 2011) with CLUSTAL and translated into amino acids. Sequences with stop codons were considered non-functional and removed from the dataset. Amino acidic MHC sequences were reversed to nucleotides and searched in Blast (<https://blast.ncbi.nlm.nih.gov/Blast.cgi>) to confirm whether they corresponded to new (unpublished) or known MHC nucleotide sequences. Blast matches lower than 100% were considered as new MHC alleles. We refer to the "MHC allele," although primers do not discern among loci and do not cover all MHC-I exon 3 and MHC-II exon 2. MHC-I and MHC-II were considered all putative sequences even they came from one PCR event.

2.5 | Phylogenetic reconstruction and selection in the peptide binding region

We combined nucleotide sequences from all individuals in two datasets (MHC-I and MHC-II), aligned with MUSCLE (Edgar, 2004), and applied redundancy sequence removal in Jalview (Waterhouse et al., 2009). A maximum likelihood phylogenetic tree for each MHC class plus other published rufous-collared sparrow MHC-I

and MHC-II sequences were obtained in [NGPhylogeny.fr](https://ngphylogeny.fr) (Lemoine et al., 2019). We selected BMGE curation and PhyML tree inference. The best model of molecular evolution was estimated by SMS (Lefort et al., 2017) and node support with aBayes. Trees were edited in iTol (Letunic & Bork, 2021), and chicken *Gallus gallus* MHC-I (GenBank accession number KM014730.1) and MHC-II (GenBank accession number [AY744349.1](https://www.ncbi.nlm.nih.gov/nuccore/AY744349.1)) were selected as outgroups.

To explore signs of positive selection operating on the MHC, we performed a selection analysis of MHC-I and MHC-II nucleotide sequences in the Datamonkey 2.0 server (Weaver et al., 2018) by means of a Bayesian approach to infer non-synonymous (dN) and synonymous (dS) substitution rates per-site with FUBAR method (Murrell et al., 2013). Excess of non-synonymous vs synonymous substitutions was considered a sign of positive selection. In addition, we calculated Tajima's D in the PBR versus the non-PBR positions under selection following Minias et al. (2018) in MEGA11 (Tamura et al., 2021).

2.6 | Individual parasite infection status

Plasmodium and *Haemoproteus* infection status were determined by amplifying the cytochrome b DNA by PCR, microscopy, or both techniques (see [Table S1](#)). Different combinations of primers were used in each laboratory of origin (Cadena, 2015; Cadena-Ortiz et al., 2019; Doussang et al., 2019; González et al., 2015; Martínez et al., 2016; Marzal et al., 2015). Seventy-three out of 93 individuals were also investigated by inspecting blood smears by microscopy (Cadena-Ortiz et al., 2019; González et al., 2015; Hussing, 2020). The combined use of both methodologies enhanced the accuracy of infection confirmation, as different primers can produce slight variations in parasite detection (Valkiūnas et al., 2006, 2008).

2.7 | Statistical analyses

2.7.1 | Associations between MHC allele diversity and latitude

Associations among MHC diversity and latitude were investigated by generalized additive mixed models (GAMM) with the `mgcv` package (Wood, 2023) in R (R Development Core Team, 2016). For each MHC class, we created a set of competitive models including MHC-I or MHC-II amino acidic allele diversity as dependent variable, and latitude, location, and year as explanatory variables. In the models, latitude was included as a smooth parameter=restricted maximum likelihood (REML). Because birds' sampling spanned different years and several individuals came from the same location, year and location were included as random effects. An AICc model selection (suited for small sample sizes) was performed over all model term combinations with family=Poisson (MHC-I) or Gaussian (MHC-II), based on previous data exploration of the dependent variable distribution. Model assumptions were checked with `gam.check`. We evaluated the goodness of fit of the final models by inspecting the dependence of the residuals, and checked their normality with

qqplots. The final best model and also the equally possible models were obtained within a threshold of $\Delta < 2$.

2.7.2 | Associations among infections and latitude/biomes

In order to explore whether infections follow the same pattern as the observed for MHC amino acidic allele diversity and latitude, we investigated the associations between infection status and latitude using logistic GAMM models. Both *Plasmodium* and *Haemoproteus* infection status (non-infected=0 and infected=1) were selected as the dependent variables and latitude, location, and year as explanatory variables. As described previously, latitude was included as a fixed factor and year and location as random effects. An AICc model selection was performed over all models obtained with method="REML" and family=binomial. Model assumptions were checked as before.

Finally, differences in *Plasmodium* and *Haemoproteus* prevalence of infection among biomes were investigated by Fisher exact tests with post hoc Bonferroni correction applied among factor levels. Biomes were obtained for every location by importing GPS data points into QGIS (QGIS.org, 2022) and loading a shapefile layer of "Biomes of the World" for each country (Dinerstein et al., 2017; Olson et al., 2001).

3 | RESULTS

3.1 | MHC allele diversity

The average number of reads \pm [SD] per sample was 1273 ± 361 (MHC-I) and 901 ± 198 (MHC-II). We obtained 48 MHC-I and 104 MHC-II nucleotide sequence variants when considering all individuals ([Tables S1](#) and [S2](#)). Individual MHC allele nucleotide diversity ranged from one to four alleles for MHC-I (mean=1.8, SD=0.99), and from one to six for MHC-II (mean=3.34, SD=1.21). When converted to amino acids, the MHC allele amino acidic diversity was similar (MHC-I: mean=1.57, SD=0.87; MHC-II mean=3.28, SD=1.16). A total of 21 MHC-I stop codon sequences were removed from the dataset, while no stop codon sequences were observed for MHC-II. Considering that MHC genes are codominantly expressed, the maximum number of MHC-I alleles found in an individual was four, indicating the presence of at least two loci (assuming all loci are heterozygous). For MHC-II, the maximum number of alleles was six, suggesting at least three loci. We excluded several individuals with low-quality MHC sequences from the analyses ([Table S1](#)).

MHC-I blast search indicated that 21 out of the 48 nucleotide sequences were new MHC alleles (GenBank accession numbers: [OR578737](https://www.ncbi.nlm.nih.gov/nuccore/OR578737)–[OR578757](https://www.ncbi.nlm.nih.gov/nuccore/OR578757)) ([Table S2](#)). The most common alleles were ZocaU*2 (60 individuals) and ZocaU*5 (10 individuals). Less than 10 individuals shared the rest of the alleles. For MHC-II blast, they resulted in 104 haplotypes (GenBank accession numbers: [OQ377810](https://www.ncbi.nlm.nih.gov/nuccore/OQ377810)–[OQ377913](https://www.ncbi.nlm.nih.gov/nuccore/OQ377913)). The most common MHC-II alleles were alleles Zoca1 (85 individuals), Zoca9 (14 individuals), and Zoca16 (13 individuals), all of them new alleles.

3.2 | Selection and phylogenetic reconstruction

We detected signs of positive selection in both MHC-I (amino acid positions 12, 48, 51) and MHC-II (amino acid positions 10, 15, 18, 27, 33, 37, 40, 47, 50, 51, 53, 57, 58). The positions identified in our analysis coincided with those previously observed under positive selection in passerine birds at rates of 9.1% for MHC-I and 69.2% for MHC-II (Minias et al., 2018) (Figure S1). And a coincidence rate of 0% for MHC-I and 35.2% for MHC-II when compared to human PBR positions (Brown et al., 1993; Saper et al., 1991). Tajima's D neutrality test confirmed a greater number of non-synonymous versus synonymous substitutions in the PBR compared with non-PBR regions for both MHC classes (Tables S3 and S4).

3.3 | Individual parasite infection status

A higher prevalence of *Plasmodium* than *Haemoproteus* was found, with 24% and 12% of birds infected, respectively. The detected parasites corresponded to *Plasmodium (Haemamoeba) cathemerium* (ZOCAP15), *Plasmodium (Novyella) homopolare* (BAEBIC02), *Plasmodium SGS1*, *Haemoproteus (P.) sp1* (ZC1), *Haemoproteus coatneyi*, *Haemoproteus erythrogravidus*, and *Haemoproteus CHLOP01* (Table S1). All these infections have been previously reported (Basto et al., 2006; Cadena, 2015; Cadena-Ortiz et al., 2019; Doussang et al., 2019; González et al., 2015; Hussing, 2020;

Martínez et al., 2016; Marzal et al., 2015), except for the new individuals from Chile to whom PCR, and microscopy was done in this study. Some infections could only be identified at the genus level for some individuals, leaving them as *Haemoproteus* sp. and *Plasmodium* sp.

3.4 | Associations between MHC allele diversity and latitude

GAMM model selection for MHC-I resulted in a best model with only the intercept as the final term, excluding models that incorporated latitude (Table 1, Figure 2). Models including latitude, year, or location were equally plausible based on $\Delta < 2$, but latitude was not statistically significant in any model. However, in two models, year was statistically significant, indicating a random effect of year on MHC-I. For MHC-II, the best GAMM model retained latitude as the final term. MHC-II amino acid allele diversity was non-linearly associated with latitude (Table 1, Table S5). Specifically, MHC-II amino acid allele diversity was low in North Chile (latitude -18° to -20°) and locations near the Equator (0° to 4°), then increased in Peru (-6° to -12°), and further increased from central to the austral region of Chile (-33° to -54°) (Figure 2). Based on $\Delta < 2$, three other models were also plausible, with latitude always being statistically significant, but not location or year, excluding any random effects of these variables. All models' assumptions were met.

		df	logLik	AICc	Delta
MHC-I	Intercept	0	-121.908	245.861	0
	Latitude+Location	2	-121.685	247.508	1.646
	Latitude	2	-121.685	247.509	1.647
	Latitude+Location+Year	2	-121.504	247.855	1.994
	Latitude+Year	2	-121.504	247.855	1.994
MHC-II	Latitude	7	-122.012	261.375	0
	Latitude+Location	7	-122.012	261.382	0.006
	Latitude+Year	8	-121.567	262.737	1.361
	Latitude+Year+Location	8	-121.567	262.738	1.362

TABLE 1 Associations among rufous-collared sparrow major histocompatibility complex (MHC) amino acid allele diversity and latitude, year, and location, investigated by generalized additive mixed models (GAMM) and AICc model selection.




FIGURE 2 Generalized additive mixed model (GAMM) associations between MHC-I and MHC-II amino acid allele diversity and latitude. 95% confidence intervals are shown.

3.5 | Associations between infections and latitude/biomes

GAMM models found no association between infection status for either *Plasmodium* or *Haemoproteus* and latitude. The best GAMM model for *Plasmodium* included latitude as an explanatory term, along with several plausible models that included combinations of year and location (Table 2). In the best model and the equally plausible models, the association with latitude were not statistically significant (Table S6, Figure S3). For *Haemoproteus*, the best model showed a marginally significant association with latitude ($p=.051$), with a pattern similar to that observed for MHC-II, with individuals from north Chile having a low prevalence of infection (Table S6, Figure S3). Another plausible model also included latitude and year as factors.

Finally, we found a statistically significant association between *Haemoproteus* infection and biome (Fisher's exact test, $p=.0018$). *Haemoproteus* prevalence was higher in the tropical biome compared with the desert biome (post-hoc Bonferroni test: tropical vs. desert, $n=69$, adjusted $p=.007$) (Figure 3). There was also a significant association between *Plasmodium* prevalence and biome (Fisher's exact test, $p=.028$); however, this result became non-significant after the Bonferroni test (post hoc Bonferroni test: tropical vs. desert, $n=69$, adjusted $p=.281$).

4 | DISCUSSION

In this study, we explored the relationships between MHC diversity and avian malaria infections (*Plasmodium* and *Haemoproteus*) across


a latitudinal gradient in a passerine bird species in South America. While avian MHC is known to respond to a diverse array of antigens, there is compelling evidence linking MHC alleles to avian malaria infections (O'Connor et al., 2019). Based on the latitudinal gradient in biological diversity (LGD) hypothesis, we expected a decrease in MHC allele diversity from the Equator to the austral region of South America, which would correspond to a decrease in the prevalence of *Plasmodium* and *Haemoproteus* parasites at higher latitudes. However, if the prevalence of infection does not follow the trend predicted by the LGD, then MHC diversity should be lower in regions where the prevalence of infection is lower or absent. We found that Individuals from north Chile exhibited the lowest MHC-II diversity, a finding that aligned with the absence of infections by both parasites. MHC diversity could be in part explained by a low prevalence of avian malaria parasites in the region. This result was supported by the observation that individuals from the deserts and xeric shrublands (intermediate latitude, north Chile) were less infected by *Haemoproteus* parasites compared to individuals from tropical and subtropical biomes.

Although LGD is observed for many plant and animal species, other taxonomic groups, like parasites, have less clear patterns (Clark et al., 2014), even the reverse, as is the case for the *Leucocytooon* parasite, a closely related species (Cuevas et al., 2020; Fecchio et al., 2020; Merino et al., 2008). In rufous-collared sparrows, in the most complete study to date, the higher prevalence of *Plasmodium* is found in central Chile and in Peru, and for *Haemoproteus* in central and northern Chile (Doussang et al., 2019), and the lowest prevalence for both parasites in Colombia and the austral region of Chile. This trends is supported by a low prevalence

TABLE 2 Associations among rufous-collared sparrow infection status with latitude, year, and location, investigated by generalized additive mixed models (GAMM) and AICc model selection.

		df	logLik	AICc	Delta
<i>Plasmodium</i>	Latitude	4	-27.494	63.549	0
	Latitude+Year	4	-27.494	63.549	0.0001
	Latitude+Location	5	-26.629	65.047	1.498
	Latitude+Year+Location	5	-26.629	65.047	1.498
<i>Haemoproteus</i>	Latitude	5	-43.298	98.424	0
	Latitude+Year	5	-43.295	98.433	0.0096

FIGURE 3 *Plasmodium* and *Haemoproteus* parasite prevalence among biomes. Des=Deserts and Xeric Shrublands, Med=Mediterranean Forests, Mon=Montane Grasslands and Shrublands, Tem=Temperate Broadleaf and Mixed Forests, Woodlands and Scrub, and Tro=Tropical and Subtropical Moist Broadleaf Forests.



for both parasites reported in an Andean community of birds in Colombia (although at elevations ranging from 2100 to 4000 m.a.s.l.) (González et al., 2015). Overall, these contrasting results highlight the variability in the patterns of avian malaria prevalence detected in South America. These differences may be better understood by considering the type of biome. North Chile is characterized by deserts and semi-arid habitats, including the presence of the Atacama Desert, the driest desert on Earth (Darack, 2008). This disrupts the expected linear decreasing latitudinal pattern for the prevalence of avian malaria. Thus, biome could be a factor that better summarizes the effect of latitude, along with other biotic and abiotic factors, on infections (Fecchio et al., 2019). In the case of avian malaria parasites, precipitation and temperature are main factors affecting the life cycle of insect vectors (Doussang et al., 2019; Martínez-de la Puente et al., 2009). Hence, desert or semi-arid habitats could limit the occurrence and their activity (Chapa-Vargas et al., 2020). On the contrary, tropical and subtropical environments may harbour a larger community of insect vectors, which is reflected in the higher prevalence of *Haemoproteus* and greater diversity of MHC-II compared to desert biomes. However, in locations close to the Equator in South America, a low prevalence of infection have been also reported, possible related to a dilution effect, where high diversity of host birds is associated with a decrease in parasite transmission among hosts of the same species (Moens & Pérez-Tris, 2016). This could explain the moderate MHC-II diversity levels observed in our study. However, an intermediate prevalence of *Plasmodium* and *Haemoproteus* (23%–50%, respectively) is also observed in Ecuador (Cadena-Ortiz et al., 2019), and in our study.

Contrary to our findings, Doussang et al. (2019) reported an elevated prevalence of infection by *Plasmodium* and *Haemoproteus* in rufous-collared sparrows in northern Chile. In the same region, another study reported *Plasmodium* infecting house sparrows (*Passer domesticus*) but not rufous-collared sparrows (Martínez et al., 2016), and *Haemoproteus* was observed infecting eared doves (*Zenaidura macroura*). This parasite belongs to the subgenus *Haemoproteus* (different from the subgenus *Parahaemoproteus* that usually infects rufous-collared sparrows), thus confirming local transmission of *Plasmodium* and at least the *Haemoproteus* subgenus. Differences in results among studies could be due to particularities among sampling localities or other factors, including the sampling period. Although arid, the region presents valleys and oases with water presence where suitable conditions could exist for vectors, favoring local transmission (González-Gómez et al., 2018). One possibility is that the birds concentrate in these oases, generating greater density in those areas and favoring the transmission of the parasite. On the contrary, the prevalence of infection varies with season, so it can confound the actual prevalence if not sampled throughout the year. Birds included in our study were sampled in this region during the austral summer (January) and winter (June–July) in 2012 and 2017, covering a broad period where transmission is expected to occur and in different years. The detection of avian malaria parasites in local birds supports active transmission, but also indicates differential transmission

depending on the host species (Clark, 2018; Doussang et al., 2021). Differences in prevalence in bird species from the same area could be related to life-history traits like nestling period, nest location, vegetation strata, or body mass that increase the exposition to vectors (Quillfeldt et al., 2011). In order to account for this variability, we included year as a random effect in the models, but no effect was observed.

Evidence of low MHC diversity in species living in environments with a low prevalence of parasites has been observed in other birds. For example, the prevalence of blood parasitic infections is low in seabirds, which have been related to the reduced occurrence of vectors in marine environments (Quillfeldt et al., 2011). Moreover, in a latitudinal study ranging from Peru to South Chile evaluating Magellanic and Humboldt penguins, only Humboldt penguins from Peru were infected with *Haemoproteus* compared with southern locations. Precisely, individuals from the infected population exhibited elevated MHC-I and MHC-II allele diversity, suggesting a greater diversity associated with the infections (Sallaberry-Pincheira et al., 2015, 2016). Interestingly, it has been observed that Magellanic penguins translocated to other parts of the world, such as zoos from the northern hemisphere, tend to succumb to infection caused by avian malaria parasites (Hernandez-Colina et al., 2021). This results supports that parasites' absence or low abundance could result in low MHC allele diversity in rufous-collared sparrows in this region (Radwan et al., 2010).

Due to the latitudinal amplitude of the work and the small sample size, we could not perform a detailed analysis of the associations between MHC diversity or specific alleles on infections. However, for both types of MHC we have detected positive selection on PBR. In the case of MHC-I, several alleles were closely related to other alleles previously associated with *Haemoproteus* infections in Perú (Jones et al., 2014, 2015). The most common *Haemoproteus* molecular lineage detected in those studies was *H. (P.) sp1* (ZC1) (KC480265). This parasite corresponds to the same lineage reported in (Cadena-Ortiz et al., 2019; Doussang et al., 2019; Merino et al., 2008) and detected in our study in individuals from Ecuador. Its high prevalence in South America is suggested possibly associated with chronic infections of low virulence (Doussang et al., 2019; Merino et al., 2008). This parasite has been related to the morpho species *H. coatneyi* parasite. *H. coatneyi* is closely related to *H. erythrogravidus* (which was also detected in our study), and both share *Z. capensis* as a host. They have a wide distribution and impact various bird species across South America. *H. coatneyi* has a broad geographical distribution, occurring in South, Central, and North America, infecting several species, whereas *H. erythrogravidus* occurs only in South America and is restricted to infecting only two species: the rufous-collared sparrows and the blue-winged mountain tanager (*Anisognathus somptuosus*) (F. Thraupidae) (de Oliveira et al., 2020; Valkiūnas, 2005). On the contrary, the other lineage found in our study, CHLOP01, was restricted to *Z. capensis* in Peru (Marzal et al., 2015); however, the virulence of these parasites is unknown. With respect to *Plasmodium* parasites infecting rufous-collared sparrows we detected *Plasmodium (Haemamoeba) cathemerium* (ZOCAP15), *Plasmodium (Novyella)*

homopolare (BAEBIC02), and *Plasmodium relictum* SGS1. In Bosque de Jerusalem in Ecuador, the prevalence of ZOCAP15 was very low, which could mean that it is either rare in the population or actually is very virulent (Cadena-Ortiz et al., 2019). Interestingly, one individual infected by this lineage had one of the highest parasitemia, suggesting high susceptibility to this lineage. On the contrary, BAEBIC02 has been reported to infect several passerines, mainly species from Emberizidae and Passerellidae families, included rufous-collared sparrows (Cadena-Ortiz et al., 2019; Rivero de Aguilar et al., 2018; Walther et al., 2014). *Plasmodium relictum* SGS1 has been recently detected for the first time in Peru and is considered an invasive species for the South American continent. Therefore, it is necessary to continue monitoring these parasites to discern the effects they may have on the host and their role on parasite-mediated selection on MHC. For example, in Bosque de Jerusalem in Ecuador, the prevalence of both genera was among the highest among all locations, suggesting that individuals cannot avoid infections, instead trying to keep infections at low intensities through their immune system, that is, MHC genes.

Our results generally support variability in MHC diversity on a latitudinal scale related to biome; however, we have also identified other factors that could explain our results. On the one hand, our results support differences both in MHC diversity and infection prevalence in a specific area coinciding with an arid biome, but also a result consistent with the separation between *Zonotrichia* control region-based molecular lineages observed between northern Chile and Peru. In South America, there is evidence of molecularly distinct clusters based on a mitochondrial control region genes (Lougheed et al., 2013). Thus, the genetic differentiation of these populations could determine the differences observed in the MHC genes. Conversely, differences in MHC diversity may reflect the distinct demographic histories of populations from various regions. Reduced MHC diversity could result from low genetic diversity in bottlenecked or inbred populations (Radwan et al., 2010). Moreover, MHC diversity variation found in our study should reflect adaptation to the local community of parasites, however, selection can be modulated or being stronger toward other pathogens (Llanos-Soto et al., 2017; Loiseau et al., 2008). Finally, bird movements could be affecting our results' accuracy. The rufous-collared sparrow is considered a sedentary species in most of the distribution, except for southern Chile, with only one subspecies *Z. c. australis* being a long-distance migrant from central Chile and Argentina to the Magallanes region. Even for a common species, such as the rufous-collared sparrow, there is still scarce information about the migratory status of many populations (Medrano et al., 2018). Nevertheless, in a broad latitudinal scale, as is our study, we should have captured latitudinal differences in the MHC related to local parasites.

AUTHOR CONTRIBUTIONS

Juan Rivero de Aguilar: Conceptualization (lead); data curation (lead); formal analysis (lead); funding acquisition (equal); investigation (lead); methodology (equal); project administration (equal); resources (equal); software (equal); supervision (equal); validation

(equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Omar Barroso:** Conceptualization (equal); data curation (equal); investigation (equal); methodology (equal); validation (equal); visualization (equal); writing – review and editing (equal). **Elisa Bonaccorso:** Conceptualization (equal); data curation (equal); formal analysis (equal); funding acquisition (equal); investigation (equal); methodology (equal); project administration (equal); resources (equal); software (equal); supervision (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Hector Cadena:** Data curation (equal); investigation (equal); writing – review and editing (equal). **Lucas Hussing:** Conceptualization (equal); data curation (equal); formal analysis (equal); investigation (equal); methodology (equal); validation (equal); writing – review and editing (equal). **Josefina Jorquera:** Data curation (equal); formal analysis (equal); investigation (equal); methodology (equal); software (equal); supervision (equal); validation (equal); writing – review and editing (equal). **Javier Martinez:** Conceptualization (equal); data curation (equal); funding acquisition (equal); investigation (equal); methodology (equal); project administration (equal); resources (equal); software (equal); supervision (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Josué Martínez-de la Puente:** Data curation (equal); investigation (equal); methodology (equal); supervision (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Alfonso Marzal:** Conceptualization (equal); data curation (equal); formal analysis (equal); investigation (equal); methodology (equal); supervision (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Fabiola León Miranda:** Conceptualization (equal); data curation (equal); formal analysis (equal); investigation (equal); methodology (equal); software (equal); supervision (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Santiago Merino:** Conceptualization (equal); funding acquisition (equal); investigation (equal); methodology (equal); project administration (equal); resources (equal); supervision (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Nubia E. Matta:** Conceptualization (equal); data curation (equal); formal analysis (equal); funding acquisition (equal); investigation (equal); methodology (equal); project administration (equal); resources (equal); software (equal); supervision (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Marilyn Ramenofsky:** Conceptualization (equal); data curation (equal); funding acquisition (equal); investigation (equal); methodology (equal); project administration (equal); resources (equal); supervision (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Ricardo Rozzi:** Funding acquisition (equal); project administration (equal); resources (equal); supervision (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Carlos E. Valeris-Chacín:** Conceptualization (equal); data curation (equal); investigation (equal); methodology (equal); supervision (equal); validation (equal); writing

– original draft (equal); writing – review and editing (equal). **Rodrigo A. Vásquez:** Conceptualization (equal); funding acquisition (equal); investigation (equal); methodology (equal); project administration (equal); resources (equal); supervision (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **Juliana A. Vianna:** Conceptualization (equal); funding acquisition (equal); investigation (equal); methodology (equal); project administration (equal); resources (equal); software (equal); supervision (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal). **John C. Wingfield:** Conceptualization (equal); data curation (equal); funding acquisition (equal); investigation (equal); methodology (equal); project administration (equal); resources (equal); supervision (equal); validation (equal); visualization (equal); writing – original draft (equal); writing – review and editing (equal).

ACKNOWLEDGMENTS

The sampling methodology was approved by the Scientific Ethics Committee of Vicerrectoría de Investigación y Postgrado, Universidad de Magallanes CERTIFICADO N°035/CEC-UMAG/2022 and University of Chile 2017, following guidelines from the Biosecurity Manual of CONICYT (version 2008) and Chilean law N°20380 about animal protection. Full permissions for sampling and animal ethics approval in Chile were granted by the Servicio Agrícola y Ganadero (SAG; 404/2017 and 3813/2018). Sequencing of samples from Ecuador was authorized by Contrato Marco de Acceso a Recursos Genéticos MAE-DNB-CM-2015-0017 granted to Universidad Tecnológica Indoamérica. The import of blood samples to Chile was approved by SAG (No 1398/2018 and No 5452/2018).

FUNDING INFORMATION

This study was funded by FONDECYT-CONICYT Postdoctorado 3,170,211; CONICYT PIA APOYO CCTE AFB170008; IEB-FB210006; ANID—Convocatoria Nacional Subvención a la Instalación en la Academia, 2020, PAI77200078; Cape Horn International Center (CHIC), ANID/BASAL FB210018; ANID—Programa Iniciativa Milenio—ICN2021_044 (CGR), ICN2021_002 (BASE), and NCN2021-050 (LiLi); Universidad San Francisco de Quito (HUBI 12424); JMP was financed by the PID2020-118205GB-I00 grant funded by MCIN/AEI/10.13039/501100011033 and by “ERDF A way of making Europe”; AM was funded by line of action LA4 (R+D+I program in the Biodiversity Area financed with the funds of the FEDER Extremadura 2021–2027 Operational Program of the Recovery, Transformation and Resilience Plan) and by “Consejería de Economía e Infraestructura of the Junta de Extremadura” and the European Regional Development Fund, a Way to Make Europe, through the research project IB20089.

CONFLICT OF INTEREST STATEMENT

The authors have no conflicts of interest to declare.

DATA AVAILABILITY STATEMENT

MHC sequencing reads have been deposited at NCBI (GenBank accession numbers: MHC-I [OR578737–OR578757](https://doi.org/10.5281/zenodo.11398506); MHC-II:

[OQ377810–OQ377913](https://doi.org/10.5281/zenodo.11398506)). Database as well as R code for stats and figures used in this publication are available on Zenodo ([10.5281/zenodo.11398506](https://doi.org/10.5281/zenodo.11398506)).

ORCID

Juan Rivero de Aguilar [ORCID](https://orcid.org/0000-0002-4746-5759) <https://orcid.org/0000-0002-4746-5759>
 Omar Barroso [ORCID](https://orcid.org/0000-0003-0565-329X) <https://orcid.org/0000-0003-0565-329X>
 Elisa Bonaccorso [ORCID](https://orcid.org/0000-0002-7262-9356) <https://orcid.org/0000-0002-7262-9356>
 Hector Cadena [ORCID](https://orcid.org/0000-0003-4653-2072) <https://orcid.org/0000-0003-4653-2072>
 Javier Martinez [ORCID](https://orcid.org/0000-0003-2657-1154) <https://orcid.org/0000-0003-2657-1154>
 Josué Martínez-de la Puente [ORCID](https://orcid.org/0000-0001-8055-4115) <https://orcid.org/0000-0001-8055-4115>
 Alfonso Marzal [ORCID](https://orcid.org/0000-0001-5872-1060) <https://orcid.org/0000-0001-5872-1060>
 Santiago Merino [ORCID](https://orcid.org/0000-0003-0603-8280) <https://orcid.org/0000-0003-0603-8280>
 Nubia E. Matta [ORCID](https://orcid.org/0000-0003-1775-0804) <https://orcid.org/0000-0003-1775-0804>
 Marilyn Ramenofsky [ORCID](https://orcid.org/0000-0002-3694-8464) <https://orcid.org/0000-0002-3694-8464>
 Ricardo Rozzi [ORCID](https://orcid.org/0000-0001-5265-8726) <https://orcid.org/0000-0001-5265-8726>
 Carlos E. Valeris-Chacín [ORCID](https://orcid.org/0009-0005-3589-570X) <https://orcid.org/0009-0005-3589-570X>
 Juliana A. Vianna [ORCID](https://orcid.org/0000-0003-2330-7825) <https://orcid.org/0000-0003-2330-7825>
 John C. Wingfield [ORCID](https://orcid.org/0000-0001-9063-7985) <https://orcid.org/0000-0001-9063-7985>

REFERENCES

- Alcaide, M., Edwards, S. V., Negro, J. J., Serrano, D., & Tella, J. L. (2008). Extensive polymorphism and geographical variation at a positively selected MHC class II B gene of the lesser kestrel (*Falco naumanni*). *Molecular Ecology*, 17(11), 2652–2665. <https://doi.org/10.1111/j.1365-294X.2008.03791.x>
- Alcaide, M., Liu, M., & Edwards, S. V. (2013). Major histocompatibility complex class I evolution in songbirds: Universal primers, rapid evolution and base compositional shifts in exon 3. *PeerJ*, 1, e86. <https://doi.org/10.7717/peerj.86>
- Asghar, M., Hasselquist, D., & Bensch, S. (2011). Are chronic avian haemosporean infections costly in wild birds? *Journal of Avian Biology*, 42, 530–537. <https://doi.org/10.1111/j.1600-048X.2011.05281.x>
- Asghar, M., Hasselquist, D., Hansson, B., Zehindjiev, P., Westerdahl, H., & Bensch, S. (2015). Hidden costs of infection: Chronic malaria accelerates telomere degradation and senescence in wild birds. *Science*, 347(6220), 436–438. <https://doi.org/10.1126/science.1261121>
- Atkinson, C. T., Dusek, R. J., Woods, K. L., & Iko, W. M. (2000). Pathogenicity of avian malaria in experimentally-infected Hawaii Amakihi. *Journal of Wildlife Diseases*, 36(2), 197–204. <https://doi.org/10.7589/0090-3558-36.2.197>
- Balakrishnan, C. N., Ekblom, R., Völker, M., Westerdahl, H., Godinez, R., Kotkiewicz, H., Burt, D. W., Graves, T., Griffin, D. K., Warren, W. C., & Edwards, S. V. (2010). Gene duplication and fragmentation in the zebra finch major histocompatibility complex. *BMC Biology*, 8(1), 29. <https://doi.org/10.1186/1741-7007-8-29>
- Basto, N., Rodriguez, O. A., Marinkelle, C. J., Gutierrez, R., & Matta, N. E. (2006). Haematozoa in birds from La Macarena National Natural Park (Colombia). *Caldasia*, 28(2), 371–377.
- Biedrzycka, A., Bielański, W., Ćmiel, A., Solarz, W., Zając, T., Migalska, M., Sebastian, A., Westerdahl, H., & Radwan, J. (2018). Blood parasites shape extreme major histocompatibility complex diversity in a migratory passerine. *Molecular Ecology*, 27(11), 2594–2603. <https://doi.org/10.1111/mec.14592>
- Borghans, J. A. M., Beltman, J. B., & De Boer, R. J. (2004). MHC polymorphism under host-pathogen coevolution. *Immunogenetics*, 55(11), 732–739. <https://doi.org/10.1007/s00251-003-0630-5>

- Brown, J. H., Jardetzky, T. S., Gorga, J. C., Stern, L. J., Urban, R. G., Strominger, J. L., & Wiley, D. C. (1993). Three-dimensional structure of the human class II histocompatibility antigen HLA-DR1. *Nature*, 364(6432), 33–39. <https://doi.org/10.1038/364033a0>
- Cadena, H. (2015). *Malaria aviar en los chingolos (Zonotrichia capensis) del Parque Recreacional-Bosque Protector Jerusalem, Pichincha, Ecuador*. Pontificia Universidad Católica del Ecuador.
- Cadena-Ortiz, H., Mantilla, J. S., Rivero de Aguilar, J., Flores, D., Bahamonde, D., Matta, N. E., & Bonaccorso, E. (2019). Avian haemosporidian infections in rufous-collared sparrows in an Andean dry forest: Diversity and factors related to prevalence and parasitaemia. *Parasitology*, 146(6), 765–773. <https://doi.org/10.1017/S0031182018002081>
- Chapa-Vargas, L., Matta, N. E., & Merino, S. (2020). Effects of ecological gradients on tropical avian hemoparasites. In D. Santiago-Alarcon & A. Marzal (Eds.), *Avian malaria and related parasites in the tropics* (pp. 349–377). Springer. https://doi.org/10.1007/978-3-030-51633-8_10
- Chapman, F. (1940). The post-glacial history of *Zonotrichia capensis*. *Bulletin of the American Museum of Natural History*, 77, 381–438.
- Clark, N. J. (2018). Phylogenetic uniqueness, not latitude, explains the diversity of avian blood parasite communities worldwide. *Global Ecology and Biogeography*, 27(6), 744–755. <https://doi.org/10.1111/geb.12741>
- Clark, N. J., Clegg, S. M., & Lima, M. R. (2014). A review of global diversity in avian haemosporidians (*Plasmodium* and *Haemoproteus*: Haemosporida): New insights from molecular data. *International Journal for Parasitology*, 44(5), 329–338. <https://doi.org/10.1016/j.ijpara.2014.01.004>
- Cuevas, E., Vianna, J. A., Botero-Delgado, E., Doussang, D., González-Acuña, D., Barroso, O., Rozzi, R., Vásquez, R. A., & Quirici, V. (2020). Latitudinal gradients of haemosporidian parasites: Prevalence, diversity and drivers of infection in the thorn-tailed Rayadito (*Aphrastura spinicauda*). *International Journal for Parasitology: Parasites and Wildlife*, 11, 1–11. <https://doi.org/10.1016/j.ijppaw.2019.11.002>
- Darack, E. (2008). The Atacama Desert: The driest land in the world. *Weatherwise*, 61(1), 14–15. <https://doi.org/10.3200/wewi.61.1.14-15>
- de Oliveira, L., Barino, G. T. M., Rossi, M. F., D'agosto, M., Dias, R. J. P., & Santos, H. A. (2020). Morphological and molecular characterization of *Haemoproteus coatneyi* and *Haemoproteus erythrogravidus* (Haemosporida: Haemoproteidae) in passeriformes in Brazil's atlantic forest. *Revista Brasileira de Parasitologia Veterinária*, 29(4), 1–13. <https://doi.org/10.1590/s1984-296120200074>
- Demas, G. E., & Nelson, R. J. (2012). *Ecoimmunology*. Oxford University Press.
- Dinerstein, E., Olson, D., Joshi, A., Vynne, C., Burgess, N. D., Wikramanayake, E., Hahn, N., Palminteri, S., Hedao, P., Noss, R., Hansen, M., Locke, H., Ellis, E. C., Jones, B., Barber, C. V., Hayes, R., Kormos, C., Martin, V., Crist, E., ... Saleem, M. (2017). An ecoregion-based approach to protecting half the terrestrial realm. *Bioscience*, 67(6), 534–545. <https://doi.org/10.1093/biosci/bix014>
- Doussang, D., González-Acuña, D., Torres-Fuentes, L. G., Loughheed, S. C., Clemente-Carvalho, R. B., Greene, K. C., & Vianna, J. A. (2019). Spatial distribution, prevalence and diversity of haemosporidians in the rufous-collared sparrow, *Zonotrichia capensis*. *Parasites & Vectors*, 12(1), 2. <https://doi.org/10.1186/s13071-018-3243-4>
- Doussang, D., Sallaberry-Pincheira, N., Cabanne, G. S., Lijtmaer, D. A., González-Acuña, D., & Vianna, J. A. (2021). Specialist versus generalist parasites: The interactions between host diversity, environment and geographic barriers in avian malaria. *International Journal for Parasitology*, 51(11), 899–911. <https://doi.org/10.1016/j.ijpara.2021.04.003>
- Durrant, K. L., Beadell, J. S., Ishtiaq, F., Graves, G. R., Olson, S. L., Gering, E., Peirce, M. A., Milensky, C. M., Schmidt, B. K., Gebhard, C., & Fleischer, R. C. (2006). Avian hematozoa in South America: A comparison of temperate and tropical zones. *Ornithological Monographs*, 60, 98–111. <https://doi.org/10.2307/40166831>
- Edgar, R. C. (2004). MUSCLE: Multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Research*, 32(5), 1792–1797. <https://doi.org/10.1093/nar/gkh340>
- Edwards, S. V., & Hedrick, P. W. (1998). Evolution and ecology of MHC molecules: From genomics to sexual selection. *Trends in Ecology & Evolution*, 13(8), 305–311. [https://doi.org/10.1016/S0169-5347\(98\)01416-5](https://doi.org/10.1016/S0169-5347(98)01416-5)
- Fecchio, A., Bell, J. A., Bosholn, M., Vaughan, J. A., Tkach, V. V., Lutz, H. L., Cueto, V. R., Gorosito, C. A., González-Acuña, D., Stromlund, C., Kvasager, D., Comiche, K. J. M., Kirchgatter, K., Pinho, J. B., Berv, J., Anciães, M., Fontana, C. S., Zyskowski, K., Sampaio, S., ... Clark, N. J. (2020). An inverse latitudinal gradient in infection probability and phylogenetic diversity for *Leucocytozoon* blood parasites in New World birds. *Journal of Animal Ecology*, 89(2), 423–435. <https://doi.org/10.1111/1365-2656.13117>
- Fecchio, A., Bell, J. A., Pinheiro, R. B. P., Cueto, V. R., Gorosito, C. A., Lutz, H. L., Gaiotti, M. G., Paiva, L. V., França, L. F., Toledo-Lima, G., Tolentino, M., Pinho, J. B., Tkach, V. V., Fontana, C. S., Grande, J. M., Santillán, M. A., Caparroz, R., Roos, A. L., Bessa, R., ... Collins, M. D. (2019). Avian host composition, local speciation and dispersal drive the regional assembly of avian malaria parasites in south American birds. *Molecular Ecology*, 28(10), 2681–2693. <https://doi.org/10.1111/mec.15094>
- García-Longoria, L., Muriel, J., Magallanes, S., Villa-Galarce, Z. H., Ricopa, L., Inga-Díaz, W. G., Fong, E., Vecco, D., Guerra-Saldaña, C., Salas-Rengifo, T., Flores-Saavedra, W., Espinoza, K., Mendoza, C., Saldaña, B., González-Blázquez, M., Gonzales-Pinedo, H., Luján-Vega, C., Del Águila, C. A., Vilca-Herrera, Y., ... Marzal, A. (2022). Diversity and host assemblage of avian haemosporidians in different terrestrial ecoregions of Peru. *Current Zoology*, 68(1), 27–40. <https://doi.org/10.1093/cz/zoab030>
- González, A. D., Lotta, I. A., García, L. F., Moncada, L. I., & Matta, N. E. (2015). Avian haemosporidians from neotropical highlands: Evidence from morphological and molecular data. *Parasitology International*, 64(4), 48–59. <https://doi.org/10.1016/j.parint.2015.01.007>
- González-Gómez, P. L., Echeverría, V., Estades, C. F., Perez, J. H., Krause, J. S., Sabat, P., Li, J., Kültz, D., & Wingfield, J. C. (2018). Contrasting seasonal and aseasonal environments across stages of the annual cycle in the rufous-collared sparrow, *Zonotrichia capensis*: Differences in endocrine function, proteome and body condition. *Journal of Animal Ecology*, 87(5), 1364–1382. <https://doi.org/10.1111/1365-2656.12846>
- Hall, T. (2011). BioEdit: An important software for molecular biology. *GERF Bulletin of Biosciences*, 2(June), 60–61.
- Handford, P. (1985). Morphological relationships among subspecies of the rufous-collared sparrow, *Zonotrichia capensis*. *Canadian Journal of Zoology*, 63(10), 2383–2388. <https://doi.org/10.1139/z85-352>
- Hasselquist, D. (2007). Comparative immunology in birds: Hypotheses and tests. *Journal of Ornithology*, 148(S2), 571–582. <https://doi.org/10.1007/s10336-007-0201-x>
- Hawley, D. M., & Fleischer, R. C. (2012). Contrasting epidemic histories reveal pathogen-mediated balancing selection on class II MHC diversity in a wild songbird. *PLoS One*, 7(1), e30222. <https://doi.org/10.1371/journal.pone.0030222>
- Hernandez-Colina, A., Gonzalez-Olvera, M., Eckley, L., Lopez, J., & Baylis, M. (2021). Avian malaria affecting penguins in zoological gardens, aquariums and wildlife parks in the UK. *Veterinary Record*, 189(9), e511. <https://doi.org/10.1002/vetr.511>
- Hess, C. M., & Edwards, S. V. (2002). The evolution of the major histocompatibility complex in birds. *Bioscience*, 52(5), 423–431. [https://doi.org/10.1641/0006-3568\(2002\)052\[0423:TEOTMH\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2002)052[0423:TEOTMH]2.0.CO;2)

- Hussing, L. (2020). Factores ecológicos y ambientales relacionados con la distribución de malaria aviar en el chincol (*Zonotrichia capensis*) a lo largo de Chile. Universidad de Chile.
- Johnson, P., & Haas, S. E. (2021). Why do parasites exhibit reverse latitudinal diversity gradients? Testing the roles of host diversity, habitat and climate. *Global Ecology and Biogeography*, 30(9), 1810–1821. <https://doi.org/10.1111/geb.13347>
- Jones, M. R., Cheviron, Z. A., & Carling, M. D. (2014). Variation in positively selected major histocompatibility complex class I loci in rufous-collared sparrows (*Zonotrichia capensis*). *Immunogenetics*, 66(12), 693–704. <https://doi.org/10.1007/s00251-014-0800-7>
- Jones, M. R., Cheviron, Z. A., & Carling, M. D. (2015). Spatially variable coevolution between a haemosporidian parasite and the MHC of a widely distributed passerine. *Ecology and Evolution*, 5(5), 1045–1060. <https://doi.org/10.1002/ece3.1391>
- Kaufman, J. (2018). Unfinished business: Evolution of the MHC and the adaptive immune system of jawed vertebrates. *Annual Review of Immunology*, 36(1), 383–409. <https://doi.org/10.1146/annurev-immunol-051116-052450>
- Lefort, V., Longueville, J. E., & Gascuel, O. (2017). SMS: Smart model selection in PhyML. *Molecular Biology and Evolution*, 34(9), 2422–2424. <https://doi.org/10.1093/MOLBEV/MSX149>
- Lemoine, F., Correia, D., Lefort, V., Doppelt-Azeroual, O., Mareuil, F., Cohen-Boulakia, S., & Gascuel, O. (2019). NGPhylogeny.fr: New generation phylogenetic services for non-specialists. *Nucleic Acids Research*, 47(W1), W260–W265. <https://doi.org/10.1093/nar/gkz303>
- Lenz, T. L. (2011). Computational prediction of MHC II-antigen binding supports divergent allele advantage and explains trans-species polymorphism. *Evolution*, 65(8), 2380–2390. <https://doi.org/10.1111/j.1558-5646.2011.01288.x>
- Letunic, I., & Bork, P. (2021). Interactive tree of life (iTOL) v5: An online tool for phylogenetic tree display and annotation. *Nucleic Acids Research*, 49(W1), W293–W296. <https://doi.org/10.1093/nar/gkab301>
- Levy, H., Fiddaman, S. R., Vianna, J. A., Noll, D., Clucas, G. V., Sidhu, J. K. H., Polito, M. J., Bost, C. A., Phillips, R. A., Crofts, S., Miller, G. D., Pistorius, P., Bonnadonna, F., Le Bohec, C., Barbosa, A., Trathan, P., Raya Rey, A., Frantz, L. A. F., Hart, T., & Smith, A. L. (2020). Evidence of pathogen-induced immunogenetic selection across the large geographic range of a wild seabird. *Molecular Biology and Evolution*, 37(6), 1708–1726. <https://doi.org/10.1093/molbev/msaa040>
- Llanos-Soto, S., Muñoz, B., Moreno, L., Landaeta-Aqueveque, C., Kinsella, J. M., Mironov, S., Cicchino, A., Barrientos, C., Torres-Fuentes, G., & González-Acuña, D. (2017). External and gastrointestinal parasites of the rufous-collared sparrow *Zonotrichia capensis* (Passeriformes, Emberizidae) in Chile. *Revista Brasileira de Parasitologia Veterinária*, 26(3), 314–322. <https://doi.org/10.1590/s1984-29612017043>
- Loiseau, C., Richard, M., Garnier, S., Chastel, O., Julliard, R., Zoorob, R., & Sorci, G. (2009). Diversifying selection on MHC class I in the house sparrow (*Passer domesticus*). *Molecular Ecology*, 18(7), 1331–1340. <https://doi.org/10.1111/j.1365-294X.2009.04105.x>
- Loiseau, C., Zoorob, R., Garnier, S., Birard, J., Federici, P., Julliard, R., & Sorci, G. (2008). Antagonistic effects of a Mhc class I allele on malaria-infected house sparrows. *Ecology Letters*, 11(3), 258–265. <https://doi.org/10.1111/j.1461-0248.2007.01141.x>
- Lougheed, S. C., Campagna, L., Dávila, J. A., Tubaro, P. L., Lijtmaer, D. A., & Handford, P. (2013). Continental phylogeography of an ecologically and morphologically diverse Neotropical songbird, *Zonotrichia capensis*. *BMC Evolutionary Biology*, 13(1), 58. <https://doi.org/10.1186/1471-2148-13-58>
- Lundie, R. J., de Koning-Ward, T. F., Davey, G. M., Nie, C. Q., Hansen, D. S., Lau, L. S., Mintern, J. D., Belz, G. T., Schofield, L., Carbone, F. R., Villadangos, J. A., Crabb, B. S., & Heath, W. R. (2008). Blood-stage plasmodium infection induces CD8 + T lymphocytes to parasite-expressed antigens, largely regulated by CD8 α + dendritic cells. *Proceedings of the National Academy of Sciences of the United States of America*, 105(38), 14509–14514. <https://doi.org/10.1073/pnas.0806727105>
- Mantilla, J. S., González, A. D., Lotta, I. A., Moens, M., Pacheco, M. A., Escalante, A. A., Valkiunas, G., Moncada, L. I., Pérez-Tris, J., & Matta, N. E. (2016). *Haemoproteus erythrogravidus* n. sp. (Haemosporida, Haemoproteidae): Description and molecular characterization of a widespread blood parasite of birds in South America. *Acta Tropica*, 159, 83–94. <https://doi.org/10.1016/j.actatropica.2016.02.025>
- Martin, R., & Kaufman, J. (2022). Some thoughts about what non-mammalian jawed vertebrates are telling us about antigen processing and peptide loading of MHC molecules. *Current Opinion in Immunology*, 77, 102218. <https://doi.org/10.1016/j.coi.2022.102218>
- Martínez, J., Vásquez, R. A., Marqués, A., Díez-Fernández, A., & Merino, S. (2016). The prevalence and molecular characterisation of blood parasites infecting the vulnerable Tamarugo Conebill (*Conirostrum tamarugense*) and other birds in the Pampa del Tamarugal, Chile. *Emu - Austral Ornithology*, 116(3), 310–314. <https://doi.org/10.1071/MU15090>
- Martínez-de la Puente, J., Merino, S., Lobato, E., Rivero-de Aguilar, J., del Cerro, S., Ruiz-de-Castañeda, R., & Moreno, J. (2009). Does weather affect biting fly abundance in avian nests? *Journal of Avian Biology*, 40(6), 653–657. <https://doi.org/10.1111/j.1600-048X.2009.04726.x>
- Martínez-de la Puente, J., Merino, S., Tomás, G., Moreno, J., Morales, J., Lobato, E., García-Fraile, S., & Belda, E. J. (2010). The blood parasite *Haemoproteus* reduces survival in a wild bird: A medication experiment. *Biology Letters*, 6(5), 663–665. <https://doi.org/10.1098/rsbl.2010.0046>
- Marzal, A., García-Longoria, L., Callirgos Cárdenas, J. M., & Sehgal, R. (2015). Invasive avian malaria as an emerging parasitic disease in native birds of Peru. *Biological Invasions*, 17(1), 39–45. <https://doi.org/10.1007/s10530-014-0718-x>
- Medrano, F., Barros, R., Norambuena, H. V., Matus, R., & Schmitt, F. (2018). *Atlas de las aves nidificantes de Chile*. Red de Observadores de Aves y Vida Silvestre de Chile.
- Merino, S., Moreno, J., Jose Sanz, J., Arriero, E., Sanz, J. J., & Arriero, E. (2000). Are avian blood parasites pathogenic in the wild? A medication experiment in blue tits (*Parus caeruleus*). *Proceedings. Biological Sciences/The Royal Society*, 267(1461), 2507–2510. <https://doi.org/10.1098/rspb.2000.1312>
- Merino, S., Moreno, J., Vásquez, R. A., Martínez, J., Sánchez-Monsálvez, I., Estades, C. F., Ippi, S., Sabat, P., Rozzi, R., & McGehee, S. (2008). Haematzoa in forest birds from southern Chile: Latitudinal gradients in prevalence and parasite lineage richness. *Austral Ecology*, 33(3), 329–340. <https://doi.org/10.1111/j.1442-9993.2008.01820.x>
- Minias, P., Pikus, E., Whittingham, L. A., & Dunn, P. O. (2018). A global analysis of selection at the avian MHC. *Evolution*, 72(6), 1278–1293. <https://doi.org/10.1111/evo.13490>
- Minias, P., Pikus, E., Whittingham, L. A., & Dunn, P. O. (2019). Evolution of copy number at the MHC varies across the avian tree of life. *Genome Biology and Evolution*, 11(1), 17–28. <https://doi.org/10.1093/gbe/evy253>
- Moens, M. A. J., & Pérez-Tris, J. (2016). Discovering potential sources of emerging pathogens: South America is a reservoir of generalist avian blood parasites. *International Journal for Parasitology*, 46(1), 41–49.
- Møller, A. P. (1998). Evidence of larger impact of parasites on hosts in the tropics: Investment in immune function within and outside the tropics. *Oikos*, 82(2), 265–270. <https://doi.org/10.2307/3546966>
- Murphy, K., & Weaver, C. (2017). *Janeway's immunobiology* (9th ed.). Taylor & Francis Group LCC.
- Murrell, B., Moola, S., Mabona, A., Weighill, T., Sheward, D., Kosakovsky Pond, S. L., & Scheffler, K. (2013). FUBAR: A fast, unconstrained

- Bayesian AppRoximation for inferring selection. *Molecular Biology and Evolution*, 30(5), 1196–1205. <https://doi.org/10.1093/molbev/mst030>
- Nikolich-Zugich, J. (2004). The role of mhc polymorphism in antimicrobial resistance. *Microbes and Infection*, 6(5), 501–512. <https://doi.org/10.1016/j.micinf.2004.01.006>
- O'Connor, E. A., Westerdahl, H., Burri, R., & Edwards, S. V. (2019). Avian MHC evolution in the era of genomics: Phase 1.0. *Cells*, 8(10), 1152. <https://doi.org/10.3390/cells8101152>
- Olson, D. M., Dinerstein, E., Wikramanayake, E. D., Burgess, N. D., Powell, G. V. N., Underwood, E. C., D'Amico, J. A., Itoua, I., Strand, H. E., Morrison, J. C., Loucks, C. J., Allnutt, T. F., Ricketts, T. H., Kura, Y., Lamoreux, J. F., Wettengel, W. W., Hedao, P., & Kassem, K. R. (2001). Terrestrial ecoregions of the world: A new map of life on earth. *Bioscience*, 51(11), 933–938. [https://doi.org/10.1641/0006-3568\(2001\)051\[0933:TEOTWA\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2001)051[0933:TEOTWA]2.0.CO;2)
- Owen-Ashley, N. T., Hasselquist, D., Råberg, L., & Wingfield, J. C. (2008). Latitudinal variation of immune defense and sickness behavior in the white-crowned sparrow (*Zonotrichia leucophrys*). *Brain, Behavior, and Immunity*, 22(4), 614–625. <https://doi.org/10.1016/j.bbi.2007.12.005>
- Piertney, S. B., & Oliver, M. K. (2006). The evolutionary ecology of the major histocompatibility complex. *Heredity*, 96(1), 7–21. <https://doi.org/10.1038/sj.hdy.6800724>
- Poblete, Y., Contreras, C., Fernández, C., Flores, C. R., Vega, P., Ávila, M., & Sabat, P. (2023). Geographic variation in the altitudinal migration patterns, body size, oxidative status and exploratory behavior in a neotropical bird. *Ecology and Evolution*, 13, e9941. <https://doi.org/10.1002/ece3.9941>
- QGIS.org. (2022). QGIS geographic information system. QGIS association. Open source geospatial foundation project.
- Quillfeldt, P., Arriero, E., Martínez, J., Masello, J. F., & Merino, S. (2011). Prevalence of blood parasites in seabirds - a review. *Frontiers in Zoology*, 8(1), 26. <https://doi.org/10.1186/1742-9994-8-26>
- R Development Core Team. (2016). *R: A language and environment for statistical computing*. R foundation for statistical computing. <https://www.r-project.org/>
- Radwan, J., Babik, W., Kaufman, J., Lenz, T. L., & Winternitz, J. (2020). Advances in the evolutionary understanding of MHC polymorphism. *Trends in Genetics*, 36(4), 298–311. <https://doi.org/10.1016/j.tig.2020.01.008>
- Radwan, J., Biedrzycka, A., & Babik, W. (2010). Does reduced MHC diversity decrease viability of vertebrate populations? *Biological Conservation*, 143(3), 537–544. <https://doi.org/10.1016/j.biocon.2009.07.026>
- Rekdal, S. L., Anmarkrud, J. A., Johnsen, A., & Lifjeld, J. T. (2018). Genotyping strategy matters when analyzing hypervariable major histocompatibility complex-experience from a passerine bird. *Ecology and Evolution*, 8(3), 1680–1692. <https://doi.org/10.1002/ece3.3757>
- Rising, J. D., & Jaramillo, A. (2020). Rufous-collared sparrow (*Zonotrichia capensis*), version 1.0. In J. del Hoyo, A. Elliott, J. Sargatal, D. A. Christie, & E. de Juana (Eds.), *Birds of the world*. Cornell Lab of Ornithology. <https://doi.org/10.2173/bow.rucspa1.01>
- Rivero de Aguilar, J., Castillo, F., Moreno, A., Peñafiel, N., Browne, L., Walter, S. T., Karubian, J., & Bonaccorso, E. (2018). Patterns of avian haemosporidian infections vary with time, but not habitat, in a fragmented neotropical landscape. *PLoS One*, 13(10), e0206493. <https://doi.org/10.1371/journal.pone.0206493>
- Rock, K. L., Reits, E., & Neeffes, J. (2016). Present yourself! By MHC class I and MHC class II molecules. *Trends in Immunology*, 37(11), 724–737. <https://doi.org/10.1016/j.it.2016.08.010>
- Sallaberry-Pincheira, N., Gonzalez-Acuña, D., Herrera-Tello, Y., Dantas, G. P. M., Luna-Jorquera, G., Frere, E., Valdés-Velasquez, A., Simeone, A., & Vianna, J. A. (2015). Molecular epidemiology of avian malaria in wild breeding colonies of Humboldt and Magellanic penguins in South America. *EcoHealth*, 12(2), 267–277. <https://doi.org/10.1007/s10393-014-0995-y>
- Sallaberry-Pincheira, N., González-Acuña, D., Padilla, P., Dantas, G. P. M., Luna-Jorquera, G., Frere, E., Valdés-Velasquez, A., & Vianna, J. A. (2016). Contrasting patterns of selection between MHC I and II across populations of Humboldt and Magellanic penguins. *Ecology and Evolution*, 6(20), 7498–7510. <https://doi.org/10.1002/ece3.2502>
- Sanchez-Mazas, A. (2020). A review of HLA allele and SNP associations with highly prevalent infectious diseases in human populations. *Swiss Medical Weekly*, 150, w20214. <https://doi.org/10.4414/smww.2020.20214>
- Santiago-Alarcon, D., & Marzal, A. (2020). *Avian malaria and related parasites in the tropics: Ecology, evolution and systematics*. Springer. <https://doi.org/10.1007/978-3-030-51633-8>
- Santiago-Alarcon, D., Palinauskas, V., & Schaefer, H. M. (2012). Diptera vectors of avian Haemosporidian parasites: Untangling parasite life cycles and their taxonomy. *Biological Reviews*, 87(4), 928–964. <https://doi.org/10.1111/j.1469-185X.2012.00234.x>
- Saper, M. A., Bjorkman, P. J., & Wiley, D. C. (1991). Refined structure of the human histocompatibility antigen HLA-A2 at 2.6 Å resolution. *Journal of Molecular Biology*, 219(2), 277–319. [https://doi.org/10.1016/0022-2836\(91\)90567-P](https://doi.org/10.1016/0022-2836(91)90567-P)
- Sato, A., Tichy, H., Grant, P. R., Grant, B. R., Sato, T., & O'hUigin, C. (2011). Spectrum of MHC class II variability in Darwin's finches and their close relatives. *Molecular Biology and Evolution*, 28(6), 1943–1956. <https://doi.org/10.1093/molbev/msr015>
- Sebastian, A., Herdegen, M., Migalska, M., & Radwan, J. (2016). AMPLISAS: A web server for multilocus genotyping using next-generation amplicon sequencing data. *Molecular Ecology Resources*, 16(2), 498–510. <https://doi.org/10.1111/1755-0998.12453>
- Spurgin, L. G., & Richardson, D. S. (2010). How pathogens drive genetic diversity: MHC, mechanisms and misunderstandings. *Proceedings. Biological Sciences/The Royal Society*, 277(1684), 979–988. <https://doi.org/10.1098/rspb.2009.2084>
- Tamura, K., Stecher, G., & Kumar, S. (2021). MEGA11: Molecular evolutionary genetics analysis version 11. *Molecular Biology and Evolution*, 38(7), 3022–3027. <https://doi.org/10.1093/molbev/msab120>
- Trowsdale, J. (1995). "Both man & bird & beast": Comparative organization of MHC genes. *Immunogenetics*, 41(1), 1–17. <https://doi.org/10.1007/BF00188427>
- Valkiūnas, G. (2005). *Avian malaria parasites and other Haemosporidia*. CRC Press.
- Valkiūnas, G., Bensch, S., Iezhova, T. A., Krizanauskiene, A., Hellgren, O., & Bolshakov, C. V. (2006). Nested cytochrome b polymerase chain reaction diagnostics underestimate mixed infections of avian blood haemosporidian parasites: Microscopy is still essential. *Journal of Parasitology*, 92(2), 418–422. <https://doi.org/10.1645/GE-3547RN.1>
- Valkiūnas, G., Iezhova, T. A., Krizanauskiene, A., Palinauskas, V., Sehgal, R. N. M., & Bensch, S. (2008). A comparative analysis of microscopy and PCR-based detection methods for blood parasites. *Journal of Parasitology*, 94(6), 1395–1401. <https://doi.org/10.1645/GE-1570.1>
- van Oosterhout, C. (2009). A new theory of MHC evolution: Beyond selection on the immune genes. *Proceedings of the Royal Society B: Biological Sciences*, 276, 657–665. <https://doi.org/10.1098/rspb.2008.1299>
- Videvall, E., Palinauskas, V., Valkiūnas, G., & Hellgren, O. (2020). Host transcriptional responses to high- and low-virulent avian malaria parasites. *The American Naturalist*, 195(6), 1070–1084. <https://doi.org/10.1086/708530>
- Walther, E. L., Valkiūnas, G., González, A. D., Matta, N. E., Ricklefs, R. E., Cornel, A., & Sehgal, R. N. M. (2014). Description, molecular characterization, and patterns of distribution of a widespread New World avian malaria parasite (Haemosporida: Plasmodiidae),

- Plasmodium (Novyella) homopolare* sp. nov. *Parasitology Research*, 113(9), 3319–3332. <https://doi.org/10.1007/s00436-014-3995-5>
- Waterhouse, A. M., Procter, J. B., Martin, D. M. A., Clamp, M., & Barton, G. J. (2009). Jalview version 2—A multiple sequence alignment editor and analysis workbench. *Bioinformatics*, 25(9), 1189–1191. <https://doi.org/10.1093/bioinformatics/btp033>
- Weaver, S., Shank, S. D., Spielman, S. J., Li, M., Muse, S. V., & Kosakovsky Pond, S. L. (2018). Datamonkey 2.0: A modern web application for characterizing selective and other evolutionary processes. *Molecular Biology and Evolution*, 35(3), 773–777. <https://doi.org/10.1093/molbev/msx335>
- Westerdahl, H., Mellinger, S., Sigeman, H., Kutschera, V. E., Proux-Wéra, E., Lundberg, M., Weissensteiner, M., Churcher, A., Bunikis, I., Hansson, B., Wolf, J. B. W., & Strandh, M. (2022). The genomic architecture of the passerine MHC region: High repeat content and contrasting evolutionary histories of single copy and tandemly duplicated MHC genes. *Molecular Ecology Resources*, 22(6), 2379–2395. <https://doi.org/10.1111/1755-0998.13614>
- White, E. M., Greiner, E. C., Bennett, G. F., & Herman, C. M. (1978). Distribution of the hematozoa of neotropical birds. *Revista de Biología Tropical*, 26(Suppl 1), 43–102.
- Willig, M. R., Kaufman, D. M., & Stevens, R. D. (2003). Latitudinal gradients of biodiversity: Pattern, process, scale, and synthesis. *Annual Review of Ecology, Evolution, and Systematics*, 34, 273–309. <https://doi.org/10.1146/annurev.ecolsys.34.012103.144032>

Wood, S. N. (2023). Package “mgcv”: Mixed GAM computation vehicle with automatic smoothness estimation. In *Generalized Additive Models: An Introduction with R* (2nd ed.). CRC Press.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Rivero de Aguilar, J., Barroso, O., Bonaccorso, E., Cadena, H., Hussing, L., Jorquera, J., Martínez, J., Martínez-de la Puente, J., Marzal, A., León Miranda, F., Merino, S., Matta, N. E., Ramenofsky, M., Rozzi, R., Valeris-Chacín, C. E., Vásquez, R. A., Vianna, J. A., & Wingfield, J. C. (2024). Associations among MHC genes, latitude, and avian malaria infections in the rufous-collared sparrow (*Zonotrichia capensis*). *Ecology and Evolution*, 14, e11634. <https://doi.org/10.1002/ece3.11634>