

UC Irvine

UC Irvine Previously Published Works

Title

Direct Construction of Quaternary Carbons from Tertiary Alcohols via Photoredox-Catalyzed Fragmentation of tert-Alkyl N-Phthalimidoyl Oxalates

Permalink

<https://escholarship.org/uc/item/2272070q>

Journal

Journal of the American Chemical Society, 135(41)

ISSN

0002-7863

Authors

Lackner, Gregory L
Quasdorf, Kyle W
Overman, Larry E

Publication Date

2013-10-16

DOI

10.1021/ja408971t

Peer reviewed



Published in final edited form as:

J Am Chem Soc. 2013 October 16; 135(41): . doi:10.1021/ja408971t.

Direct Construction of Quaternary Carbons from Tertiary Alcohols via Photoredox-Catalyzed Fragmentation of *tert*-Alkyl *N*-Phthalimidoyl Oxalates

Gregory L. Lackner[†], Kyle W. Quasdorf[†], and Larry E. Overman^{*}

Department of Chemistry, 1102 Natural Sciences II, University of California, Irvine, California 92697-2025

Abstract

A convenient method for the direct construction of quaternary carbons from tertiary alcohols by visible-light photoredox coupling of *tert*-alkyl *N*-phthalimidoyl oxalate intermediates with electron-deficient alkenes is reported.

Forming sterically demanding quaternary carbons by the reaction of nucleophilic carbon radicals with electron-deficient alkenes is attractive for several reasons: the forming bond is unusually long in calculated transition structures (2.2–2.5 Å);¹ the rates of addition of tertiary radicals to electron-deficient alkenes are faster than those of Me, primary and secondary radicals;² and stereoselectivity in the addition of tertiary radicals to prochiral alkenes is typically larger than that of primary or secondary radicals.^{2a,3,4} Although these appealing features have been recognized for many years,^{2a,5} this method is not mentioned in a recent comprehensive survey of methods to construct quaternary carbons.⁶ The considerable potential such bond constructions hold was suggested in our recent formal total synthesis of (–)- aplyviolene, wherein the stereoselective coupling of a tertiary carbon radical, generated by visible-light photoredox decarboxylative fragmentation of *N*-(acyloxy)phthalimide **1**, to α -chlorocyclopentenone **2** was the central step (eq 1).^{7a} This result represented the first utilization of such substrates in a C–C bondforming reaction since their initial disclosure by Okada.⁸ We surmise that the reason tertiary carbon radicals have not heretofore played a significant role in assembling quaternary carbons is the lack of convenient methods for generating these intermediates from widely available tertiary alcohols.⁴ Further-more, despite the pronounced advancement of visible-light photoredox-catalyzed transformations in recent years, methods that generate tertiary carbon radicals remain largely unexplored.^{7b,9} In this communication, we report such a method.


Corresponding Author: leoverman@uci.edu.

[†]Author Contributions

G.L.L. and K.W.Q. contributed equally

Supporting Information

Experimental details, characterization data, and CIF files for **6j** and **7d**. This material is available free of charge via the Internet at <http://pubs.acs.org>.



(1)

The method we developed was inspired by Barton's pioneering introduction of *tert*-alkyl *N*-hydroxypyridine-2-thionyl oxalates for generating carbon radicals from alcohols.¹⁰ Although Barton oxalate intermediates can be formed from tertiary alcohols, the instability of these intermediates, which prevents their isolation, and their light sensitivity are likely responsible for their limited use in the formation of quaternary carbons.^{11–15} Anticipating that related *tert*-alkyl *N*-phthalimidoyl oxalates would be more convenient precursors of tertiary carbon radicals, we initially explored conditions to access these compounds.

After examining several potential methods for preparing these mixed oxalate diesters, we found that tertiary alcohols smoothly underwent acylation with chloro *N*-phthalimidoyl oxalate (**4**, generated *in situ* from oxalyl chloride and *N*-hydroxyphthalimide) at room temperature in the presence of Et₃N and catalytic DMAP to afford *tert*-alkyl *N*-phthalimidoyl oxalates in high yields (**5**→**6**, Figure 1). In contrast to *N*-(acyloxy)phthalimides, which are generally quite stable to aqueous workup and silica gel chromatography, oxalates **6** proved to be much more sensitive. However, we found that the byproducts of their synthesis typically could be removed by simple filtration to furnish the desired product in acceptable purity. *tert*-Alkyl *N*-phthalimidoyl oxalates **6a–6k** are stable to ambient light and can be easily prepared on multigram scale. Confirmation that this sequence did provide the previously unknown alkyl *N*-phthalimidoyl oxalates was secured by single-crystal X-ray analysis of oxalate diester **6j**.

With a general route to *tert*-alkyl *N*-phthalimidoyl oxalates in hand, we examined the coupling of oxalate **6a** with methyl vinyl ketone. Employing conditions similar to those used in our earlier studies with *N*-(acyloxy)phthalimides⁷ resulted largely in decomposition of the oxalate, furnishing coupled product **7a** in only small amounts (Table 1, entry 1). Omission of *i*-Pr₂NEt provided an improvement in yield (entry 2), reflecting the instability of **6a** to the presence of this amine. The use of *i*-Pr₂NEt•HBF₄ resulted in improved yields of **7a** (entries 3–9). Whereas using an excess or equal amount of methyl vinyl ketone relative to **6a** proved detrimental (entries 1–5),¹⁶ optimal yields were obtained when oxalate **6a** was present in slight excess (entries 6–10). It was also found that a 1:1 THF-CH₂Cl₂ solvent mixture was superior to either CH₂Cl₂ or THF alone (entries 8 and 9). The *i*-Pr₂NEt•HBF₄ additive was found to be beneficial even under optimized conditions, as its omission led to lower product yields (entry 10). Finally, [Ru(bpy)₃](BF₄)₂ or the commercially available [Ru(bpy)₃](PF₆)₂ performed comparably in the reaction (entries 8 and 9). Under the optimized reaction conditions, the coupling of oxalate **6a** with methyl vinyl ketone gave ketone product **7a** in 82% yield.

The results of our initial survey of the scope of the coupling of *tert*-alkyl *N*-phthalimidoyl oxalates with methyl vinyl ketone are summarized in Table 2. In most cases, yields of the coupled products were excellent, ranging from 68–85% (entries 1–7). Chiral oxalates **6c**, **6d** and **6e** coupled with high diastereoselectivity (>20:1) from the sterically most accessible face (entries 3, 4 and 5). The coupling of the estrone-derived precursor **6e** is significant as it

demonstrates the construction of vicinal-quaternary carbon centers (entry 5). Two nitrogen-containing heterocyclic substrates were also investigated (entries 7 and 8). Piperidinone-derived **6h** furnished coupled product **7h** in 82% yield, whereas the coupling of indole-containing oxalate **6i** proceeded in diminished yield. As expected from our exploratory studies, carrying out the coupling of **6d** and **6e** with equal equivalents of the oxalate precursor and methyl vinyl ketone gave the coupled products in somewhat diminished yields (4 and 5). The reaction of adamantyl oxalate **6j** led to the expected product **7j** in low yield, with the major product deriving from coupling of the intermediate alkoxycarbonyl radical with methyl vinyl ketone (entry 9). To no surprise, homoallylic oxalate **6k** coupled with methyl vinyl ketone to give butyrolactone **7k** (entry 7).^{15a,17}

The scope of this new coupling reaction with respect to the conjugate acceptor is summarized in Table 3. Acceptors possessing a terminal double bond generally performed best in the reaction, with acrylonitrile, phenyl vinyl sulfone, and methyl acrylate providing the coupled products in excellent yield (entries 1–3); the yield was somewhat lower with dimethyl acrylamide (entry 4). Dimethyl fumarate also served as an excellent coupling partner producing **9e** in 85% yield (entry 5).

Cyclic acceptors could be utilized also, although the yields of coupled products were slightly lower (entries 6–9). Use of a butenolide acceptor (entry 8) furnished the desired product in 72% yield, with addition occurring exclusively from the face opposite the methoxy substituent.¹⁸ Finally, the activated trisubstituted acceptor 2-carbomethoxycyclopropane-2-ene-1-one coupled with *tert*-alkyl *N*-phthalimidoyl oxalate **6a** to afford trans product **9i** in 62% yield (entry 9).

A plausible mechanism for the visible-light photoredox coupling reported herein is outlined in Scheme 1. As proposed by Okada⁸ for the fragmentation of *N*-(acyloxy)phthalimides, single-electron transfer from Ru(bpy)₃⁺ to the *tert*-alkyl *N*-phthalimidoyl oxalate, followed by homolytic cleavage of the N–O bond and subsequent decarboxylation generates alkoxycarbonyl radical **12**. A second slower decarboxylation leads to the formation of tertiary radical **13**,^{10,19} which after addition to an electron-deficient alkene provides α-acyl radical **14**. Hydrogen atom abstraction from Hantzsch ester radical cation **10b** would furnish the final product (**14**→**7**). An alternative pathway (not shown), involving reduction of α-acyl radical **14** to the corresponding enolate and subsequent protonation, could also lead to the formation of **7**. While the presence of Hantzsch ester **10a** is necessary to realize catalytic turnover, the role of the ammonium additive is not fully understood at this time. However, it could potentially serve to protonate intermediate radical anion **11** and also act as a source of BF₄[−] to undergo anion exchange with [Ru(bpy)₃](PF₆)₂ leading to the formation of the more soluble [Ru(bpy)₃](BF₄)₂ complex.

In conclusion, we have developed a convenient method for the direct construction of quaternary carbons from tertiary alcohols by visible-light photoredox coupling of *tert*-alkyl *N*-phthalimidoyl oxalate intermediates with electron-deficient alkenes. In three examples, in which the intermediate tertiary carbon radical is chiral and sterically biased, diastereoselection in forming the new quaternary carbon stereocenter was excellent. Additional synthetic applications of *tert*-alkyl *N*-phthalimidoyl oxalates, as well as mechanistic studies of their reactivity in photoredox-mediated processes, are currently under investigation and will be reported in due course.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Support was provided by the National Science Foundation (CHE1265964). NMR and mass spectra were determined at UC Irvine using instruments purchased with the assistance of NSF and NIH shared instrumentation grants. We thank Dr. Joseph Ziller and Dr. John Greaves, Department of Chemistry, UC Irvine, for their assistance with X-ray and mass spectrometric analyses.

References

1. (a) Damm W, Giese B, Hartung J, Hasskerl T, Houk KN, Hüter; Zipse H. *J Am Chem Soc.* 1992; 114:4067. Arnaud R, Postlethwaite H, Barone V. *J Phys Chem.* 1994; 98:5913.
2. (a) Giese B. *Angew Chem, Int Ed.* 1983; 22:753. (b) Beckwith AL, Poole JS. *J Am Chem Soc.* 2002; 124:9489. [PubMed: 12167045]
3. Hayen A, Koch R, Saak W, Haase D, Metzger JO. *J Am Chem Soc.* 2000; 122:12458.
4. For recent reviews that discuss the addition of carbon radicals to alkenes, see: Renaud P, Sibi MP. *Radicals in Organic Synthesis.* Wiley-VCH Weinheim 2001; Zard SZ. *Radical Reactions in Organic Synthesis.* Oxford New York 2003 Srikanth GSC, Castle SL. *Tetrahedron.* 2005; 61:10377. Rowlands GJ. *Tetrahedron.* 2009; 65:8603.
5. Barton DHR, Sas W. *Tetrahedron.* 1990; 46:3419.
6. Christoffers, A Baro. *Quaternary Stereocenters—Challenges and Solutions for Organic Synthesis.* Wiley-VCH; Weinheim: 2005.
7. For use of visible-light photoredox catalysis in complex natural product synthesis see: Schnermann MJ, Overman LE. *Angew Chem, Int Ed.* 2012; 51:9576. For use of a tertiary halide as a radical precursor see: Furst L, Narayanam JMR, Stephenson CRJ. *Angew Chem, Int Ed.* 2011; 50:9655.
8. For the pioneering development of N-(acyloxy)phthalimides for forming carbon radicals, see: Okada K, Okamoto K, Morita N, Okubo K, Oda M. *J Am Chem Soc.* 1991; 113:9401.
9. For recent reviews, see: Tucker JW, Stephenson CRJ. *J Org Chem.* 2012; 77:1617. [PubMed: 22283525] Prier CK, Rankic DA, MacMillan DWC. *Chem Rev.* 2013; 113:5322. [PubMed: 23509883]
10. (a) Barton DHR, Crich D. *Tetrahedron Lett.* 1985; 26:757. (b) Barton DHR, Crich D, Kretzschmar G. *J Chem Soc, Perkin Trans 1.* 1986:39.
11. Elimination is a serious complication,^{10b} see also: Aggarwal VK, Angelaud R, Bihan D, Blackburn P, Fieldhouse R, Fonquerna SJ, Ford GD, Hynd G, Jones E, Jones RVH, Jubault P, Palmer MJ, Ratcliffe PD, Adams H. *J Chem Soc, Perkin Trans 1.* 2001:2604.
12. Tertiary bromides and iodides are the most widely employed precursors of tertiary radicals.^{4,7b} However, as a result of competing rearrangement and elimination reactions, these halides will not be available from many structurally complex tertiary alcohols.^{7a}
13. Although many alcohol derivatives have found use in the deoxygenation of tertiary alcohols,¹⁴ to our knowledge there are only scattered reports of the bimolecular trapping of tertiary radicals generated from these precursors to construct quaternary carbons.^{10,15}
14. McCombie SW, Motherwell WB, Tozer MJ. *Org React.* 2012; 77:161.
15. For examples, see: Togo H, Fujii M, Yokoyama M. *Bull Chem Soc Jpn.* 1991; 64:57. Togo H, Matsubayashi S, Yamazaki O, Yokoyama M. *J Org Chem.* 2000; 65:2816. [PubMed: 10808463] Blazejewski J-C; Diter P, Warchol T, Wakselman C. *Tetrahedron Lett.* 2001; 42:859. Sunazuka T, Yoshida K, Kojima N, Shirahata T, Hirose T, Handa M, Yamamoto D, Harigaya Y, Kuwajima I, Omura S. *Tetrahedron Lett.* 2005; 46:1459.
16. Excess methyl vinyl ketone contributes to diminished product yields by the addition of intermediate 14 (Scheme 3) to a second equivalent of methyl vinyl ketone. See reference 8 for similar observations.
17. Togo H, Yokoyama M. *Heterocycles.* 1990; 31:437.
18. Morikawa T, Washio Y, Harada S, Hanai R, Kayashita T, Nemoto H, Shiro M, Taguchi T. *J Chem Soc, Perkin Trans 1.* 1995:271.
19. Simakov PA, Martinez FN, Horner JH, Newcomb M. *J Org Chem.* 1998; 63:1226.





Figure 1. Synthesis of *tert*-alkyl *N*-phthalimidoyl oxalates

^aIsolated yield, ^bPhth = *N*-phthalimido



Scheme 1.

Table 1

Optimization for the coupling of 6a with methyl vinyl ketone.

Entry	oxalate equiv	MVK equiv	Hantzsch ester equiv	additive equiv	solvent ^d	X	Yield (%) ^{b,c}
1	1.0	1.5	1.0	<i>i</i> -Pr ₂ NEt•Hantzsch ester methyl vinyl ketone additive	CH ₂ Cl ₂	BF ₄	16 ^b
2	1.0	1.5	1.0	none	CH ₂ Cl ₂	BF ₄	36 ^b
3	1.0	1.5	1.0	<i>i</i> -Pr ₂ NEt•HBF ₄ (2.2)	CH ₂ Cl ₂	BF ₄	40 ^b
4	1.0	5.0	1.0	<i>i</i> -Pr ₂ NEt•HBF ₄ (2.2)	CH ₂ Cl ₂	BF ₄	22 ^b
5	1.0	1.0	1.0	<i>i</i> -Pr ₂ NEt•HBF ₄ (2.2)	CH ₂ Cl ₂	BF ₄	56 ^b
6	1.5	1.0	1.5	<i>i</i> -Pr ₂ NEt•HBF ₄ (2.2)	CH ₂ Cl ₂	BF ₄	77 ^b
7	1.5	1.0	1.5	<i>i</i> -Pr ₂ NEt•HBF ₄ (1.0)	CH ₂ Cl ₂	BF ₄	82 ^b
8	1.5	1.0	1.5	<i>i</i> -Pr ₂ NEt•HBF ₄ (1.0)	THF/CH ₂ Cl ₂ ^a	BF ₄	92 ^b
9	1.5	1.0	1.5	<i>i</i> -Pr ₂ NEt•HBF ₄ (1.0)	THF/CH ₂ Cl ₂ ^a	PF ₆	92 ^b 82 ^c
10	1.5	1.0	1.5	none	THF/CH ₂ Cl ₂ ^a	PF ₆	65 ^c

^a 1:1 mixture of THF-CH₂Cl₂; Hantzsch ester = diethyl 1,4-dihydro-2,6-dimethyl-3,5-pyridinedicarboxylate.

^b Yield determined by ¹H NMR.

^c Isolated yield after silica gel chromatography.





Table 2

Coupling of tert-alkyl *N*-phthalimidoyl oxalates with methyl vinyl ketone.

Entry	Oxalate	Product	Yield (%) ^{b,c}
1	 6a	 7a	82 ^b
2	 6b	 7b	85 ^b
3	 6c	 7c	82 ^b
4	 6d	 7d	85 ^b 63 ^c
5	 6e	 7e	68 ^b 43 ^c
6	 6f R = TBS	 7f	81 ^b
	 6g R = PMB	 7g	78 ^b



Entry	Oxalate	Product	Yield (%) ^{b,c}
7			82 ^b
8			36 ^b
9			22 ^b
			65 ^b
10			43 ^b

^a 1:1 mixture of THF: CH_2Cl_2 ,

^b Isolated yield after silica gel chromatography (average of two experiments),


^c Isolated yield after silica gel chromatography with 1:1 ratio of oxalate precursor to acceptor,

^d Phth = *N*-phthalimido

Table 3

Coupling of 6a with various acceptors.

Entry	Acceptor	Product	Yield (%) ^b
1			92 ^b
2			89 ^b
3			84 ^b
4			64 ^b
5			85 ^b
6			54 ^b
7			52 ^b
8			72 ^b



Entry	Acceptor	Product	Yield (%) ^b
9			62 ^b

^a 1:1 mixture of THF:CH₂Cl₂.

^b Isolated yield after silica gel chromatography (average of two experiments)