

UC Agriculture & Natural Resources Farm

Title

Pesticide Selection to Reduce Impacts on Water Quality

Permalink

<https://escholarship.org/uc/item/22n4f423>

Authors

Ferruzzi, Giulio

Gan, Jay

Publication Date

2004-09-01

DOI

10.3733/ucanr.8119

Peer reviewed


UNIVERSITY OF CALIFORNIA

Division of Agriculture
and Natural Resources
<http://anrcatalog.ucdavis.edu>

In partnership with



<http://www.nrcs.usda.gov>

**Farm Water
Quality Planning**
*A Water Quality and
Technical Assistance
Program for California
Agriculture*

This Reference Sheet is part of the **Farm Water Quality Planning (FWQP)** series, developed for a short course that provides training for growers of irrigated crops who are interested in implementing water quality protection practices. The short course teaches the basic concepts of watersheds, nonpoint source pollution (NPS), self-assessment techniques, and evaluation techniques. Management goals and practices are presented for a variety of cropping systems.



Reference:

Pesticide Selection to Reduce Impacts on Water Quality

GIULIO FERRUZZI is Agronomist, USDA–NRCS, San Luis Obispo County, and **JAY GAN** is Associate Professor and Water Quality Specialist, UC Riverside.

This publication's purpose is to help individual growers make their pesticide-use decisions with water quality in mind. There are several factors that influence a pesticide's potential to affect water quality, including soil properties (e.g., soil texture, organic matter content), pesticide properties (e.g., sorption to soil, half-life), climatic conditions (e.g., rainfall, temperature), and management practices or techniques (e.g., application method, irrigation). **Table 1** summarizes how a number of these factors can affect water quality. Along with other site-specific factors, you should take the factors listed in **Table 1** into consideration every time you choose a pesticide for application.

Despite all of these factors that can affect a pesticide's potential to impact water quality, though, it is the chemical properties of the pesticide that determine whether it is likely to impact surface water or ground water. In **Tables 2, 3, and 4** we have ranked the potential of some common pesticides to impact surface water bodies and ground water according to three pesticide properties: K_{OC} , water solubility, and soil half-life.

K_{OC} is the soil/organic carbon partitioning coefficient. It is highly dependent both on the pesticide's fat solubility and on the organic carbon content of a soil. The larger the K_{OC} , the more strongly the pesticide will sorb to the soil and the less likely it is to migrate to ground water. Water solubility is reported in ppm (parts per million), which is equivalent to milligrams of chemical (in this case, pesticide) that can be dissolved in a liter of water (1 ppm \approx 1 mg/L). The more soluble a pesticide is, the more likely it is to migrate to ground water or move offsite in surface runoff, although there are exceptions (e.g., glyphosate). A pesticide's soil half-life is the number of days it takes for half of the pesticide to degrade in the soil. The longer the half-life, the more persistent a pesticide is and thus the more probable it is that it will move into ground water or surface water.

Figure 1 gives examples of how K_{OC} and solubility affect the movement of pesticides in soils. Note that K_{OC} and solubility have a strong influence on pesticide leachability. The pesticides in **Tables 2, 3, and 4** are reported as having a small, medium, or large potential to impact the surface or ground water. Precautionary actions should be taken above and beyond basic best management practices for pesticides that rank as medium or large potential.

It is important to note that, while the impact of pesticides on groundwater quality is mainly a human health concern (because of its effect on the potability of well water), the effect on surface water quality is often a concern for aquatic organisms or wildlife. This is especially important for pesticides that can run off from fields into streams and lakes, where the pesticides may harm aquatic invertebrates and fish or cause long-term harm to wildlife that feeds on those fish and invertebrates. **Tables 2, 3, and 4**, list the toxicity of some common pesticides to fish, birds, and other wildlife. By taking appropriate measures (e.g., modifying application methods, application




Figure 1. Comparative leaching of several herbicides in soils. USDA photo by C.S. Helling; previously published in Brady 1984.

has minimal effect on natural enemies, among other desirable characteristics. You can find more information on some common pesticides' toxicity toward certain natural enemies in the Natural Enemies Handbook (Flint and Dreistadt 1998) (UC ANR Publication 3386).

In summary, your choice of pesticides should be based on several factors. For example, when you are applying pesticides to a location where leaching is a major concern, you should choose a pesticide with a short half-life, high K_{OC} , low solubility, and low toxicity toward aquatic organisms, wildlife, beneficial insects, and other nontarget species. In contrast, when you are applying pesticides to a location where runoff is a major concern, pesticide properties (half-life, K_{OC} , solubility) become less important and management and pesticide toxicity become more important. You should still choose a pesticide with a small half-life, high K_{OC} , low solubility, and low toxicity to aquatic organisms, wildlife, beneficial insects, and other nontarget species, but you also need to pay close attention to field management practices. Erosion should be controlled, for instance, to keep pesticides that are associated with field sediment from making their way into streams and waterways. Pesticides should be incorporated during application if possible and any sediment generated from the field during rainfall or irrigation events should be retained, for instance, in a sediment pond or vegetated filter strip.

In selecting pesticides that are appropriate for a specific crop and pest issue, refer to the University of California IPM Pest Management Guidelines (<http://www.ipm.ucdavis.edu>), UC Cooperative Extension IPM Advisors, or a certified Pest Control Advisor (PCA). You can use Tables 1 through 4 along with other site-specific information to make an educated decision about which pesticide would be most appropriate in terms of water quality and fish or wildlife toxicity. If the pesticide you are interested in is not in Tables 2 through 4, consult the USDA–NRCS Windows Pesticide Screening Tool (<http://www.wcc.nrcs.usda.gov/pestmgt/winpst.html>) or the University of California's Pesticide Wise Web site (<http://www.pw.ucr.edu>), which contains a more comprehensive listing of pesticides and their properties.

rates, timing, sediment/erosion control structures, tail-water recovery systems, vegetative buffers, etc.), you can reduce the potential impact to these species.

An additional concern is the potential effect of pesticides on the natural enemies of the target pests. If pesticide applications significantly reduce natural enemy populations, you may have to make more-frequent applications to suppress the resurgent pest populations. More applications, in turn, increase the potential to affect water quality. One of your goals is to protect water quality, so you should select a pesticide that

Table 1. Water quality impact potential as influenced by water, pesticides, and soil properties (H = High and L = Low).

| | Pesticide Properties | | | | | Soil Properties | | | | | | | | Rainfall/Irrigation Events | | Management Practices | |
|-------------------------------------|-----------------------|----------------------|-------------------------------------------------|-----------------------------------------------|------------|----------------------|--------------------|------------------------|-----------------------|---------------------------------|------------------------------------|---------------------|------------------|---------------------------------------------|--------------------------------------------|----------------------|----------------|
| | High water solubility | Low water solubility | Large K _{OC} (strongly sorbed to soil) | Small K _{OC} (weakly sorbed to soil) | Persistent | Coarse-textured soil | Fine-textured soil | High in organic matter | Low in organic matter | Many large connected soil pores | Few small discontinuous soil pores | Shallow water table | Deep water table | Small volumes not extending below root zone | Large volumes exceeding evapotranspiration | Broad-casting | Incorporating |
| Risk of ground-water impact | H | L | L | H | H | H | L | L | H | H | L | H | L | L | H* | L [†] | — [‡] |
| Risk of surface-water impact | L [#] | H | H | L [#] | H | L | H | H | L | L | H | — [#] | — [#] | L [#] | H | H | L |

* can be L if pesticide solubility is low or K_{OC} is large and organic matter is high
 † can be H if pesticide has high solubility, low K_{OC} and excessive rainfall/irrigation exists
 ‡ dependent on pesticide properties, soil properties, and rainfall/irrigation events
 # can be H if excessive runoff exists

Table 2. Water quality impact potential and toxicity information of some common herbicides.

| Herbicide Common Name | Herbicide Trade Name | Soil Sorption Index (K _{OC}) | Water Solubility (ppm) | Soil Half-Life (days) | Leaching Potential* | Solution Runoff Potential [†] | Adsorbed Runoff Potential [‡] | Toxicity to Fish [#] | Toxicity to Birds and Other Wildlife [§] | References |
|-----------------------------------------|----------------------|----------------------------------------|------------------------|-----------------------|---------------------|----------------------------------------|----------------------------------------|--------------------------------|---------------------------------------------------|------------|
| <i>Phenoxy and Benzoic Acids</i> | | | | | | | | | | |
| 2,4-D dimethylamine | 2,4-D amine | 20 | 890 | 10 | Medium | Medium | Small | Moderate to slight | High to Slight | 1, 2 |
| DCPA | Dacthal | 5,000 | 0.5 | 100 | Small | Medium | Large | Slight to practically nontoxic | Moderate to practically nontoxic | 1, 2 |
| Dicamba | Banvel | 2 | 4x10 ⁵ | 14 | Large | Medium | Small | Practically nontoxic | Practically nontoxic | 1, 2 |
| <i>Triazines</i> | | | | | | | | | | |
| Atrazine | Aatrex | 100 | 33 | 60 | Large | Large | Medium | Slight | Practically nontoxic | 1, 2 |
| Cyanazine | Bladex | 190 | 170 | 14 | Medium | Medium | Small | Slight to moderate | Slight to moderate | 1, 2 |
| Hexazinone | Velpar | 54 | 33,000 | 90 | Large | Large | Medium | Slight | Slight to practically nontoxic | 1, 2 |
| Metribuzin | Sencor | 60 | 1,220 | 40 | Large | Large | Small | Slight | Moderate to slight | 1, 2 |
| Prometon | Pramitol | 150 | 720 | 500 | Large | Large | Medium | Practically nontoxic | Slight | 1, 3 |
| Prometryn | Promet | 400 | 33 | 60 | Medium | Large | Medium | Moderate | Practically nontoxic | 1, 2 |
| Simazine | Simazine | 130 | 6.2 | 60 | Large | Large | Medium | Slight to practically nontoxic | Practically nontoxic | 1, 2 |
| <i>Substituted Ureas</i> | | | | | | | | | | |
| Chlorsulfuron | Glean | 40 | 7,000 | 160 | Large | Large | Medium | Practically nontoxic | Practically nontoxic | 1, 4 |
| Diuron | Karmex | 480 | 42 | 90 | Medium | Large | Medium | Moderate | Slight | 1, 3 |
| Linuron | Lorox | 400 | 75 | 60 | Medium | Large | Medium | Slight | Slight | 1, 2 |
| Sulfometuron-methyl | Oust | 78 | 70 | 20 | Medium | Large | Small | Slight | Practically nontoxic | 1, 2 |
| Tebuthiuron | Spike | 80 | 2,500 | 360 | Large | Large | Medium | Slight to practically nontoxic | Practically nontoxic | 1, 2 |
| <i>Thiocarbamates</i> | | | | | | | | | | |
| Butylate | Sutan | 400 | 44 | 13 | Small | Large | Small | Moderate | Practically nontoxic | 1, 2 |
| Cycloate | Ro-Neet | 430 | 95 | 30 | Medium | Large | Small | Moderate | Practically nontoxic | 1, 4 |

* The potential for the pesticide to be lost via leaching.

[†] The potential for the pesticide to be lost by being transported away in surface runoff in the solution phase.[‡] The potential for the pesticide to be lost by being transported away in surface runoff while adsorbed to soil particles.[#] The toxicity categories are defined in Table 5. Most toxicities are reported for fish, although some include aquatic invertebrates also.[§] The toxicity categories are defined in Table 5. Most toxicities are reported for birds, although some include rabbits and other wildlife also.

Table 2. Water quality impact potential and toxicity information of some common herbicides (con't).

| Herbicide Common Name | Herbicide Trade Name | Soil Sorption Index (K_{OC}) | Water Solubility (ppm) | Soil Half-Life (days) | Leaching Potential* | Solution Runoff Potential† | Adsorbed Runoff Potential‡ | Toxicity to Fish# | Toxicity to Birds and Other Wildlife§ | References |
|--------------------------------|----------------------|----------------------------------|------------------------|-----------------------|---------------------|----------------------------|----------------------------|----------------------------------|---------------------------------------|------------|
| EPTC | Eradicane | 200 | 344 | 6 | Small | Medium | Small | Slight | Slight to practically nontoxic | 1, 2 |
| Molinate | Molinate | 190 | 970 | 21 | Medium | Medium | Small | High to slight | Practically nontoxic | 1, 2 |
| Pebulate | Tillam | 430 | 100 | 14 | Small | Medium | Small | Moderate | Practically nontoxic | 1, 4 |
| Triallate | Far-Go | 2,400 | 4 | 82 | Small | Large | Large | High | Practically nontoxic | 1, 2 |
| <i>Bipyridyliums</i> | | | | | | | | | | |
| Diquat dibromide | Diquat | 1×10^6 | 7.2×10^5 | 1,000 | Very small | Small | Large | Moderate to practically nontoxic | Slight to moderate | 1, 2 |
| Paraquat | Gramoxone | 1×10^6 | 6.2×10^5 | 1,000 | Very small | Small | Large | Moderate to slight | Moderate | 1, 2 |
| <i>Chloroacetamides</i> | | | | | | | | | | |
| Alachlor | Lasso | 170 | 240 | 15 | Medium | Medium | Small | Moderate | Practically nontoxic | 1, 2 |
| Metolachlor | Dual | 200 | 530 | 90 | High | High | Medium | Moderate | Slight to practically nontoxic | 1, 2 |
| <i>Dinitroanilines</i> | | | | | | | | | | |
| Oryzalin | Surflan | 600 | 2.5 | 20 | Small | Medium | Small | High | Slight to practically nontoxic | 1, 2 |
| Pendimethalin | Prowl | 5,000 | 0.28 | 90 | Small | Medium | Large | High | Slight | 1, 2 |
| Trifluralin | Treflan | 8,000 | 0.3 | 60 | Small | Medium | Large | Very high | Practically nontoxic | 1, 2 |
| <i>Nitriles</i> | | | | | | | | | | |
| Bromoxynil | Buctril | 192 | 0.8 | 8 | Small | Small | Medium | Very high to moderate | High to moderate | 1, 2 |
| Dichlobenil | Casoron | 400 | 21.2 | 60 | Medium | Large | Medium | Moderate to slight | Slight to practically nontoxic | 1, 4 |

* The potential for the pesticide to be lost via leaching.

† The potential for the pesticide to be lost by being transported away in surface runoff in the solution phase.

‡ The potential for the pesticide to be lost by being transported away in surface runoff while adsorbed to soil particles.

The toxicity categories are defined in Table 5. Most toxicities are reported for fish, although some include aquatic invertebrates also.

§ The toxicity categories are defined in Table 5. Most toxicities are reported for birds, although some include rabbits and other wildlife also.

Table 2. Water quality impact potential and toxicity information of some common herbicides (con't).

| Herbicide Common Name | Herbicide Trade Name | Soil Sorption Index (K_{OC}) | Water Solubility (ppm) | Soil Half-Life (days) | Leaching Potential* | Solution Runoff Potential† | Adsorbed Runoff Potential‡ | Toxicity to Fish# | Toxicity to Birds and Other Wildlife§ | References |
|-------------------------|----------------------|----------------------------------|------------------------|-----------------------|---------------------|----------------------------|----------------------------|--------------------------------|---------------------------------------|------------|
| <i>Other Herbicides</i> | | | | | | | | | | |
| Acrolein | Magnacide-H | 1 | 2.1x10 ⁵ | 14 | Large | Medium | Small | High to slight | Very high to high | 1, 4 |
| Bensulide | Prefar | 1,000 | 5.6 | 120 | Medium | Large | Large | Moderate to high | Slight | 1, 2 |
| Bentazon | Basagran | 35 | 2.3x10 ⁶ | 20 | Large | Large | Small | Practically nontoxic | Slight | 5, 1, 2 |
| Bromacil | Hyvar | 32 | 700 | 60 | Large | Large | Medium | Slight to practically nontoxic | Practically nontoxic | 1, 2 |
| Clopyralid | Stinger | 6 | 1,000 | 30 | Large | Medium | Small | Practically nontoxic | Slight to practically nontoxic | 1, 5 |
| Diethyl-ethyl | Antor | 1,400 | 105 | 21 | Small | Medium | Medium | Moderate | N/A | 1, 6 |
| Ethofumesate | Norton | 340 | 50 | 30 | Medium | Large | Small | Slight to practically nontoxic | Practically nontoxic | 1, 4 |
| Glyphosate | Roundup | 24,000 | 12,000 | 47 | Very small | Large | Large | Practically non-Toxic | Slight | 1, 2 |
| Imazethapyr | Pursuit | 10 | 2x10 ⁵ | 90 | Large | Large | Medium | Practically nontoxic | Practically nontoxic | 1, 4 |
| Isoxaben | Snapshot TG | 1,400 | 1 | 100 | Small | Large | Large | High to moderate | Practically nontoxic | 1, 4 |
| Napropamide | Devrinol | 400 | 74 | 70 | Medium | Large | Medium | Moderate to slight | Practically nontoxic | 1, 2 |
| Norflurazon | Evital | 600 | 28 | 90 | Medium | Large | Medium | Moderate | Slight to practically nontoxic | 1, 4 |
| Oxyfluorfen | Goal | 1x10 ⁵ | 0.1 | 35 | Very small | Large | Medium | High | Practically nontoxic | 1, 2 |
| Propyzamide | Kerb | 200 | 15 | 60 | Large | Large | Medium | Slight to practically nontoxic | Practically nontoxic | 1, 4 |
| Pyrazon | Pyramin | 120 | 400 | 21 | Large | Large | Medium | Slight | Slight | 1, 3 |
| Rimsulfuron | Matrix | 47 | 7,300 | 10 | Medium | Medium | Small | Practically nontoxic | Slight to moderate | 1, 3 |

* The potential for the pesticide to be lost via leaching.

† The potential for the pesticide to be lost by being transported away in surface runoff in the solution phase.

‡ The potential for the pesticide to be lost by being transported away in surface runoff while adsorbed to soil particles.

The toxicity categories are defined in Table 5. Most toxicities are reported for fish, although some include aquatic invertebrates also.

§ The toxicity categories are defined in Table 5. Most toxicities are reported for birds, although some include rabbits and other wildlife also.

Table 3. Water quality impact potential of common insecticides, nematicides, and other pesticides.

| Pesticide Common Name | Pesticide Trade Name | Soil Sorption Index (K_{OC}) | Water Solubility (ppm) | Soil Half-Life (days) | Leaching Potential | Solution Runoff Potential | Adsorbed Runoff Potential | Toxicity to Fish | Toxicity to Birds and Other Wildlife | References |
|------------------------------------------|----------------------|----------------------------------|------------------------|-----------------------|--------------------|---------------------------|---------------------------|----------------------------------|--------------------------------------|------------|
| <i>Pyrethroids and Botanicals</i> | | | | | | | | | | |
| Esfenvalerate | Asana | 5,300 | 0.002 | 35 | Small | Medium | Medium | Very high | Slight | 1, 2 |
| Fenvalerate | Pydrin | 5,300 | 0.002 | 35 | Small | Medium | Medium | High to moderate | Practically nontoxic | 1, 4 |
| Fluvalinate | Mavrik | 1×10^6 | 0.005 | 30 | Very small | Small | Medium | Very high | Slight | 1, 2 |
| Permethrin | Ambush | 1×10^5 | 0.006 | 30 | Very small | Small | Medium | Very high | Practically nontoxic | 1, 2 |
| Resmethrin | Crossfire | 1×10^5 | 0.01 | 30 | Very small | Small | Medium | Very high | Practically nontoxic | 1, 2 |
| <i>Carbamates</i> | | | | | | | | | | |
| Aldicarb | Temik | 30 | 6,000 | 30 | Large | Medium | Small | Moderate | Very high | 1, 2 |
| Carbaryl | Sevin | 300 | 120 | 10 | Small | Medium | Small | Moderate | Practically nontoxic | 1, 2 |
| Carbofuran | Furadan | 22 | 351 | 50 | Large | Large | Medium | High | High | 1, 2 |
| Formetanate | Carzol | 1×10^6 | 5×10^5 | 100 | Very small | Small | Large | Moderate to slight | High | 1, 4 |
| Methiocarb | MesuroI | 300 | 24 | 30 | Medium | Large | Small | Moderate to high | Very high to high | 1, 4 |
| Methomyl | Lannate | 72 | 58,000 | 30 | Large | Medium | Small | High to moderate | High | 1, 2 |
| Oxamyl | Vydate-L | 2.8×10^5 | 25 | 4 | Small | Medium | Small | Moderate to slight | Very high | 1, 2 |
| Thiodicarb | Larvin | 350 | 19.1 | 7 | Small | Large | Small | High to moderate | Practically nontoxic | 1, 4 |
| <i>Organophosphates</i> | | | | | | | | | | |
| Acephate | Orthene | 2 | 818,000 | 3 | Small | Medium | Small | Slight to practically nontoxic | Moderate to slight | 1, 4 |
| Azinphos-methyl | Guthion | 1,000 | 29 | 10 | Small | Medium | Medium | Very high to moderate | Slight to moderate | 1, 2 |
| Chlorpyrifos | Lorsban | 6,070 | 0.4 | 30 | Small | Small | Medium | Very high | Very high to moderate | 1, 2 |
| Chlorpyrifos-methyl | Dursban methyl | 3,000 | 4 | 7 | Small | Medium | Medium | Moderate to practically nontoxic | Slight | 1, 4 |
| Diazinon | D.Z.N. | 1,000 | 60 | 40 | Small | High | High | High to slight | Very high to high | 1, 2 |
| Dimethoate | Cygon | 20 | 39,800 | 7 | Medium | Medium | Small | Moderate | Very high to moderate | 1, 2 |
| Disulfoton | Di-Syston | 600 | 25 | 30 | Medium | Large | Small | High | Moderate | 1, 2 |
| Fenamiphos | Nemacur | 100 | 400 | 50 | Large | Large | Medium | High to moderate | Very high | 1, 2 |

Table 3. Water quality impact potential of common insecticides, nematicides and other pesticides (con't).

| Pesticide Common Name | Pesticide Trade Name | Soil Sorption Index (K_{OC}) | Water Solubility (ppm) | Soil Half-Life (days) | Leaching Potential | Solution Runoff Potential | Adsorbed Runoff Potential | Toxicity to Fish | Toxicity to Birds and Other Wildlife | References |
|----------------------------------------------------------|----------------------|----------------------------------|------------------------|-----------------------|------------------------|---------------------------|---------------------------|----------------------------------|--------------------------------------|------------|
| Fonofos | Dyfonate | 870 | 16.9 | 40 | Small | Large | Small | High | Extremely high | 1, 2 |
| Malathion | Cythion | 1,800 | 130 | 1 | Small | Small | Small | Very high to slight | Moderate | 1, 2 |
| Methyl parathion | Penncap-M | 5,100 | 60 | 5 | Small | Medium | Medium | Very high to high | Extreme | 1, 2 |
| Naled | Dibrom | 180 | 2,000 | 1 | Small | Medium | Small | High to moderate | High to moderate | 1, 2 |
| Parathion | Phoskil | 5,000 | 24 | 14 | Small | Medium | Medium | Very high to moderate | Very high to high | 1, 4 |
| Phorate | Thimet | 1,000 | 22 | 60 | Small | Large | Large | Very high | Very high to high | 1, 2 |
| Terbufos | Counter | 500 | 5 | 5 | Small | Medium | Small | High | Very high | 1, 2 |
| Trichlorfon | Dylox | 10 | 1.2x10 ⁵ | 10 | Large | Medium | Small | Very high | High to moderate | 1, 2 |
| <i>Organochlorides (Chlorinated hydrocarbons)</i> | | | | | | | | | | |
| Dicofol | Kelthane | 1.8x10 ⁵ | 1 | 60 | Very small | Small | Large | High | Slight | 1, 2 |
| Dienochlor | Pentac | 1,000 | 25 | 300 | Medium | Large | Large | Very high to high | Practically nontoxic | 1, 2 |
| Endosulfan | Thiodan | 12,400 | 0.32 | 50 | Very small | Medium | Large | Very high | High to moderate | 1, 2 |
| Lindane | Isotox | 1,100 | 7 | 400 | Medium | Large | Large | Very high to high | Moderate to practically nontoxic | 1, 2 |
| <i>Other</i> | | | | | | | | | | |
| Abamectin | Avid | 5,000 | 5 | 28 | Small | Medium | Medium | Very high | Practically nontoxic | 1, 2 |
| <i>Bacillus thuringiensis</i> | Dipel | N/A | N/A | 120 | Very small (estimated) | N/A | N/A | Practically nontoxic | Practically nontoxic | 2 |
| Bifenthrin | Talstar | 2.4x10 ⁵ | 0.1 | 26 | Very small | Small | Medium | Very high | Slight to practically nontoxic | 1, 4 |
| Cryolite | Kryocide | 10,000 | 420 | 3,000 | Small | Large | Large | Slight to practically nontoxic | Practically nontoxic | 1, 4 |
| Diflubenzuron | Dimilin | 10,000 | 0.08 | 10 | Small | Small | Medium | Practically nontoxic | Practically nontoxic | 1, 2 |
| Ethoprop | Mocap | 70 | 750 | 25 | Large | Medium | Small | Very high to slight | Very high to slight | 1, 4 |
| Imidacloprid | Admire | 440 | 580 | 127 | Large | Large | Medium | Slight to practically nontoxic | High to slight | 1, 4 |
| Metaldehyde | Metaldehyde | 240 | 230 | 10 | Small | Medium | Small | Moderate to practically nontoxic | Moderate to slight | 1, 4 |
| Oxydemeton-methyl | Metasystox-R | 10 | 1x10 ⁶ | 10 | Large | Medium | Small | High to slight | High to slight | 1, 4 |

Table 4. Water quality impact potential of common fungicides.

| Fungicide Common Name | Fungicide Trade Name | Soil Sorption Index (K_{OC}) | Water Solubility (ppm) | Soil Half-Life (days) | Leaching Potential | Solution Runoff Potential | Adsorbed Runoff Potential | Toxicity to Fish | Toxicity to Birds and Other Wildlife | Reference |
|---------------------------------------------------|----------------------|----------------------------------|------------------------|-----------------------|------------------------|---------------------------|---------------------------|--------------------------------|--------------------------------------|-----------|
| <i>Dithiocarbamates</i> | | | | | | | | | | |
| Mancozeb | Dithane | 2,000 | 6 | 70 | Small | Large | Large | High to moderate | Slight | 1, 2 |
| Maneb | Maneb | 2,000 | 6 | 70 | Small | Large | Large | High | Practically nontoxic | 1, 2 |
| <i>Dicarboximides</i> | | | | | | | | | | |
| Iprodione | Rovral | 700 | 13.9 | 14 | Small | Large | Small | Moderate | Slight | 1, 2 |
| Vinclozolin | Ronilan | 100 | 1,000 | 20 | Medium | Medium | Small | Moderate to slight | Practically nontoxic | 1, 2 |
| <i>Organochlorides (Chlorinated hydrocarbons)</i> | | | | | | | | | | |
| Chlorothalonil | Bravo | 1,380 | 0.6 | 30 | Small | Medium | Medium | High | Practically nontoxic | 1, 2 |
| PCNB (Quintozene) | Terraclor | 5,000 | 0.44 | 21 | Small | Small | Medium | High | Practically nontoxic | 1, 2 |
| Dichloropropene | Telone II | 32 | 2,250 | 10 | Medium | Medium | Small | Moderate | Moderate to practically nontoxic | 1, 4 |
| <i>Other Fungicides</i> | | | | | | | | | | |
| <i>Bacillus subtilis</i> | Serenade | N/A | N/A | N/A | Very small (estimated) | Small | Small | Practically nontoxic | Practically nontoxic | 3, 7 |
| Benomyl | Benlate | 1,900 | 2 | 67 | Very small | Small | Small | Very high to high | Moderate to practically nontoxic | 1, 2 |
| Captan | Captan | 200 | 5.1 | 3 | Small | Medium | Small | Very high | Practically nontoxic | 1, 2 |
| Carboxin | Vitavax | 260 | 195 | 7 | Small | Medium | Small | High to slight | Slight to practically nontoxic | 1, 4 |
| Chloropicrin | Chlor-O-Pic | 62 | 2,270 | 1 | Small | Medium | Small | Very high to high | N/A | 1, 4 |
| Dicloran | Botran | 1,000 | 7 | 10 | Small | Medium | Medium | High to slight | Slight to practically nontoxic | 1, 4 |
| Fosetyl-Al technical | Aliette | 20 | 1.2x10 ⁵ | 1 | Very small | Medium | Small | Practically nontoxic | Practically nontoxic | 1, 4 |
| Metalaxyl | Ridomil | 70 | 8,400 | 50 | Large | Large | Medium | Practically nontoxic | Practically nontoxic | 1, 2 |
| Triadimefon | Bayleton | 300 | 71.5 | 26 | Medium | Large | Small | Slight to practically nontoxic | Practically nontoxic | 1, 2 |
| Triflumizole | Procure | 40 | 12,500 | 14 | Medium | Medium | Small | High to moderate | Practically nontoxic | 1, 4 |

Table 5. Definition of toxicity categories used in Tables 2, 3, and 4.

| Toxicity Rating | Bird Acute Oral LD ₅₀ (mg/kg) | Fish water LC ₅₀ (mg/L) |
|----------------------|------------------------------------------|------------------------------------|
| Very high | <10 | <0.1 |
| High | 10–50 | 0.1–1 |
| Moderate | >50–500 | >1–10 |
| Slight | >500–2000 | >10–100 |
| Practically nontoxic | >2000 | >100 |

SOURCE: Modified from Kamrin, 1997, Lewis Publishers (an imprint of CRC Press).

REFERENCES FOR TABLES 2 THROUGH 4

Note: These references are numbered, and they are referenced by number in Tables 2 through 4.

1. Plotkin, Steven. 2004. USDA–NRCS Windows pesticide screening tool (WIN–PST) pesticide properties database. Amherst, MA: USDA–NRCS National Water and Climate Center. Available at <http://www.wcc.nrcs.usda.gov/pestmgmt/winpst.html>
2. Kamrin, Michael A. 1997. Pesticide profiles: Toxicity, environmental impact, and fate. Boca Raton: CRC Press LLC. 676 pp.
3. Material Safety Data Sheets for specific chemicals.
4. Office of Pesticide Programs. 2001. Environmental effects database (EEDB). Environmental Fate and Effects Division. Washington, DC: US Environmental Protection Agency.
5. Waskom, Reagan M. 1995. Best management practices for agricultural pesticide use. Fort Collins, CO: Colorado State University Cooperative Extension, Bulletin #XCM-177.
6. Hunn, J. B., E. P. Multer, and M. S. DeFelice. 1993. Fish and agricultural chemicals: Safeguarding your ponds. Columbia, MO: University of Missouri Cooperative Extension, publication G1912.
7. Marrone, Pamela. 2004. Personal communication.

ADDITIONAL INFORMATION ON MANAGING PESTICIDES IN AGRICULTURE

- Brady, N. C. 1984. The nature and properties of soils. 9th edition. New York: Macmillan.
- Brady, N. C., and R. R. Weil. 2002. The nature and properties of soils. 13th edition. Upper Saddle River, NJ: Prentice Hall.
- Dreistadt, Steve H. 2001. Integrated pest management for floriculture and nurseries. Oakland: University of California Division of Agriculture and Natural

- Resources, Publication 3402.
- Flint, Mary Louise, and Steve H. Dreistadt. 1998. *Natural enemies handbook*. Oakland: University of California Division of Agriculture and Natural Resources, Publication 3386.
- Forster, R., J. Trumbo, and T. Yargeau. 1997. *California wildlife and pesticides: A guide to using pesticides in and near fish and wildlife habitat*. Sacramento: California Department of Fish and Game. PAPA.
- Gan, Jay, and Peter Pang. no date. *Pesticide wise online database*. Riverside: University of California. Available at <http://www.pw.ucr.edu>
- IPM Manual Group. 1992. *Integrated pest management for cole crops and lettuce*. Oakland: University of California Division of Agriculture and Natural Resources, Publication 3307.
- IPM Manual Group. 1994. *Integrated pest management for strawberries*. Oakland: University of California Division of Agriculture and Natural Resources, Publication 3351.
- IPM Manual Group. 1999. *Integrated pest management for stone fruits*. Oakland: University of California Division of Agriculture and Natural Resources, Publication 3389.
- Kamrin, Michael A. 1997. *Pesticide profiles: Toxicity, environmental impact, and fate*. Boca Raton: CRC Press LLC.
- Mahler, Robert L., Hugh W. Homan, and Gene P. Carpenter. 1998. *Pesticides and their movement in soil and water*. Moscow, ID: University of Idaho Cooperative Extension, 5M 5-95, Current Information Series No. 865.
- UC IPM Program. various dates. *University of California integrated pest management guidelines*. Oakland: University of California Division of Agriculture and Natural Resources. Available at <http://www.ipm.ucdavis.edu/PMG/crops-agriculture.html>
- USDA. 2000. *USDA-NRCS Pest Management Standard 595*. Available from <http://www.ca.nrcs.usda.gov/rts/secr.htm>
- USDA. 2004. *USDA-NRCS Windows pesticide screening tool (WIN-PST), version 2.0062b*. Amherst, MA: USDA-NRCS National Water and Climate Center. <http://www.wcc.nrcs.usda.gov/pestmgt/winpst.html>

Figure 1 in this publication was adapted from *The Nature and Properties of Soils* (Brady 1984), courtesy of C. S. Helling, USDA. The information in this publication was drawn from the above listed sources, the sources referenced in the tables, and discussions with qualified professionals. Contact your local NRCS office or visit <http://www.nrcs.usda.gov> for further information.

FOR MORE INFORMATION

You'll find detailed information on many aspects of resource conservation in these titles and in other publications, slide sets, CD-ROMs, and videos from UC ANR:

Developing a Nonpoint Source Pollution Evaluation Program, publication 8087
Nonpoint Sources of Pollution in Irrigated Agriculture, publication 8055
The Farm Water Quality Plan, publication 9002

To order these products, visit our online catalog at <http://anrcatalog.ucdavis.edu>. You can also place orders by mail, phone, or FAX, or request a printed catalog of publications, slide sets, CD-ROMs, and videos from

University of California
Agriculture and Natural Resources
Communication Services
6701 San Pablo Avenue, 2nd Floor
Oakland, California 94608-1239

Telephone: (800) 994-8849 or (510) 642-2431, FAX: (510) 643-5470
e-mail inquiries: danrcs@ucdavis.edu

An electronic version of this publication is available on the ANR Communication Services Web site at <http://anrcatalog.ucdavis.edu>.

Publication 8119

© 2004 by the Regents of the University of California, Division of Agriculture and Natural Resources. All rights reserved.

The University of California prohibits discrimination or harassment of any person on the basis of race, color, national origin, religion, sex, gender identity, pregnancy (including childbirth, and medical conditions related to pregnancy or childbirth), physical or mental disability, medical condition (cancer-related or genetic characteristics), ancestry, marital status, age, sexual orientation, citizenship, or status as a covered veteran (covered veterans are special disabled veterans, recently separated veterans, Vietnam era veterans, or any other veterans who served on active duty during a war or in a campaign or expedition for which a campaign badge has been authorized) in any of its programs or activities.

University policy is intended to be consistent with the provisions of applicable State and Federal laws.

Inquiries regarding the University's nondiscrimination policies may be directed to the Affirmative Action/Staff Personnel Services Director, University of California, Agriculture and Natural Resources, 300 Lakeside Drive, 6th Floor, Oakland, CA 94612-3550, (510) 987-0096. For information about obtaining this publication, call (800) 994-8849. For downloading information, call (530) 754-5112.

pr-9/04-WJC/VJG



This publication has been anonymously peer reviewed for technical accuracy by University of California scientists and other qualified professionals. The review process was managed by the ANR Associate Editor for Natural Resources.