

UC Davis

UC Davis Previously Published Works

Title

Overexpression of an ABA biosynthesis gene using a stress-inducible promoter enhances drought resistance in petunia

Permalink

<https://escholarship.org/uc/item/2300w8ns>

Journal

Horticulture Research, 2(1)

ISSN

2662-6810

Authors

Estrada-Melo, Alejandro C
Chao
Reid, Michael S
et al.

Publication Date

2015-12-01

DOI

10.1038/hortres.2015.13

Peer reviewed

ARTICLE

Overexpression of an ABA biosynthesis gene using a stress-inducible promoter enhances drought resistance in petunia

Alejandro C Estrada-Melo¹, Chao Ma¹, Michael S Reid¹ and Cai-Zhong Jiang²

The response of plants to drought stress includes reduced transpiration as stomates close in response to increased abscisic acid (ABA) concentrations. Constitutive overexpression of 9-*cis*-epoxycarotenoid dioxygenase (*NCED*), a key enzyme in ABA biosynthesis, increases drought resistance, but causes negative pleiotropic effects on plant growth and development. We overexpressed the tomato *NCED* (*LeNCED1*) in petunia plants under the control of a stress-inducible promoter, *rd29A*. Under water stress, the transgenic plants had increased transcripts of *NCED* mRNA, elevated leaf ABA concentrations, increased concentrations of proline, and a significant increase in drought resistance. The transgenic plants also displayed the expected decreases in stomatal conductance, transpiration, and photosynthesis. After 14 days without water, the control plants were dead, but the transgenic plants, though wilted, recovered fully when re-watered. Well-watered transgenic plants grew like non-transformed control plants and there was no effect of the transgene on seed dormancy.

Horticulture Research (2015) 2, 15013; doi:10.1038/hortres.2015.13; Published online: 8 April 2015

INTRODUCTION

Drought, the major abiotic stress constraining agricultural production,¹ is a complex stress. Plants have evolved diverse strategies to respond to drought, depending on the timing, severity, and duration of drought in their native habitat.² Drought stress can be defined as the lack of the minimum amount of water to supply the basal requirements of otherwise well-adapted plants and drought resistance simply as the ability of the plant to survive under drought conditions and to recover following rehydration. Global climate change is expected to increase the occurrence of drought periods in some parts of the world; improving plant growth under water-limited conditions will therefore become one of the most challenging tasks for growers of all plant crops.³

Breeding for drought resistance would seem to be straightforward. Growing an outcrossed population under water-stressed conditions and then selecting the best lines for subsequent rounds of crossing should provide genotypes that are more resistant to drought. Unfortunately, such plants seldom perform well under favorable conditions. This suggests the potential for using a molecular genetic approach, either by increasing transcription of native genes that provide drought resistance or by introducing such genes from other species.

Obvious targets for this approach are genes involved in the synthesis and response to abscisic acid (ABA), the plant hormone that mediates the response of plants to stresses such as drought, cold, and salinity.⁴ Drought stress triggers increased tissue ABA concentrations, and the ABA signaling cascade results in stomatal closure, and reduced transpiration.^{5,6} When the water stress is removed, ABA concentrations fall, through catabolism and conjugation.⁷ The ability of plants to control endogenous ABA levels and regulate the stomatal aperture during water stress has been a useful tool for experiments in improving drought resistance.⁸

Prolonged exposure to osmotic stress that results from dehydration is the likely cause of tissue death in drought-stressed plants. In addition to stomatal closure, plants respond to increased ABA concentrations by activating osmotic-responsive genes. These genes encode proteins that help plants maintaining water uptake under drought stress by accumulating solutes, thus lowering water potential, preserving cell turgor, and minimizing water loss.⁹ There are two major mechanisms for osmotic adjustment, the accumulation of inorganic ions such as K⁺, Cl⁻, and Na⁺ or the accumulation of organic solutes (also called compatible solutes) that reduce the osmotic potential of the tissue. The mechanism employed varies among species, cultivars, and even by position within the same plant.¹⁰

The most common compatible solutes accumulated during osmotic adjustment are sugar alcohols such as mannitol, quaternary ammonium compounds such as betaine, and amino acids such as glycine and proline. Proline is also considered to protect plant cells from toxic by-products produced in response to drought. Yamada *et al.*¹¹ demonstrated in transgenic petunia plants that overexpression of *pyrroline-5-carboxylate synthetase* (*P5CS*), which is a key gene in proline biosynthesis, increased proline levels and improved plant resistance to drought stress.

ABA also acts as a signal for the induction of dehydration resistance genes,¹² which, in addition to protecting the cells from stress, regulate other genes involved in the response to drought stress. These genes include the *late embryogenesis abundant* (*LEA*) genes that provide desiccation tolerance during seed maturation, water transporters such as aquaporins, detoxification enzymes like superoxide dismutase and catalase, protein kinases, phosphatases and transcription factors that act as regulatory molecules.¹³

ABA biosynthesis is initiated in plastids where the C₄₀ carotenoid xanthophyll zeaxanthin is synthesized from isopentenyl diphospho-

¹Department of Plant Sciences, University of California Davis, Davis, CA, USA and ²Crops Pathology and Genetic Research Unit, United States Department of Agriculture, Agricultural Research Service, Davis, CA, USA

Correspondence: CZ Jiang, (cjiang@ucdavis.edu), MS Reid, (msreid@ucdavis.edu)

Received: 12 January 2015; Revised: 23 February 2015; Accepted: 26 February 2015

sphate (IPP). Epoxidation of zeaxanthin by zeaxanthin epoxidase (ZEP)¹⁴ results in another C₄₀ compound, violaxanthin, which is converted into neoxanthin through isomerization. Neoxanthin is oxidatively cleaved by 9-*cis*-epoxycarotenoid dioxygenase (NCED) to yield xanthoxin, the first C₁₅ intermediate in ABA biosynthesis.¹⁵ Xanthoxin is exported from the plastid to the cytosol and is converted in a two-step reaction to ABA with ABA aldehyde as the intermediate.^{16,17} NCED is considered to be the rate-limiting step in ABA biosynthesis; the gene encoding the enzyme was first cloned from the ABA-deficient maize mutant *vp14*.¹⁸

Overexpression of *NCED* has been shown to result in ABA accumulation and increased drought resistance in tomato (*Solanum lycopersicum*),¹⁹ *Arabidopsis thaliana*,^{20,21} and tobacco (*Nicotiana tabacum*).^{22,23} These studies all used a constitutive promoter *CaMV-35S* to drive expression of *NCED* genes, with negative pleiotropic effects on plant growth and development, including stunted growth, delayed germination, and seed dormancy.^{19,24}

Overexpression of *LeNCED1* in tomato under the control of the tissue-specific promoter *rbc53C* generated transgenic plants with much higher ABA levels in leaves and sap while the effect on seed dormancy was reduced. However in non-stressed conditions, these plants showed reduced stomatal conductance and CO₂ assimilation, increased photobleaching in young seedlings, reduction in chlorophyll and carotenoid content, interveinal water soaking, and greatly reduced growth.²⁴

We sought a strategy that would stimulate transcription of an *NCED* transgene only under stress conditions. For this purpose we selected the stress-inducible promoter *rd29A* from *Arabidopsis* to drive expression of a tomato *LeNCED1* in petunia. The *cis*-acting region of the *rd29A* promoter contains three dehydration-responsive elements and one abscisic acid responsive element. The promoter functions in almost all vegetative organs and tissues during water stress.²⁵

We hypothesized that under water stress conditions, the overexpression of *LeNCED1* using the stress-inducible promoter *rd29A* would increase the ABA and proline concentration in leaves, leading to stomatal closure, reducing the osmotic potential and enhancing drought resistance in petunia, while under non-stress conditions, there would be no negative effects on plant growth and development.

MATERIALS AND METHODS

Plant materials and growth conditions

Petunia (*Petunia hybrida* 'Mitchell Diploid'), obtained from Goldsmith Seeds (Gilroy, CA, USA), was used as a control and as the material for petunia *Agrobacterium*-mediated transformations. Petunia seeds were sterilized and germinated on a solid MS medium²⁶ in a growth chamber at 22 °C with light intensity of 40 μmol/m²/s. Seedlings with two true leaves were then carefully transferred into pots containing the same amounts of UC Mix (1/3 Peat, 1/3 Sand, 1/3 redwood compost), and grown in a greenhouse at University of California Davis under natural conditions without supplement lights at 25–30 °C.

Plasmid constructs

For the cloning of the *rd29A* promoter, a 1626 bp fragment of *rd29A* (GenBank Accession No. CS191722.1) in the upstream 5' noncoding region was amplified from wild-type *Arabidopsis* genomic DNA using the primers (Supplementary Table S1). The 2222 bp coding region of *LeNCED1* cDNA (GenBank Accession No. AJ439079.2) was amplified from tomato (*Solanum lycopersicum* Mill. L. cv. New Yorker) using the cloning primers (Supplementary Table S1). The PCR products were cloned into pGEM-T easy vector (Promega, Madison, WI, USA) for sequence confirmation by the sequencing service of the College of Biological Sciences at University of California Davis.

After sequence confirmation, the *LeNCED1* gene was introduced into a modified pGSA1403 binary expression vector²⁷ in which the *CaMV-35S* promoter was replaced with the *rd29A* promoter (Supplementary Figure 1). For transgenic plants that were used as a constitutively expressed control, the

LeNCED1 gene was introduced into the pGSA1403 binary vector downstream of the *CaMV-35S* promoter. The expression constructs, containing the kanamycin resistance gene for selection, were used for plant transformation.

Plant transformation

Petunia (*Petunia hybrida* 'Mitchell Diploid') was transformed via *Agrobacterium tumefaciens* strain LBA4404. Plants were transformed at the UC Davis transformation facility.²⁸ At the fourth to sixth leaf stage, the kanamycin-resistant primary transformants (T₀) were screened using polymerase chain reaction (PCR) to determine whether the plants contained the transgene.^{28,29}

Lines containing the transgene were then transferred into pots containing same amounts of UC Mix (1/3 Peat, 1/3 Sand, 1/3 redwood compost) and grown in a greenhouse under natural conditions without supplement lights at 25–30 °C until flowering. Plants then were self-pollinated to prevent cross-pollination T1 seeds that displayed a 3 : 1 ratio of survival on kanamycin were collected at maturity and used for further analysis.

Seeds of T1 transgenic lines were sterilized, germinated at room temperature under controlled illumination (14 h of light) on a solid MS medium,²⁶ and supplemented with 100 mg L⁻¹ kanamycin for selection of segregated progeny. Seeds from non-transformed control plants were germinated and grown on a solid MS medium without kanamycin. To identify homozygous lines, seedlings from selected lines were transferred to pots filled with same amounts of UC Mix and grown in a greenhouse with natural illumination at 25–30 °C until they set seed. One hundred seeds from each line were germinated again on MS medium containing 100 mg L⁻¹ kanamycin. Three transgenic *rd29A:LeNCED1* lines (8F, 7D, and 3D) showing 100% survival were selected as homozygous and used for molecular analysis and physiological assessments. In addition, one *CaMV-35S:LeNCED1* line 12-F was selected to serve as a control. Due to low germination, stunted growth and leaf photobleaching, this line was only used for limited experiments.

Semi-quantitative RT-PCR

To determine the expression of the *LeNCED1* and *PhP5CS* genes in non-transformed and transgenic petunia plants, RNA extracted from the leaves of T3 plants was analyzed using semi-quantitative RT-PCR. Prior to the extraction, plants were drought stressed by withholding water for 3 days. RNA was extracted from harvested leaves using TRIzol reagent (Invitrogen, CA, USA) according to the manufacturer's instructions. After DNase I treatment to remove contaminating DNA, 1 μg of total RNA was used for reverse transcription using the Superscript II Reverse Transcriptase Kit (Invitrogen, CA, USA) according to the manufacturer's protocol. The synthesized cDNA was used for analysis of transcript abundance using semi-quantitative RT-PCR and the primers shown in Supplementary Table S1. Gene expression levels were compared to the expression levels of 26S ribosomal RNA.²⁸ Semi-quantitative PCR reactions were carried out in 25 μL volumes containing 1 μL reverse transcription reaction products as templates. PCR products were analyzed by 1% (w/v) agarose gel electrophoresis. Three biological replications were used for the analysis.

The gel images from the semi-quantitative PCR analyses were quantified using the Gel Logic 200 software (Kodak, Rochester, NY). The mean area value of the three biological replicates was used to determine relative expression levels. Graphs were generated and statistically analyzed by one-way analysis of variance (ANOVA) using JMP (SAS Institute Inc., Cary, NC, USA). Student's *t*-test was used to examine significance of differences between treatments using a *P* value of 0.05.

ABA analysis

To determine ABA concentrations, two fully expanded leaves nearest to the apical meristem were collected and immediately frozen in liquid nitrogen. The tissues were lyophilized and placed at -80 °C until extraction. Approximately 100 mg samples were ground in liquid nitrogen and homogenized for 24 h in the dark at 4 °C in 80% chilled methanol containing 100 mg l⁻¹ butylated hydroxytoluene. After centrifugation, the supernatant was collected, purified through a Sep-Pak C18 cartridge column (Waters Inc., Milford, MA, USA) and eluted with 70% methanol. ABA content was determined using the enzyme-linked immunosorbent assay (ELISA) test kit (Aglia/Phytodetek, Elkhart, IN, USA) according to the manufacturer's instructions. All measurements were made from three individual plants as replicates and the results are expressed as nanomole ABA per gram fresh weight of tissue.

Proline determination

Proline content was measured spectrophotometrically based on proline's reaction with ninhydrin as described by Bates *et al.*³⁰ with modifications. The fully expanded leaves from transgenic and wild-type plants were sampled. Fresh leaves (100 mg) were frozen in liquid nitrogen and ground in a mortar and pestle in 0.5 ml of 3% aqueous sulfosalicylic acid. The resulting supernatant was added to the reaction mixture of 100 μ L 3% sulfosalicylic acid, 200 μ L glacial acetic acid and 200 μ L acidic ninhydrin. The reaction mix was incubated at 100 $^{\circ}$ C for 60 min, then extracted with 1 ml toluene. Absorbance at 520 nm was determined in a NanoDrop 2000c UV-Vis spectrophotometer (Thermo Fisher Scientific Inc., Waltham, MA USA) using toluene as a reference. The concentration of proline was determined by reference to a standard concentration curve.

Drought response

Eighteen *rd29A:LeNCED1-8F* and non-transformed petunia seedlings (2 weeks old) were transplanted into 4-inch pots containing same amounts of UC Mix. Pots were covered with aluminum foil to prevent any loss of water by evaporation from the soil surface. The plants were grown for 3 weeks in a greenhouse (natural illumination at 25–30 $^{\circ}$ C) under well watered conditions (pot capacity). At this point (day 0), the pots were weighed, and this value was considered to be 100% water content. Drought stress was then imposed by withholding water for 9 days, at which point most leaves of the non-transformed plants were wilted. The pots were weighed every morning (9:00–10:00 a.m.) for determination of percentage water loss.

Recovery after severe drought

After the drought treatment (14 days), plants were re-watered, and 10 days later the plants that displayed full turgor and resumed their growth were counted as recovered plants.

Effect of drought stress on ABA and proline content

Transgenic and non-transformed Petunia seedlings were transplanted into 4-inch pots containing UC Mix. Four weeks after transplanting, second and third leaves of transgenic and non-transformed plants were harvested to serve as controls. The remaining plants were drought stressed by withholding water. The second and third leaves of transgenic and non-transformed plants were harvested from each plant for determination of ABA and proline content 3 days after treatment.

Seed germination

To evaluate the effect of the *rd29A:LeNCED1* construct on seed germination, seeds of non-transformed and transgenic Petunia plants were sterilized and sown (100 seeds per assay) on Petri plates containing standard MS agar medium. Plates were held at 24 $^{\circ}$ C in 16 to 8-h light/dark cycles. Germination was scored when two fully expanded green cotyledons had developed. The experiment was repeated three times.

Gas-exchange measurements

We analyzed the effects of the *LeNCED1* transgene on net photosynthesis rate (A), stomatal conductance (g_s), and transpiration rate (E) using a portable photosynthesis system (model LI-6400, LI-COR Biosciences, Lincoln, NE, USA). Measurements were made on the uppermost fully expanded leaves of five individual transgenic plants using a photon flux density of 1000 μ mol.m⁻².s⁻¹ and a CO₂ concentration of 400 μ mol.mol⁻¹. Non-transformed and *CaMV-35S:LeNCED1* plants served as controls.

Stem water potential measurement

Stem water potential was measured using the bagged leaf technique in a Scholander-type pressure chamber.³¹ Fully expanded leaves near the stem at similar positions were used. Measurements were conducted at midday.³¹

Stomatal density

To measure leaf stomatal density a leaf imprint method was used. Three fully developed leaves for each plant and three mature plants were sampled. A drop of methacrylate glue (Crazy Glue) was applied to the adaxial and abaxial side of three different areas of each leaf, avoiding the midvein, and the leaf was pressed to a glass slide for 30 s. The leaf was removed and the imprint on the glass slide was observed using a light microscope.

Flower senescence

Ten flowers from plants of each transgenic line and from control plants were harvested when flowers were fully open but before the anthers had dehisced. Flowers were placed in 2 ml tubes containing a standard vase solution (Chrysal, Walterboro, SC, USA). Flowers were monitored daily and were scored as senesced when the corolla wilted. To determine the longevity of attached flowers, 10 fully open flowers of each transgenic line and of control plants were labeled, then monitored daily until their corollas were completely wilted.

Data analysis

All measurements were repeated at least three times from different individual plants. All data were subject to ANOVA according to the model for completely randomized design using Proc GLM (generalized linear model) with LSD means separation (least significant difference test; $P \leq 0.05$). Values were analyzed by JMP Version 11 (SAS Institute Inc., Cary, NC, USA).

RESULTS

Screening and characteristics of *rd29A:LeNCED1* lines

Following self-pollination and kanamycin selection of 12 T₀ lines obtained from the transformation facility, we selected three homozygous T₂ transgenic *rd29A:LeNCED1* lines 8F, 7D, and 3D. These plants exhibited normal growth and development under non-stress growth conditions (Figure 1). Anatomically, too, the plants were indistinguishable; we found no significant differences in stomatal density or stomatal index on abaxial and adaxial leaf surfaces between the transgenic *rd29A:LeNCED1* and control plants (data not shown). T₃ seedlings of these lines were used for further experiments. One homozygous line *CaMV-35S:LeNCED1-12F*, was used as a constitutive expression control in limited experiments. Under non-stressed conditions, the seeds from *CaMV-35S:LeNCED1* plants showed low germination rate and the plants grew more slowly than the non-transformed control and *rd29A:LeNCED1* plants (Figure 1), and older plants developed a severe photobleaching phenotype (Supplementary Figure 4).

In the germination assay, 95% of seeds from non-transformed plants germinated within 10 days. Seeds of the *rd29A:LeNCED1* lines, 8F, 7D, and 3D, gave germination rates of 95, 90, and 92% respectively. Only 50% of the seeds from the *CaMV-35S:LeNCED1* plants had germinated after 10 days (Supplementary Figure 2b). Germination was restored to the levels of the control seeds when *CaMV-35S:LeNCED1* seeds were treated with 500 ppm GA (Supplementary Figure 2c).

Drought stress response of vegetative plants

After 6 weeks in the greenhouse under normal conditions there were no visible differences in phenotype between the *rd29A:LeNCED1* transgenic plants and control plants (Figure 2). When water was then withheld for 1 week, there was a dramatic difference. The control plants were visibly wilted, while plants of the three transgenic lines were still turgid (Figure 2).

Recovery from drought stress

The recovery experiment compared control and *rd29A:LeNCED1-8F* plants. Control plants started to wilt at 3 days of drought (Figure 3) and were severely wilted by 10 days. At that time the transgenic plants showed wilting and yellowing only of the older (basal) leaves (Figure 3). After 2 weeks without water, the control plants were completely desiccated, and some transgenic plants were severely wilted, although their apices remained turgid. The plants were re-watered after 2 weeks of drought stress (Figure 3). Only 11% of the control plants survived, whereas 94.4% of the *rd29A:LeNCED1* transgenic plants recovered completely.

Drought stress response of flowering plants

After 12 weeks in the greenhouse, the *rd29A:LeNCED1* and control plants were flowering and were visually indistinguishable




Figure 1. Phenotypes of *LeNCED*-overexpressed petunia. The seedlings from non-transformed control, transgenic *rd29A:LeNCED1* lines 8F, 7D, 3D, and *CaMV-35S:LeNCED1-12F* were grown in UC Mix soil in greenhouse. Daily irrigations were provided to maintain optimal non-stress growth conditions. Photographs were taken at different stages of development (4, 7, 10, and 13 weeks after sowing).




Figure 2. Phenotypes of control and *rd 29A:LeNCED1* plants under non-stress and stress conditions. Plants of non-transformed control and transgenic *rd29A:LeNCED1* (8F, 7D, and 3D) lines were grown in UC mix soils in greenhouse for 6 weeks under non-stress conditions (a) and were subjected to drought treatment by withholding irrigation for 7 days (b). Photographs taken at 7 days after drought treatment clearly showed wilting phenotype for the non-transformed control while the *rd29A:LeNCED1* lines showed no apparent negative impact on the growth and development.




Figure 3. Drought stress resistance of *rd29A::LeNCED1* petunia. Plants of non-transformed control and transgenic *rd29A::LeNCED1* line 8F were grown in pots filled with same amounts of UC Mix under non-stress conditions for 5 weeks and were subjected to drought stress by withholding water for 14 days. The control plants started wilting at day 3 and were severely wilted by day 10. In comparison, transgenic plants started to show wilting at day 10. The stressed plants were rehydrated to allow recovery and photographs were taken 10 days after re-watering. The number of surviving plants (out of the total tested) was indicated on the upper right of the image.

(Supplementary Figure 3). Withholding water for 10 days resulted in severe wilting of leaves and flowers in the control plants while the transgenic plants were only slightly wilted, and most of their flowers were still turgid (Supplementary Figure 3). As with the vegetative plants, most of the *rd29A::LeNCED1* plants at the flowering stage recovered if they were re-watered after 15 days of drought stress; the control plants were unable to recover (Supplementary Figure 4). Transgenic plants constitutively expressing *LeNCED1* also completely recovered (Supplementary Figure 4).

Effect on transpiration

Determination of water loss by measuring the pot weight of 5-week-old plants subjected to drought stress demonstrated a significant difference between non-transformed and *rd29A::LeNCED1-8F* plants. For the first day after stopping irrigation, water loss was similar in all plants. Thereafter there was a dramatic difference in water loss between the control and transgenic plants. Water loss from non-transformed control and from transgenic plants was almost linear for the first week of the drought stress period; the rate of water loss in the transgenic plants was half that in the controls (Figure 4). The time to visible wilting in the controls was less than half of that in the transgenic plants.

LeNCED1 expression results in an increase of ABA content

LeNCED1 transcripts could not be detected in non-stressed leaves from the three *rd29A::LeNCED1* transgenic lines (0 h) but were abundant in leaves from the *CaMV-35S::LeNCED1* line (Figure 5a). After 3 days of drought, *LeNCED1* transcript abundance had increased dramatically in leaves from the three *rd29A::LeNCED1* transgenic lines and was similar to that in the *CaMV-35S::LeNCED1* transgenic plants (Figure 5a).

ABA concentrations in non-stressed leaves were low (Figure 5c); the concentration in leaves of the *CaMV-35S::LeNCED1* line was twice that in the other lines. After 3 days of drought stress, ABA concentrations had increased dramatically in all plants, especially in the *rd29A::LeNCED1* line 8F and in the *CaMV-35S::LeNCED1* line 12F (Figure 5c).

Physiological parameters

In well-watered plants, the photosynthesis rate, transpiration rate, stomatal conductance, and stem water potential of control and *rd29A::LeNCED1* plants were very similar (Figure 6). In contrast, these parameters were all significantly lower in plants of the *CaMV-35S::LeNCED1* line. Drought stress reduced these physiological parameters in all plants. Photosynthesis, stomatal conductance, and




Figure 4. Water loss of *rd29A::LeNCED1* petunia under drought stress. Plants of non-transformed control and transgenic *rd29A::LeNCED1-8F* were grown in the pots filled with same amounts of UC mix soils under well-irrigated conditions for 5 weeks and were subjected to drought stress treatment by withholding water over a period of 9 days. The weights of each pot were measured daily in the morning and water loss was calculated. The arrows mark the points when all the plants showed wilting. Each time point is presented as the mean \pm SD ($n = 6$). Different letters denote significant differences at the 0.05 level.

transpiration in the stressed transgenic plants were lower than in the stressed non-transformed controls, but stem water potential of the controls was much lower than that in the transgenic lines (Figure 6).

Effects of drought stress on proline synthesis

Under non-stressed conditions, *P5CS* transcript abundance was low, but detectable in transgenic *rd29A::LeNCED1* and control plants (Figure 7a). Abundance was much higher in the non-stressed *CaMV-35S::LeNCED1* plants. After 3 days of drought stress, *P5CS* transcript abundance had increased substantially in all plants, but much more in the transgenic plants than in the controls (Figure 7b). The proline content of the plants mirrors the transcript abundance (Figure 7c); concentrations were low in non-stressed plants, although somewhat higher in the *CaMV-35S::LeNCED1* plants (Figure 7c). Stressed plants all contained much higher concentrations of proline, but concentrations in the transgenic lines were double that in the controls (Figure 7c).

DISCUSSION

Our data strongly support the hypothesis that overexpression of *LeNCED1* using a stress-specific promoter would enhance drought resistance in petunia without negative effects on plant growth and development under non-stress conditions. Previous researchers overexpressed *NCED* using a constitutive promoter such as *CaMV-35S*. Although they found increased ABA accumulation, faster stomatal closure and enhanced drought resistance in several plants including tomato,³² *Arabidopsis*,^{33,34,20} *Nicotiana plumbaginifolia*,²³ and tobacco,^{22,19} the transgenic plants showed defective growth and development, increased seed dormancy and delayed germination.¹⁹ In the *CaMV-35S::LeNCED1* petunia lines that we generated, similar effects, including a marked photobleaching phenotype in leaves of mature plants, were observed (Supplementary Figure 5).

To avoid these pleiotropic effects, we generated petunia plants overexpressing the *LeNCED1* gene under the control of the stress-responsive *rd29A* promoter. Inducible promoters are a very powerful tool used in plant genetic engineering to regulate the expression




Figure 5. *LeNCED1* expression and ABA content in *LeNCED1*-overexpressed petunia. Plants of non-transformed control, transgenic *rd29A::LeNCED1* lines 8F, 7D, 3D, and *CaMV-35S::LeNCED1-12F* line were grown in UC mix soils in greenhouse for 6 weeks under non-stress conditions and were subjected to drought treatment by withholding irrigation for 3 days. Transcript abundances of the *LeNCED1* were determined with semi-quantitative RT-PCR using intact leaves after 0- and 3-day drought treatment (a, b). Error bars show SE of the means of three biological replicates; different letters denote significant differences using *t*-test at $P > 0.05$. ABA contents were measured using ELISA methods as described in M & M (c). Leaf samples were taken from the non-transformed control, transgenic *rd29A::LeNCED1* and *CaMV-35S::LeNCED1* lines at 0 and 3 days after drought treatment. Data are means \pm SD ($n = 6$). Different letters denote significant differences at the 0.05 level.




Figure 6. Leaf gas exchange and water potential in *LeNCED1*-overexpressed petunia. Plants of non-transformed control, transgenic *rd29A:LeNCED1* lines 8F, 7D, 3D, and *CaMV-35S:LeNCED1-12F* line were grown in UC mix soils in greenhouse for 6 weeks under non-stress conditions and were subjected to drought treatment by withholding irrigation for 3 days. CO₂ assimilation rate (a), leaf conductance (b), transpiration rate (c), and stem water potential (d) were measured using intact leaves. Error bars represent standard deviation of the mean where *n* = 6. Different letters denote significant differences at the 0.05 level.

of genes in a particular tissue or at specific stages of development.³⁵ In normal growth conditions, the transgenic plants were indistinguishable from the control plants (Figure 1), indicating that the transgene was tightly controlled. Even in line 7D, whose stomatal conductance was significantly reduced under non-stress conditions (Figure 6b), suggesting some 'leakiness,' there were only minor negative effects on plant growth and development. Seed germination rate (Supplementary Figure 2), flower longevity (Supplementary Figure 6), and stomatal density (data not shown) of the transgenic plants grown under non-stressed conditions were also similar to those of control plants.

The photobleaching symptom observed in the mature leaves of the *CaMV-35S:LeNCED1* plants might be attributed to reduced production of photoprotective carotenoids³⁶ resulting from diversion of precursors to ABA synthesis. Since there was no photobleaching in the *rd29A:LeNCED1* plants, photobleaching in the constitutive *NCED* plants cannot be explained in this way, and must instead be the result of other correlated effects, such as stomatal closure and limited CO₂ fixation under non-stressed conditions.

One of the more remarkable responses to drought stress of the *rd29A:LeNCED1* transgenic plants was their ability to recover from severe water stress. After 14 days without water, most of the transgenic plants recovered, whereas almost all of the control plants died when re-watered. This difference may simply reflect the reduction in water loss and accompanying physiological stresses in the transgenic plants, but it may also be the result of the greatly increased concentration of proline and perhaps other osmoregulatory compounds in the cells of the stressed transgenic plants. We found that the relatively low concentration of ABA in the leaves of constitutive *CaMV-35S:LeNCED1* plants under non-stressed conditions had about the same effect on gas exchange as the much higher levels in the drought-stressed *rd29A:LeNCED1* transgenic

plants. This suggests that mechanisms other than stomatal closure are key to the improved stress resistance resulting from stress-induced overexpression of *NCED*.

In response to drought stress, many plants reduce their water potential to maintain turgor by accumulating compatible solutes such as proline, mannitol, or trehalose in the cytoplasm.^{37,38} Proline, one of the most commonly reported compatible solutes, is important not only in osmoregulation,³⁹ but also in protecting cells by enhancing the stability of proteins and membranes.^{40,41} The increased transcript abundance of *P5CS*, considered to be the rate limiting enzyme in proline biosynthesis,⁴² was correlated with a significantly higher accumulation of free proline in *rd29A:LeNCED1* transgenic plants during drought stress (Figure 7). Other researchers have shown correlations between endogenous increase⁴³ and exogenous application of ABA and proline concentrations in drought-stressed plants, but the relationship between proline and ABA in stress survival requires further examination.

Our results demonstrated that overexpression of a target gene related to ABA biosynthesis pathway such as *NCED* using the inducible promoter *rd29A* is an effective strategy for improving the drought resistance and avoiding the negative effects resulting from a constitutive promoter. Drought is a complex stress. Drought resistance can vary widely among crops; accurately assessing the drought resistance of different crops and varieties requires extensive field trials over a number of growing seasons in varied environments. Whether the approach described in this study would be widely applicable to other plants or agricultural field crops or the mechanism based on the conditionally enhanced ABA biosynthesis would be useful under field conditions requires further study. In addition, it would be informative to examine whether this approach could enhance water use efficiency and affect biomass production in the future.




Figure 7. *P5CS* expression and proline accumulation in *LeNCED1*-overexpressed petunia. Plants of non-transformed control, transgenic *rd29A:LeNCED1* lines 8F, 7D, 3D, and *CaMV-35S:LeNCED1-12F* line were grown in UC mix soils in greenhouse for 6 weeks under non-stress conditions and were subjected to drought treatment by withholding irrigation for 3 days. Relative expression levels of *P5CS* in petunia leaves at 0 and 3 days after drought treatment were measured by semi-quantitative RT-PCR (a, b). Proline accumulation in the control and *rd29A:LeNCED1* petunia intact leaves exposed to 0 and 3 days under drought (c). Mean SD values of three independent experiments are shown. Different letters denote significant differences at the 0.05 level.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

ACKNOWLEDGMENTS

This work was funded in part by United States Department of Agriculture (USDA) Floriculture Initiative (5306-21000-019-00D and 5306-13210-001-02S) and by the Plant Science Department at the University of California, Davis through the Jastro Scholarship. We thank Linda Donnelly and Lee Ann Richmond for technical assistance. Goldsmith Seeds (Gilroy, CA, USA) generously donated seeds of the Mitchell petunia cultivar.

REFERENCES

- Boyer JS. Plant productivity and environment. *Science* 1982; **218**: 443–448.
- Gurian-Sherman D. *Why genetic engineering is not solving agriculture's drought problem in a thirsty world*. Report. Cambridge, MA; 2012.
- Salinger MJ, Sivakumar MVK, Motha R. Reducing vulnerability of agriculture and forestry to climate variability and change: workshop summary and recommendations. *Climatic Change* May 2005; **70**: 341–362.
- Davies WJ, Zhang JH. Root signals and the regulation of growth and development of plants in drying soil. *Annu Rev Plant Phys* 1991; **42**: 55–76.
- Blatt MR. Cellular signaling and volume control in stomatal movements in plants. *Annu Rev Cell Dev Bi* 2000; **16**: 221–241.
- Malladi A, Burns JK. Communication by plant growth regulators in roots and shoots of horticultural crops. *Hortscience* August 2007; **42**: 1113–1117.
- Dietz KJ, Sauter A, Wichert K, Messdaghi D, Hartung W. Extracellular beta-glucosidase activity in barley involved in the hydrolysis of ABA glucose conjugate in leaves. *J Exp Bot* May 2000; **51**: 937–944.
- Wang Y, Ying JF, Kuzma M, Chalifoux M, Sample A et al. Molecular tailoring of far-nylation for plant drought tolerance and yield protection. *Plant J* August 2005; **43**: 413–424.
- Gagneul D, Ainouche A, Duhaze C, Lugan R, Larher FR et al. A reassessment of the function of the so-called compatible solutes in the halophytic *Plumbaginaceae* *Limonium latifolium*. *Plant Physiol* July 2007; **144**: 1598–1611.
- Ashraf M, Bashir A. Salt stress induced changes in some organic metabolites and ionic relations in nodules and other plant parts of two crop legumes differing in salt tolerance. *Flora* 2003; **198**: 486–498.
- Yamada M, Morishita H, Urano K, Shiozaki N, Yamaguchi-Shinozaki K, Shinozaki K, et al. Effects of free proline accumulation in petunias under drought stress. *J Exp Bot* July 2005; **56**: 1975–1981.
- Xiong LM, Ishitani M, Lee H, Zhu JK. The Arabidopsis *LOS5/ABA3* locus encodes a molybdenum cofactor sulfuryase and modulates cold stress- and osmotic stress-responsive gene expression. *Plant Cell* September 2001; **13**: 2063–2083.
- Roychoudhury A, Paul S, Basu S. Cross-talk between abscisic acid-dependent and abscisic acid-independent pathways during abiotic stress. *Plant Cell Rep* July 2013; **32**: 985–1006.
- Marin E, Nussaume L, Quesada A, Gonneau M, Sotta B et al. Molecular identification of zeaxanthin epoxidase of *Nicotiana glauca*, a gene involved in abscisic acid biosynthesis and corresponding to the ABA locus of *Arabidopsis thaliana*. *Embo J* 15 May 1996; **15**: 2331–2342.
- Schwartz SH, Tan BC, Gage DA, Zeevaart JAD, McCarty DR. Specific oxidative cleavage of carotenoids by VP14 of maize. *Science* 20 Jun 1997; **276**: 1872–1874.
- Schwartz SH, Qin X, Zeevaart JA. Elucidation of the indirect pathway of abscisic acid biosynthesis by mutants, genes, and enzymes. *Plant Physiol* April 2003; **131**: 1591–1601.
- Taylor IB, Sonneveld T, Bugg TDH, Thompson AJ. Regulation and manipulation of the biosynthesis of abscisic acid, including the supply of xanthophyll precursors. *J Plant Growth Regul* December 2005; **24**: 253–273.
- Tan BC, Schwartz SH, Zeevaart JAD, McCarty DR. Genetic control of abscisic acid biosynthesis in maize. *Proc Nat Acad Sci USA* 28 October 1997; **94**: 12235–12240.
- Thompson AJ, Jackson AC, Symonds RC, Mulholland BJ, Dadswell AR et al. Ectopic expression of a tomato 9-cis-epoxycarotenoid dioxygenase gene causes overproduction of abscisic acid. *Plant J* August 2000; **23**: 363–374.
- Wan XR, Li L. Regulation of ABA level and water-stress tolerance of *Arabidopsis* by ectopic expression of a peanut 9-cis-epoxycarotenoid dioxygenase gene. *Biochem Bioph Res Co* 8 September 2006; **347**: 1030–1038.
- Iuchi S, Kobayashi M, Taji T, Naramoto M, Seki M et al. Regulation of drought tolerance by gene manipulation of 9-cis-epoxycarotenoid dioxygenase, a key enzyme in abscisic acid biosynthesis in *Arabidopsis*. *Plant J* 2001; **27**: 325–333.
- Zhang Y, Yang J, Lu S, Cai J, Guo Z. Overexpressing *SgNCE1* in *Tobacco* Increases ABA Level, Antioxidant Enzyme Activities, and Stress Tolerance. *J Plant Growth Regul* 2008; **27**: 151–158.
- Qin XQ, Zeevaart JAD. Overexpression of a 9-cis-epoxycarotenoid dioxygenase gene in *Nicotiana glauca* increases abscisic acid and phaseic acid levels and enhances drought tolerance. *Plant Physiol* February 2002; **128**: 544–551.

- 24 Tung SA, Smeeton R, White CA, Black CR, Taylor IB *et al*. Over-expression of LeNCED1 in tomato (*Solanum lycopersicum* L.) with the *rbc53C* promoter allows recovery of lines that accumulate very high levels of abscisic acid and exhibit severe phenotypes. *Plant Cell Environ* July 2008; **31**: 968–981.
- 25 Yamaguchishinozaki K, Koizumi M, Urao S, Shinozaki K. Molecular-cloning and characterization of 9 Cdnas for genes that are responsive to desiccation in *Arabidopsis-Thaliana* - sequence-analysis of one Cdna clone that encodes a putative transmembrane channel protein. *Plant Cell Physiol* April 1992; **33**: 217–224.
- 26 Murashige T, Skoog F. A revised medium for rapid growth and bio assays with tobacco tissue cultures. *Physiologia plantarum* 1962; **15**: 473–497.
- 27 Chang XX, Donnelly L, Sun DY, Rao JP, Reid MS *et al*. A petunia homeodomain-leucine zipper protein, PhHD-Zip, plays an important role in flower senescence. *Plos One* 14 February 2014; **9**(2): e88320.
- 28 Wang H, Stier G, Lin J, Liu G, Zhang Z *et al*. Transcriptome changes associated with delayed flower senescence on Transgenic Petunia by inducing expression of *etr1-1*, a mutant ethylene receptor. *Plos One* 9 July 2013; **8**(7): e65800.
- 29 Liang YC, Reid MS, Jiang CJ. Controlling plant architecture by manipulation of gibberellic acid signalling in petunia. *Hortic Res* 2014; **1**: 14061.
- 30 Bates LS, Waldren RP, Teare ID. Rapid determination of free proline for water-stress studies. *Plant Soil* 1973; **39**: 205–207.
- 31 McCutchan H, Shackel KA. Stem-water potential as a sensitive indicator of water stress in prune trees (*Prunus domestica* L. cv. French). *Amer Soc Hort Sci* 1992; **117**(4): 607–611.
- 32 Thompson AJ, Andrews J, Mulholland BJ, McKee JM, Hilton HW *et al*. Overproduction of abscisic acid in tomato increases transpiration efficiency and root hydraulic conductivity and influences leaf expansion. *Plant Physiol* April 2007; **143**: 1905–1917.
- 33 Iuchi SK, Kobayashi M, Taji T, Naramoto M, Seki M *et al*. Regulation of drought tolerance by gene manipulation of 9-cis-epoxycarotenoid dioxygenase, a key enzyme in abscisic acid biosynthesis in *Arabidopsis*.-2001.pdf. *Plant J* 2001; **27**: 325–333.
- 34 Lefebvre V, North H, Frey A, Sotta B, Seo M *et al*. Functional analysis of *Arabidopsis* NCED6 and NCED9 genes indicates that ABA synthesized in the endosperm is involved in the induction of seed dormancy. *Plant J* February 2006; **45**: 309–319.
- 35 Dutt M, Dhekney SA, Soriano L, Kandel R, Grosser JW. Temporal and spatial control of gene expression in horticultural crops. *Hortic Res* 2014 09/ 24/online; **1**: 14047: doi:14010.11038.
- 36 Demmigadams B, Adams WW. Photoprotection and other responses of plants to high light stress. *Annu Rev Plant Phys* 1992; **43**: 599–626.
- 37 Csonka LN. Physiological and genetic responses of bacteria to osmotic-stress. *Microbiol Rev* March 1989; **53**: 121–147.
- 38 Chen THH, Murata N. Enhancement of tolerance of abiotic stress by metabolic engineering of betaines and other compatible solutes. *Curr Opin Plant Biol* June 2002; **5**: 250–257.
- 39 Samaras Y, Bressan RA, Csonka LN, Garcia-Rios MG, Paino D'urzo MP *et al*. Proline accumulation during drought and salinity. In: Smirnoff N, ed. Environment and plant metabolism: flexibility and accumulation. Oxford, UK: BIOS Scientific Publishers 1995; p161–187.
- 40 Hare PD, Cress WA. Metabolic implications of stress-induced proline accumulation in plants. *Plant Growth Regul* February 1997; **21**: 79–102.
- 41 Kishor PBK, Sangam S, Amrutha RN, Laxmi PS, Naidu KR *et al*. Regulation of proline biosynthesis, degradation, uptake and transport in higher plants: its implications in plant growth and abiotic stress tolerance. *Curr Sci India* 10 February 2005; **88**: 424–438.
- 42 Hu CAA, Delauney AJ, Verma DPS. A bifunctional enzyme (delta-1-pyrroline-5-carboxylate synthetase) catalyzes the 1st 2 steps in proline biosynthesis in plants. *Proc Nat Acad Sci USA* 1 October 1992; **89**: 9354–9358.
- 43 Zhou L, Xu H, Mischke S, Meinhardt LW, Zhang D *et al*. Exogenous abscisic acid significantly affects proteome in tea plant (*Camellia sinensis*) exposed to drought stress. *Hortic Res* June 2014; **1**: 14029.



This work is licensed under a Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by-nc-sa/3.0/>

Supplemental Information for this article can be found on the *Horticulture Research* website (<http://www.nature.com/hortres>).