

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

THE MOLECULAR AND ELECTRONIC STRUCTURE OF PHOSPHONIUM CYCLOPROPYLIDE  
H<sub>3</sub>P=C(CH<sub>2</sub>)<sub>2</sub>: A THEORETICAL STUDY

### Permalink

<https://escholarship.org/uc/item/23g3f7g5>

### Author

Vincent, M.A.

### Publication Date

1982-12-01



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

## Materials & Molecular Research Division

RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

MAY 17 1983

LIBRARY AND  
DOCUMENTS SECTION

Submitted to the Journal of the American Chemical  
Society


THE MOLECULAR AND ELECTRONIC STRUCTURE OF  
PHOSPHONIUM CYCLOPROPYLIDE  $H_3P=C(CH_2)_2$ :  
A THEORETICAL STUDY

Mark A. Vincent, Henry F. Schaefer III,  
Annette Schier, and Hubert Schmidbauer

December 1982

### TWO-WEEK LOAN COPY

*This is a Library Circulating Copy  
which may be borrowed for two weeks.  
For a personal retention copy, call  
Tech. Info. Division, Ext. 6782.*



LBL-15399  
c-2

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

The Molecular and Electronic Structure of Phosponium

Cyclopropylide  $\text{H}_3\text{P}=\text{C}(\text{CH}_2)_2$ : A Theoretical Study

Mark A. Vincent and Henry F. Schaefer III

Department of Chemistry and Lawrence Berkeley Laboratory  
University of California  
Berkeley, California, 94720 U.S.A.

and

Annette Schier and Hubert Schmidbaur

Anorganisch-chemisches Institut  
Technische Universität München  
8046 Garching, West Germany

Abstract


Theoretical studies have been carried out for a phosponium cyclopropylide model  $\text{H}_3\text{P}=\text{C}(\text{CH}_2)_2$ . The molecule was shown to have a ground state equilibrium geometry (Structure I) with a pyramidal carbanion center, the details being in satisfactory agreement with experimental findings for the triphenylphosphine cyclopropylide homologue. One of the  $\text{PH}_3$  hydrogen atoms is affected by the carbanion (partial) negative charge, and its position shows substantial distortion away from the idealized tetrahedral phosphorus geometry. In the energy profile for the carbanion inversion the stationary point of maximum energy (two imaginary vibrational frequencies) occurs not for the planar carbanion configuration, but well beyond in a situation with a dihedral angle  $\text{HOPClC2}$  of  $123^\circ$  instead of  $90^\circ$ . The rotated

configuration (a genuine transition state) is virtually the eclipsed analogue of Structure I. The barriers to inversion and rotation are predicted to be 6.3 and 5.8 kcal, respectively. Calculated bond distances, bond angles, dihedral angles, total energy values and total electron populations on atoms are given for all three conformations. Atomic orbital components and energies of frontier orbitals of the ground state are also discussed.

### Introduction

Chemical bonding and the molecular structure of phosphorus ylides are classical problems of longstanding interest, which is reflected in an ever increasing number of publications on pertinent experimental or theoretical investigations. In the last two decades about 35 molecular structures of ylides have been determined by X-ray<sup>1</sup> or electron diffraction.<sup>2</sup> The ylidic electronic structure has been probed by dipole moment studies,<sup>3</sup> PES measurements,<sup>4</sup> ESR experiments,<sup>5</sup> multi-nuclei NMR,<sup>6</sup> vibrational<sup>7</sup> and UV<sup>8</sup> spectroscopy, and mass spectrometry.<sup>9</sup> Following the rapid advancement of theoretical chemistry, computational methods at various levels of sophistication were applied to simple model systems,<sup>10</sup> particularly of the type  $R_3P=CH_2$ <sup>4a,4b,6h,8a,11</sup> and led to a number of mostly complimentary descriptions of the bonding situation in ylides of phosphorus, be they real or hypothetical (like  $H_3P=CH_2$ ).

It appears to be generally accepted that phosphorus ylides ("phosphonium ylides", "alkylene phosphoranes") should be referred to as molecules (or ions) in which a phosphonium center of nearly tetrahedral geometry is directly attached to a carbanionic center, as implied by one of Georg Wittig's original canonical formulae



The nature of the electronic interaction between phosphorus and the ylidic carbon, however, is already a point of controversy, not the least because of uncertainties about the actual geometry of the carbanion, and hence its hybridisation. The significance of d-orbital

participation is closely associated with this problem, since a planar  $sp^2$  geometry for the carbanion is believed to optimize ( $p \rightarrow d$ )  $\pi$  interactions. This argument is questionable, since the short "P=C" distances found in ylides can be accounted for simply by a polarisation effect of the d-orbitals and even by considering solely the heteropolar (electrostatic) forces without any d-orbital contributions.<sup>12</sup>

For  $H_3P=CH_2$  a trigonally planar carbanion geometry has been predicted by both semi-empirical<sup>10a</sup> and ab initio calculations,<sup>10</sup> but deviations from planarity are not associated with a great increase in energy. Rotation about the ylidic bond is virtually unrestricted, with activation barriers smaller than 1 kcal/mole. Structural studies seemed to bear out these results, as all molecular structures determined until recently showed indeed planar carbanion geometry.<sup>1,2</sup>

In this situation it was of great interest that the X-ray diffraction analysis of the triphenylphosphoniumcyclopropylide provided unambiguous proof for a pyramidal carbanion geometry.<sup>13</sup> (Figure 1). In fact, the angle between the ylidic bond and the three-membered ring is close to  $60^\circ$  and thus in keeping with the standard ligand arrangements of  $\sigma$ -bonded substituents of cyclopropane ( $60^\circ$ ).

This result prompted a reinvestigation of the molecular orbital (MO) description of the prototype  $H_3P=C(CH_2)_2$ , which had been considered previously in two other studies, with a planar carbanion geometry at the Extended Hückel MO level<sup>11a</sup> and with both planar and pyramidal geometries in CNDO/2 and MINDO/3 type calculations.<sup>6h</sup>

A deviation from planarity was derived from these studies and from NMR correlations. The latter were subsequently found not to be meaningful being based on erroneous data.<sup>13</sup>

A temperature dependent NMR investigation of  $(C_6H_5)_3P=C(CH_2)_2$  has shown that inversion at the ylidic carbon is rapid on the NMR time scale even at  $-110^\circ C$ , indicating a very low activation energy for the inversion process in solution. Apart from a calculation of the optimized ground state geometry of the  $H_3P=C(CH_2)_2$  model, it was therefore one of the goals of the work presented in this report to determine the energy differences relative to the expected transition state and the invertsomer obtained in the carbanion inversion.

#### Theoretical Approach

Unfortunately, previous experience has shown that the reliable description of carbanions requires large basis sets, including diffuse s- and p- functions. This is especially true for the study of inversion barriers.<sup>14</sup> However, the presence of the positively charged phosphorus next to the carbon anion in ylides means that the charge buildup will not be one full electron, but considerably less. Thus a set of orbitals similar to those used to describe inversion in a molecule like ammonia should be suitable. Carlsen, et al.,<sup>15</sup> have found that the smallest set of orbitals to give a reasonable barrier for ammonia is a double zeta plus polarization set. Hence a Dunning double zeta basis<sup>16a</sup> was used with a set of six d-functions (exponent 0.75). Due to the size of the phosphonium cyclopropylide it was inadvisable to use such a large basis set on the non-ylidic carbons. Thus only a double zeta basis was used for them.<sup>16a</sup> This inequality

between the carbons is not expected to have any major effects. However, research by Dixon and coworkers<sup>17</sup> suggests that the omission of d functions on the other two carbon atoms may result in some overestimation of the inversion barrier.

Test calculations with and without one set of more diffuse p functions on the ylidic carbon were also carried out. However the energy lowering relative to the straight double zeta basis was only 0.00067 hartrees, and such functions were thereafter omitted. In contrast the energy lowering due to the addition of ylidic carbon d functions is substantial, 0.02487 hartrees.

A double zeta<sup>16b</sup> plus polarization (exponent 0.6) basis set was employed to describe phosphorus, as Lischka has shown that such a basis set is necessary to describe the quantitative features of phosphorus bonding in the related model ylide  $H_3PCH_2$ .<sup>10d</sup> For hydrogen it was decided, because of the size of the ylide, to use only a Dunning double zeta basis,<sup>16</sup> scaled by a factor of 1.2. The complete technical designation of this contracted gaussian basis set is thus P(11s 7p 1d/6s 4p 1d), C(9s 5p 1d/4s 2p 1d), H(4s/2s). Determination of the geometrical structures was carried out via analytic gradient techniques, which guarantee stationarity with respect to variations of the coordinates of the system.<sup>18</sup> All geometries were determined assuming a plane of symmetry for the molecule. This plane contained the unique hydrogen bonded to phosphorus, the phosphorus atom itself and the ylidic carbon. Vibrational analyses later proved this assumption to be valid.

Pictorial Results

The final results for the ground state equilibrium geometry of the molecule (Structure I), and the charge distribution (as indicated by the Mulliken population analysis) are summarized in Tables 1 and 2. These Tables also contain the corresponding data for the conformation, which represents the energy maximum in the profile of the carbanion inversion process, referred to as Structure III, and finally the configuration of the stationary point (Structure II) reached as the cyclopropyl ring folds to the opposite side of the original ground state equilibrium geometry. Note that the latter conformation is not geometrically identical with the ground state, because the cyclopropyl ring is in a different rotatory position relative to the  $\text{PH}_3$  group: Structure I is a staggered, Structure II an eclipsed conformation. The molecular

geometries are represented by the cartesian coordinates of the 11 atoms and the meaningful atomic distances, bond angles and dihedral angles. The perspective drawings of the three configurations in Figures 2a-2c are complementary to the three projections parallel to the P-C<sub>ylide</sub> bond shown in Figures 3a-3c.

From the latter figures the different rotamers are again immediately obvious. Structure II would require a 60° rotation of the PH<sub>3</sub> group around the P-C<sub>ylide</sub> axis to relax to the configuration of Structure I. A compilation of the energies of the individual molecular orbitals and their individual atomic orbital components are available from the authors on request. Some of these data are used in the discussion below where appropriate. Energy gradients for specific stretching and bonding movements were also calculated. They are useful to characterize the various energy profiles associated with distortions of the individual geometries.

#### The Ground State Equilibrium Configuration (Structure I)

The calculations have clearly shown that phosphonium cyclopropylide indeed has an equilibrium geometry with a carbanion center C1 of a pronounced pyramidal array of the neighboring atoms (P, C2, C3). With valence angles PC1C2 = PC1C3 = 127.5° and C2C1C3 = 61.1° the orientation of the ylidic bond relative to the cyclopropyl plane is 44.9°. These values are in reasonable agreement with the experimental data for the triphenyl (!) homologue (117.1°, 117.8° and 58.8°, respectively).<sup>13</sup> The agreement is even better for the PC1 bond length of 168.3 pm in the model as compared to 169.6 pm in the triphenyl homologue, and for the

ring distances C1C2 = C1C3 = 150.4 pm, C2C3 = 152.8 pm [152.5 and 149.9 pm in  $(C_6H_5)_3P=C(CH_2)_2$ ].

In the triphenylphosphonium group of the cyclopropylide, one phenyl group is unique in that it forms valence angles at phosphorus that differ markedly from the remaining bond angles at phosphorus (see C3PC4 in Table 1). This situation is met again for the unique hydrogen HO in the model with its longer P-HO distance (141.3 pm as compared to 138.1 pm for P-H10 and P-H11) and its angle HOPC1 = 129.0°. Obviously this hydrogen is interacting most strongly with the carbanion lone pair, and the resulting repulsive forces lead to a substantial distortion away from the idealized tetrahedral phosphorus geometry.

The Mulliken population analysis (Table 2) ascribes a significant negative charge to HO and, of course, to C1 (-0.05 and -0.51 electrons, respectively), suggesting again repulsive Coulomb forces. The remaining Mulliken population analysis data characterize phosphorus as an onium center (+0.54), as expected. Cyclopropyl hydrogens are also positive, cyclopropyl carbons negative. The dipole moment of the ground state molecule is  $\mu = 3.34$  Debye. No experimental value is available for comparison, but estimations based on data for non-cyclic ylides<sup>3</sup> could lead to a value of around 4 Debye.

The molecule has 20 occupied MO's with a frontier HOMO of ionization potential  $IP(20) = -0.2755$  a.u., according to Koopman's theorem. Surprisingly, this orbital is largely a 2p orbital (of C1) with an out-of-phase contribution from HO. The second highest occupied MO is a ring-bonding orbital comprised mainly of p-functions on C1, C2 and C3 [ $IP(19) = -0.4035$ ]. The population analysis for the highest three occupied orbitals is shown in Table 3. The molecular orbital picture

does not impose a hybridization upon the ylid carbon, C1. Nevertheless, one can deduce from the orbitals an approximate hybridization midway between  $sp^2$  and  $sp^3$ .

#### The Pathway for Inversion (Structure III)

The energy difference between Structure I and Structure III leads to a calculated barrier to carbanion inversion of 6.3 kcal/mole. Note that Structure III has two imaginary vibrational frequencies, i.e., it is a maximum with respect to both inversion and rotation. It is very important to note that this energy maximum does not correspond to the situation where the ylidic carbon atom C1 is in a planar configuration (PC1C2C3). The conformational constraints caused by the relative orientation of the  $PH_3$  group apparently shift the maximum towards larger dihedral angles HO-P-C1-C2. For the almost planar situation (HO-P-C1-C2 =  $92.5^\circ$ ) the energy is calculated to be -458.21798 a.u., or 1.2 kcal lower than for Structure III (-458.21613 a.u.). In the projectional drawing in Figure 2c this means that the C2-C3 horizontal is well above the P-C1 axis, whereas P-C1 would point towards the center of C2-C3 for the fully planar configuration at C1.

In Structure III, characterized by a dihedral angle HO-P-C1-C2 =  $123.2^\circ$  with its inverted PC1C2C3 pyramid, the P-C1 ylide bond distance has been shortened to 163.4 pm. This value is extremely low and in fact lower than any of the experimental values available to date. It may indicate, however, that "P=C bonding" is indeed an optimum at a virtually planar ylidic carbon, were it not that other factors destabilize such a configuration, as it is evidently the case for  $H_3PC(CH_2)_2$ . The angle between the PC1 bond and the cyclopropane ring is  $-23.4^\circ$  for Structure III. Note however that this initially surprising

result is consistent with Hammond's postulate. That is, for the exothermic (by 5.8 kcal) inversion of II to I, the stationary point structurally resembles Structure II. Most other of its structural and charge distribution data follow more or less expected trends.

#### The Transition State (Structure II)

As pointed out above, the transition state for rotation is fixed in an eclipsed conformation regarding the position of the  $\text{PH}_3$  group relative to the cyclopropane moiety (Figure 2b). It is therefore as expected that the energy of this conformation is higher (by 5.8 kcal) than that of the equilibrium geometry (Structure I) with its relaxed staggered conformation. The barrier is of course much greater than that predicted ( $\sim 1$  kcal) previously<sup>10</sup> for the model ylid  $\text{H}_3\text{PCH}_2$ . Calculated bond lengths are similar to those already found for Structure I. The dihedral angle  $\text{HO-P-Cl-C2} = 141.2^\circ$  is about complementary to the  $39.8^\circ$  value for Structure I, showing that the dihedral angle is indeed a reasonable indicator for the state of the inversion process. Accordingly, the angle of the ylidic P-Cl bond relative to the cyclopropane ring is  $-45.6^\circ$ , very similar to the arrangement in Structure I ( $+44.9^\circ$ ).

The three hydrogen atoms attached to phosphorus show very similar P-H distances, HPH or ClPH angles, and net atomic charges (from Mulliken population analysis), and there is accordingly no evidence for a unique bonding situation for HO as compared to Structure I. No doubt this finding is again the consequence of the eclipsed conformation. The cyclopropane ring is a nearly equilateral triangle in all three stationary point structures (I, II, and III). The Mulliken population analysis also points to a roughly similar charge distribution for the cyclopropane part of the models.

### Vibrational Analyses

In order to precisely ascertain the nature of the stationary point geometries reported above, it was necessary to carry out vibrational analyses of Structures I and III. Structure I was found to be a genuine equilibrium geometry, that is, all vibrational frequencies are real. Structure III, lying 6.3 kcal higher in energy, was found to have two imaginary vibrational frequencies. This means that Structure III represents a maximum on the potential energy hypersurface with respect to both inversion and rotation. It follows that Structure II, lying 5.8 kcal above the equilibrium geometry, is the true transition state.

Predicted vibrational frequencies are given in Table 4. However it should be noted that frequencies predicted at this level of theory are typically  $\sim 10-12\%$  higher than the observed fundamentals.<sup>22</sup> The assignment of the phosphorous-carbon stretching frequency is not entirely unambiguous, but the normal modes at  $1428\text{ cm}^{-1}$  (equilibrium) and  $1550\text{ cm}^{-1}$  (Structure III, two imaginary vibrational frequencies) seem to best fit this description.

### Discussion

This theoretical study is consistent with the experimental finding<sup>13</sup> that ylidic carbon atoms can adopt a pyramidal ground state configuration at least in the special situation of the cyclopropylides, where drastic ring strain and a unique state of bonding have to be assumed. The predicted barrier to rotation about the P-C bond is 5.8 kcal, in harmony with the experimental finding that this barrier is  $\leq 8$  kcal. The results parallel previous reports on the unusual configurational stability of aziridines as compared to amines,<sup>19</sup> both of which are isoelectronic with the ylidic carbanionic fragments. Spectroscopic and mechanistic data for phosphonium<sup>13</sup> and sulfonium<sup>20</sup> cyclopropylides had also indicated a preference of these systems for a pyramidal geometry.

There are other implications associated with the details of the theoretical predictions, however, which should be considered. As judged from the equilibrium bond distance P-Cl the ylide bond originating from a pyramidal carbon appears not to be very different from bonds observed with planar carbon atoms. A very flat energy profile for the carbanion inversion is thus suggested and in fact observed both in the calculations<sup>10</sup> and experimentally.<sup>13a</sup> Also, conformational effects are perhaps often underestimated in the discussion of ylidic bonding. The relatively short P-Cl contact and the partial negative charge at Cl clearly can cause distortions at the phosphane moiety, as also detected experimentally in a study of hindered rotation in ylides and their complexes.<sup>21</sup> The above examples are illustrative of the importance of an inversion process associated with a rotational movement to reach again a true energy minimum state.

Unfortunately, the present unavailability of the model compound  $\text{H}_3\text{P}=\text{C}(\text{CH}_2)_2$  precludes further crosschecking of the calculated data by experiments. Photoelectron spectroscopy and dipole moment studies would be the obvious means of control. Nevertheless data from simple homologues may well be sufficient to probe some of the expected or unexpected details.

#### Acknowledgments

This research was supported by Fonds der Chemischen Industrie, Frankfurt, Main and by Deutsche Forschungsgemeinschaft, Bonn-Bad Godesberg. A travel grant to A.S., by Proctor & Gamble, European Office, Brussels is gratefully acknowledged. Dr. D. Neugebauer is thanked for establishing the computer drawings presented in the Figures. MAV and HFS were supported by the Director, Office of Energy Research, Office of Basic Energy Sciences, Chemical Sciences Division of the U.S. Department of Energy under Contract Number DE-AC03-76SF00098. The calculations were carried out on a Harris H800 minicomputer supported by the National Science Foundation, Grant CHE-8009320.

References

- (1) (a) Mak, T.C.W., Trotter, J., Acta Cryst. 1965, 18, 81-88.  
(b) Stephens, F.S., J. Chem. Soc. 1965, 5640-5650.  
(c) Stephens, F.S., J. Chem. Soc. 1965, 5658-5678.  
(d) Wheatley, P.J., J. Chem. Soc. 1965, 5785-5800.  
(e) Speziale, A.J., Ratts, K.W., J. Am. Chem. Soc. 1965, 87, 5603-5606. (f) Daly, J.J., Wheatley, P.J., J. Chem. Soc. A, 1966 1703-1706. (g) Daly, J.J., J. Chem. Soc. A, 1967, 1913-1917.  
(h) Chioccola, G., Daly, J.J., J. Chem. Soc. A, 1968, 568-577.  
(i) Bart, J.C.J., J. Chem. Soc. B, 1969, 350-365. (j) Cameron, T.S., Prout, C.K., J. Chem. Soc. C, 1969, 2292-2295.  
(k) Bailey, A.S., Cameron, T.S., Evans-Peach, J.M., Prout, C.K., J. Chem. Soc. C, 1969, 2295-2298. (l) Ross, F.K., Hamilton, W.C., Ramirez, F., Acta Cryst. 1971, 27B, 2331-2334.  
(m) Kennard, O., Motherwell, W.D.S., Coppola, J.C., J. Chem. Soc. C, 1971, 2461-2464. (n) Buckle, J., Harrison, P.G., King, T.J., Richards, J.A., J. Chem. Soc. Chem. Comm. 1972, 1104-1105. (o) Vincent, A.T., Wheatley, P.J., J. Chem. Soc. Dalton Trans. 1972, 617-622. (p) Ross, F.K., Manojlovic-Muir, L., Hamilton, W.C., Ramirez, F., Pilot, J.F., J. Am. Chem. Soc. 1972, 94, 8734-8738. (qu) Gilardi, R.D., Karle, I.L., Acta Cryst. 1972, 28B, 3420-3425. (r) Goldberg, S.Z., Duesler, E.N., Raymond, K.N., Inorg. Chem. 1972, 11, 1397-1401. (s) Goldberg, S.Z., Raymond, K.N., Inorg. Chem. 1973, 12, 2923-2927.  
(t) Dreissig, W., Hecht, H.J., Plieth, K., Z. Kristallogr. 1973,

132-145. (u) Howells, M.A., Howells, R.D., Baenziger, N.C.,  
Burton, D.J., J. Am. Chem. Soc. 1973, 95, 5366-5370. (v) Ammon,  
H.L., Wheeler, G.H., Watts, P.L., J. Am. Chem. Soc. 1973, 95,  
6158-6163. (w) Lingner, U., Burzlaff, H., Acta Cryst. 1974  
30B, 1715-1722. (x) Burzlaff, H., Voll, U., Bestmann, H.J.,  
Chem. Ber. 1974, 107, 1949-1956. (y) Cameron, A.F., Duncanson,  
F.D., Freer, A.A., Armstrong, V.W., Ramage, R., J. Chem. Soc.  
Perkin Trans. II 1975, 1030-1036. (z) Carrol, P.J., Titus, D.D.,  
J. Chem. Soc. Dalton Trans. 1977, 824-829. (aa) Burzlaff, H.,  
Wilhelm, E., Bestmann, H.J., Chem. Ber. 1977, 110, 3168-3171.  
(bb) Hardy, G.E., Zink, J.I., Kaska, W.C., Baldwin, J.C.,  
J. Am. Chem. Soc. 1978, 100, 8001-8002. (cc) Karsch, H.H.,  
Zimmer-Gasser, B., Neugebauer, D., Schubert, U., Angew. Chem.  
1979, 91, 519-520. Angew. Chem. Int. Ed. Engl. 1979, 18, 484-485.  
Z. Naturforsch. 1979, 34b, 1267-1269. (dd) Schmidbaur, H.,  
Haßlberger, G., Deschler, U., Schubert, U., Kappenstein, C.,  
Frank, A., Angew. Chem. 1979, 91, 437-438. Angew. Chem. Int. Ed.  
Engl. 1979, 18, 408-409. (ee) Schmidbaur, H., Costa, T., Milewski-  
Mahrla, B., Schubert U., Angew. Chem. 1980, 92, 557-558. Angew.  
Chem. Int. Ed. Engl. 1980, 19, 555-556. (ff) Baldwin, J.C.,  
Keder, N.L., Strouse, C.E., Kaska, W.C., Z. Naturforsch. 1980,  
35b, 1289-1297. (gg) Schubert, U., Kappenstein, C., Milewski-  
Mahrla, B., Schmidbaur, H., Chem. Ber. 1981, 114, 3070.  
(hh) Weiss, J., Nuber, B., Z. Anorg. Allg. Chem. 1981, 473,  
101-106. (ii) Cramer, R.E., Maynard, R.B., Paw, J.C., Gilje, J.W.,  
J. Am. Chem. Soc. 1981, 103, 3589-3591.

- (2) (a) Ebsworth, E.A.V., Frazer, T.E., Rankin, D.W.H., Chem. Ber. 1977, 110, 3494-3500. (b) Ebsworth, E.A.V., Frazer, T.E., Rankin, D.W.H., Gasser, O., Schmidbaur, H., Chem. Ber. 1977, 110, 3508-3515. (c) Ebsworth, E.A.V., Rankin, D.W.H., Zimmer-Gasser, B., Schmidbaur, H., Chem. Ber. 1980, 113, 1637-1645.
- (3) (a) Ramirez, F., Dershowitz, S., J. Org. Chem. 1957, 22, 41-45. (b) Ramirez, F., Levy, S., J. Am. Chem. Soc. 1957, 79, 67-69. (c) Johnson, A.W., J. Org. Chem. 1959, 24, 282-284. (d) Lumbroso, H., Pigenet, C., Arcoria, A., Scarlata, G., Bull. Soc. Chim. Fr. 1971, 38, 3838-3846. (e) Lumbroso, H., Bertin, D.M., Froyen, P., Bull. Soc. Chim. Fr. 1974, 41, 819-822 (f) Boldesbul, I.E., Loktionova, R.A., Repina, L.A., Lysenko, V.P., Zh. Obshch. Khim. 1977, 47, 2735-2740. Engl. Trans 1977, 47, 2490-2494. (g) Lumbroso, H., Curé, J., Bestmann, H.J., J. Organomet. Chem. 1978, 161, 347-360.
- (4) (a) Ostoja-Starzewski, K. A., tom Dieck, H., Bock, H., J. Organomet. Chem. 1974, 65, 311-325. (b) Ostoja-Starzewski, K. A., Bock, H., J. Am. Chem. Soc. 1976, 98, 8486-8494. (c) Perry, W.B., Schaaf, T.F., Jolly, W.L., J. Am. Chem. Soc. 1975, 97, 4899-4905.

- (5) (a) Lucken, E.A.C., Mazeline, C., J. Chem. Soc. A 1966, 1074-1077. *ibid.* A, 1967, 439-442. (b) Schipper, P., Jansen, E.H.J., Buck, H.M, in Crayson, E.J. and M. (Ed.), "Topics in Phosphorus Chemistry", Vol. 9, Wiley-Interscience, New York 1977, S. 419 ff.
- (6) (a) Schmidbaur, H., Tronich, W., *Angew. Chem.* 1967, 79, 412-413. *Angew. Chem. Int. Ed. Engl.* 1967, 6, 448-449. (b) Schmidbaur, H., Tronich, W., *Chem. Ber.* 1968, 595-603 and 604-610. (c) Grim, S.O., McFarlane, W., Marks, T.J., *Chem. Comm.* 1967, 1191-1192. (d) Schmidbaur, H., Buchner, W., Scheutzow, D., *Chem. Ber.* 1975, 106, 1251-1255. (e) Hildenbrand, K., Dreeskamp, H., *Z. Naturforsch.* 1973, 28b, 226-228. (f) Gray, G.A., *J. Am. Chem. Soc.*, 1973, 95, 5092-5094 and 7736-7742. (g) Schmidbaur, H., Richter, W., Wolf, W., Köhler, F.H., *Chem. Ber.* 1975, 108, 2649-2655. (h) Albright, T.A., Gordon, M.D., Freeman, W.J., Schweizer, E.E., *J. Am. Chem. Soc.* 1976, 98, 6249-6252. (i) Köster, R., Simić, D., Grassberger, M.A., *Justus Liebigs Ann. Chem.* 1970, 739, 211-219.

- (7) (a) Lüttke, W., Wilhelm, K., *Angew. Chem.* 1965, 77, 867.  
(b) Sawodny, W., *Z. Anorg. Allg. Chem.* 1969, 368, 284-292.  
(c) Thomas, L.C., "Interpretation of the Infrared Spectra of Organophosphorus Compounds", Heyden, New York, 1974, 87.
- (8) (a) Iwata, K., Yoneda, S., Yoshida, Z., *J. Am. Chem. Soc.* 1971, 93, 6745-6750. (b) Fischer, H., Fischer, H., *Chem. Ber.* 1966, 99, 658-669. (c) Grim, S.O., Ambrus, J.H., *J. Org. Chem.* 1968, 33, 2993-2994. (d) Zhmurova, I.N., Yurchenko, R.I., Tukkar, A.A., Yurchenko, V.G., Voitsekhovskaya, O.M., *Zh. Obshch. Khim.* 1979, 49, 2401-2405, *Engl. Trans* 1979, 49, 2119-2124.
- (9) (a) Tökés, L., Jones, G.H., *Org. Mass. Spektrom.* 1975, 10, 241-249. (b) Alexander, R.G., Bigley, D.B., Todd, J.F.J., *J. Chem. Soc. Chem. Comm.* 1972, 553-554.
- (10) (a) Hoffmann, R., Boyd, D.B., Goldberg, S.Z., *J. Am. Chem. Soc.* 1970, 92, 3929-3936. (b) Absar, I., Van Wazer, J.R., *J. Am. Chem. Soc.* 1972, 94, 2382-2387. (c) Whangbo, M.-H., Wolfe, S., *Can. J. Chem.* 1976, 54, 949-962. (d) Lischka, H., *J. Am. Chem. Soc.* 1977, 99, 353-360. (e) Bernardi, F., Schlegel, H.B., Whangbo, M.-H., Wolfe, S., *J. Am. Chem. Soc.* 1977, 99, 5633-5636. (f) Strich, A., *Nouv. J. Chim.* 1979, 3, 105-108. (g) Trinquier, G., Malrieu, J.P., *J. Am. Chem.*

- Soc. 1979, 101, 7169-7172. (h) Eades, R.A., Gassman, P.G., Dixon, D.A., J. Am. Chem. Soc. 1981, 103, 1066-1068.
- (i) Mitchell, D.J., Wolfe, S., Schlegel, H.B., Can. J. Chem. 59, 3280 (1981).
- (11) (a) Boyd, D.B., Hoffmann, R., J. Am. Chem. Soc. 1971, 93 1064-1066. (b) Dijk, J.M.F., Buck, H.M., Recu. Trav. Chim. Pays-Bas 1974, 93, 155-156. (c) Klabuhn, B., Tetrahedron 1974, 30, 2327-2330. (d) Ostoja-Starzewski, K.-H.A., Bock, H., tom Dieck, H., Angew. Chem. 1975, 87, 197-198., Angew. Chem. Int. Ed. Engl. 1975, 14, 173. (e) Ostoja-Starzewski, K.-H.A., Richter, W., Schmidbaur, H., Chem. Ber. 1976, 109, 473-481. (f) Albright, T.A., Hofmann, P., Rossi, A.R., Z. Naturforsch. 1980, 35b, 343-351.
- (12) Whangbo, M.-H., Wolfe, S., Bernardi, F., Can. J. Chem. 1975, 53, 3040-3043.
- (13) (a) Schmidbaur, H., Schier, A., Milewski-Mahrla, B., Schubert, U., Chem. Ber. 1982, 115, 722-731. (b) Schmidbaur, H., Schier, A., Chem. Ber. 1981, 114, 3385-3394.
- (14) Dykstra, C.E., Hereld, M., Lucchese, R.R., Schaefer, H.F., Meyer, W., J. Chem. Phys. 1977, 4071-4075.
- (15) Carlsen, N.R., Radom, L., Riggs, N.V., Rodwell, W.R., J. Am. Chem. Soc. 1979, 101, 2233-2234.

- (16). (a) Dunning, T. H., J. Chem. Phys. 1970, 53, 2823, (b) Dunning, T. H., and Hay P. J., in Schaefer, H. F. (Ed.), "Modern Theoretical Chemistry", Vol. 3, Plenum, New York 1977, 1ff.
- (17). Eades, R. A., Weil, D. A., Dixon, D. A. and Douglass, C. H., J. Phys. Chem. 85, 976 (1981).
- (18). Pulay, P., Mol. Phys. 1969, 17, 197-204.
- (19). (a) Lehn, J. M., Fortsch. Chem. Forsch. 1970, 15, 311-377.  
(b) Kessler, H., Angew. Chem. 1970, 82, 237-253. Angew. Chem. Int. Ed. Engl. 1970, 9, 219-235. (c) Rauk, A., Allen, L.C., Mislou, K., Angew. Chem. 1970, 82, 453-468. Angew. Chem. Int. Ed. Engl. 1970, 9, 400-415. (d) Carter, R. E., Drakenberg, T., Bergmann, N.-A., J. Am. Chem. Soc. 1975, 97, 6990-6996.
- (20). Ratajczak, A., Anet, F.A.L., Cram, D.J., J. Am. Chem. Soc. 1967, 89, 2072-2077.
- (21). (a) Schmidbaur, H., Müller, G., Blaschke, G., Chem. Ber. 1980, 113, 1480-1485. (b) Schmidbaur, H., Müller, G., Milewski-Mahrla, B., Schubert, U., Chem. Ber. 1980, 113, 2575-2578.
- (22). Yamaguchi, Y., Schaefer, H. F., J. Chem. Phys. 1980, 73, 2310.

Figure Captions

1. Molecular structure of triphenylphosphonium cyclopropylide as determined by X-ray diffraction.<sup>13</sup>
2. Perspective drawing of the phosphonium cyclopropylide model with molecular dimensions calculated for
  - (a) the Ground State (Structure I)
  - (b) the Transition State for Rotation (Structure II).
  - (c) Structure III, a stationary point having two imaginary vibrational frequencies. (For details see text and Figures 3a-c.)
3. Projection of the phosphonium cyclopropylide model along the P-Cl axis ("ylide bond") with the molecular dimensions calculated for
  - (a) the Ground State (Structure I)
  - (b) the Transition State for Rotation (Structure II).
  - (c) Structure III, a stationary point having two imaginary vibrational frequencies. (For details see text and Figures 2a-c.)

Table 1. Calculated bond distances (in nm), bond angles (in degrees) and dihedral angles in the phosphonium cyclopropylide model as determined for the equilibrium ground state (Structure I), the stationary point for carbanion inversion (Structure III), and the transition state (Structure II) reached after carbanion inversion (for details see text). The experimental data for triphenylphosphonium cyclopropylide (by X-ray diffraction)<sup>13</sup> are given for comparison. See Figure for number of atoms.

<u>Triphenylphosphonium Cyclopropylide</u>		<u>Structure I (Equilibrium)</u>	<u>Structure II (Transition State)</u>	<u>Structure III (Two Imaginary Vibrational Frequencies)</u>
Energy (hartrees)	— —	-458.22613	-458.21686	-458.21613
Energy (kcal)	— —	0.0	5.8	6.3
PC1	181.5	PHO 141.3	138.6	139.8
PC2	180.0	PH10 138.1	139.4	139.2
PC3	181.6	PH11 138.1	139.4	139.2
PC4	169.6	PC1 168.2	168.0	163.4
C4C41	152.6	C1C2 150.4	150.6	148.6
C4C42	152.6	C1C3 150.4	150.6	148.6
C41C42	149.9	C2C3 152.8	152.5	153.7
C1PC4	109.7	H10PC1 110.9	118.6	116.4
C2PC4	110.2	H11PC1 110.9	118.6	116.4
C1PC2	107.1	H10PH11 104.6	101.9	103.8
C3PC4	117.0	HOPC1 129.0	115.4	120.3
C1PC3	105.3	HOPH10 99.3	99.5	98.3
C2PC3	107.0	HOPH11 99.3	99.5	98.3

Table 1 continued

<u>Triphenylphosphonium Cyclopropylide</u>	<u>Structure I (Equilibrium)</u>	<u>Structure II (Transition State)</u>	<u>Structure III (Two Imaginary Vibrational Frequencies)</u>
C41C4P 117.1	C2C1P 127.6	127.1	141.8
C42C4P 117.8	C3C1P 127.6	127.1	141.8
C42C4C41 58.8	C2C1C3 61.1	60.9	62.3
C42C41C4 60.6	C1C2C3 59.5	59.6	58.9
C41C42C4 60.6	C1C3C2 59.5	59.6	58.9
Inclination 58.0 (P-Cylid vs. triangle)	44.9	45.6	23.4
	HOPC1C2 39.9	140.6	123.3
	PC1C2H30 134.5	12.1	35.3
	PC1C2H20 12.7	134.7	112.9

Table 2. Total Gross Population on Atoms as determined from the Mulliken Population Analysis of the three phosphonium cyclopropylide model configurations. (For atom numbering see Figures 2a-c).

<u>Atom</u>	<u>Structure I (Equilibrium)</u>	<u>Structure II (Transition State)</u>	<u>Structure III (Two Imaginary Vibrational Frequencies)</u>
1 Carbon 1	6.512	6.458	6.560
2 Phosphorous	14.458	14.427	14.419
3 Hydrogen 0	1.050	0.988	1.023
4 Hydrogen 10	0.972	0.998	1.000
5 Hydrogen 11	0.972	0.998	1.000
6 Carbon 2	6.374	6.366	6.349
7 Carbon 3	6.374	6.366	6.349
8 Hydrogen 20	0.835	0.841	0.829
9 Hydrogen 21	0.835	0.841	0.829
10 Hydrogen 30	0.809	0.813	0.821
11 Hydrogen 31	0.809	0.813	0.821


Table 3. Breakdown of the three highest occupied molecular orbitals for the equilibrium geometry of  $\text{H}_3\text{P}=\text{C}(\text{CH}_2)_2$  according to contributions from different atomic orbitals.

<u>Orbital</u>		<u>14a'</u>	<u>6a''</u>	<u>13a'</u>
Orbital Energy (hartrees)		0.2755	0.4035	0.4070
Carbon 1	s	0.07	0.00	0.02
	p	0.63	0.43	0.10
	d	0.00	0.01	0.01
Phosphorous	s	0.01	0.00	0.01
	p	0.05	0.00	0.07
	d	0.04	0.01	0.00
Hydrogen 0	s	0.08	0.00	0.00
Hydrogen 10	s	0.01	0.02	0.01
Carbon 2	s	0.00	0.00	0.00
	p	0.03	0.21	0.37
Hydrogen 20	s	0.01	0.02	0.02
Hydrogen 30	s	0.02	0.02	0.00

Table 4. Predicted vibrational frequencies (in  $\text{cm}^{-1}$ ) for  
 $\text{H}_3\text{P}=\text{C}(\text{CH}_2)_2$ .

Structure I (Equilibrium)		Structure III	
A'	3372	A'	3356
A''	3356	A''	3340
A'	3280	A'	3267
A''	3269	A''	3257
} C-H		} C-H	
A'	2726	A'	2635
A''	2720	A''	2628
A'	2450	A'	2525
} P-H		} P-H	
A'	1652	A'	1658
A''	1624	A''	1625
A'	1428	A'	1550
A'	1339	A'	1340
A''	1281	A''	1338
A''	1274	A''	1264
A''	1247	A''	1247
A'	1196	A'	1184
A'	1179	A'	1160
A'	1142	A'	1116
A''	1044	A''	1086
A'	1014	A''	1021
A''	997	A'	1012
A'	887	A'	904
A'	795	A''	828
A''	761	A'	810
A'	623	A'	626
A''	271	A''	247
A'	212	A'	158 i Ring inversion
A''	180	A''	229 i $\text{PH}_3$ internal rotation





.....

.....

Figure 2(a)

2b




Figure 2(b)

2c




Figure 2(c)

3a




Figure 3(a)

3b.




Figure 3(b)




Figure 3(c)

This report was done with support from the Department of Energy. Any conclusions or opinions expressed in this report represent solely those of the author(s) and not necessarily those of The Regents of the University of California, the Lawrence Berkeley Laboratory or the Department of Energy.

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Department of Energy to the exclusion of others that may be suitable.

TECHNICAL INFORMATION DEPARTMENT  
LAWRENCE BERKELEY LABORATORY  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720