UC Irvine UC Irvine Previously Published Works

Title

DUAL AND OPPOSING ROLES OF MIR-124 IN EPILEPTOGENESIS ARE MEDIATED THROUGH NRSF AND INFLAMMATORY PATHWAYS

Permalink

https://escholarship.org/uc/item/2450b2x0

Authors

Brennan, G Dey, D Chen, Y <u>et al.</u>

Publication Date

2016

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at https://creativecommons.org/licenses/by/4.0/

Peer reviewed

Abstracts

Platform Session 4: Basic Science 1 Monday 12th September

0008

DUAL AND OPPOSING ROLES OF MIR-124 IN EPILEPTOGENESIS ARE MEDIATED THROUGH NRSF AND INFLAMMATORY PATHWAYS

Brennan G^{1,2}, Dey D², Chen Y², Patterson K³, Magnetta E², Hall A², Dube C², Mei YT², Henshall D¹, Baram TZ^{2,3} ¹Physiology and Medical Physics, Royal College of Surgeons in Ireland, Dublin, Ireland, ²Pediatrics, University of California, Irvine, CA, United States, ³Anatomy and Neurobiology, University of California, Irvine, CA, United States

Insult-provoked transformation of neuronal networks into epileptic ones involves multiple mechanisms. Intervention studies have identified both dysregulated inflammatory pathways and NRSF-mediated repression of crucial neuronal genes as contributors to epileptogenesis. However, it remains unclear how epilepsy-provoking insults (e.g., prolonged seizures) induce both inflammation and NRSF and whether common mechanisms exist. We examined miR-124 as a candidate dual regulator of NRSF and inflammatory pathways. Status epilepticus (SE) induced by

kainic acid led to reduced miR-124 expression via SIRT1 – and, in turn, miR-124 repression – via C/EBP α upregulated NRSF. We tested whether augmenting miR-124 using miR-124 mimics/agomirs after SE would abort epileptogenesis by preventing both inflammation and NRSF upregulation. SE-sustaining animals developed epilepsy as detected by 24-video EEG monitoring, but supplementing miR-124 did not modify epileptogenesis. Examining this result further, we found that synthetic miR-124 not only effectively blocked NRSF upregulation and rescued NRSF target genes, but also augmented microglia activation and inflammatory cytokines. Thus, miR-124 attenuates epileptogenesis via NRSF while promoting epilepsy via inflammation.

Epilepsia, 57(Suppl. 2):8-9 2016 doi: 10.1111/epi.13609