

UCSF

UC San Francisco Previously Published Works

Title

Isolation of a Complete Circular Virus Genome Sequence from an Alaskan Black-Capped Chickadee (*Poecile atricapillus*) Gastrointestinal Tract Sample

Permalink

<https://escholarship.org/uc/item/2622004b>

Journal

Microbiology Resource Announcements, 3(5)

ISSN

2576-098X

Authors

Hanna, Zachary R
Runckel, Charles
Fuchs, Jérôme
[et al.](#)

Publication Date

2015-10-29

DOI

10.1128/genomea.01081-15

Peer reviewed

Isolation of a Complete Circular Virus Genome Sequence from an Alaskan Black-Capped Chickadee (*Poecile atricapillus*) Gastrointestinal Tract Sample

Zachary R. Hanna,^{a,b} Charles Runckel,^{c,d*} Jérôme Fuchs,^{a*} Joseph L. DeRisi,^{c,d} David P. Mindell,^{c*} Caroline Van Hemert,^e Colleen M. Handel,^e John P. Dumbacher^a

California Academy of Sciences, San Francisco, California, USA^a; Department of Integrative Biology and Museum of Vertebrate Zoology, University of California Berkeley, Berkeley, California, USA^b; Department of Biochemistry and Biophysics, University of California San Francisco, San Francisco, California, USA^c; Howard Hughes Medical Institute, Bethesda, Maryland, USA^d; Alaska Science Center, U.S. Geological Survey, Anchorage, Alaska, USA^e

* Present address: Charles Runckel, Runckel & Associates, Portland, Oregon, USA; Jérôme Fuchs, UMR 7205 Institut de Systématique, Evolution, Biodiversité, CNRS, MNHN, UPMC, EPHE, Sorbonne Universités, Département Systématique et Evolution, Muséum National d'Histoire Naturelle, Paris, France; David P. Mindell, Division of Environmental Biology, National Science Foundation, Arlington, Virginia, USA.

We report here the genome sequence of a circular virus isolated from samples of an Alaskan black-capped chickadee (*Poecile atricapillus*) gastrointestinal tract. The genome is 2,152 bp in length and is most similar (30 to 44.5% amino acid identity) to the genome sequences of other single-stranded DNA (ssDNA) circular viruses belonging to the gemycircularvirus group.

Received 5 August 2015 Accepted 11 August 2015 Published 24 September 2015

Citation Hanna ZR, Runckel C, Fuchs J, DeRisi JL, Mindell DP, Van Hemert C, Handel CM, Dumbacher JP. 2015. Isolation of a complete circular virus genome sequence from an Alaskan black-capped chickadee (*Poecile atricapillus*) gastrointestinal tract sample. *Genome Announc* 3(5):e01081-15. doi:10.1128/genomeA.01081-15.

Copyright © 2015 Hanna et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Zachary R. Hanna, zachanna@berkeley.edu, or John P. Dumbacher, jdumbacher@calacademy.org.

We isolated a circular viral genome from buccal and cloacal swabs sampled from an Alaskan black-capped chickadee (*Poecile atricapillus*) with avian keratin disorder (1) in Anchorage, AK, on 21 December 2010 and preserved it in RNAlater (Life Technologies). We extracted DNA and RNA using a ZR viral RNA kit (Zymo Research). We split the extracted nucleic acids into three separate 3- μ l aliquots, leaving one aliquot unmodified. We treated the second aliquot with T7 exonuclease and ExoI (New England BioLabs [NEB]) for 1 h at 37°C, and then 20 min at 65°C to eliminate noncircular DNA. We trapped circular DNA present in the third aliquot by adding 50 μ l of 0.8% low-melting-point agarose gel melted in Tris-acetate-EDTA buffer, cooling and solidifying the gel, running under a 100-V current for 1 h, and extracting DNA from the gel (2). Next, we performed a multiple displacement amplification (MDA) using Phi29 polymerase (NEB) on each aliquot and precipitated DNA in isopropanol. We pooled equal masses of the three aliquots and then enzymatically cut the DNA with six different 6-cutter restriction enzymes, separated the digested DNA on an agarose gel, and cut out and extracted DNA from the sharpest visualized band, an ~2-kb fragment from the AvrII (NEB) digest. We cloned the gel-extracted DNA, amplified colonies, and sequenced them using an ABI 3130xl.

We assembled sequences and produced subsequent assemblies using the Geneious *de novo* assembler [version 8.1.5; Biomatters (<http://www.geneious.com>)]. Using the alignment tool BLASTx, we searched the NCBI nonredundant protein sequences collection (3–5), finding matches of our assemblies to circular virus genome capsid and replication gene sequences. We designed primers with reference to the circular virus gene sequences, used them to amplify the remaining circular genomic sequences from the exonu-

lease and untreated aliquot MDA reaction products, and sequenced them on an ABI 3130xl. We assembled a complete circular genome of length 2,152 bp with two large bidirectionally oriented open reading frames (ORFs). One ORF (protein ID AKU89609) encodes a viral capsid protein and the other (AKU89610) a replicase protein. Using primers designed from the genome, we unsuccessfully attempted to amplify the virus from samples from 17 additional Alaskan *P. atricapillus* birds with avian keratin disorder.

We searched for publicly available sequences most similar to the assembled genome using the alignment tool BLASTx with a word size of 3 to search the NCBI nonredundant protein sequences collection (3–6). Ranked by bit score and discarding hits to hypothetical and unverified bacterial proteins, the top hits were capsid- and replication-associated proteins from viruses in the gemycircularvirus group (7). We performed a MUSCLE (version 3.8.425) (8) alignment of our genome against the 32 gemycircularvirus group complete genomes in NCBI GenBank (3, 6) on 27 May 2015. The alignment resulted in 30 to 44.5% identity with the genomes of the gemycircularvirus group at the amino acid level. Our genome sequence contributes to the resource of circular viral sequences isolated from vertebrate fecal samples (9–11).

Nucleotide sequence accession numbers. The genome has been deposited in GenBank under the accession no. **KT309029**. The version described in this paper is the first version, KT309029.1.

ACKNOWLEDGMENTS

Funding provided by Michael and Katalina Simon and the Howard Hughes Medical Institute made this work possible. The U.S. Geological Survey's Ecosystems Mission Area and Wildlife Program funded sample

collection field work. This material is based upon work supported by the National Science Foundation Graduate Research Fellowship under grant no. DGE 1106400 to Z.R.H.. The funders had no role in the study design, data collection, and interpretation, or the decision to submit the work for publication.

We thank Anna Sellas for assistance with lab work and Lisa Pajot for help with sample collection. We generated genetic data at the Center for Comparative Genomics, California Academy of Sciences.

Any opinion, findings, and conclusions or recommendations expressed in this material are those of the authors and do not necessarily reflect the views of the National Science Foundation. Any use of trade, product, or firm names in this publication is for descriptive purposes only and does not imply endorsement by the U.S. Government.

REFERENCES

- Handel CM, Pajot LM, Matsuoka SM, Hemert CV, Terenzi J, Talbot SL, Mulcahy DM, Meteyer CU, Trust KA. 2010. Epizootic of beak deformities among wild birds in Alaska: an emerging disease in North America? *Auk* 127:882–898. <http://dx.doi.org/10.1525/auk.2010.10111>.
- Schindler CW, Krolewski JJ, Rush MG. 1982. Selective trapping of circular double-stranded DNA molecules in solidifying agarose. *Plasmid* 7:263–270. [http://dx.doi.org/10.1016/0147-619X\(82\)90007-5](http://dx.doi.org/10.1016/0147-619X(82)90007-5).
- Acland A, Agarwala R, Barrett T, Beck J, Benson DA, Bollin C, Bolton E, Bryant SH, Canese K, Church DM, Clark K, DiCuccio M, Dondoshansky I, Federhen S, Feolo M, Geer LY, Gorenkov V, Hoepfner M, Johnson M, Kelly C, Khotomlianski V, Kimchi A, Kimelman M, Kitts P, Krasnov S, Kuznetsov A, Landsman D, Lipman DJ, Lu Z, Madden TL, Madej T, Maglott DR, Marchler-Bauer A, Karsch-Mizrachi I, Murphy T, Ostell J, O'Sullivan C, Panchenko A, Phan L, Pruitt DPKD, Rubinstein W, Sayers EW, Schneider V, Schuler GD, Sequeira E, Sherry ST, Shumway M, Sirotkin K, Siyan K, Slotta D, et al. 2014. Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res* 42:D7–D17.
- Boratyn GM, Camacho C, Cooper PS, Coulouris G, Fong A, Ma N, Madden TL, Matten WT, McGinnis SD, Merezuk Y, Raytselis Y, Sayers EW, Tao T, Ye J, Zaretskaya I. 2013. BLAST: a more efficient report with usability improvements. *Nucleic Acids Res* 41:W29–W33. <http://dx.doi.org/10.1093/nar/gkt282>.
- Johnson M, Zaretskaya I, Raytselis Y, Merezuk Y, McGinnis S, Madden TL. 2008. NCBI blast: a better Web interface. *Nucleic Acids Res* 36:W5–W9. <http://dx.doi.org/10.1093/nar/gkn201>.
- Benson DA, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, Sayers EW. 2015. GenBank. *Nucleic Acids Res* 43:D30–D35. <http://dx.doi.org/10.1093/nar/gku1216>.
- Rosario K, Dayaram A, Marinov M, Ware J, Kraberger S, Stainton D, Breitbart M, Varsani A. 2012. Diverse circular ssDNA viruses discovered in dragonflies (Odonata: Epiprocta). *J Gen Virol* 93:2668–2681. <http://dx.doi.org/10.1099/vir.0.045948-0>.
- Edgar RC. 2004. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 32:1792–1797. <http://dx.doi.org/10.1093/nar/gkh340>.
- Reuter G, Boros Á, Delwart E, Pankovics P. 2014. Novel circular single-stranded DNA virus from turkey faeces. *Arch Virol* 159:2161–2164. <http://dx.doi.org/10.1007/s00705-014-2025-3>.
- Sikorski A, Argiello-Astorga GR, Dayaram A, Dobson RCJ, Varsani A. 2013. Discovery of a novel circular single-stranded DNA virus from porcine faeces. *Arch Virol* 158:283–289. <http://dx.doi.org/10.1007/s00705-012-1470-0>.
- Blinkova O, Victoria J, Li Y, Keele BF, Sanz C, Ndjango J-N, Peeters M, Travis D, Lonsdorf EV, Wilson ML, Pusey AE, Hahn BH, Delwart EL. 2010. Novel circular DNA viruses in stool samples of wild-living chimpanzees. *J Gen Virol* 91:74–86. <http://dx.doi.org/10.1099/vir.0.015446-0>.