

UC San Diego

UC San Diego Previously Published Works

Title

Wearable Sensor/Device (Fitbit One) and SMS Text-Messaging Prompts to Increase Physical Activity in Overweight and Obese Adults: A Randomized Controlled Trial

Permalink

<https://escholarship.org/uc/item/27p8k0wt>

Journal

Telemedicine Journal and e-Health, 21(10)

ISSN

1530-5627

Authors

Wang, Julie B  
Cadmus-Bertram, Lisa A  
Natarajan, Loki  
et al.

Publication Date

2015-10-01

DOI

10.1089/tmj.2014.0176

Peer reviewed

# Wearable Sensor/Device (Fitbit One) and SMS Text-Messaging Prompts to Increase Physical Activity in Overweight and Obese Adults: A Randomized Controlled Trial

Julie B. Wang, PhD, MPH,<sup>1-3</sup> Lisa A. Cadmus-Bertram, PhD,<sup>1</sup> Loki Natarajan, PhD,<sup>1</sup> Martha M. White, MS,<sup>1</sup> Hala Madanat, PhD,<sup>2</sup> Jeanne F. Nichols, PhD, FACSM,<sup>4</sup> Guadalupe X. Ayala, PhD, MPH,<sup>2</sup> and John P. Pierce, PhD<sup>1</sup>

<sup>1</sup>Moores Cancer Center and <sup>4</sup>Center for Wireless Population Health Systems, University of California, San Diego, La Jolla, California.

<sup>2</sup>Graduate School of Public Health, San Diego State University, San Diego, California.

<sup>3</sup>Cardiovascular Research Institute, University of California, San Francisco, California.

This study is registered at [ClinicalTrials.gov](http://ClinicalTrials.gov) with clinical trial registration number NCT02188875.

## Abstract

**Background:** Studies have shown self-monitoring can modify health behaviors, including physical activity (PA). This study tested the utility of a wearable sensor/device (Fitbit<sup>®</sup> One<sup>™</sup>; Fitbit Inc., San Francisco, CA) and short message service (SMS) text-messaging prompts to increase PA in overweight and obese adults. **Materials and Methods:** Sixty-seven adults wore a Fitbit One tracker for 6 weeks; half were randomized to also receive three daily SMS-based PA prompts. The Fitbit One consisted of a wearable tracker for instant feedback on performance and a Web site/mobile application (app) for detailed summaries. Outcome measures were objectively measured steps and minutes of PA by intensity using two accelerometers: Actigraph<sup>™</sup> (Pensacola, FL) GT3X+ (primary measure) at baseline and Week 6 and Fitbit One (secondary measure) at baseline and Weeks 1, 2, 3, 4, 5, and 6. **Results:** Mixed-model repeated-measures analysis of primary measures indicated a significant within-group increase of +4.3 (standard error [SE]=2.0) min/week of moderate- to vigorous-intensity PA (MVPA) at 6-week follow-up ( $p=0.04$ ) in the comparison group (Fitbit only), but no study group differences across PA levels. Secondary measures indicated the SMS text-messaging effect lasted for only 1 week: the intervention group increased by +1,266 steps (SE=491;  $p=0.01$ ), +17.8 min/week MVPA (SE=8.5;  $p=0.04$ ), and +38.3 min/week total PA (SE=15.9;  $p=0.02$ ) compared with no changes in the comparison

group, and these between-group differences were significant for steps ( $p=0.01$ ), fairly/very active minutes ( $p<0.01$ ), and total active minutes ( $p=0.02$ ). **Conclusions:** These data suggest that the Fitbit One achieved a small increase in MVPA at follow-up and that the SMS-based PA prompts were insufficient in increasing PA beyond 1 week. Future studies can test this intervention in those requiring less help and/or test strategies to increase participants' engagement levels.

**Key words:** behavioral health, e-health, mobile health, sensor technology, technology

## Introduction

The combination of excess weight and lack of physical activity (PA) is associated with several chronic conditions, including diabetes, cardiovascular disease, and many cancers.<sup>1-3</sup> With over a third of U.S. adults obese<sup>4,5</sup> and half not meeting recommended levels of 150 min/week of moderate- to vigorous-intensity PA (MVPA),<sup>6,7</sup> developing low-cost interventions to increase PA levels is a public health priority.

Interventions to increase PA have involved several modalities, including in-person (individual or group), telephone, and Web-based counseling/coaching approaches.<sup>8-11</sup> More recently, there is growing interest among health behavior researchers in mobile health interventions that use mobile devices.<sup>12</sup>

In the United States, mobile phone usage is ubiquitous, and in 2012, approximately 86% of subscribers reported using short message service (SMS) text-messaging.<sup>13</sup> Researchers agree that text messaging has the potential to reach large audiences, including traditionally underserved populations,<sup>14</sup> and possibly serve as an inexpensive intervention modality.<sup>15</sup> Previous studies have shown that text messaging as a primary mode of communication can be successful for diabetes management,<sup>15-17</sup> smoking cessation,<sup>18-23</sup> and diet and/or PA for weight loss.<sup>24-27</sup> In these studies, text-messaging components were used in a variety of ways, ranging from simple reminders for medication adherence to rapid feedback on PA performance. Several studies with text messaging as the main intervention

component to promote PA have reported higher levels at follow-up compared with their respective comparison groups,<sup>28-35</sup> with a few showing no improvement.<sup>36-38</sup> It is unclear, however, whether these study effects were associated with the content (and/or intensity) of the text messages or simply because participants were responding to behavioral cues<sup>39</sup> as they received text messages that were reminding them to increase their activity levels. There is evidence that simple cues or prompts such as signs can be sufficient to increase the use of stairwells.<sup>40-43</sup> We hypothesize that prompts delivered as text messages could be equally effective at increasing daily PA.

An analysis of different behavioral strategies across 122 dietary and PA intervention studies concluded that the greatest behavioral change effects were achieved using self-monitoring plus at least one other self-regulatory technique (i.e., intention formation, specific goal setting, review of behavioral goals, and feedback on performance).<sup>44</sup> New commercially available wearable sensors/devices with integrated Web sites and mobile applications (apps) such as the Fitbit<sup>®</sup> One™ (Fitbit Inc., San Francisco, CA) offer a user-friendly tool for enhanced self-monitoring of PA (compared with traditional recordkeeping using diaries) that can help users to deploy their self-regulatory skills. These trackers allow users to collect objective measures of their own PA levels with a wearable device/sensor (an accelerometer), upload these data onto a personal Web site/mobile app account, view daily summary data to obtain more detailed feedback on their PA performance, and set/review goals. Wearable sensors/devices like the Fitbit One possess the technology that allows users to collect and monitor large amounts of their own PA data. However, there is a scarcity in the published literature on the usability of these devices and their effects on increasing PA.

The primary objective of this study was to test the effects on PA level of a technology-based intervention that delivered simple prompts using SMS text messaging in conjunction with the Fitbit One for self-monitoring. The study sample consisted of overweight and obese adults (mostly women) who were interested in increasing their PA.<sup>45</sup> A secondary objective was to examine the usability and effects of a wearable device/sensor (the Fitbit One) on PA levels. Specifically, the 6-week study tested and compared the effects of daily SMS-based PA prompts plus self-monitoring with the Fitbit One (intervention group) versus self-monitoring with the Fitbit One only (comparison group). Outcome measures were number of steps and minutes of PA by

intensity level using two accelerometers: the Actigraph (Pensacola, FL) GT3X+ (primary measure) and Fitbit One (secondary measure). The Actigraph GT3X+ is a valid and reliable measure of PA among adults<sup>46,47</sup> and thus provided primary measures of PA change from the baseline week to Week 6. The Fitbit One's technology allowed collection of additional days of PA measurement throughout the entire study period (up to 49 days). A study aim was to test the feasibility of the Fitbit One as a daily measure of PA without limitations in the number of days. A *priori* power and sample size estimates were calculated to compare likely differences in change in steps (primary outcome) between study groups. We hypothesized that SMS-based PA prompts plus the Fitbit One would show a greater increase in PA levels than the Fitbit One alone at 6-week follow-up.

## Materials and Methods

### STUDY DESIGN AND PARTICIPANTS

A two-group design was used to test the effects of daily text messaging as simple prompts to increase PA in a sample of overweight and obese adults (Fig. 1). Initial eligibility criteria were screened over the telephone and included being a non-smoker, 18-69 years old, overweight or obese (body mass index [BMI]  $\geq 25$  kg/m<sup>2</sup>), not meeting recommended levels of PA (<150 min/week of MVPA),<sup>48</sup> ability to safely increase PA,




Fig. 1. Participant flow (CONSORT diagram).

which was assessed using the Physical Activity Readiness Questionnaire,<sup>49</sup> text-messaging capabilities on a personal mobile phone, and meeting operation systems' requirements for the Fitbit One on a personal computer. Additionally, participants were eligible if they indicated willingness to increase their PA levels within 1 month of screening.

The study recruitment pool consisted of 177 subjects, of whom approximately 69% were women who had consented during their mammography appointments at the University of California at San Diego (UCSD) to be contacted for future research opportunities. Additionally, 19% were recruited via word-of-mouth and 12% from flyers that were posted throughout the community, including the UCSD and San Diego State University campuses. In total, 117 participants completed the initial telephone eligibility screening. The UCSD Institutional Review Board approved (on January 3, 2013) the study protocol and consent, and all participants provided written informed consent.

## INTERVENTION

*Baseline clinic visit and run-in period (prior to randomization).* Eligible participants were inactive and overweight/obese adults with varying history of PA and invited to a 1-h baseline clinic visit at UCSD Moores Cancer Center. The visit included a baseline questionnaire and measure of height (Seca<sup>®</sup> [Chino, CA] stadiometer) and weight (Scale-Tronix [White Plains, NY] medical scale) to verify self-reported height and weight taken at telephone screening. To set the PA agenda for all participants, study personnel provided participants with a brief 5-min intervention to review motivation, set goals (i.e., toward 10,000 steps/day), and plan for challenging situations. They also provided print materials from the U.S. Department of Health and Human Services (the 2008 *Physical Activity Guidelines for Americans*).<sup>48</sup> Participants were randomized to one of two study groups: Fitbit One alone or Fitbit One plus SMS text messaging.

Study personnel demonstrated how to wear the Actigraph GT3X+ (on an elastic belt clipped at the hip) and Fitbit One (clipped at the pocket, hip, or bra), as well as other functions of the Fitbit One (e.g., charging the tracker, wirelessly uploading data, and navigating the Fitbit Web site and/or app). They also demonstrated how to access personal Fitbit accounts for daily summaries of PA levels (i.e., steps, minutes of PA by "lightly active," "fairly active," and "very active" minutes) and highlighted the importance of charging and uploading the tracker almost every day to minimize missing data. Participants wore the Actigraph GT3X+ and Fitbit One devices concurrently for 7 days to assess baseline PA levels and to demonstrate their ability to use the Fitbit One. A "valid" day of measure was

defined as wearing both devices concurrently for a minimum of 600 min/day. Only those who provided at least 5 valid days with at least 1 weekend day from both devices met the eligibility criteria for randomization.

*Intervention group: self-monitoring with Fitbit One plus SMS text-messaging prompts.* Intervention participants were asked to indicate three preferred times of the day to receive text message prompts to engage in PA. The study used a commercial text-messaging Web site (EzTexting.com) to program automatic delivery of messages according to participants' pre-specified times. Participants were asked to contact the study if they wanted to change their schedules. Messages were limited to 150 characters, typically stated the time of delivery, and prompted participants to do PA (Example: "Good morning [name]! This is your 9AM reminder to do at least a 10-minute bout of moderate-to-vigorous intensity physical activity."). In total, 42 text messages were delivered sequentially, in which three messages were delivered every day within a 14-day cycle, and this pattern was repeated every 2 weeks throughout the 6-week study period. All participants were asked to continue wearing the Fitbit One tracker and upload data every day for the duration of the study.

*Comparison group: self-monitoring with Fitbit One only.* Participants who were randomly assigned to the comparison group were also asked to continue wearing the Fitbit One tracker and upload data every day for the duration of the study.

## OUTCOMES

*PA assessments: Actigraph GT3X+ and Fitbit One.* This study objectively measured PA using two types of triaxial accelerometers: (1) the Actigraph GT3X+, a well-validated assessment tool<sup>46,47</sup> that does not provide feedback to the individual, and (2) the Fitbit One, a more recent assessment tool that has not been well validated but *does* provide feedback on both the tracker and Web site/mobile app. The comparative validity of the Fitbit One measure will be addressed in a separate article. In this study, we examined number of steps (primary outcome) and minutes of PA by intensity levels.

*Actigraph GT3X+.* The maximum recording time of an Actigraph GT3X+ accelerometer is approximately 19 days before it needs to be returned to the study site for recharging. Therefore we provided participants with these accelerometers to wear for two weekly periods: at baseline (Week 0) and at the end of the intervention (Week 6). A minimum of 600 min/day was used as the cutoff for a valid day of measurement.<sup>50</sup> Data were processed using ActiLife version 6.10 software (Actigraph) (using Troiano default settings<sup>50</sup>) for nonwear bouts, spike

tolerance, and days with less than 600 min of measurement. Changes from baseline to Week 6 were calculated for steps per week and minutes per week of MVPA and total PA.

*Fitbit One.* The Fitbit One accelerometer can be recharged by participants using a USB cable, and it also wirelessly uploads data to the participant's personal computers or mobile devices. An initial validation report on Fitbit One (based on treadmill PA) has been published.<sup>51</sup> The Fitbit offers an added advantage to the Actigraph GT3X+ in that it can provide continuous measurement of PA across the entire study period. As this is a relatively novel device, one of our study aims focused on the usability of the Fitbit One as a self-monitoring system. Fitbit One measures of steps, fairly/very active minutes, and total active minutes were collected for Weeks 0, 1, 2, 3, 4, 5, and 6.

Study personnel accessed participants' Fitbit.com accounts (with consent) and recorded daily summary data. They were trained to identify days with nontypical wear patterns by visually scanning Fitbit graphs and flagging wear periods of zero movement for 4 or more h. These days were marked as "nontypical" to suggest that the tracker may not have been consistently worn throughout the day and/or data were not recorded possibly due to a depleted battery. The number of such nontypical days, however, was rare, particularly during baseline and follow-up weeks, and ranged from 5% to 9% of all observation points for all participants across all days for 5 weeks. Nonetheless, these days were excluded in the final analysis of daily Fitbit One data.

*Baseline questionnaire.* Participants completed a brief self-administered questionnaire during the baseline clinic visit that included items on demographics (i.e., age, sex, race, and education), text-messaging use, previous Web and/or app use for PA, personal and environmental factors associated with PA including motivation, and attitudes pertaining to self-monitoring and text messaging. Participants were also asked to rate on a 4-point scale (from "Very Confident" to "Not at All Confident") their answers to the question "How confident are you in your ability to increase your current physical activity levels to 150 min/wk of moderate-to-vigorous intensity physical activity in the next 6 weeks?" Three items with Likert-type responses were used to calculate a composite index score to assess baseline text-messaging use: (1) number of days used in a typical week, (2) average number of messages received per day, and (3) average number of messages sent per day. Participants' scores were categorized around the median split to determine whether their baseline text-messaging use was "frequent" or "infrequent."

*Follow-up questionnaire items.* Participants were asked to complete a 5–10-min telephone questionnaire at follow-up that assessed attitudes/behaviors pertaining to each intervention component (i.e., Fitbit tracker, Web site, and/or text messages). Participants were asked to rate on a 5-point scale from "Very Often" to "Never": "On a typical day, I checked the Fitbit tracker to see (a) how many steps I've taken (b) how much distance I've travelled and (c) if the flower grew taller (for intensity)." They were also asked "In a typical week, I logged onto my Fitbit account..." and to rate their response on a 5-point scale from "Everyday" to "Never." Items on text messaging included "The three daily text messages that prompted me to be physically active were..." with responses on a 3-point scale of "Too many" to "Too few," as well as an open-ended item, "Please describe in your own words how the text messages were useful or not useful in increasing your physical activity levels."

#### SAMPLE SIZE

Power and sample size estimates were calculated *a priori* to test the primary hypothesis that the group provided with the Fitbit One plus SMS-based PA prompts would have a greater increase in number of steps at follow-up than the group provided the Fitbit One only. From the literature<sup>52–55</sup> we assumed that the combined intervention effect on the standardized mean difference in steps would be at least 17% higher at follow-up than for the Fitbit only group. We assessed power at 80% with an alpha level of 0.05 for a two-sided two-sample *t* test. Allowing for 10% attrition, these calculations required us to enroll a minimum of 54 participants in the study.

#### RANDOMIZATION

A study personnel member who was not involved in baseline clinic visits used a permuted-block randomization procedure to allocate participants into study groups. Participants were contacted by e-mail to notify them of their group assignments.

#### STATISTICAL ANALYSIS

To assess if baseline demographic and lifestyle factors were comparable between randomized groups, we applied two-sided *t* tests for continuous variables (i.e., age, BMI, PA levels defined as steps and minutes by intensity level measured by the Actigraph GT3X+) and chi-squared tests for categorical variables (i.e., sex, education, race/ethnicity, text-messaging use, previous Web and/or app use for PA). A mixed-model repeated-measures analysis<sup>56</sup> was conducted to test and compare PA effects between intervention and comparison groups. An important advantage of this modeling paradigm is

that subjects with partially missing data can still be included in the models, thus potentially avoiding selection biases that would result from including only subjects with complete data (i.e., all 7 days of Actigraph GT3X+) (see Cnaan et al.<sup>56</sup> and Holzapfel et al.<sup>57</sup>). The outcome in the models was daily estimates of PA from the Actigraph GT3X+ at pre- (baseline), and postintervention (6-week follow-up) with up to seven measures per time point. A random subject-specific intercept was included to model between-subject variability, and fixed effects were time (i.e., pre- and postintervention), group, and the group by time interactions. A statistically significant group by time interaction indicated whether pre- to post-intervention changes in PA differed by study groups. All analyses were adjusted for daily wear-time minutes of the accelerometer. Mixed-models were for three outcomes: (a) steps (b) minutes of MVPA, and (c) minutes of total PA. Adherence to modeling assumptions was tested using residual plots (e.g., Q-Q plots to examine if residuals followed a Gaussian distribution).

To examine trajectories of activity over the 6-week period, mixed-model repeated-measure analysis were conducted for Fitbit One measures of PA levels at Weeks 0, 1, 2, 3, 4, 5, and 6 for (a) steps, (b) minutes of fairly/very active minutes, and (c) minutes of total active minutes. These analyses were also adjusted for wear-time minutes. All reported *p* values were considered statistically significant at an alpha level of <0.05. Analyses were conducted using SAS version 9.3 software (SAS Institute Inc., Cary, NC).

## Results

### STUDY SAMPLE

In total, 67 participants were randomized from January 2013 to January 2014 (see Fig. 1 for the CONSORT participant flow). Thirty-three participants were allocated to the SMS-based intervention group and 34 in the comparison group. Two participants were lost to follow-up in each study group. Additionally, two comparison participants indicated they were too busy and withdrew from the study within 1 week of randomization. All results were based on an intent-to-treat analysis and included all 67 subjects in the mixed models.

The study sample was 91% female, 61% college graduates, and 67% non-Hispanic white, with a mean (standard deviation) age of 48.2 (11.7) years (range, 19–66 years) and a BMI of 31.0 (3.7) kg/m<sup>2</sup>; 49% were overweight (BMI 25–29 kg/m<sup>2</sup>), and 51% were obese (BMI ≥30 kg/m<sup>2</sup>) (Table 1). At baseline, 50% reported that they frequently used SMS text messaging. Thirty-nine percent reported previously using a Web and/or mobile app for PA. Randomization achieved

**Table 1. Participants' Baseline Characteristics**

	<i>N</i>	INTERVENTION ( <i>N</i> =33)	COMPARISON ( <i>N</i> =34)	<i>P</i> VALUE
Age (years)	67	49.3 (11.5)	47.1 (11.9)	0.45
Sex				0.38
Female	61	88	94	
Male	6	12	6	
Education				0.37
< College	26	33	67	
≥ College graduate	41	44	56	
Race/ethnicity				0.83
White	45	67	68	
Hispanic	11	18	15	
African-American	3	12	9	
Asian	2	3	3	
Other	2	0	6	
BMI (kg/m <sup>2</sup> )				
25–29	33	52	47	0.72
≥30	34	48	53	
PA (Actigraph GT3X+)				
Steps ( <i>n</i> /day)	67	6,909 (415)	6,732 (401)	0.58
MVPA (min/week)	67	34.6 (3.0)	32.7 (2.9)	0.46
Total PA (min/week)	67	154.6 (5.3)	149.9 (6.8)	0.30
Wear time (min/day)	67	847.7 (122.2)	835.0 (119.1)	0.26
Text-messaging use				0.12
Frequent	34	52	47	
Infrequent <sup>a</sup>	33	48	53	
Web or app use <sup>b</sup>				0.26
Yes	27	41	37	
No	40	59	63	
Confidence change PA				
Very confident	31	38	53	<0.0001 <sup>c</sup>
Confident/somewhat	36	62	47	

Data are mean (standard deviation) values or percentages.

<sup>a</sup>A three-item composite index score assessed "frequent" and "infrequent" text-messaging use: (1) number of days text messaging used in a typical week, (2) average number of text messages received per day, and (3) average number of text messages sent per day.

<sup>b</sup>Previous Web and/or application (app) use specifically to monitor physical activity (PA).

<sup>c</sup>Chi-squared or *t* tests, alpha level *p*<0.05.

BMI, body mass index.

comparable study groups except for baseline confidence level in meeting recommended MVPA by the end of the study period. Baseline PA levels indicated significant group differences in steps ( $p=0.05$ ) and MVPA ( $p=0.04$ ) between those who were “Very Confident” versus “Confident” or “Somewhat Confident” (data not shown). Accordingly, overall and stratified analyses by baseline confidence level were conducted using the primary PA data.

#### ACTIGRAPH GT3X+: PA CHANGE FROM BASELINE TO WEEK 6

Primary assessment of PA was measured using the Actigraph GT3X+ at baseline (Week 0) and 6-week follow-up. Device wear times were comparable across assessment periods and group, which suggests results were not skewed by more or less PA that was collected depending on the amount of time devices were worn: baseline medians were 7 days (range, 5–7 days) and 843.8 min/day (range, 601.0–1,178.3 min/day), and 6-week follow-up medians were 7 days (range, 5–7 days) and 872.5 min/day (range, 607.3–1,110.3 min/day) (Table 2). There were no between-group differences in changes for steps or minutes of PA by intensity level (group by time interactions,  $p>0.1$ ). However, there was a significant within-group increase of +4.3 (standard error [SE]=2.0) min/week of MVPA from baseline to Week 6 ( $p=0.04$ ) in the comparison group.

A significant difference in baseline confidence levels in achieving PA goals suggested the need for stratified analyses. In the overall analyses, three-way interaction terms were tested in each model that included baseline confidence level (group by time by baseline confidence level) and were not significant for any of the outcomes. Nonetheless, we conducted stratified analyses by baseline confidence level,

and the results indicated no between-group differences (group by time interactions,  $p\geq 0.2$ ) (data not shown). In summary, the findings suggest that participants in the comparison group achieved a small increase in MVPA at 6-week follow-up (within-group difference from baseline to Week 6) and that baseline confidence level did not moderate this effect.

#### FITBIT ONE: PA LEVELS AT BASELINE AND WEEKS, 1, 2, 3, 4, 5, AND 6

Fitbit One measures of PA were collected at baseline (Week 0) and Weeks 1, 2, 3, 4, 5, and 6 (Fig. 2). When comparing the pattern of PA over the 6-week period, there were statistically significant group by time interactions for (a) steps ( $p=0.02$ ), (b) fairly/very active minutes ( $p<0.001$ ), and (c) total active minutes ( $p=0.04$ ), with the intervention group having on average higher activity levels over the 6 weeks compared with the comparison group.

To further investigate the intervention effects across time, group differences in PA were examined at each week compared with baseline in the mixed models: group by time interactions indicated significant between-group differences in PA change from baseline to week 1 for steps ( $p=0.01$ ), fairly/very active minutes ( $p<0.01$ ), and total active minutes ( $p=0.02$ ). These PA changes, however, were short term and not maintained through Weeks 2–6. In a further examination of these results within each study group, at Week 1, the intervention participants significantly increased their steps by +1,266 (SE=491;  $p=0.01$ ), fairly/very active minutes/week by +17.8 (SE=8.5;  $p=0.04$ ), and total active minutes/week by +38.3 (SE=15.9;  $p=0.02$ ) (Fig. 2). During the same period, there were no significant changes in PA levels in the comparison group: steps, -48 (SE=240,

**Table 2. Change in Physical Activity Levels Measured by Actigraph GT3X+ from Baseline to 6-Week Follow-Up, Adjusted for Wear Time**


	INTERVENTION GROUP			COMPARISON GROUP			P VALUE <sup>a,b</sup>
	BASELINE (N=33)	WEEK 6 (N=30)	CHANGE	BASELINE (N=34)	WEEK 6 (N=29)	CHANGE	
Steps (n/day)	6,885 (638)	6,909 (415)	24 (276)	7,165 (417)	6,732 (401)	-433 (222)	0.20
PA by intensity level (minutes/week)							
Moderate to vigorous	34.6 (3.0)	35.7 (2.5)	-1.1 (2.4)	32.7 (2.9)	36.9 (3.4)	4.3 (2.0) <sup>c</sup>	0.33
All intensity	154.6 (5.3)	153.0 (6.5)	-1.6 (4.5)	149.9 (6.8)	157.7 (6.9)	7.8 (4.2)	0.13

Data are mean (standard error) values.

<sup>a</sup>Mixed-model repeated-measures (group by time), alpha level  $p<0.05$ .

<sup>b</sup>Mixed-model repeated-measures three-way interactions (group by time by baseline confidence) for steps:  $p=0.63$ ; moderate to vigorous physical activity (PA),  $p=0.60$ ; all intensity PA,  $p=0.67$ .

<sup>c</sup>Significant, within-group increase,  $p=0.04$ .



**Fig. 2.** Weekly physical activity levels measured by the Fitbit One from baseline (Week 0) to Week 6, adjusted for baseline wear time: **(A)** number of steps, **(B)** fairly active and very active minutes, and **(C)** total active minutes. Group by time interactions were significant for steps ( $p=0.02$ ), fairly/very active minutes ( $p<0.001$ ), and total active minutes ( $p=0.04$ ). Cntl, control; Intv, intervention.

$p=0.84$ ); fairly/very active minutes/week,  $+2.3$  ( $SE=4.1$ ,  $p=0.57$ ); and total active minutes/week,  $-6.7$  ( $SE=11.7$ ,  $p=0.55$ ) (Table 3).

In summary, these data suggest that the Fitbit One (comparison group) was able to achieve a small within-group increase in MVPA at the 6-week follow-up among a sample of overweight and obese adults. This small within-group effect was significant in the Actigraph GT3X+ measures of PA. In the present study, it was feasible to collect daily objective measures of PA using the Fitbit One for up to 49 days throughout the study period. Analyses of these data suggest that a combination of daily SMS-based PA prompts and the Fitbit One (intervention group) increased PA levels for steps and minutes of PA by intensity, although only for a short-term period of 1 week. Therefore, daily SMS-based PA prompts in combination

with a Fitbit One device were not able to achieve sustained (i.e., 6-week) PA change. Instead, Fitbit One alone may help to increase MVPA at 6 weeks in overweight and obese adults. However, further research is needed with larger sample sizes and longer study periods to elucidate these findings.

#### FOLLOW-UP ATTITUDES AND BEHAVIORS ON THE FITBIT ONE AND SMS PA PROMPTS

At follow-up, a greater proportion of comparison (versus intervention) participants reported that, on a typical day, they viewed their Fitbit trackers “Very Often” or “Often” for steps (90% versus 71%) and distance (70% versus 55%). Those who self-reported that they frequently viewed their tracker were associated with greater increases in PA (even more so compared with the Web site). In the intervention group, approximately half indicated that the three daily text messages were “Too Many.” Additionally, in an open-ended question about the text-messaging intervention, a common response among participants was that they had stopped reading them altogether when they noticed that the messages were “automated.” Other notable phrases were that the messages were “inconvenient,” “annoying,” and “impersonal.”

#### Discussion

This study focused on a technology-based intervention to increase PA in a sample consisting mostly of inactive and overweight/obese women. Providing a technology-based self-monitoring device (Fitbit One) led to a small and statistically significant (within-group) increase in MVPA (in minutes/week) from baseline to Week 6. The intervention group included three daily SMS-based prompts to undertake PA, which were associated with increased PA over the first week, but this effect was lost by Week 2 of the 6-week intervention.

#### SELF-MONITORING (FITBIT ONE) ONLY

Results from the follow-up questionnaire indicated that participants in the comparison group, compared with the intervention group, was more engaged in using the Fitbit tracker, which might help to explain their small increase in MVPA at follow-up. These findings support several studies that have shown a positive association between self-



**Table 3. Summary of Changes in Physical Activity Levels Measured Using the Actigraph GT3X+ and Fitbit One from Baseline to Week 1 (Short-Term Effect) and Baseline to 6-Week Follow-Up, Adjusted for Wear Time**

	ACTIGRAPH GT3X+			FB		
	TEXTS + FB (INTERVENTION)	FB ONLY (COMPARISON)	P VALUE <sup>a</sup>	TEXTS + FB (INTERVENTION)	FB ONLY (COMPARISON)	P VALUE <sup>a</sup>
Change from baseline to Week 1						
Steps ( <i>n/day</i> )	–	–	–	1,266 (491)	– 48 (240)	0.01 <sup>b</sup>
PA by intensity level (minutes/week)						
MVPA or FA+VA	–	–	–	17.8 (8.5)	2.3 (4.1)	<0.01 <sup>b</sup>
All intensity	–	–	–	38.3 (15.9)	–6.7 (11.7)	0.02 <sup>b</sup>
Change from baseline to Week 6						
Steps ( <i>n/day</i> )	24 (276)	– 433 (222)	0.20	44 (292)	495 (257)	0.44
PA by intensity level (minutes/week)						
MVPA or FA+VA	– 1.1 (2.4)	4.3 (2.0) <sup>c</sup>	0.33	– 4.4 (5.0)	4.5 (5.2)	0.72
All intensity	– 1.6 (4.5)	7.8 (4.2)	0.13	– 16.1 (8.4)	– 19.6 (12.1)	0.10

<sup>a</sup>Mixed-model repeated-measures (group by time), alpha level  $p < 0.05$ , adjusted for baseline wear time (in minutes/week).

<sup>b</sup>Between-group significance.

<sup>c</sup>Within-group significance ( $p = 0.04$ ).

FA+VA, fairly active+very active; FB, Fitbit One; MVPA, moderate- to vigorous-intensity physical activity; PA, physical activity.

monitoring and PA change.<sup>58–62</sup> In the present study, the Fitbit tracker allowed participants to access quick readings of their PA performance throughout the day. It is unclear as to why comparison participants were more engaged compared with those in the intervention. A possible explanation might be that, although the randomization procedure achieved group comparability on most variables, there was a difference in baseline confidence level in their ability to increase PA. Therefore, it is possible that the comparison group had higher baseline confidence or self-efficacy to increase their PA levels.<sup>63</sup> However, further analyses also indicated that this small increase in MVPA within the comparison group was probably not moderated by baseline confidence.

Another possible explanation for the higher level of engagement in the comparison group might be that, because they only had the Fitbit One, they relied solely on that device. In contrast, the intervention group also received daily SMS-based prompts, which at follow-up they indicated were too frequent and automated. Thus, the text messages could have distracted them from further engagement in the study. It is important to note that, although the improvement in MVPA was significant in the comparison group, the effect was small and not accompanied by an increase in daily steps. Therefore, these findings need to be replicated before we can consider

them as evidence to support that simply providing a wearable sensor/sensor for self-monitoring would be sufficient to increase PA in inactive overweight/obese adults.

#### SHORT-TERM EFFECTS OF SMS PA PROMPTS

In this study, adding automated daily text messages as simple prompts for PA was not associated with increased PA at follow-up. However, using the Fitbit data, we were able to analyze change throughout the study period. During the first week, there was a significant increase in PA, which suggests these messages were able to serve as cues (or reminders); however, this effect was not maintained by the second week and into the remainder of the study. Indeed, many participants reported that they quickly stopped reading the daily texts when they discovered that the messages were not tailored to their performance. In addition, three messages a day as reminder messages were perceived too frequent to be helpful. Clearly, these SMS-based prompts were not effective in motivating change in PA levels for more than the first week.

Previous studies have used more intensive messaging strategies than simple prompts to promote PA.<sup>28,30–35</sup> A 9-week study used automated messages to help participants identify/reduce barriers and identify motivating benefits.<sup>29</sup> However, this more intensive messaging intervention reported similar results to the present study—an increase in MVPA that

was not maintained after the first week (assessed using a wrist-worn accelerometer).<sup>29</sup> The similarity in results between the two studies suggests that it might be the automated nature of text messaging rather than the content and/or frequency of messages associated with the studies' inability to maintain study effects for longer than a week. In the present study, it is possible that participants no longer felt accountable once they realized that the text messages were automated, which might explain the loss of PA effects. Other studies have reported longer-term effects but did not include objective measures of PA<sup>32,34,35</sup>; self-reported measures can easily be biased in many trial settings. In future studies, we recommend reducing the frequency of messages if they are used as basic cues/prompts. However, in order for these cues to be motivational, the messages might need to include some level of individually tailored (and/or adaptive) feedback on PA performance. For example, wearable sensors/devices like the Fitbit One could include push notifications with PA feedback to cue users to increase their PA levels.

#### STUDY LIMITATIONS

The study sample consisted of overweight and obese adults, mostly women who were participating in breast cancer screening within a clinic in a tertiary teaching hospital and indicated an interest in a PA study. Therefore, generalizability of these results is limited. This intervention might have had better success with a sample of adults that was more representative of the general population and among those with higher motivation to increase their PA levels. Participants were asked to concurrently wear the Actigraph GT3X+ and Fitbit One at baseline. Therefore, participants were inadvertently exposed to some PA intervention with the Fitbit One during baseline measure of PA and prior to the start of the study, which could have diminished the effects of either the Fitbit One and/or text messaging. A strength of this study was the use of the Actigraph GT3X+, which is a valid and reliable measure of PA.<sup>46,47</sup> A validation study has compared Fitbit One measures with the Actigraph GT3X+ for steps that were taken on a treadmill,<sup>51</sup> and validation in a real-world setting is currently underway. More generally, improvements are needed in these technologies for more accurate measures across an array of activities beyond steps (e.g., cycling and swimming).

#### Conclusions

Results from this study suggest that simply providing a wearable sensor/device for self-monitoring of PA was insufficient in achieving increases in target PA levels in a sample of overweight and obese adults consisting mostly of women. Future studies on wearable sensors/devices may require closer

examination of engagement levels with the technology and level of help needed to achieve target PA levels. The addition of daily automated text messages as simple reminders in conjunction with the wearable sensor/device did not help to increase PA levels. These data suggest that text-messaging interventions likely require more individualized and/or adaptive strategies such as feedback on PA performance. More research is needed to investigate an optimal intervention package that feature wearable sensors/devices to effectively promote PA change. Text messages (or even push notifications via mobile apps) may be part of such an intervention, but results from this study suggest that messages should be more responsive to participants' individual PA performance.

#### Acknowledgments

This research was supported by a gift from the Carol Vasiliadis family and in part by grant CA-113710 from the National Cancer Institute. The study would like to acknowledge UCSD undergraduate interns Quynh Nguyen and Amy Nham for their assistance in study implementation.

#### Disclosure Statement

No competing financial interests exist.

#### REFERENCES

1. Bassuk SS, Manson JE. Epidemiological evidence for the role of physical activity in reducing risk of type 2 diabetes and cardiovascular disease. *J Appl Physiol* **2005**;99:1193–1204.
2. Ehemann C, Henley SJ, Ballard-Barbash R, et al. Annual Report to the Nation on the status of cancer, 1975–2008, featuring cancers associated with excess weight and lack of sufficient physical activity. *Cancer* **2012**;118:2338–2366.
3. Malnick SDH, Knobler H. The medical complications of obesity. *QJM* **2006**;99:565–579.
4. Flegal KM, Carroll MD, Ogden CL, Curtin LR. Prevalence and trends in obesity among US adults, 1999–2008. *JAMA* **2010**;303:235–241.
5. Ogden CL, Carroll MD, Kit BK, Flegal KM. Prevalence of obesity in the United States, 2009–2010. *NCHS Data Brief* **2012**; (82):1–8.
6. Prevalence of self-reported physically active adults—United States, 2007. *MMWR Morb Mortal Wkly Rep* **2008**;57:1297–1300.
7. Troiano RP, Berrigan D, Dodd KW, Mâsse LC, Tilert T, McDowell M. Physical activity in the United States measured by accelerometer. *Med Sci Sports Exerc* **2008**;40:181–188.
8. Jakicic JM, Marcus BH, Lang W, Janney C. Effect of exercise on 24-month weight loss maintenance in overweight women. *Arch Intern Med* **2008**;168:1550–1559; discussion 1559–1560.
9. Irwin ML, Alvarez-Reeves M, Cadmus L, et al. Exercise improves body fat, lean mass, and bone mass in breast cancer survivors. *Obesity* **2009**;17:1534–1541.
10. Rock CL, Flatt SW, Sherwood NE, Karanja N, Pakiz B, Thomson CA. Effect of a free prepared meal and incentivized weight loss program on weight loss and weight loss maintenance in obese and overweight women: A randomized controlled trial. *JAMA* **2010**;304:1803–1810.
11. Pierce JP, Natarajan L, Caan BJ, et al. Influence of a diet very high in vegetables, fruit, and fiber and low in fat on prognosis following treatment for breast cancer: The Women's Healthy Eating and Living (WHEL) randomized trial. *JAMA* **2007**;298:289–298.

12. Norman GJ, Zabinski MF, Adams MA, Rosenberg DE, Yaroch AL, Atienza AA. A review of eHealth interventions for physical activity and dietary behavior change. *Am J Prev Med* **2007**;33:336–345.
13. The Nielsen Company. The mobile consumer: A global snapshot. **2013**. Available at <http://www.nielsen.com/content/dam/corporate/uk/en/documents/Mobile-Consumer-Report-2013.pdf> (last accessed May 1, 2014).
14. Fjeldsoe BS, Marshall AL, Miller YD. Behavior change interventions delivered by mobile telephone short-message service. *Am J Prev Med* **2009**;36:165–173.
15. Cole-Lewis H, Kershaw T. Text-messaging as a tool for behavior change in disease prevention and management. *Epidemiol Rev* **2010**;32:56–69.
16. Hanauer DA, Wentzell K, Laffel N, Laffel LM. Computerized Automated Prompt Diabetes System (CARDS): E-mail and SMS cell phone text-messaging prompts to support diabetes management. *Diabetes Technol Ther* **2009**;11:99–106.
17. Franklin V, Waller A, Pagliari C, Greene S. "Sweet Talk": Text-messaging support for intensive insulin therapy for young people with diabetes. *Diabetes Technol Ther* **2003**;5:991–996.
18. Rodgers A, Corbett T, Bramley D, et al. Do u smoke after txt? Results of a randomised trial of smoking cessation using mobile phone text-messaging. *Tob Control* **2005**;14:255–261.
19. Free C, Knight R, Robertson S, et al. Smoking cessation support delivered via mobile phone text-messaging (txt2stop): A single-blind, randomised trial. *Lancet* **2011**;378:49–55.
20. Obermayer JL, Riley WT, Asif O, Jean-Mary J. College smoking-cessation using cell phone text-messaging. *J Am Coll Health* **2004**;53:71–78.
21. Riley W, Obermayer J, Jean-Mary J. Internet and mobile phone text-messaging intervention for college smokers. *J Am Coll Health* **2008**;57:245–248.
22. Bramley D, Riddell T, Whittaker R, et al. Smoking cessation using mobile phone text-messaging is as effective in Maori as non-Maori. *N Z Med J* **2005**;118:U1494.
23. Haug S, Meyer C, Schorr G, Bauer S, John U. Continuous individual support of smoking cessation using text-messaging: A pilot experimental study. *Nicotine Tob Res* **2009**;11:915–923.
24. Patrick K, Raab F, Adams MA, et al. A text message-based intervention for weight loss: Randomized controlled trial. *J Med Internet Res* **2009**;11:e1.
25. Gerber BS, Stolley MR, Thompson AL, Sharp LK, Fitzgibbon ML. Mobile phone text-messaging to promote healthy behaviors and weight loss maintenance: A feasibility study. *Health Inform J* **2009**;15:17–25.
26. Napolitano MA, Hayes S, Bennett GG, Ives AK, Foster GD. Using Facebook and text-messaging to deliver a weight loss program to college students. *Obesity* **2013**;21:25–31.
27. Shapiro JR, Bauer S, Hamer RM, Kordy H, Ward D, Bulik CM. Use of text-messaging for monitoring sugar-sweetened beverages, physical activity, and screen time in children: A pilot study. *J Nutr Educ Behav* **2008**;40:385–391.
28. Cheung PP, Chow BC, Parfitt G. Using environmental stimuli in physical activity intervention for school teachers: A pilot study. *Glob J Heal Educ Promot* **2008**;11:47–56.
29. Hurling R, Catt M, Boni MD, et al. Using internet and mobile phone technology to deliver an automated physical activity program: Randomized controlled trial. *J Med Internet Res* **2007**;9:e7.
30. Prestwich A, Perugini M, Hurling R. Can the effects of implementation intentions on exercise be enhanced using text messages? *Psychol Health* **2009**;24:677–687.
31. Prestwich A, Perugini M, Hurling R. Can implementation intentions and text messages promote brisk walking? A randomized trial. *Health Psychol* **2010**;29:40–49.
32. Conroy MB, Yang K, Elci OU, et al. Physical activity self-monitoring and weight loss: 6-month results of the SMART trial. *Med Sci Sports Exerc* **2011**;43:1568–1574.
33. Antypas K, Wangberg SC. An Internet- and mobile-based tailored intervention to enhance maintenance of physical activity after cardiac rehabilitation: Short-term results of a randomized controlled trial. *J Med Internet Res* **2014**;16:e77.
34. Fjeldsoe BS, Miller YD, Marshall AL. MobileMums: A randomized controlled trial of an SMS-based physical activity intervention. *Ann Behav Med* **2010**;39:101–111.
35. King AC, Ahn DK, Oliveira BM, Atienza AA, Castro CM, Gardner CD. Promoting physical activity through hand-held computer technology. *Am J Prev Med* **2008**;34:138–142.
36. Haapala I, Barengo NC, Biggs S, Surakka L, Manninen P. Weight loss by mobile phone: A 1-year effectiveness study. *Public Health Nutr* **2009**;12:2382–2391.
37. Nguyen HQ, Gill DP, Wolpin S, Steele BG, Benditt JO. Pilot study of a cell phone-based exercise persistence intervention post-rehabilitation for COPD. *Int J Chron Obstruct Pulmon Dis* **2009**;4:301–313.
38. Gasser R, Brodbeck D, Degen M, Luthiger J, Wyss R, Reichlin S. Persuasiveness of a mobile lifestyle coaching application using social facilitation. In: IJsselsteijn WA, Kort YAW, Midden C, Eggen B, Hoven E, eds. *Persuasive technology*. Berlin: Springer, **2006**:27–38.
39. Abraham C, Michie S. A taxonomy of behavior change techniques used in interventions. *Health Psychol* **2008**;27:379–387.
40. Anderson RE, Franckowiak SC, Snyder J, Barlett SJ, Fontaine KR. Physical activity promotion by the encouraged used of stairs. *Ann Intern Med* **1998**;129:363–369.
41. Blamey A, Mutrie N, Aitchison, T. Promoting active living: A step in the right direction. *J Inst Health Educ* **1996**;34:5–9.
42. Brownell KD, Stunkard AJ, Albaum JM. Evaluation and modification of exercise patters in the natural environment. *Am J Psychiatry* **1980**;137:1540–1545.
43. Kerr J, Eves FF, Carroll D. The influence of poster prompts on stair use: The effects of setting, poster size and content. *Br J Health Psychol* **2001**;6:397–405.
44. Michie S, Abraham C, Whittington C, McAteer J, Gupta S. Effective techniques in healthy eating and physical activity interventions: A meta-regression. *Health Psychol* **2009**;28:690–701.
45. Williams AD. Use of a text-messaging program to promote adherence to daily physical activity guidelines: A review of the literature. *Bariatr Nurs Surg Patient Care* **2012**;7:13–16.
46. Le Masurier GC, Lee SM, Tudor-Locke C. Motion sensor accuracy under controlled and free-living conditions. *Med Sci Sports Exerc* **2004**;36:905–910.
47. Bassett DR, Dinesh J. Use of pedometers and accelerometers in clinical populations: Validity and reliability issues. *Phys Ther Rev* **2010**;15:135–142.
48. U.S. Department of Health and Human Services. *Physical activity guidelines for Americans*. Washington, DC: U.S. Department of Health and Human Services, **2008**:21–25.
49. Scott T, Reading J, Shephard RJ. Revision of the Physical Activity Readiness Questionnaire (PAR-Q). *Can J Sport Sci* **1992**;17:338–345.
50. Cain KL, Geremia GM. Accelerometer data collection and scoring manual for adult and senior studies. San Diego, CA: San Diego State University, **2012**. Available at <http://www.drjamesallis.sdsu.edu/>, accessed May 1, 2014.
51. Takacs J, Pollock CL, Guenther JR, Bahar M, Napier C, Hunt MA. Validation of the Fitbit One activity monitor device during treadmill walking. *J Sci Med Sport* **2014**;17:496–500.
52. Friedenreich C, Woolcott C, McTiernan A, et al. Adiposity changes after a 1-year aerobic exercise intervention among postmenopausal women: A randomized controlled trial. *Int J Obes* **2010**;35:427–435.
53. Napolitano MA, Borradaile KE, Lewis BA, et al. Accelerometer use in a physical activity intervention trial. *Contemp Clin Trials* **2010**;31:514–523.
54. Rogers LQ, Hopkins-Price P, Vicari S, et al. Physical activity and health outcomes three months after completing a physical activity behavior change intervention: Persistent and delayed effects. *Cancer Epidemiol Biomarkers Prev* **2009**;18:1410–1418.
55. Rogers LQ, Hopkins-Price P, Vicari S et al. A randomized trial to increase physical activity in breast cancer survivors. *Med Sci Sports Exerc* **2009**;41:935.

56. Cnaan A, Laird NM, Slasor P. Using the general linear mixed model to analyse unbalanced repeated measures and longitudinal data. *Stat Med* **1997**;16:2349–2380.
57. Holzapfel C, Cresswell L, Ahern AL, et al. The challenge of a 2-year follow-up after intervention for weight loss in primary care. *Int J Obes* **2014**;38:806–811.
58. Carels RA, Darby LA, Rydin S, Douglass OM, Cacciapaglia HM, O'Brien WH. The relationship between self-monitoring, outcome expectancies, difficulties with eating and exercise, and physical activity and weight loss treatment outcomes. *Ann Behav Med* **2005**;30:182–190.
59. Gleeson-Kreig JM. Self-monitoring of physical activity: Effects on self-efficacy and behavior in people with type 2 diabetes. *Diabetes Educ* **2006**;32:69–77.
60. Napolitano MA, Fotheringham M, Tate D, et al. Evaluation of an internet-based physical activity intervention: A preliminary investigation. *Ann Behav Med* **2003**;25:92–99.
61. Tudor-Locke C, Bassett DR, Swartz AM, et al. A preliminary study of one year of pedometer self-monitoring. *Ann Behav Med* **2004**;28:158–162.
62. Tate DF. Using internet technology to deliver a behavioral weight loss program. *JAMA* **2001**;285:1172.
63. Bandura A. *Self-efficacy: The exercise of self-control*. Gordonsville, VA: W.H. Freeman & Co., **1997**.

Address correspondence to:  
*Julie B. Wang, PhD, MPH*  
*University of California, San Francisco*  
*530 Parnassus Avenue*  
*San Francisco, CA 94143-1390*

*E-mail: julie.wang@ucsf.edu*

*Received: August 26, 2014*

*Revised: December 4, 2014*

*Accepted: December 4, 2014*