

UCSF

UC San Francisco Previously Published Works

Title

Phospholipase C gamma 1 is required for activation of store-operated channels in human keratinocytes

Permalink

<https://escholarship.org/uc/item/280114jn>

Journal

Journal of Investigative Dermatology, 124(1)

ISSN

0022-202X

Authors

Tu, Chia-Ling L
Chang, W H
Bikle, D D

Publication Date

2005

Peer reviewed

Phospholipase C γ 1 Is Required for Activation of Store-operated Channels in Human Keratinocytes

Chia-Ling Tu*, Wenhan Chang and Daniel D. Bikle

Endocrine Unit, Veteran Affairs Medical Center and Department of Medicine, University of California, San Francisco, California 94121, USA

*Corresponding author: Dr. Chia-Ling Tu

Endocrine Unit (111N), 4150 Clement Street, San Francisco, CA 94121, USA

Telephone: 415-7502089

Fax: 415-750-6929

E-mail: ctu@itsa.ucsf.edu

Running title: Store-operated Ca^{2+} entry in keratinocytes

The abbreviations used are; CCE, capacitative calcium entry; PLC, phospholipase C; DAG, diacylglycerol; IP_3 , inositol 1,4,5-trisphosphate; IP_3R , inositol 1,4,5-trisphosphate receptor; ER, endoplasmic reticulum; ROC, receptor-operated channel; SOC, store-operated channel; TRPC, canonical transient receptor potential channel; SH3, Src homology 3 domain; 2-APB, 2-aminoethoxydiphenylborane; $[\text{Ca}^{2+}]_i$, cytosolic Ca^{2+} concentration; OAG, 1-oleoyl-2-acetyl-*sn*-glycerol; PKC, protein kinase C.

Keywords: SOC, keratinocyte, PLC γ 1, TRPC, IP_3R

ABSTRACT

Store-operated calcium entry depicts the movement of extracellular Ca^{2+} into cells through plasma membrane Ca^{2+} channels activated by depletion of intracellular Ca^{2+} stores. The members of the canonical subfamily of transient receptor potential channels (TRPC) have been implicated as the molecular bases for store-operated channels (SOC). Here we investigate the role of phospholipase C (PLC) in regulation of native SOC and the expression of endogenous TRPC in human epidermal keratinocytes. Calcium entry in response to store depletion with thapsigargin was reversibly blocked by 2-aminoethoxydiphenyl borane, an effective SOC inhibitor, and suppressed by the diacylglycerol analogue, 1-oleoyl-2-acetyl-sn-glycerol. Inhibition of PLC with U73122 or transfection of a PLC γ 1 antisense cDNA construct completely blocked SOC activity, indicating a requirement for PLC, especially PLC γ 1, in the activation of SOC. RT-PCR and immunoblotting analyses showed that TRPC1, 3, 4, 5 and 6 are expressed in keratinocytes. Knockdown of the level of endogenous TRPC1 or TRPC4 inhibited store-operated calcium entry, indicating they are part of the native SOC. Co-immunoprecipitation studies demonstrated that TRPC1, but not TRPC4, interacts with PLC γ 1 and the inositol 1,4,5-trisphosphate receptor (IP $_3$ R). The association of TRPC1 with PLC γ 1 and IP $_3$ R decreased in keratinocytes with higher intracellular Ca^{2+} , coinciding with a down-regulation in SOC activity. Our results indicate that the activation of SOC in keratinocytes depends, at least partly, on the interaction of TRPC with PLC γ 1 and IP $_3$ R.

INTRODUCTION

In most non-excitable cells, release of calcium from intracellular stores triggers entry of extracellular calcium through channels in the plasma membrane, a process known as capacitative calcium entry (CCE) (Putney, et al, 2001). CCE provides an important means for mediating long-term Ca^{2+} signals and replenishment of Ca^{2+} stores (Parekh and Penner, 1997; Venkatachalam, et al, 2002). Activation of CCE can occur through various mechanisms (Putney, et al, 2001; Venkatachalam, et al, 2002). Binding of ligands to plasma membrane receptors activates phospholipase C (PLC), which catalyzes the hydrolysis of phosphatidyl inositol 4,5-bisphosphate to generate diacylglycerol (DAG) and inositol 1,4,5-trisphosphate (IP_3). DAG has been known for its ability to directly activate ion channels (Vazquez et al, 2001; Venkatachalam, et al, 2001) as well as protein kinase C (PKC), whereas IP_3 triggers Ca^{2+} release from intracellular Ca^{2+} stores, e.g. the endoplasmic reticulum (ER), through binding to IP_3 receptors (IP_3R) (Berridge, 1993). The resulting reduction of Ca^{2+} content in the ER lumen activates Ca^{2+} entry through ion channels in the plasma membrane (Putney, 1986). The term “agonist-induced Ca^{2+} entry” was used to refer to the Ca^{2+} entry mechanism induced by receptor activation (Patterson et al, 2002). In addition, agents such as the Ca^{2+} pump inhibitor thapsigargin and the calcium ionophore ionomycin, which deplete calcium stores independently of receptor activation or IP_3 generation, also activate Ca^{2+} entry (Putney, 1999). Therefore, the CCE channels can be loosely classified as receptor-operated channels (ROC) and store-operated channels (SOC), referring to Ca^{2+} entry channels activated in response to activation of PLC-coupled receptors and to Ca^{2+} store depletion, respectively (Berridge, 1997). However, it is difficult to clearly define Ca^{2+} entry channels as either ROCs or SOCs, since both mechanisms converge on Ca^{2+} release from internal stores. The canonical subfamily of transient receptor potential (TRPC) channels

has long been considered the candidates for ROCs and/or SOCs (Clapham et al, 2001; Montell, 2001). TRPC channels can be activated by a number of mediators involved in the PLC signaling cascade, including Ca^{2+} (Zitt et al, 1997), DAG (Kiselyov and Muallem, 1999; Venkatachalam et al, 2003) and activated IP_3R (Boulay et al, 1999; Kiselyov et al, 1998). Mammalian TRPC channels include 7 members: TRPC1, TRPC2, the DAG-responsive subgroup of TRPC3/6/7 (Trebak et al, 2003) and DAG-insensitive subgroup of TRPC4/5 (Kiselyov and Muallem, 1999; Venkatachalam et al, 2003). Many studies demonstrate that mammalian TRPC channels can be activated in response to store-depletion with calcium pump blockers or ionophores (Clapham et al, 2001), even though studies on overexpressed mammalian TRPC channels suggest that receptor-mediated stimulation of PLC is the major activation mechanism for TRPCs (Hofmann et al, 2000). Nevertheless, gene knockdown (Bough et al, 2001; Philipp et al, 2000) or knockout (Freichel et al, 2001; Mori et al, 2002) studies indicate that endogenously expressed TRPC channels contribute to native SOC. In particular, TRPC1, TRPC3 and TRPC4 have been shown to be part of endogenous SOC in human submandibular gland cells (Liu et al, 2000), neurons (Li et al, 1999), and adrenal cortex cells (Philipp et al, 2000), respectively.

Though the underlying mechanisms of agonist-induced and store-operated Ca^{2+} entry are unclear, the favored conformational coupling model (Irvine, 1990; Putney, 1999) proposes that a reduction in ER luminal calcium induces a conformational change of IP_3R , which directly activates SOC in the plasma membrane through direct interaction. Studies of overexpressed TRPC demonstrated that IP_3R interacts with TRPC1 (Lockwich et al, 2000; Rosado and Sage, 2000a), TRPC3 (Kiselyov et al, 1998; Kiselyov et al, 1999), TRPC4 (Tang et al, 2001) and TRPC6 (Boulay et al, 1999). Overexpression of TRPC peptides containing the IP_3R -interacting domain reduced the activity of endogenous SOC, suggesting that native SOC is activated via

IP₃Rs (Boulay et al, 1999). In several cell types store-depletion induced Ca²⁺ entry was blocked when PLC activity was inhibited or polyphosphoinositides were depleted, indicating that besides activated IP₃ receptors, activation of SOC requires functional PLC activity (Broad et al, 2001; Rosado and Sage, 2000b). It has been suggested that the IP₃R-SOC complex may locate in close proximity to a PLC to respond to low levels of IP₃ required for channel activation (Broad et al, 2001; Putney, 1999). Recently Patterson et al (Patterson et al, 2002) demonstrated that PLC γ plays a structural role in regulating agonist-induced calcium entry, since the action of PLC γ is dependent on its Src homology 3 (SH3) domain but independent of its lipase activity. It is unclear whether PLC γ regulates the function of SOC, although overexpression of PLC γ 1 has been shown to augment both agonist-induced and store-operated Ca²⁺ entry (Patterson et al, 2002).

In human keratinocytes, emptying the intracellular calcium store has been shown to activate SOC (Csernoch et al, 2000; Karvonen et al, 2000). In this study, we investigated the role of PLC γ 1 in regulation of store-operated Ca²⁺ entry and the molecular identities of endogenous SOCs in normal epidermal keratinocytes. We demonstrate that SOC function is sensitive to disruption of basal PLC activity, and that activation of SOC requires PLC γ 1. Our results indicate that TRPC1 and TRPC4 are part of the native SOC subunits in keratinocytes and that TRPC1 physically associates with PLC γ 1 and IP₃R. In addition, we demonstrate that the inhibition of SOC activity by calcium parallels the attenuated interactions of TRPC1 with PLC γ 1 and IP₃R. Our findings indicate that the coupling of native SOC to signaling molecules in the PLC/IP₃ pathway provides a gating mechanism for activation of calcium entry by store depletion.

EXPERIMENTAL PROCEDURES

Materials- Ionomycin, thapsigargin, and 2-APB (2-aminoethoxydiphenylborane) were purchased from Calbiochem-Novabiochem Corp. (La Jolla, CA). Membrane permeable DAG analogues, 1-oleoyl-2-acetyl-sn-glycerol (OAG) and 1, 2-dioctanoyl-sn-glycerol (DOG) and the phospholipase C inhibitor U-73122 and its inactive analog U-73343 were purchased from Sigma Chemicals (St. Louis, MO). Stock solutions of these compounds were prepared in DMSO or ethanol. All DNA constructs used in the transfection were prepared using Qiagen Maxi-prep columns (Chatsworth, CA) according to the manufacturer's protocol.

Cell Culture- Normal human keratinocytes (NHK) were isolated from newborn foreskin and cultured as described (Gibson et al, 1996). Cells were maintained in serum free keratinocyte growth medium (KGM; Clonetics Inc., San Diego, CA) supplemented with 0.03 mM CaCl_2 .

Measurement of Cytosolic Ca^{2+} - Keratinocytes cultured on glass coverslips were loaded with 7.5 μM Fura-2/AM (Molecular Probes, Eugene, OR) in 0.1% Pluronic F127 in buffer A (20 mM Hepes, 120 mM NaCl, 5 mM KCl, 1 mM MgCl_2 , 1 mg/ml sodium pyruvate, 1 mg/ml glucose) containing 0.07 mM calcium. Cells were then washed with Ca^{2+} -free buffer A before Ca^{2+} measurements were made. The fluorescence of Fura2-loaded cells was monitored using a Dual-wavelength Fluorescence Imaging System (Intracellular Imaging Inc., Cincinnati, OH). The cells were alternately illuminated with 340 nm and 380 nm light, and the fluorescence at emission wavelength 510 nm was recorded. Ca^{2+} stores were depleted by 1 μM thapsigargin in 0.1 mM EGTA, and Ca^{2+} entry in Fura2-loaded keratinocytes was determined in the presence of 2 mM calcium, unless indicated otherwise. All experiments were performed at room temperature. The signals from 30 to 50 single cells for each measurement were recorded. Each sample was calibrated by the addition of 20 μM ionomycin (R_{max}) followed by 20 mM

EGTA/Tris, pH 8.3 (R_{min}). Cytosolic Ca^{2+} concentration ($[Ca^{2+}]_i$) was calculated from the ratio of emission at the two excitation wavelengths based on the formula $[Ca^{2+}]_i = K_d Q(R - R_{min}) / (R_{max} - R)$, $R = F_{340} / F_{380}$, $Q = F_{min} / F_{max}$ at 380 nm, and K_d for Fura-2 for Ca^{2+} is 224 nM.

RT-PCR Analysis of TRPC Transcripts: The expression of the TRPC channel messages was determined by RT-PCR using sets of primers that spanned the pore-forming region of the TRPC channels. Total RNA isolated from confluent keratinocytes cultured in KGM containing 0.03 mM Ca^{2+} was reverse transcribed into cDNA by reverse transcriptase (Superscript II, Life Science Technology, Goat Island, NY), using an oligo-dT primer. Those cDNAs were then used as templates in PCR amplification by Expand Long Template PCR system (Roche Molecular Biochemicals, Indianapolis, IN). The primer pairs used in PCR were: TRPC1 forward primer 5'-GATTTTGGAAAATTTCTTGGGATGT-3', reverse primer 5'-TTTGTCTTCATGATTTGCTATCA-3'; TRPC3 forward primer 5'-GACATATTCAAGTTCATGGTCCTC-3', reverse primer 5'-ACATCACTGTCATCCTCAATTTTC-3'; TRPC4 forward primer 5'-TCTGCAAATATCTCTGGGAAGAATGCT-3', reverse primer 5'-AAGCTTTGTTTCGTGCAAATTTCCATTC-3'; TRPC5 forward primer 5'-ATCTACTGCCTGGTACTACTAGC-3', reverse primer 5'-TCAGCATGATCGGCAATAAGCTG-3'; TRPC6 forward primer 5'-AAAGACATCTTCAAGTTCATGGTC-3', reverse primer 5'-CACATCAGCGTCATCCTCAATTTTC-3'. The expected size of the PCR products for the TRPCs was between 323 to 415 bp. The PCR DNA products were subcloned into a pCR II vector (Invitrogen, Carlsbad, CA), and their sequences were verified by double-stranded DNA sequencing (Biomedical Resource Center, University of California, San Francisco).

TRPC Anti-sense cDNA Vector Construction and Transfection of Keratinocytes – To construct a vector expressing antisense RNA for human TRPCs, a 374 bp TRPC1 cDNA fragment, a 315 bp TRPC3 cDNA fragment and a 415 bp TRPC4 cDNA fragment were subcloned in an antisense orientation into a mammalian expression vector pcDNA3.1 (Invitrogen) which contains a hygromycin resistance gene. Keratinocytes cultured on glass coverslips were transfected with these human TRPC anti-sense cDNA constructs in KGM containing 0.03 mM calcium using TransIT keratinocyte transfection reagent (Mirus Corp., Madison, WI) according to the manufacturer's protocol. This results in an initial transfection efficiency of 30-40%. Two days after transfection, transfected cells were selected with 100 µg/ml of hygromycin (Roche Molecular Biochemicals) for 24 hours to enrich the transfected cells. In the case of knockdown of PLCγ1, an expression vector containing the full-length PLCγ1 gene in an antisense orientation (Xie and Bikle, 1999) was used for transfection. Transfected cells were selected with 200 µg/ml of neomycin analogue G418 for 48 hours. These selection protocols kill essentially all non-transfected cells. Afterwards, transfected cells were loaded with Fura-2/AM and measured for intracellular Ca²⁺ as described above. For the immunoblotting analyses in the antisense knockdown studies, keratinocytes were grown in 100-mm tissue culture dishes until 35-50% confluent. Cells were then transfected with the desired antisense cDNA construct and selected with G418 or hygromycin B as described above. Cells were then lysed in RIPA buffer and protein content in total lysates was determined by BCA assay. Equivalent amounts of protein (75 µg) were loaded per lane in 5% polyacrylamide gels for immunoblotting analyses as described in the following section.

Immunoblotting Analysis of Membrane Proteins- Preconfluent keratinocytes were cultured in KGM containing 0.03 mM CaCl₂. Crude plasma membranes were isolated from these cultures as described. Briefly, the cells were rinsed twice and scraped into PBS. Cells were then collected by centrifugation and the cell pellet was sonicated in homogenization buffer (containing 20 mM Tris-HCl, pH 7.4, 0.25 M sucrose, 4 mM MgCl₂, 5 mM EGTA, and protease inhibitor cocktail (Complete protease inhibitors, Roche Molecular Biochemicals)). The homogenate was centrifuged at 100,000x g for 30 min, and the resultant pellet was extracted with RIPA buffer (50 mM Hepes, pH 7.4, 1% deoxycholate acid, 1% Triton X-100, 0.1% SDS, 150 mM NaCl, 1 mM EDTA, and Complete protease inhibitor cocktail) for 30 min at 4°C. The soluble membrane protein fraction was separated from the insoluble cytoskeleton fraction by centrifugation at 100,000x g for 30 min. All steps were carried out at 4°C. Protein concentration in these membrane preparations was determined by the BCA Protein Assay Kit (Pierce Corp., Rockford, IL) and equivalent amounts (75 µg) per sample were analyzed as follows. Membrane protein samples were electrophoresed through 5% polyacrylamide gels and electroblotted onto polyvinylidene fluoride membranes (Immobilon-P, 0.45 µm; Millipore Corp., Bedford, MA). After blocking with 5% milk in TBS (20 mM Tris- HCl, pH 7.4, 150 mM NaCl, 1 mM EDTA), the blots were incubated with 1 µg/ml of primary polyclonal antibodies at 4°C overnight. Subsequently, the blots were incubated with 0.1 µg/ml HRP conjugated anti-rabbit IgG antibody (Amersham Pharmacia Biotech, Piscataway, NJ) for 1 h at room temperature. The bound antibody was visualized using the SuperSignal West Dura Chemiluminescent Kit (Pierce Corp.) and subsequent exposure to x-ray film. The data presented are representative of three independent experiments. Anti-TRPC antibodies were purchased from Chemicon International Inc. (Temecula, CA). The corresponding sequences of the peptides used to raise antibodies for

TRPC proteins are as follows: human TRPC1 (Accession P48995) residues 557-571, mouse TRPC3 (Accession Q9QXC1) residues 822-835, mouse TRPC4 (Accession AAC05179) residues 943-958, human TRPC5 (Accession Q9UL62) residues 959-973, mouse TRPC6 (Accession Q61143) residues 24-38. All epitopes are specific for their designated TRPC proteins and not present in any other known proteins. Antibodies against PLC γ 1 and PLC β 1 were obtained from Santa Cruz Biotechnology, Inc. (Santa Cruz, CA).

Co-immunoprecipitation - Preconfluent keratinocytes were scraped and lysed in ice-cold lysis buffer (1% NP-40, 50 mM Tris, pH 7.4, 150 mM NaCl, 5 mM EDTA supplemented with Complete protease inhibitor cocktail) with gentle tumbling for 2 h. The lysates were then centrifuged at 10,000x g for 10 min, and the supernatants were collected and used in immunoprecipitation experiments. All steps were carried out at 4°C. Protein concentration in lysates was determined by the BCA assay. 500 μ g of total proteins was immunoprecipitated by 5 μ g of monoclonal antibody against PLC γ 1 (Santa Cruz Biotechnology, Inc.) or IP $_3$ R (Calbiochem-Novabiochem Corp.), followed by Sepharose-conjugated protein G (ImmunoLink Immobilized Protein G, Pierce Corp.) in 0.5 ml of lysis buffer with gentle tumbling at 4°C overnight. Immunoprecipitants were collected, washed, eluted, and separated by SDS-PAGE. In subsequent immunoblotting analyses, polyclonal antibodies were used to detect the presence of TRPC1, TRPC3 and TRPC4 in the immunoprecipitants.

Fluorescence Immunolocalization of TRPC in Keratinocytes- Keratinocytes were cultured on coverslips and were fixed with 4% paraformaldehyde for 20 min at 37°C. Cells were permeablized with 1% NP-40 in PBS. After blocking with 5% goat serum in PBS with 0.01% Tween-20, cells were incubated with 10 μ g/ml of polyclonal antibodies against TRPC proteins (Chemicon) at 4°C for overnight. Subsequently cells were incubated with fluorescein-

conjugated anti-rabbit IgG antibody (Molecular Probes) at room temperature for 1 h. Finally, coverslips were washed in PBS, mounted on glass slides using Gel-Mount (Biomedica, Foster City, CA) and examined with a Leica TCS NT/SP confocal microscope (Leica Microsystems, Heidelberg, Germany).

RESULTS

SOC in Keratinocytes requires PLC Activity – To activate the SOC in human epidermal keratinocytes, we treated confluent cells grown in 0.03 mM Ca^{2+} with a Ca^{2+} -ATPase inhibitor thapsigargin (TG), and monitored SOC activity by measuring the $[\text{Ca}^{2+}]_i$. In the presence of 0.1 mM EGTA, 1 μM TG elicited an initial increase in $[\text{Ca}^{2+}]_i$ due to passive release from ER Ca^{2+} stores. Subsequent addition of 2 mM extracellular Ca^{2+} induced a second phase of increased $[\text{Ca}^{2+}]_i$ (from 83 ± 10 nM to 256 ± 32 nM, mean \pm SD; n=50) resulting from calcium influx through SOC (**Fig 1a**). As shown in **Fig 1b**, this store-operated Ca^{2+} entry is blocked by application of 75 μM 2-aminoethoxydiphenyl borate (2-APB) (**Fig 1b**), an effective SOC inhibitor independent from its action on IP_3R (Bootman et al, 2002; Ma et al, 2002). This inhibition of SOC activity was reversible. After 2-APB induced blockade of SOC, removal of 2-APB resulted in a slow return of SOC activity over several minutes ($[\text{Ca}^{2+}]_i$ increased from 86 ± 5 nM to 154 ± 23 nM; n=42). Subsequent re-addition of 2-APB while SOC-mediated Ca^{2+} entry was maximally active resulted in rapid termination of SOC activity ($[\text{Ca}^{2+}]_i$ decreased to 94 ± 11 nM) (**Fig 1b**).

To investigate the role of PLC in the activation of SOC in keratinocytes, we blocked the activation of PLC with a membrane-permeable PLC inhibitor, U73122, and examined its impact on SOC activity. We previously found that 15-min pretreatment of keratinocytes with 10 μM U73122 fully prevented the PLC-mediated cellular response to extracellular calcium (Xie and Bialek, 1999). A 15 min preincubation of cells with 10 μM U73122 had no effect on the ability of thapsigargin to release intracellular calcium stores but fully prevented the rise in $[\text{Ca}^{2+}]_i$ upon the addition of extracellular Ca^{2+} ($[\text{Ca}^{2+}]_i$ remained at 103 ± 10 nM; n=37) (**Fig 2a**). On the contrary, pretreatment of cells with 10 μM U73343, an inactive analog of U73122, had no effect on

thapsigargin-induced SOC activity ($[Ca^{2+}]_i$ increased from 100 ± 5 nM to 284 ± 31 nM; n=46) (**Fig 2a**). However, the inhibitory effect of U73122 on Ca^{2+} entry was not due to a direct action on SOC channels. Unlike the SOC inhibitor 2-APB, application of 10 μ M U73122 after induction of the Ca^{2+} entry by thapsigargin and subsequent addition of calcium did not affect the SOC activity. As shown in **Fig 2b**, 7 min after application of 10 μ M U73122, $[Ca^{2+}]_i$ remained at 252 ± 15 nM (n=29), compared with cells treated with vehicle (273 ± 25 nM; n=34). Therefore, the specific blocking effect of U73122 on PLC action prevents store depletion induced Ca^{2+} entry, indicating that PLC is essential for the activation but not for the continuance of SOC activity.

Activation of SOC Is Prevented by Transfection of Human PLC γ 1 Antisense cDNA- Our previous studies had demonstrated that PLC γ 1 is a critical mediator in regulation of calcium signaling in keratinocytes. Knockdown of PLC γ 1 expression blocked the rise in $[Ca^{2+}]_i$ in response to extracellular Ca^{2+} (Xie and Bikle, 1999) or 1,25-dihydroxyvitamin D3 (Xie and Bikle, 2001). To elucidate the role played by PLC γ 1 in SOC function, we blocked the expression of endogenous PLC γ 1 by transfecting confluent keratinocytes with a full-length human PLC γ 1 antisense cDNA construct. The transfectants were selected with the neomycin analog G418 in KGM containing 0.03 mM calcium for 2 days to enrich transfected cells. Protein extracts were prepared and subjected to immunoblotting analyses to examine the level of endogenous PLC γ 1 protein. The expression of PLC γ 1 in the keratinocytes transfected with antisense PLC γ 1 cDNA was fully blocked compared with that in the cells transfected with the control vector (**Fig 3a**). In contrast, the protein level of PLC β 1 was not affected by the antisense PLC γ 1 cDNA (**Fig 3a**), confirming the specificity of the PLC γ 1 antisense construct. We next evaluated the effects of the PLC γ 1 knockdown on SOC activity by comparing the Ca^{2+} entry evoked by TG induced store emptying in the keratinocytes transfected with antisense PLC γ 1

cDNA and in cells transfected with control vector. As shown in **Fig 3b**, the ability of thapsigargin to release intracellular calcium stores was not affected, but the rise in $[Ca^{2+}]_i$ upon the addition of extracellular Ca^{2+} ($[Ca^{2+}]_i$ increased from 107 ± 7 nM to 203 ± 12 nM in cells transfected with vector, n=21) was completely blocked by transfection of the PLC γ 1 antisense cDNA ($[Ca^{2+}]_i$ decreased from 104 ± 3 nM to 92 ± 5 nM; n=12). Our results indicate that PLC γ 1 is involved in the activation of SOC.

PLC activation generates IP $_3$ and DAG. The latter has been reported to directly activate calcium entry mediated by certain TRPC channels (Kiselyov and Muallem, 1999). Therefore, the requirement for PLC in SOC function may signal a need for DAG in addition to an activated IP $_3$ R. To assess the involvement of DAG, we tested the effect of the membrane permeable DAG analog, 1-oleoyl-2-acetyl-*sn*-glycerol (OAG), on SOC activity in keratinocytes. After TG induced store depletion, subsequent addition of 1 mM extracellular Ca^{2+} induced a moderate SOC activity that was not affected by the addition of 0.1% DMSO, the vehicle for OAG (**Fig 4a**). 7 min after application of 10 μ M U73122, $[Ca^{2+}]_i$ remained at 210 ± 20 nM (n=50). However, the store depletion-induced Ca^{2+} entry was gradually blocked by the application of 100 μ M OAG ($[Ca^{2+}]_i$ decreased from 185 ± 8 nM to 101 ± 6 nM; n=50) (**Fig 4b**). Similar results were obtained with 100 μ M 1,2-dioctanoyl-*sn*-glycerol (DOG), another DAG analogue (data not shown). These results indicate that DAG does not promote but rather inhibits the coupling of SOC activity to PLC function.

Expression of TRPC channels in keratinocytes- Substantial evidence has shown that TRPC channels mediate SOC activity (Clapham et al, 2001; Philipp et al, 2000). To investigate the molecular identities of the native SOC, we first examined the expression of TRPCs in keratinocytes by RT-PCR and immunoblotting. Total RNA isolated from confluent

keratinocytes cultured in KGM containing 0.03 mM Ca²⁺ was reverse transcribed into cDNA then used as templates. The expression of the TRPC messages was determined by RT-PCR using sets of primers that spanned the pore-forming region of the channels. The expected size of the PCR product was 374 bp for the TRPC1, 323 bp for TRPC3, 415 bp for TRPC4, 341 bp for TRPC5 and 328 bp for TRPC6. No template (no RT) controls were run for all experiments (**Fig 5a**). Our results showed that TRPC1, TRPC3, TRPC4, TRPC5 and TRPC6 were expressed in preconfluent keratinocytes (**Fig 5a**), whereas no TRPC2 or TRPC7 messages were detected (data not shown). The sequence of each amplified TRPC cDNA was verified by double-stranded DNA sequencing. The primers for TRPC3 also amplified a 450 bp fragment, the identity of which has not been established. To examine the expression of TRPC proteins, membrane proteins were prepared from keratinocytes. Immunoblotting analyses using antibodies recognizing various TRPC proteins detected endogenous TRPC1, TRPC3, TRPC4, TRPC5 and TRPC6 proteins (**Fig 5b**). Their estimated molecular weights were 91 kDa for TRPC1, 135 kDa for TRPC3, 96 kDa for TRPC4, 150 kDa for TRPC5, 125 and 195 kDa for the major and minor band of TRPC6, respectively. No immunoreactive bands were detected when the blots were incubated with the same antibodies preabsorbed with the specific peptides against which they were raised (**Fig 5b**).

Role of TRPC1 and TRPC4 in SOC activity- In addition to store-depletion and receptor activation, DAG has been shown to directly activate members of the TRPC3/6/7 subgroup, but has no effect on other TRPCs, and sometimes prevents the activation of members of TRPC4/5 subgroup (Kiselyov and Muallem, 1999; Venkatachalam et al, 2003). Our results showed that the endogenous SOC activity in keratinocytes could not be enhanced by the application of the DAG analogue (**Fig 4**), suggesting that the members of the TRP3/6/7 subgroup are not involved in mediating SOC activity. Furthermore, we have found that a decrease in SOC activity with

time in culture coincided with a sharp decline in the protein levels of TRPC1 and TRPC4 but not in that of TRPC3, 5, and 6 (data not shown). Therefore we turned our attention to TRPC1 and TRPC4 channels as the likely mediators of store-operated Ca^{2+} entry. We performed fluorescence immunostaining to detect and localize endogenous TRPC1 and TRPC4 proteins in preconfluent keratinocytes cultured in 0.03 mM calcium. Examination by confocal microscopy detected extensive fluorescence signal of TRPC1 and TRPC4 at the plasma membrane and the submembrane area. However, a substantial amount of TRPC protein also localized within an intracellular compartment(s) near the perinuclear region (**Fig 6**). Minimal fluorescence signal was detected when the antibodies were preabsorbed with the antigenic peptides, confirming the specificity of the immunoreaction (**Fig 6**).

To determine whether TRPC1 and TRPC4 mediate SOC activity in keratinocytes, we knocked down the expression of endogenous TRPC1 and TRPC4 by transfection with antisense cDNA constructs, and examined the impact on SOC activity. Preconfluent keratinocytes were transfected with antisense cDNAs and then selected with hygromycin in KGM containing 0.03 mM calcium for 24h to enrich transfected cells. Protein extracts were prepared and subjected to immunoblotting analyses to examine the level of the endogenous TRPC channels. The level of TRPC1 and TRPC4 proteins was greatly reduced in the keratinocytes transfected with antisense cDNA for TRPC1 (anti-TRPC1) and TRPC4 (anti-TRPC4), respectively, as compared with that in the cells transfected with the control vector (**Fig 7a**). In contrast, transfection of antisense TRPC1 or antisense TRPC4 cDNA had minimal effect on the expression of other TRPC proteins (**Fig 7a**). The Ca^{2+} entry in response to TG induced store emptying was reduced as a result of the hygromycin selection of transfected cells (**Fig 7b**, as compared to the Ca^{2+} entry in untransfected cells shown in **Fig 1a**). Nevertheless, knockdown of either TRPC1 or TRPC4

further inhibited SOC activity. As shown in **Fig 7b**, 6 min after application of 2 mM extracellular Ca^{2+} , $[\text{Ca}^{2+}]_i$ increased from 103 ± 2 nM (mean \pm S.D.) to 154 ± 5 nM in the cells transfected with vector (n=14). Transfection of antisense cDNA for TRPC1 and TRPC4 resulted in reduction in SOC activity to ~26% ($[\text{Ca}^{2+}]_i$ increased from 112 ± 2 nM to 125 ± 3 nM; n=11) and ~40% ($[\text{Ca}^{2+}]_i$ increased from 112 ± 6 nM to 132 ± 5 nM; n=10), respectively, of that in cells transfected with vector. Knockdown of both channels led to a complete blockade of SOC activity ($[\text{Ca}^{2+}]_i$ remained unchanged; n=13) (**Fig 7b**). On the other hand, transfection of antisense TRPC3 cDNA decreased the endogenous level of TRPC3 (Fig 7c) but had no effect on the store-depletion induced Ca^{2+} entry (Fig 7d). After application of 2 mM extracellular Ca^{2+} , $[\text{Ca}^{2+}]_i$ increased from 95 ± 5 nM to 206 ± 10 nM in cells transfected with TRPC3 antisense cDNA (n=12), which is comparable with that in cells transfected with vector (92 ± 3 nM to 210 ± 21 nM, n=14). Our results indicate that TRPC1 and TRPC4, but not TRPC3, are the major mediators of SOC activity in these cells.

TRPC1, but not TRPC4, forms protein complexes with PLC γ 1 and IP $_3$ R- Several studies have shown that TRPC proteins can interact with IP $_3$ R through binding to its ligand-binding domain (Boulay et al, 1999; Lockwich et al, 2000; Tang et al, 2001). Furthermore, a recent report demonstrated binding of PLC γ to TRPC3 and TRPC4 (Patterson et al, 2002). In order to investigate possible direct interactions of endogenous TRPC channels with PLC γ and IP $_3$ R, we performed co-immunoprecipitation experiments to test whether antibodies against IP $_3$ R and PLC γ 1 could also bring down TRPC1 and TRPC4. Protein extracts were prepared from confluent keratinocytes cultured in 0.03 mM Ca^{2+} , incubated with monoclonal antibodies against PLC γ 1 or IP $_3$ R, and precipitated with protein G-conjugated beads. Immunoprecipitants were collected and analyzed by immunoblotting to detect the presence of TRPC1 and TRPC4.

Both anti-PLC γ 1 and anti-IP $_3$ R antibodies co-immunoprecipitated TRPC1, but not TRPC4 (**Fig 8a**). In addition, an agarose-conjugated peptide containing the Src homology domains (SH2SH2SH3) of PLC γ 1 was sufficient to bind TRPC1 (**Fig 8a**), as well as TRPC3 (data not shown), indicating that PLC γ 1 interacts with TRPC through its SH2SH2SH3 domain. No band was detected when the blots were incubated with the same antibodies preabsorbed with their respective antigenic peptides, confirming the specificity of the immunoreactions (**Fig 8a**).

We then performed fluorescence immunolocalization of endogenous TRPC1, PLC γ 1 and IP $_3$ R proteins to assess the associations among these calcium regulatory molecules. Examination by confocal microscopy confirmed the association of TRPC1 with PLC γ 1 and IP $_3$ R. As shown in **Fig 8b**, substantial amounts of PLC γ 1 and IP $_3$ R colocalized with TRPC1 near the plasma membrane. Interestingly, TRPC1 protein exhibited little interaction with either IP $_3$ R or PLC γ 1 in the perinuclear compartment, even though all three molecules were present in this region (**Fig 8b**). Our findings indicate that these calcium regulators form a signaling complex at or near the plasma membrane to regulate the SOC activity.

Ca $^{2+}$ has been known to be a factor regulating TRPC function (Zitt et al, 1997) in part by regulating the interaction of TRP channels and IP $_3$ R (Singh et al, 2002; Tang et al, 2001; Zhang et al, 2001). To assess this possibility we compared the interaction of TRPC1 with PLC γ 1 and IP $_3$ R in keratinocytes grown in 0.03 or 1.2 mM extracellular Ca $^{2+}$. As shown in **Fig 9a**, the interactions between TRPC1 and PLC γ 1 or IP $_3$ R were attenuated in the cells treated with 1.2 mM Ca $^{2+}$ for 24h as compared to cells maintained in 0.03 mM Ca $^{2+}$. This reduction in association among these molecules was not due to a decrease in the levels of TRPC1 (**Fig 9a**), PLC γ 1 or IP $_3$ R (**Fig 9b**). In fact, the level of PLC γ 1 was increased by extracellular calcium (**Fig 9b**), consistent with previous studies (Xie and Bikle, 1999). We next compared the thapsigargin

activated SOC activity in these cells. As shown in **Fig 9c**, the basal $[Ca^{2+}]_i$ was higher in cells cultured in 1.2 mM Ca^{2+} for 24h (149 ± 11 nM; n=36) than the cells grown in 0.03 mM Ca^{2+} (101 ± 7 nM; n=28). Keratinocytes cultured in 0.03 mM Ca^{2+} responded to store depletion and subsequent Ca^{2+} application with an active calcium entry ($[Ca^{2+}]_i$ increased from 105 ± 10 nM to 236 ± 25 nM; n=36), whereas the SOC was markedly down regulated in the cells grown in 1.2 mM Ca^{2+} ($[Ca^{2+}]_i$ increased from 99 ± 10 nM to 127 ± 19 nM; n=28) (**Fig 9c**). These results support the conclusion that Ca^{2+} inhibits SOC channel function by down regulation of the interactions between TRPC1, PLC γ 1 and IP $_3$ R, although the mechanism for this remains under investigation.

DISCUSSION

As found for most non-excitabile cells, emptying of intracellular calcium stores in human keratinocytes activates calcium influx across the plasma membrane through the operationally defined SOC (Csernoch et al, 2000; Karvonen et al, 2000). In this study we investigated the molecular basis for the native SOC activity in these cells. We placed our attention on TRPC1 and TRPC4 because their expression patterns correspond to a strong SOC activity in the proliferating keratinocytes (data not shown), though TRPC3, 5 and 6 channels are also expressed in these cells. TRPC1 plays a role in the store-operated Ca^{2+} entry pathway and is ubiquitously expressed in many cell types. Numerous studies have demonstrated that store-operated Ca^{2+} entry was inhibited when the expression of TRPC1 was attenuated or prevented (overview see Beech et al, 2003). The abolition of a store-operated current in adrenal cortical cells by antisense TRPC4 (Philipp et al, 2000) and the lack of store-operated Ca^{2+} entry in endothelial cells from TRPC4 knockout mice (Freichel et al, 2001) indicate that TRPC4 is also one of the SOC components. Similarly, inhibiting the expression of either TRPC1 or TRPC4 attenuated the store-operated Ca^{2+} entry in keratinocytes (**Fig 7a , 7b**), whereas knockdown of the expression of TRPC3 had no effect on SOC function (Fig 7c, 7d), indicating TRPC1 and TRPC4, but not TRPC3, are the subunits of the endogenous SOC in these cells. TRPC1 may not act alone, since it can heteromultimerise with TRPC4 or TRPC5 (but not TRPC3, TRPC6 or TRPC7) in brain, vascular smooth muscle and overexpression systems (Goel et al, 2002; Hofmann et al, 2002). However, the interaction between TRPC1 and TRPC4 or other TRPCs in keratinocytes needs further investigation.

The conformational coupling model (Irvine, 1990; Putney, 1999) for store-operated Ca^{2+} entry hypothesizes that the communication between the ER and plasma membrane involves a

direct protein-protein interaction. Kiselyov et al. demonstrated that the C-terminus of TRPC3 directly interacts with IP₃R, and identified the N-terminal IP₃-binding domain of IP₃R as the minimal portion required to activate TRPC3 (Kiselyov et al, 1998; Kiselyov et al, 1999). Our studies have showed that TRPC1 physically associates with IP₃R (**Fig 8**). These data support the notion that the direct relay of a signal from depleted intracellular calcium stores to calcium channels in the plasma membrane is mediated through protein-protein interactions.

The finding that SOC activation requires both IP₃ and IP₃R (Kiselyov et al, 1998; Zubov et al, 1999) seems counterintuitive to the observation that store-depletion alone in the absence of IP₃ generation is able to activate SOC. However, Putney et al. (Putney et al, 2001) proposed that the SOC-IP₃R complex is located in close proximity to a PLC molecule, perhaps through an organized signaling complex, and the latter produces basal levels of IP₃ sufficient to fulfill the requirement for IP₃ in SOC activation. In epithelial cells and in the mast cell line BRL, inhibition of PLC with the PLC inhibitor U73122 or depletion of phosphatidylinositol-4-phosphate (PIP) with wortmannin abrogated both agonist-induced and store depletion-induced Ca²⁺ entry (Broad et al, 2001; Rosado and Sage, 2000b). Likewise, in keratinocytes when U73122 was administrated at the same concentration reported to inhibit PLC-dependent signaling (Xie and Bikle, 1999), this drug but not the inactive isomer U73343, completely blocked thapsigargin-induced Ca²⁺ entry (**Fig 2a**). A direct inhibitory effect on SOC channels is unlikely, since U73122 failed to block Ca²⁺ entry after the activation of SOC by store depletion (Fig 2b). These results indicate that basal PLC activity is necessary for the activation but not for the maintenance of SOC. While our findings implicate a role for PLC in SOC activation, it appears that this is not due to a requirement for DAG, since application of the DAG analogues, OAG and DOG, inhibited SOC in keratinocytes (**Fig 3**). Although other studies on TRPC

channels indicate that the inhibitory action of DAG on TRPC4- and TRPC5-mediated calcium entry is PKC-dependent (Venkatachalam et al, 2003), it is unclear whether the inhibitory effect of DAG on SOC activity in keratinocytes is mediated by endogenous PKCs.

Recently Patterson et al reported a structural role for PLC γ in mediating agonist-induced calcium entry (Patterson et al, 2002). Overexpression of PLC γ 1 augmented calcium entry induced either by a G protein-coupled receptor agonist or by thapsigargin. This action of PLC γ 1 is independent of its lipase activity but dependent on its Src homology 3 (SH3) domain (Patterson et al, 2002), indicating that PLC γ functions through its ability to associate with other signaling molecules. Yeast two-hybrid analyses and co-immunoprecipitation experiments show that the PLC γ SH3 domain is able to interact with the N-terminus of overexpressed TRPC3 and TRPC4 (Patterson et al, 2002). In this study, co-immunoprecipitation results showed that in keratinocytes the endogenous TRPC1 physically interacts with PLC γ 1 as well as IP $_3$ R (**Fig 8**). Consistent with the finding that the SH3 domain of PLC γ 1 mediates the interaction with SOC, a peptide containing the SH2SH2SH3 domain of PLC γ 1 was able to interact with endogenous TRPC1 (**Fig 8**). The association of native SOC with PLC γ 1 and IP $_3$ R is in agreement with the notion that the SOC-IP $_3$ R complex and PLC γ are located in close proximity via an organized signaling complex. Fluorescence immunolocalization of endogenous TRPC1, PLC γ 1 and IP $_3$ R proteins verified that colocalization of TRPC1 with PLC γ 1 and IP $_3$ R was restricted to the plasma membrane, whereas these proteins showed little interaction in other intracellular locations. In PC12 cells, HEK293 cells and DT40 lymphocytes, knockdown or knockout of PLC γ expression diminished or abrogated agonist-induced calcium entry, whereas store-depletion activated calcium entry was not affected (Patterson et al, 2002). On the contrary, knockdown of PLC γ 1 expression in keratinocytes by anti-sense transfection completely abolished store-depletion

activated calcium entry (**Fig 4**), indicating a requirement for PLC γ 1 in coupling store emptying to activation of SOC. The explanation for these cell specific differences is unclear. Although PLC γ 1 is capable of directly interacting with SOC, we must consider the possibility that an intermediary target may mediate the actions of PLC γ 1. The N-terminus of TRPC proteins contains several ankyrin repeats, a putative interaction site with cytoskeletal proteins (Montell, 2001). PLC γ isoforms have been known to associate with the actin cytoskeleton (Rebecchi and Pentylala, 2000; Rhee, 2001) and have close structural and functional relations to the rho/rac family of GTPases (Hong-Geller and Cerione, 2000; Zenget et al, 2000), which are involved in cytoskeletal reassembly and trafficking (Arriemerlou et al, 2000; Zeng et al, 2000). Since the activation of SOC is sensitive to cytoskeletal rearrangement (Ma et al, 2000; Patterson et al, 1999; Rosado et al, 2000), PLC γ may activate SOC by altering the cytoskeletal-dependent organization of ER and plasma membrane.

Although both TRPC1 and TRPC4 contributed to SOC activity, only TRPC1 directly interacted with PLC γ 1 and IP $_3$ R (**Fig 8**). Many studies support the idea that TRPC1 functions within a signaling complex (overview see Beech et al, 2003). TRPC1 associates with PLC-related signaling proteins such as G $_{\alpha q/11}$, IP $_3$ R, PLC, plasma membrane Ca $^{2+}$ -ATPase, caveolin-1 (Lockwich et al, 2000) and calmodulin (Singh et al, 2002). Since TRPC1 has the ability to hetermultimerise with TRPC4, it may also function as a regulator for TRPC4. It is possible that TRPC4 activity is gated by PLC γ 1 and IP $_3$ R via TRPC1. This hypothesis would explain the complete abolition of store-operated Ca $^{2+}$ entry by antisense PLC γ 1 (**Fig 4**), and the observation that antisense TRPC1 inhibited store-operated Ca $^{2+}$ entry more effectively than antisense TRPC4 (**Fig 7b**). In this study we also demonstrated that TRPC3, which is strongly expressed in keratinocytes, is not involved in mediating store-depletion activated calcium entry (**Fig 7d**), even

though a weak interaction of TRPC3 with a peptide containing the PLC γ 1 SH2SH2SH3 domain was detected (data not shown). The physiologic significance of this observation in keratinocytes is unclear but is consistent with the findings of Patterson et al. (Patterson et al, 2002) in other cells, showing an interaction between PLC γ and TRPC3.

Many studies have demonstrated the high sensitivity of SOC to inhibition by $[Ca^{2+}]_i$, a negative feedback mechanism for controlling $[Ca^{2+}]_i$ (Krause et al, 1999; Parekh and Penner, 1997). Activation of SOC occurs maximally when $[Ca^{2+}]_i$ is buffered to a low level, and reduction in $[Ca^{2+}]_i$ is sufficient to activate the channels (Kerschbaum and Cahalan, 1999; Krause et al, 1999). In this study we also observed an inhibition of SOC activity by elevated $[Ca^{2+}]_i$ in keratinocytes and demonstrated that the effect of $[Ca^{2+}]_i$ on SOC was due to decreased association of TRPC1 with PLC γ 1 and IP $_3$ R (**Fig 9**). Although the mechanism(s) by which $[Ca^{2+}]_i$ alters these interactions remains unclear, our findings support the conclusion that the activity of SOC depends on the interactions of TRPC channels with PLC γ and IP $_3$ R.

ACKNOWLEDGMENTS

We thank Dr. Zhongjian Xie for providing the PLC γ 1 anti-sense cDNA construct and Dr. Theodora M. Mauro for helpful discussions. We appreciate the technical help from Sally Pennypacker and the Cell Culture and Tissue Preparation Core of our program project for preparing human epidermal keratinocytes. This work was supported by a Merit Review Award from the Department of Veterans Affairs and grants PO1-AR39448, RO1-AR38386 and RO1-AG21353 from the National Institutes of Health.

REFERENCES

1. Arrieumerlou C, Randriamampita C, Bismuth G, Trautmann A: Rac is involved in early TCR signaling. *J Immunol* 165: 3182-3189, 2000
2. Beech DJ, Xu SZ, McHugh D, Flemming MR: TRPC1 store-operated cationic channel subunit. *Cell Calcium* 33: 433-440, 2003
3. Berridge MJ: Inositol trisphosphate and calcium signaling. *Nature* 361: 315-325, 1993
4. Berridge MJ: Elementary and global aspects of calcium signalling. *J Physiol* 499: 291-306, 1997
5. Bootman MD, Collins TJ, Mackenzie L, Roderick HL, Berridge MJ, Peppiatt CM: 2-Aminoethoxydiphenyl borate (2-APB) is a reliable blocker of store-operated Ca^{2+} entry but an inconsistent inhibitor of InsP_3 -induced Ca^{2+} release. *FASEB J* 16: 1145-1150, 2002
6. Boulay G, Brown D, Qin N, et al: Modulation of Ca^{2+} entry by polypeptides of the inositol 1,4,5-trisphosphate receptor (IP_3R) that bind transient receptor potential (TRP): evidence for roles of TRP and IP_3R in store depletion-activated Ca^{2+} entry. *Proc Natl Acad Sci U S A* 96: 14955-14960, 1999
7. Broad LM, Braun FJ, Lievremont JP, Bird JG, Kurosaki T, Putney JW: Role of the phospholipase C-inositol 1,4,5-trisphosphate pathway in calcium release-activated calcium current and capacitative calcium entry. *J Biol Chem* 276: 15945-15952, 2001
8. Brough GH, Wu S, Cioffi D, Moore TM, Li M, Dean N, Stevens T: Contribution of endogenously expressed Trp1 to a Ca^{2+} -selective, store-operated Ca^{2+} entry pathway. *FASEB J* 15: 1727-1738, 2001
9. Clapham DE, Runnels LW, Strubing C: The TRP ion channel family. *Nature Rev Neurosci* 2: 387-396, 2001

10. Csernoch L, Hunyadi J, Kovacs L: Calcium release activated calcium entry in a human skin derived cell line (HaCaT). *Exp Dermatol* 9: 200-205, 2000
11. Freichel M, Suh SH, Pfeifer A, et al: Lack of an endothelial store-operated Ca^{2+} current impairs agonist-dependent vasorelaxation in $\text{TRP4}^{-/-}$ mice. *Nature Cell Biol* 3: 121-127, 2001
12. Gibson DF, Ratnam AV, Bikle DD: Evidence for separate control mechanisms at the message, protein, and enzyme activation levels for transglutaminase during calcium-induced differentiation of normal and transformed human keratinocytes. *J Invest Dermatol* 106: 154-161, 1996
13. Goel M, Sinkins WG, Schilling WP: Selective association of TRPC channel subunits in rat brain synaptosomes. *J Biol Chem* 277: 48303-48310, 2002
14. Hofmann T, Schaefer M, Schultz G, Gudermann T: Transient receptor potential channels as molecular substrates of receptor-mediated cation entry. *J Mol Med* 78: 14-25, 2000
15. Hofmann T, Schaefer M, Schultz G, Gudermann T: Subunit composition of mammalian transient receptor potential channels in living cells. *Proc Natl Acad Sci USA* 90: 7461-7466, 2002
16. Hong-Geller E, Cerione RA: Cdc42 and Rac stimulate exocytosis of secretory granules by activating the IP_3 /calcium pathway in RBL-2H3 mast cells. *J Cell Biol* 148: 481-494, 2000
17. Irvine RF: "Quantal" Ca^{2+} release and the control of Ca^{2+} entry by inositol phosphates- a possible mechanism. *FEBS Lett* 263: 5-9, 1990
18. Kanki H, Kinoshita M, Akaike A, Satoh M, Mori Y, Kaneko S: Activation of inositol 1,4,5-trisphosphate receptor is essential for the opening of mouse TRP5 channels. *Mol Pharmacol* 60: 989-998, 2001

19. Karvonen SL, Korkiamaki T, Yla-Qutinen H, et al: Psoriasis and altered calcium metabolism: downregulated capacitative calcium influx and defective calcium-mediated cell signaling in cultured psoriatic keratinocytes. *J Invest Dermatol* 114: 693-700, 2000
20. Kerschbaum HH, Cahalan MD: Single-channel recording of a store-operated Ca^{2+} channel in Jurkat T lymphocytes. *Science* 283: 836-839, 1999
21. Kiselyov KI, Mignery GA, Zhu MX, Muallem S: The N-terminal domain of the IP_3 receptor gates store-operated hTrp3 channels. *Mol Cell* 4: 423-429, 1999
22. Kiselyov K, Muallem S: Fatty acids, diacylglycerol, $\text{Ins}(1,4,5)\text{P}_3$ receptors and Ca^{2+} influx. *Trends Neurosci* 22: 334-337, 1999
23. Kiselyov K, Xu X, Mozhayeva G, et al: Functional interaction between InsP_3 receptors and store-operated hTrp3 channels. *Nature* 396: 478-482, 1998
24. Krause E, Schmid A, Gonzalez A, Schulz I: Low cytoplasmic Ca^{2+} activates I(CRAC) independently of global Ca^{2+} store depletion in RBL-1 cells. *J Biol Chem* 274: 36957-36962, 1999
25. Li HS, Xu XZ, Montell C: Activation of a TRPC3-dependent cation current through the neurotrophin BDNF. *Neuron* 24: 261-273, 1999
26. Liu X, Wang W, Singh BB, et al: Trp1, a candidate protein for the store-operated Ca^{2+} influx mechanism in salivary gland cells. *J Biol Chem* 275: 3403-3411, 2000
27. Lockwich TP, Liu X, Singh BB, Jadlowiec J, Weiland S, Ambudkar IS: Assembly of Trp1 in a signaling complex associated with caveolin-scaffolding lipid raft domains. *J Biol Chem* 275: 11934-11942, 2000

28. Ma HT, Patterson RL, van Bossum DB, Birnbaumer L, Mikoshiba K, Gill DL: Requirement of the inositol trisphosphate receptor for activation of store-operated Ca^{2+} channels. *Science* 287: 1647-1651, 2000
29. Ma HT, Venkatachalam K, Parys JB, Gill DL: Modification of store-operated channel coupling and inositol trisphosphate receptor function by 2-aminoethoxydiphenyl borate in DT40 lymphocytes. *J Biol Chem* 277: 6915-6922, 2002
30. Montell C: Physiology, phylogeny, and functions of the TRP superfamily of cation channels. Science's STKE, <http://stke.sciencemag.org/cgi/content/full/sigtrans;2001/90/re1>, 2001
31. Mori Y, Wakamori M, Miyakawa T, et al: Transient receptor potential 1 regulates capacitative Ca^{2+} entry and Ca^{2+} release from endoplasmic reticulum in B lymphocytes. *J Exp Med* 195: 673-681, 2002
32. Parekh AB, Penner R: Store depletion and calcium influx. *Physiol Rev* 77: 901-930, 1997
33. Patterson RL, van Rossum DB, Ford DL, et al: Phospholipase C- γ is required for agonist-induced Ca^{2+} entry. *Cell* 111: 529-541, 2002
34. Patterson RL, van Rossum D B, Gill DL: Store operated Ca^{2+} entry: evidence for a secretion-like coupling model. *Cell* 98: 487-499, 1999
35. Philipp S, Trost C, Warnat J, et al: TRP4 (CCE1) is part of native Ca^{2+} release-activated Ca^{2+} -like channels in adrenal cells. *J Biol Chem* 275: 23965-23972, 2000
36. Putney JW Jr: A model for receptor-regulated calcium entry. *Cell Calcium* 7: 1-12, 1986
37. Putney JW Jr: "Kissin' cousins": intimate plasma membrane-ER interactions underlie capacitative calcium entry. *Cell* 99: 5-8, 1999
38. Putney JW Jr, Broad LM, Braun FJ, Lievremont JP, Bird JG: Mechanisms of capacitative calcium entry. *J Cell Sci* 114: 2223-2229, 2001

39. Rebecchi M.J, Pentylala SN: Structure, function, and control of phosphoinositide-specific phospholipase C. *Physiol Rev* 80: 1291-1335, 2000
40. Rhee SG: Regulation of phosphoinositide-specific phospholipase C. *Annu Rev Biochem* 70: 281-312, 2001
41. Rosado JA, Jenner S, Sage SO: A role for the actin cytoskeleton in the initiation and maintenance of store-mediated calcium entry in human platelets. Evidence for conformational coupling. *J Biol Chem* 275: 7527-7533, 2000
42. Rosado JA, Sage SO: Coupling between inositol 1,4,5-trisphosphate receptors and human transient receptor potential channel 1 when intracellular Ca²⁺ stores are depleted. *Biochem J* 350: 631-635, 2000
43. Rosado JA, Sage SO: Phosphoinositides are required for store-mediated calcium entry in human platelets. *J Biol Chem* 275: 9110-9113, 2000
44. Singh BB, Liu X, Tang J, Zhu MX, Ambudkar IS: Calmodulin regulates Ca²⁺-dependent feedback inhibition of store-operated Ca²⁺ influx by interaction with a site in the C terminus of TrpC1. *Mol Cell* 9: 739-750, 2002
45. Tang J, Lin Y, Zhang Z, Tikunova S, Birnbaumer L, Zhu MX: Identification of common binding sites for calmodulin and inositol 1,4,5-trisphosphate receptors on the carboxyl termini of Trp channels. *J Biol Chem* 276: 21303-21310, 2001
46. Trebak M, Vazquez G, Bird JG, Putney JW: The TRPC3/6/7 subfamily of cation channels. *Cell Calcium* 33: 451-461, 2003
47. Vazquez G, Lievremont JP, Bird JG, Putney JW Jr: Human Trp3 forms both inositol trisphosphate receptor-dependent and receptor-independent store-operated cation channels in DT40 avian B lymphocytes. *Proc Natl Acad Sci U S A* 98: 11777-11782, 2001

48. Venkatachalam K, Ma HT, Ford DL, Gill DL: Expression of functional receptor-coupled TRPC3 channels in DT40 triple InsP₃ receptor-knockout cells. *J Biol Chem* 276: 33980-33985, 2001
49. Venkatachalam K, van Bossum DB, Patterson RL, Ma HT, Gill DL: The cellular and molecular basis of store-operated calcium entry. *Nat Cell Biol* 4: E263-E272, 2002
50. Venkatachalam K, Zheng F, Gill DL: Regulation of canonical transient receptor potential (TRPC) channel function by diacylglycerol and protein kinase C. *J Biol Chem* 278: 29031-29040, 2003
51. Xie Z, Bikle DD: Phospholipase C-gamma1 is required for calcium-induced keratinocyte differentiation. *J Biol Chem* 274: 20421-20424, 1999
52. Xie Z, Bikle DD: Inhibition of 1,25-dihydroxyvitamin D-induced keratinocyte differentiation by blocking the expression of phospholipase C-γ1. *J Invest Dermatol* 117: 1250-1254, 2001
53. Zeng L, Sachdev P, Yan L, et al: Vav3 mediates receptor protein tyrosine kinase signaling, regulates GTPase activity, modulates cell morphology, and induces cell transformation. *Mol Cell Biol* 20: 9212-9224, 2000
54. Zhang Z, Tang J, Tikunova S, et al: Activation of Trp3 by inositol 1,4,5-trisphosphate receptors through displacement of inhibitory calmodulin from a common binding domain. *Proc Natl Acad Sci USA* 98: 3168-3173, 2001
55. Zitt C, Obukhov AG, Strubing C, Zobel A, Kalkbrenner F, Luckhoff A, Schultz G: Expression of *TRPC3* in Chinese hamster ovary cells results in calcium-activated cation currents not related to store depletion. *J Cell Biol* 138: 1333-1341, 1997

56. Zubov AI, Kaznacheeva EV, Alexeeno VA, Kiselyov K, Muallem S, Mozhayeva G:
Regulation of the miniature plasma membrane Ca^{2+} channel I_{min} by inositol 1,4,5-
trisphosphate receptors. *J Biol Chem* 274: 25983-25985, 1999

FIGURE LEGENDS

Figure 1. The SOC inhibitor 2-APB blocked store-operated Ca^{2+} entry. Fura-2 loaded epidermal keratinocytes on glass coverslips were treated with 1 μM thapsigargin (TG) to deplete intracellular calcium stores in the presence of 0.1 mM EGTA. After $[\text{Ca}^{2+}]_i$ returned to baseline, 2 mM extracellular Ca^{2+} was applied to induce Ca^{2+} entry (a). 75 μM 2-APB was added after application of thapsigargin for the time indicated by the bars (b). The trace shown in the figure represents the average $[\text{Ca}^{2+}]_i$ of 42-50 individual keratinocytes during recording. 2-APB abolished the rise in $[\text{Ca}^{2+}]_i$ in response to 2 mM extracellular Ca^{2+} after store depletion. Calcium entry resumed after 2-APB was removed and was blocked by re-addition of 2-APB. The results are representative of three experiments.

Figure 2. The phospholipase C inhibitor U73122 blocked the activation of store-operated Ca^{2+} entry. Keratinocytes, loaded with Fura-2, were treated with 1 μM thapsigargin (TG) to deplete intracellular calcium stores and to activate SOC. Nominally calcium-free bath was replaced with buffer containing 2 mM calcium as indicated. (a) Cells were pretreated with a PLC inhibitor U73122 (10 μM , n=37) or its inactive analogue U73343 (10 μM , n=46) for 15 min before $[\text{Ca}^{2+}]_i$ measurement. U73122 abolished the activation of SOC by store depletion. (b) Seven minutes after application of 2 mM calcium, 0.5% ethanol (vehicle, n=34) or U73122 (10 μM , n=29) was added to cells. The trace shown in the figure represents the average $[\text{Ca}^{2+}]_i$ of 29-46 individual keratinocytes during recording. The results are representative of three experiments.

Figure 3. Transfection of a PLC γ 1 antisense cDNA construct decreased the expression of endogenous PLC γ 1 and prevented SOC activation. Keratinocytes were cultured, transfected with a full-length PLC γ 1 antisense cDNA construct (anti-PLC γ 1) or pcDNA3.1 (vector), and

selected by G418 as described under “Experimental Procedures”. Total proteins were isolated from transfected cells and analyzed by immunoblotting (**a**). The expression of PLC γ 1, but not PLC β 1, was decreased by the PLC γ 1 antisense construct (anti-PLC γ 1). Transfected cells were loaded with Fura-2, and calcium entry activated by thapsigargin (TG) induced store emptying was measured (**b**). The trace shown represents the average $[Ca^{2+}]_i$ of 12-21 individual keratinocytes during recording. Transfection of PLC γ 1 antisense construct (anti-PLC γ 1) abolished the activation of calcium entry by store depletion. Data are representative of three different experiments.

Figure 4. Inhibition of SOC by the diacylglycerol analogue OAG. Calcium entry was activated by thapsigargin (TG) induced store emptying. Subsequent addition of 1 mM extracellular Ca^{2+} was not affected by application of 0.1% DMSO (vehicle, n=50) (**a**), but was blunted by application of 100 μ M of 1-oleoyl-2-acetyl-sn-glycerol (OAG, n=50) (**b**).

Figure 5. Expression of TRP channels in human keratinocytes. RT-PCR (**a**) and immunoblotting (**b**) analyses were performed to detect TRPC transcripts and proteins, respectively. Total RNA was collected from confluent human keratinocytes grown in medium containing 0.03 mM, reverse-transcribed into cDNA and used as template in PCR amplification (**a**). Specific primer pairs were used to amplify TRPCs. No template control (no RT) was run for all experiments. To examine TRPC protein expression, the membrane proteins were isolated from keratinocytes grown in medium containing 0.03 mM Ca^{2+} and subjected to immunoblotting analyses (**b**). The expression of TRPC proteins was detected by specific antibody against each TRPC as indicated. The specificity of the immunoreactions was confirmed by incubation of the blots with the antibodies preabsorbed with the specific peptides against which they were raised.

Figure 6. Fluorescence immunolocalization of the TRPC1 and TRPC4 protein endogenously expressed in human keratinocytes. Keratinocytes were fixed, permeabilized and incubated with polyclonal antibody against TRPC1 or TRPC4, followed by fluorescein-conjugated anti-rabbit IgG. Fluorescent signals were detected with a confocal microscope. TRPC1 and TRPC4 proteins were detected on the plasma membrane as well as within the cytoplasm in the perinuclear region. Data are representative of five independent experiments.

Figure 7. Transfection of anti-sense cDNA constructs for TRPC1 and TRPC4 inhibited SOC activity in keratinocytes. Keratinocytes were cultured, transfected with the control vector or the anti-sense cDNA constructs for TRPC1 (anti-TRPC1), TRPC4 (anti-TRPC4) (**a, b**) and TRPC3 (anti-TRPC3) (**c, d**), and selected by hygromycin as described under “Experimental Procedures”. Total proteins were isolated from transfected cells and analyzed by immunoblotting (**a, c**). Note that the expression of endogenous TRPC1, TRPC4 and TRPC3 was reduced by their respective antisense constructs, whereas the levels of other TRPC proteins were minimally affected. Transfected cells were loaded with Fura-2, and thapsigargin-induced store-operated calcium entry was measured (**b, d**). The trace shown represents the average $[Ca^{2+}]_i$ of 12-14 individual keratinocytes during recording. Knockdown of either TRPC1 or TRPC4 partially suppressed, while knockdown of both channels completely abrogated the activation of calcium entry by store depletion. Knockdown of TRPC3 had no effect on SOC activity. The results are representative of three experiments.

Figure 8. TRPC1 forms a protein complex with PLC γ 1 and IP $_3$ R. Co-immunoprecipitation (**a**) revealed direct interaction of TRPC1 with PLC γ 1 and IP $_3$ R. Protein extracts collected from confluent keratinocytes grown in 0.03 mM Ca^{2+} were immunoprecipitated (IP) with monoclonal antibodies against PLC γ 1 or IP $_3$ R, followed by sepharose-conjugated protein G, or

with an agarose-conjugated peptide containing the SH2SH2SH3 domain of PLC γ 1. Immunoprecipitants were then analyzed by immunoblotting using polyclonal antibodies against TRPC1 and TRPC4. One fifth of the amount of protein used in immunoprecipitations was included in the blots as a control (input). The same antibodies preabsorbed with their specific antigenic peptides were used to confirm the specificity of the immunoreactivities of these bands. Fluorescence immunolocalization (**b**) demonstrated colocalization of TRPC1 with PLC γ 1 and IP $_3$ R. Preconfluent keratinocytes cultured in 0.03 mM Ca $^{2+}$ were incubated with a polyclonal antibody for TRPC1 and a monoclonal antibody against PLC γ 1 or IP $_3$ R, followed by fluorescein-conjugated anti-rabbit IgG and Texas Red-conjugated anti-mouse IgG, respectively. Fluorescent signals were detected with a confocal microscope. TRPC1 colocalized with PLC γ 1 and IP $_3$ R near the plasma membrane. TRPC1, PLC γ 1 and IP $_3$ R were also detected in perinuclear locations, but little interaction was found among these molecules in this region. Data are representative of three different experiments.

Figure 9. Calcium decreased SOC activity and complex formation of TRPC1, PLC γ 1 and IP $_3$ R. Preconfluent keratinocytes were cultured in 1.2 mM Ca $^{2+}$ or maintained in 0.03 mM Ca $^{2+}$ for 24h. Protein extracts were collected and immunoprecipitated as described in the Fig 8 legend. Immunoprecipitants were analyzed by immunoblotting for TRPC1 (**a**). One fifth of the amount of protein used in the immunoprecipitations was included in the blots as a control (input). The levels of endogenous PLC γ 1 and IP $_3$ R were also examined by immunoblotting of protein extracts from these cells (**b**). The interactions of TRPC1 with PLC γ 1 and IP $_3$ R were markedly reduced in cells treated with 1.2 mM Ca $^{2+}$, even though the levels of TRPC1, PLC γ 1 and IP $_3$ R were not decreased in these cells. Thapsigargin (TG)-induced SOC was measured in

keratinocytes cultured in 1.2 mM Ca²⁺ (n=28) or in 0.03 mM Ca²⁺ (n=36) for 24h (c). Results are representative of two independent experiments.




Figure 1





Figure 2




Figure 3





Figure 4

a



b




Figure 5




Figure 6




Figure 7




Figure 8




Figure 9