

Lawrence Berkeley National Laboratory

Recent Work

Title

The Excess of Negative Over Positive Mesons Produced by High Energy Photons

Permalink

<https://escholarship.org/uc/item/2831v85j>

Authors

Brueckner, K.A.
Goldberger, M.L.

Publication Date

1949-10-03

UNIVERSITY OF
CALIFORNIA

*Radiation
Laboratory*

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

BERKELEY, CALIFORNIA

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UNCLASSIFIED

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

THE EXCESS OF NEGATIVE OVER POSITIVE MESONS
PRODUCED BY HIGH ENERGY PHOTONS

K. A. Brueckner and M. L. Goldberger

October 3, 1949

Berkeley, California

-2-

THE EXCESS OF NEGATIVE OVER POSITIVE MESONS
PRODUCED BY HIGH ENERGY PHOTONS

K. A. Brueckner and M. L. Goldberger*

Radiation Laboratory
University of California
Berkeley, California

October 3, 1949

Mesons produced by the high energy photon beam from the University of California Radiation Laboratory 330 Mev synchrotron are found to show an excess of negatives over positives¹. With a carbon target, observing

¹ E. M. McMillan, J. M. Peterson, R. S. White, to be published in Science

mesons in the energy range 30-130 Mev at 90° to the photon beam, the ratio of negative to positive mesons is 1.7 ± 0.2 with no significant energy dependence.

A simple classical argument can be made to give an understanding of the reason for the negative excess. The photon can interact directly with the meson and proton through the current coupling

$$(\vec{j} \cdot \vec{A}) (\text{meson}) + (\vec{j} \cdot \vec{A}) (\text{proton})$$

The meson contribution is symmetrical for the production of positive and negative mesons. However, when positive mesons are produced, the proton is the initial nucleon at rest, and its current is zero. When negative mesons are produced, the proton is the final recoil nucleon giving a current contribution. Therefore, the cross-sections for the production of positive and negative mesons are in the ratio

$$\frac{\sigma(\text{positives})}{\sigma(\text{negatives})} = \left[\frac{(\vec{j} \cdot \vec{A}) (\text{meson})}{(\vec{j} \cdot \vec{A}) (\text{meson}) + (\vec{j} \cdot \vec{A}) \text{recoil proton}} \right]^2$$

* Now at Dept. of Physics, Mass. Inst. of Technology, Cambridge, Mass.

-3-

The current interaction is

$$\vec{j} \cdot \vec{A} = \frac{e \vec{v} \cdot \vec{A}}{1 - \frac{v}{c} \cos \theta}$$

\vec{v} = velocity of particle

\vec{A} = vector potential

$\cos \theta$ = angle between direction of particle and photon

This differs from the non-relativistic expression, $e \vec{v} \cdot \vec{A}$, the factor $1 - \frac{v}{c} \cos \theta$ in the denominator taking account of the retardation effects in the interaction of charge with the electromagnetic field. Inserting this current expression and using overall energy and momentum conservation, the positive to negative ratio can be written

$$\frac{\sigma \text{ (positives)}}{\sigma \text{ (negatives)}} = \left[1 - \frac{\epsilon}{mc^2} \left(1 - \frac{v}{c} \cos \theta \right) \right]^2$$

ϵ = meson energy including rest energy

v = meson velocity

θ = angle between direction of meson and photon

m = nucleon rest mass

Further calculations have been carried out using standard perturbation theory for scalar and pseudoscalar mesons to the lowest order in the coupling constants g and e , treating the nucleons as Dirac particles and taking into account the effects of the nucleon recoil. The result of these calculations, for the ratio of the cross-sections for positive and negative mesons, is exactly the same as that derived by the above simple considerations.

Similar calculations for vector mesons with vector coupling to the nucleon field are complicated by the strong magnetic moment interaction of the vector particle with the e.m. field. The ratio of negative to positive mesons is similar to that for the scalar meson fields but is somewhat larger.

The effects of the Coulomb field of the nucleus on the production of mesons have also been investigated and found to be less than 5 percent for mesons with energies above 30 Mev.

The ratio of the cross-sections for the production of negative and positive mesons given by (1) varies from 1.55 at 40 Mev to 1.83 at 100 Mev, at 90° to the photon beam. This agrees, within the probable error with the experimentally observed ratio. Since the positive-negative ratio depends in a quite direct way on the currents carried by the mesons and nucleons, a more accurate determination of the ratio and its energy dependence could provide valuable evidence concerning the magnetic moments of the particles.