

UC Riverside

Journal of Citrus Pathology

Title

A grower question: So we are controlling the Asian Citrus Psyllid, but are we doing it well enough?

Permalink

<https://escholarship.org/uc/item/29j5p26z>

Journal

Journal of Citrus Pathology, 1(1)

Authors

Irey, M.
Gast, T.
Hou, H.

Publication Date

2014

DOI

10.5070/C411025087

Copyright Information

Copyright 2014 by the author(s). This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

6.11

A grower question: So we are controlling the Asian Citrus Psyllid, but are we doing it well enough?

Irey, M.¹, Gast, T.², and Hou, H.¹

¹United States Sugar Corporation, Clewiston, FL

²Southern Gardens Citrus, Clewiston, FL

Since Huanglongbing (HLB) was first found in Florida, growers have recognized the importance of controlling the Asian citrus psyllid (ACP) that vectors the disease. ACP management programs have been developed and applied over large acreages, yet in many instances, the incidence of the disease continues to increase. When asked, the growers invariably report that they are controlling ACP and it is obvious that the control programs are reducing the level of ACP in the groves. However there is an outstanding question as to whether the levels of ACP are being reduced enough to limit the spread of HLB. In most cases, growers have not implemented ACP scouting programs so the level of infestation is based on rough grove surveys and feelings. Even if scouting programs are in place, there are no established data-based thresholds to trigger additional applications of pesticides resulting in a “chasing” approach where applications are made after an infestation is found. Southern Gardens has had an ACP and HLB scouting program in place for many years. By comparing the annual ACP levels to resulting HLB infection succeeding years, it may be possible to elucidate thresholds and then design ACP control programs that effectively limit the spread of HLB. Data will be presented for three years of ACP scouting and four years of HLB scouting in an attempt to show the level of control that must be achieved in order to limit the spread of HLB. The bottom line is that control thresholds are very low and the level of ACP control that must be achieved in order to limit spread HLB is much lower than most growers realize.