

UC Irvine

UC Irvine Previously Published Works

Title

Clinical features in children with nevoid basal cell carcinoma syndrome.

Permalink

<https://escholarship.org/uc/item/29v8c5xm>

Journal

PEDIATRIC RESEARCH, 39(4)

ISSN

0031-3998

Authors

Kimonis, VE
Goldstein, AM
Pastakia, B
[et al.](#)

Publication Date

1996

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Clinical Features In Children With Nevoid Basal Cell Carcinoma Syndrome. • 863

Virginia E Kimonis, Alisa M Goldstein, Behram Pastakia, Michael L Yang, Ronald Kase, John J DiGiovanna, Allen E Bale & Sherri J Bale
Pediatric Research volume 39, page146 (1996)

Nevoid Basal Cell Carcinoma Syndrome (NBCC), an autosomal dominant disorder linked to 9q22.3-q31, is characterized by multiple basal cell carcinomas (BCCs), keratocysts of the jaw, palmar/plantar pits and calcification of the falx cerebri. Twenty-six families including 105 persons with NBCC participated in a linkage and clinical study at the NIH since 1985. Of these, 30 were children aged from 4 mo. to 19 yr. (mean 11.3 yr., M:F ratio 1:1). On comparing the frequency of the major clinical features, 55% of children and 91% of adults had at least one BCC. The number of BCCs ranged from 1- > 1000 (median 1) & 1-500 (median 31.5) respectively. Jaw cysts occurred in 60% of children and 81% of adults, the no. of cysts ranging from 1-6 (median 3) and 1-28 (median 5) respectively. Lifetable analysis of adults and children indicated that 50% developed their first BCCs by the age of 21.5 yr. and their first jaw cyst by the age of 15 yr. Palmar/plantar pits were observed at approximately the same frequency in children (89%) and adults (87%). Calcification of the falx cerebri was seen in 37% children and 78% adults. On comparison with their unaffected siblings (N=25) features seen exclusively in affected children were frontal bossing (44.8%), hypertelorism (38.5%), Sprengels deformity (17.9%), pectus deformity (14.8%), and cleft lip/palate (6.6%). Sixty percent of affected and 8% of unaffected children had macrocephaly (HC > 2SD) (p < 0.01). Radiological features seen exclusively in affected children were calcification of the tentorium cerebellum (13%), bifid ribs (29%), hemivertebrae (9%), fusion of the vertebral bodies (9%), and flame shaped lucencies of the hand (16%). This study indicates that many of the major features (jaw cysts, BCCs, and falx calcification) develop over time. Therefore a diagnosis can be established in young persons by the presence of palmar pits, and evaluation of the above mentioned clinical and radiological features of NBCC.

Author information

Author notes: Virginia E Kimonis: Dept. of Pediatrics, Southern Illinois University School of Medicine, Springfield, IL.

Authors and Affiliations: IRP, NIAMS.

Virginia E Kimonis, Michael L Yang, John J DiGiovanna & Sherri J Bale

Genetic Epidemiology Branch, NCI, Bethesda, MD

Alisa M Goldstein

Diagnostic Radiology, Clinical Center, NIH & Veterans Administration Hospital, Washington, DC

Behram Pastakia

Westat Inc., Rockville, MD

Ronald Kase

Yale University School of Medicine, New Haven, CT

Allen E Bale

Additional information

(Spon. by Donald N. Medearis, M.D.)

Rights and permissions

Reprints and Permissions

About this article

Cite this article

Kimonis, V., Goldstein, A., Pastakia, B. et al. CLINICAL FEATURES IN CHILDREN WITH NEVOID BASAL CELL CARCINOMA SYNDROME. • 863. *Pediatr Res* 39, 146 (1996).

<https://doi.org/10.1203/00006450-199604001-00885>

Download citation

Issue Date

01 April 1996

DOI

<https://doi.org/10.1203/00006450-199604001-00885>