

Lawrence Berkeley National Laboratory

LBL Publications

Title

Spontaneous Fission of U234, Pu236, Cm240, and Cm244

Permalink

<https://escholarship.org/uc/item/2cc6x15q>

Authors

Ghiorso, A

Higgins, G H

Larsh, A E

et al.

Publication Date

1952-04-01

UCRL-1772
Unclassified-Chemistry Distribution

UNCLASSIFIED

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

SPONTANEOUS FISSION OF U^{234} , Pu^{236} , Cm^{240} , AND Cm^{244}

A. Ghiorso, G. H. Higgins, A. E. Larsh, G. T. Seaborg, and S. G. Thompson

April 21, 1952

Berkeley, California

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

SPONTANEOUS FISSION OF U^{234} , Pu^{236} , Cm^{240} , AND Cm^{244}

A. Ghiorso, G. H. Higgins, A. E. Larsh, G. T. Seaborg, and S. G. Thompson
Radiation Laboratory and Department of Chemistry
University of California, Berkeley, California

April 21, 1952

In a recent communication commenting on the mechanism of fission we called attention to the simple exponential dependence of spontaneous fission rate on Z^2/A and to the effect of an odd nucleon in slowing the fission process.¹ Since it is of interest to test further the simple correlation of the spontaneous fission rate for even-even nuclides with Z^2/A , a further number of such rates have been determined.


The spontaneous fission rates were measured by placing the chemically purified samples on one electrode of a parallel plate ionization chamber, filled with a mixture of argon and carbon dioxide, which was connected with an amplifier followed by a register and a stylus recorder. The results are summarized in Table I.

Table I.* Spontaneous fission rates of
 U^{234} , Pu^{236} , Cm^{242} , and Cm^{244} .


Nuclide	Fissions/gm-hr	"Half-life" (years)
U^{234}	13 ± 6	$2 \pm 1 \times 10^{16}$
Pu^{236}	$5.8 \pm 2 \times 10^7$	$3.5 \pm 1 \times 10^9$
Cm^{240}	$1.0 \pm 0.2 \times 10^{11}$	$1.9 \pm 0.4 \times 10^6$
Cm^{244}	$1.4 \pm 0.2 \times 10^{10}$	$1.4 \pm 0.2 \times 10^7$

*The errors indicated are statistical only and do not include any estimate for possible systematic errors.

The U^{234} was a sample of high isotopic purity obtained by the electro-magnetic concentration process, the Pu^{236} was prepared by bombarding highly enriched U^{235} with 18 Mev deuterons according to the reactions



the Cm^{240} came from the bombardment of Pu^{239} with 38 Mev helium ions according to the reaction $Pu^{239}(\alpha,3n)Cm^{240}$, and the Cm^{244} came from the pile neutron bombardment of Am^{243} (containing Am^{241}) by the reactions



By the nature of their methods of production, the Cm^{240} and Cm^{244} contained some Cm^{242} whose spontaneous fission had to be subtracted from the total rate in each case. The Cm^{240} also contained some Cm^{241} , but since the fission rate seemed to decay with the half-life of Cm^{240} , the contribution of the Cm^{241} must have been small. This observation on Cm^{241} would agree with the lower rate expected for nuclides having odd nucleons. The result for U^{234} is consistent with the earlier observation of Segre² who reported an upper limit of 30 spontaneous fissions/gram-hour.

These data are included in Fig. 1 which is otherwise identical with the plot in the previous report¹ (where references are given), with the exception that odd nucleon nuclides, which apparently all fall above the line, are not included. As can be seen, the new even-even nuclides fit in fairly well with the correlation. However, some even-even nuclides such as U^{234} , and possibly also U^{232} and Th^{230} , exhibit substantial deviations in the direction of slower rates. It is apparent

that more data are needed in order to establish the pattern for even-even nuclides in detail. Nevertheless, it can be definitely stated that the spontaneous fission rates for even-even nuclides seem to define a certain limiting rate, and it seems especially significant that the extrapolation of the line (in Fig. 1) representing this rate to the region of instantaneous rate (that is, half-life of the order of 10^{-20} seconds) gives a value of about 47 for Z^2/A , which corresponds with the predicted limiting value for Z^2/A .

Similar considerations in regard to spontaneous fission rates have recently been published by Whitehouse and Galbraith.³

We wish to express our appreciation to Professor J. G. Hamilton, T. M. Putnam, Jr., G. B. Rossi, and the operating crew of the 60-inch cyclotron of the Crocker Laboratory for their help in the bombardments. We would also like to thank the Y-12 Area of the Oak Ridge National Laboratory for supplying the highly purified U^{234} sample. This work was performed under the auspices of the U. S. Atomic Energy Commission.

¹G. T. Seaborg, Phys. Rev. 85, 157 (1951).

²E. Segrè, U. S. Atomic Energy Commission Declassified Document LADC-975 (May 8, 1951).

³W. J. Whitehouse and W. Galbraith, Nature 169, 494 (1952).




Fig. 1. Plot of spontaneous fission rates of even-even nuclides (δ signifies lower limit to half-life).