

UC San Diego

UC San Diego Previously Published Works

Title

Evaluating psychometric properties of the Emotional Eating Scale Adapted for Children and Adolescents (EES-C) in a clinical sample of children seeking treatment for obesity: a case for the unidimensional model

Permalink

<https://escholarship.org/uc/item/2fb016jx>

Journal

International Journal of Obesity, 43(12)

ISSN

0307-0565

Authors

Kang Sim, D Eastern
Strong, David R
Manzano, Michael
et al.

Publication Date

2019-12-01

DOI

10.1038/s41366-019-0427-8

Peer reviewed

1 Evaluating Psychometric Properties of the Emotional Eating Scale Adapted for Children
2 and Adolescents (EES-C) in a Clinical Sample of Children Seeking Treatment for
3 Obesity: A Case for the Unidimensional Model
4

5 (Contributors; orders can be rearranged)

6 D. Eastern Kang Sim^a

7 David R. Strong^b

8 Michael Manzano^a

9 Dawn M. Eichen^a

10 Kyung E. Rhee^a

11 Marian Tanofsky-Kraff^c

12 Kerri N. Boutelle^{a,b,d}

13 ^a Department of Pediatrics, UC San Diego, 9500 Gilman Drive, MC 0874, La Jolla CA
14 92093, USA (dkangsim@ucsd.edu; m3manzano@ucsd.edu; deichen@ucsd.edu;
15 k1rhee@ucsd.edu; kboutille@ucsd.edu)
16

17 ^b Department of Family Medicine and Public Health, UC San Diego, 9500 Gilman Drive,
18 La Jolla CA 92093-0725, USA (dstrong@ucsd.edu; kboutelle@ucsd.edu)
19

20 ^c Department of Medical and Clinical Psychology, Uniformed Services University, 4301
21 Jones Bridge Road, Bethesda MD 20814, USA (marian-tanofsky-kraff@usuhs.edu)
22

23 ^d Department of Psychiatry, UC San Diego, 9500 Gilman Drive, MC 0874, La Jolla CA
24 92093, USA (kboutelle@ucsd.edu)

25

26 Correspondence concerning this article should be addressed to D. Eastern Kang Sim,
27 the CHEAR, UC San Diego, 9500 Gilman Drive, La Jolla, CA 92093. E-mail:
28 dkangsim@ucsd.edu

29

30

31 **Abstract:**

32 *Background:* The Emotional Eating Scale adapted for children and adolescents (EES-C)
33 assesses food-seeking behavior and overeating in response to a range of mood or
34 affects. Despite the fact that prior psychometric studies have demonstrated high
35 reliability, concurrent validity, and test-retest reliability of theoretically defined
36 subconstructs, no prior studies of the EES-C have focused on a clinical sample of
37 children with overweight or obesity. The purpose of this study was to assess construct
38 validity of a single-construct and a proposed scoring of two sub-constructs.

39 *Method:* Using a hierarchical bi-factor approach, we evaluated the EES-C's validity in
40 assessing a single general construct, a set of two separate correlated subconstructs, or
41 hierarchical arrangement of two constructs, and determine reliability in a clinical sample
42 of treatment-seeking overweight or obese children aged 8 to 12.9 years (N=150).

43 *Results:* The present study demonstrated that rigorous factor-extraction methods
44 suggest a one-factor solution. The bi-factor indices provided clear evidence that most of
45 the reliable variance in the total score (90.8 for bi-factor model with three grouping
46 factors and 95.2 for bi-factor model with five grouping factors) was attributed to the
47 general construct. Correlated subconstructs that are currently identified in the clinical
48 sample were unreliable after the variance explained by the single general construct.

49 *Conclusion:* Results suggest that the primary interpretive emphasis of the EES-C
50 among treatment-seeking children with overweight or obesity should be placed on a
51 single general construct, not at the subscale level.

52

53 **Keywords:** emotional eating; scale development

54 **Introduction**

55 The Emotional Eating Scale (EES) for adults was designed to assess food-seeking
56 behavior and overeating in relation to a range of moods or affects ¹. The EES was
57 adapted for use with children and adolescents to determine whether similar behaviors
58 occur at this younger age ². Prior psychometric research on the Emotional Eating Scale
59 Adapted for Children and Adolescents (EES-C) has shown strong internal consistency
60 reliability, concurrent validity with general indices of disordered eating and general
61 emotional problems, and test-retest reliability of theoretically defined constructs on
62 separate subscales ²⁻⁵. Although theoretically defined subscales of the EES-C have
63 been useful tools to investigate the relationship between various affectivity and
64 overeating in children ^{1,2}, there are discrepancies in the proposed number of subscales
65 for children depending on the context in which it is used ^{2,4,5}. For instance, the original
66 validation study by Tanofsky-Kraff ² proposed three subscales ('Anger, anxiety,
67 frustration', 'depression', and 'unsettledness'). Using the Spanish version of the EES-C,
68 Perpina et al. ⁵ suggested five subscales ('anger', 'anxiety', 'depression', 'restlessness',
69 and 'helplessness'). It is not surprising, thus, that Vannucci and colleagues found that a
70 total score (the sum of all items, other than eating in response to feeling "happy")
71 showed construct validity with negative mood and energy intake ³.

72 However, in a clinical setting, the differentiation of subscales may not be
73 apparent among children with overweight or obesity who are likely to generate high
74 levels of emotional eating across all domains ^{6,7}. As children participating in the
75 previous studies were predominantly healthy weight, with only one-third of children
76 having overweight or obesity, it will be valuable to evaluate the psychometrics of the

77 EES-C and its true dimensionality between a single-construct and a sub-constructs in a
78 clinical sample ².

79 To the best of our knowledge, no prior psychometric study of the EES-C has
80 assessed a mix of single and subscale evaluations. The hierarchical bi-factor model,
81 which concurrently describes the common traits such as emotional eating scale and the
82 set of subscales (e.g., eating in response to anger, depression, etc.) may supplement
83 empirical evidence that prior psychometrics studies were unable to contribute ⁸⁻¹¹. By
84 adopting a higher-order factor analysis, we can begin to partition whether responses to
85 items were more likely to arise from smaller correlated subconstructs or if item
86 responses were reflective of a single general dimension. Thus, this study aims to
87 evaluate the validity of the EES-C in a clinical sample of children seeking treatment for
88 overweight or obesity by assessing a single general construct, a set of two separate
89 sub-constructs, or a hierarchical arrangement of the two using a bi-factor approach.

90

91 **Materials and methods:**

92 The Family, Responsibility, Education, Support and Health (FRESH) study was a
93 randomized clinical non-inferiority trial, conducted between July 2011 and July 2015 in
94 San Diego, California (Clinical Trial: NCT01197443), and evaluated two 6-month
95 treatments for childhood obesity. Detailed recruitment methods are described elsewhere
96 ^{12, 13}. Briefly, eligibility criteria included children aged 8 to 12.9 years, child body mass
97 index (kg/m^2 , BMI) from 85th to 99.9th percentile, a parent in the household with a BMI of
98 at least $25 \text{ kg}/\text{m}^2$, and availability to participate in the study on designated evenings.
99 Children with medical or psychiatric conditions that could interfere with participation in
100 the treatment were excluded. In total, 150 children who meet the inclusion criteria and
101 their parents were recruited through local advertisement, school listservs, and local
102 pediatric clinics. The current study uses measures completed by these children at
103 baseline, prior to starting any treatment. The institutional review boards of the University
104 of California San Diego and Rady Children's Hospital, San Diego, California approved
105 the study. Written consent and assent were obtained from parents and children,
106 respectively.

107 Emotional Eating Scale Adapted to Use in Children and Adolescents (EES-C):

108 The EES-C is a 25-item questionnaire that assesses eating when confronted with 25
109 negative emotions (e.g., resentful, discouraged, etc.) on a 5-point Likert scale (from "no
110 desire" to "very strong desire to eat") ². Summing the individual EES-C items generates
111 an EES-C total score. To test the convergent and discriminant validity of the scale, we
112 used the median score of the EES-C total score and dichotomized the results into two
113 groups: high in emotional eating (High-EE) and low in emotional eating (Low-EE).

114 Alternative factor models derived from prior studies ^{2, 5} in non-clinical samples have
115 been replicated to provide context and described in the analysis section.

116 Child Eating Disorder Examination (ChEDE): The ChEDE is a semi-structured
117 interview that assesses eating disorder features in children ¹⁴. The overeating section
118 was administered to evaluate the number of objective bulimic episodes (i.e., objectively
119 large amount of food with loss of control over eating) or subjective bulimic episodes (i.e.,
120 smaller amount of food but viewed as excess to participant with loss of control over
121 eating) in the past 3 months. To test the convergent validity, we dichotomized children
122 into two groups, 'any experience of loss of control eating' or 'no experience of loss of
123 control eating' respectively ¹⁵.

124 Child Behavior Checklist (CBCL): The CBCL is a parent-report questionnaire that
125 assesses children's behavioral problems ¹⁶. The CBCL yields standardized T scores
126 and age-adjusted scores on internalizing, externalizing, and total behavioral difficulties,
127 which were used to test the discriminant validity of the EES-C. The CBCL has been
128 evaluated in clinical and community populations with good inter-rater and intra-rater
129 reliability ¹⁷.

130 **Statistical analysis**

131 All analyses were conducted using the R statistical programming language (version
132 3.4) ¹⁸ and SPSS (version 23, IBM) ¹⁹. Polychoric correlations were used where
133 appropriate ²⁰. Prior to the bi-factor analysis, we replicated the methods used in prior
134 studies to help define multiple EES sub-constructs for the clinical sample. In brief, these
135 methods used Kaiser-one for class enumeration and principal component or exploratory
136 factor analysis with varimax rotation. We found lack of agreement of exploratory models

137 (e.g. 'excited/uneasy/resentful', 'loneliness', 'depression' for the three-factor model;
138 'anxiety', 'agitated', 'guilty', 'upset', and 'loneliness' for the five-factor model), which in
139 turn suggests need to examine in clinical samples. For the current study, we focus on
140 the hierarchical bifactor model which simultaneously evaluate a mix of single construct
141 and subscales. *Construct validity*

142 The optimal solution for the number of factors to be retained was determined by
143 the Kaiser-one criterion ²¹. The following procedures were also tested: 1) Velicer's
144 minimum average partial (MAP) criteria ²²; 2) Horn's parallel analysis (PA) ²³; 3) the
145 optimal coordinates (OC) ²⁴; 4) the acceleration factor (AF) ²⁴; 5) the Very Simple
146 Structure (VSS) ²⁵; and 6) Ruscio and Roche's Comparison Data (CD) ²⁰. Summing the
147 factored items generated the scores for each EES-C subscale.

148 *Convergent and Discriminant Validity*

149 To assess convergent validity, differences between the groups (High-EE and
150 Low-EE) and all variables of interest were measured using a t-test, and p-values < .05
151 were considered significant. To assess discriminant validity, Spearman's correlations
152 were used to determine whether the total and subscale scores for the EES-C were
153 significantly related to the corresponding CBCL internalizing, externalizing and total
154 behavior problems.

155 *Bi-factor model indices*

156 Hierarchical bifactor models were examined to simultaneously evaluate the
157 strength of support for a primary single factor underlying the responses and the degree
158 to which additional group factors suggested the multidimensionality of the remaining

159 variability among items after adjustment was made for relationships with the primary
160 construct^{8, 11}.

161 *Explained common variance (ECV):* ECV was used to estimate the degree to
162 which a general construct and correlated subconstructs could be used to explain and
163 organize item responses^{8, 9, 26}.

164 *Percent of uncontaminated correlations (PUC):* PUC, a bifactor-specific index,
165 presents information on the percentage of correlation that is not contaminated by
166 multidimensionality²⁷.

167 *Reliability coefficients:* Cronbach's coefficient alpha (α) was used to estimate the
168 internal scale reliability coefficient²⁸. McDonald's coefficient omega (ω) was used to
169 compliment the alpha coefficient, which estimates the proportion of variance in the unit-
170 weighted total score attributable to all sources of common variance²⁹. Omega
171 hierarchical (ω_H) and Omega hierarchical subscale (ω_{HS}) were used to estimate the
172 variance that is attributable to a single general construct and/or correlated
173 subconstructs³⁰⁻³².

174 *Scalability (Coefficient H):* Coefficient H was used to evaluate how well a set of
175 items' scalability represented the latent variable²⁶.

176 **Results**

177 The mean age of child participants was 10.4 years, and 33.3% (n=50) were males.

178 Almost one-third of the subjects were Hispanic. See Table 1 for participant
179 demographics and characteristics.

180 *Convergent and Discriminant Validity of the EES-C*

181 Table 1 presents support for convergent validity and strong relationships
182 between EES-C total scores and levels of self-reported LOC eating behavior. The
183 median for the EES-C total score was 9.5 (range: 0–74). Therefore, children with EES-C
184 scores <9.5 were categorized as low in emotional eating, and those with total scores ≥
185 9.5 were categorized as High-EE. Participants with High-EE did not differ on
186 demographic or anthropometric variables, with the exception that Hispanic children
187 were more likely to be classified in the Low-EE group when compared with their peers in
188 the High-EE group. Children in the High-EE group were more likely to endorse LOC
189 eating than the Low-EE group and BMI-z score did not differ between groups (Table 1).

190 Table 2 presents an examination of discriminant validity of the EES-C total and
191 subscale scores. The correlation coefficients between the percentile of internalizing,
192 externalizing, and total behavior problems on the CBCL with the EES-C total score or
193 subscales (formed with either three or five grouping factors) were all small (range = -
194 0.08 to 0.08). No statistically significant differences were noted, suggesting the EES-C
195 reliably assess a construct of emotional eating that was distinct from general emotional
196 or behavior problems.

197 *Exploring Construct Validity*

198 Figure 1 presents scree plot of indices for determining the number of factors to
199 be retained. While the Kaiser-one approach suggested that five factors to be retained,
200 Velicer's MAP criteria provided minimum squared average partial correlations of 0.02 for
201 the first and second steps, suggesting one or two factors. The remaining four methods
202 (three are displayed in figure 1) suggested that one factor be retained.

203 *Applying the Bi-factor Model*

204 Table 3 presents summary results of standardized factor loadings and bi-factor
205 reliability indices of the three-grouping factor. The single general factor loadings ranged
206 from .57 to .79 across all items and most were within the DeVellis's common criteria for
207 an acceptable range³³. All subscales item-loadings for correlated factors were poor with
208 the exception of emotional eating in response to feeling 'furious' (.79). Across all factor
209 extractions, the single general factor of the bi-factor model accounted for 90% of reliable
210 variance with 10% of the residual variance spread across subscales. After accounting
211 for the variance due to the general factor, the subscales for the correlated factors
212 accounted for a small proportion of the total variance ($\omega_{HS} = .17, .11, .37$). The
213 remaining 3% of the ω total is estimated to be due to random error. With a coefficient H
214 of .94, the general factor presents near perfect construct replicability. None of the
215 indices of the three grouping factors show strong construct replicability.

216 Table 4 presents summary results of standardized factor loadings and indices of
217 a bi-factor model with five grouping factors. The single general factor loadings remained
218 strong and ranged from .61 to .80 across all items. Within the bi-factor model with
219 subscale for the correlated five grouping factors, item loadings were all less than 0.50
220 with the exception of the item 'furious' (.81). The single general factor accounted for 95%

221 of the reliable variance, implying only 5% of the residual variance is distributed to
222 subscales. After accounting for the variance due to the general factor, the subscale
223 grouping factors accounted for a small proportion of the total variance (ω^2
224 = .11, .14, .18, .33, .14). The coefficient H of .95 suggests strong construct replicability
225 of the general factor, whereas none of the indices of the five grouping factors show
226 strong construct reliability. The fourth grouping factor in this model (FFS-F4; table 4)
227 had an H index of .66 which meets the recommended cutoff for favored construct
228 replicability but had only two items ('furious' and 'angry'), suggesting a set of closely
229 related items strongly defined by eating in response to feeling furious.

230

231 **Discussion**

232 This study evaluated the construct validity and psychometric properties of the EES-C
233 using hierarchical bi-factor approach among children seeking weight-loss treatment.
234 Nearly all of the reliable variance of the EES-C was captured by a single general
235 construct underlying the responses, and multiple bi-factor indices supported the general
236 factor's unidimensionality. Results suggested that the single general factor of emotional
237 eating directly influenced responses on each of the subscales from the correlated
238 factors rather than simply reflecting an accumulation or indirect influence of separately
239 assessed constructs. Scores from the general factor demonstrated good convergent
240 validity with a measure of LOC eating behavior, and good discriminant validity with no
241 evidence of significant relationships with competing measures of general emotional or
242 behavioral problems from the CBCL.

243 There are several reasons why it may be useful to use a single general construct for
244 emotional eating in children rather than distinguish between several different constructs
245 of emotions related to eating among treatment-seeking children who are overweight or
246 obese. First, children between the ages of 8 and 12 years old are still developing the
247 cognitive and emotional awareness needed to distinguish between different affective
248 states that are represented in the EES-C³⁴. Second, children in this age range may
249 best relate their eating behaviors to overall levels of arousal (e.g., furious vs. calm) or
250 general valence of affect (e.g., positive vs. negative) rather than discrete emotions (e.g.,
251 lonely).

252 In terms of applied methodology, our study utilized several newer approaches that
253 move the previous psychometric work conducted on the EES-C forward. One of the

254 greatest challenges in factor analysis is choosing the correct number of factors to retain.
255 The traditional Kaiser one approach suggested that five factors exist in the EES-C. Of
256 the six alternative factor extraction methods tested (OC, AF, PA, CD, VSS, and MAP),
257 five suggested that one factor be retained and the sixth (VSS) suggested that one or
258 two factors should be retained. This implies that, while multiple sources of variability in
259 item responses within the EES-C could be scored separately, the identification of items
260 or relative importance of extracted subscales may not be stable or replicable across
261 studies. Rather, a more stable and parsimonious solution may be to organize all items
262 using the single primary construct, a solution supported by multiple indices that suggest
263 the unidimensionality of this scale.

264 Another stabilizing methodological approach addresses decisions around which test
265 of correlation to use that would best reflect the ordered categorical response process for
266 these items^{2, 4, 5}. The EES-C, which uses a five-point Likert scale, has a strong
267 skewedness or kurtosis, and using the Pearson's correlation may produce factors that
268 are based solely on item distribution similarity and can cause items to appear as
269 multidimensional when, in fact, they are not³⁵. In the present study, we have
270 implemented the polychoric correlation approach, which leads to more robust
271 estimations of dimensionality than factor analyses using Pearson's.

272 Furthermore, our study utilized several modern coefficients to evaluate internal
273 consistency. Prior psychometrics studies of the EES-C have extensively used
274 coefficient alpha (α), which demonstrated strong internal consistency; however, high α
275 values from previous studies may be partly attributable to the many redundant items
276 within the scale, which inflate correlations within the group factor. The reliance on α

277 alone has been criticized as an exclusive indicator of scale reliability because it
278 underestimates true reliability and is not sensitive to violations of assumptions of the
279 unidimensional nature of the scale^{36, 37}. By implementing a bi-factor approach, we have
280 partitioned single general and correlated group factor variance to better understand the
281 strength of a single primary factor underlying the EES-C. Upon evaluating the percent of
282 total score variance attributable to a single general factor, ω_H provided clear evidence
283 that most of the reliable variance in the total score is attributed to the general factor, not
284 to the subscales. We also provided a coefficient H, which is interpreted as a replicability
285 coefficient. Only the general factor passed the threshold of coefficient H (.7); not all
286 subscales met this criterion. The low coefficient H of all the subscales leads one to be
287 suspicious of construct reliability because they are likely to differ from one study to
288 another and in different contexts. The total score, however, had loadings greater
289 than .90, indicating high construct reliability between studies.

290 One major strength of this study is its use of newer empirical approaches that have
291 been absent from previous validation studies. These methods provide a more robust
292 evaluation of the psychometric properties of the EES-C and a more complete picture of
293 scale performance. Furthermore, this study examined psychometric properties using a
294 population that had never been evaluated: overweight children seeking to lose weight.
295 Several limitations, however, must be considered. As this was a randomized control
296 clinical trial with a population of children seeking to lose weight, self-report bias may
297 have possibly influenced our participants' responses with regards to their emotional
298 eating behaviors. For instance, the median score of the EES-C of our clinical sample
299 was nominally lower (8-12 years; median 9.5) compared to the previous validation study

300 with 151 youths (8-18 years; median 13)³. Including only treatment seeking children do
301 not necessarily generalize to other children with overweight/obesity and not to healthy
302 weight children. Future studies should test the reliability of this scale in other
303 populations while using a similar bi-factor approach.

304 In summary, these results suggest that for a clinical sample of children with
305 overweight or obesity, the EES-C should be implemented with a unidimensional scale
306 and supports the construct validity of the scale in non-treatment seeking children using
307 a total score³. Thus, recommendations to use a single total score should be applied to
308 both treatment-seeking and non-treatment seeking children. Future studies are needed
309 to determine whether the single general factor as manifested in the total score is
310 clinically important.

311 Acknowledgements: This work was supported by the National Institute of Health [grant
312 numbers R01DK075861 and K02HL1120242, PI: Boutelle]. The opinions and assertions
313 expressed herein are those of the authors and are not to be construed as reflecting the
314 views of the Uniformed Services University or the U.S. Department of Defense.

315

316 Competing Interest: The authors declare that they have no conflict of interest.

317 **References**

- 318 1. Arnow B, Kenardy J, Agras W. The Emotional Eating Scale: the development of
319 a measure to assess coping with negative affect by eating. *Int J Eat Disord* 1995;
320 **18**(1): 79-90.
- 321
- 322 2. Tanofsky-Kraff M, Theim K, Yanovski S, Bassett A, Burns N, Ranzenhofer L *et al.*
323 Validation of the emotional eating scale adapted for use in children and
324 adolescents (EES-C). *Int J Eat Disord* 2007; **40**(3): 232-240.
- 325
- 326 3. Vannucci A, Tanofsky-Kraff M, Shomaker L, Ranzenhofer L, Matheson B,
327 Cassidy O *et al.* Construct validity of the emotional eating scale adapted for
328 children and adolescents. *Int J Obes (Lond)* 2012; **36**(7): 938-943.
- 329
- 330 4. Bektas M, Bektas I, Selekoglu Y, Kudebes A, Altan S, Ayar D. Psychometric
331 properties of the Turkish version of the Emotional Eating Scale for children and
332 adolescents. *Eat Behav* 2016; **22**: 217-221.
- 333
- 334 5. Perpina C, Cebolla A, botella C, Lurbe E, Torro M. Emotional Eating Scale for
335 children and adolescents: psychometric characteristics in a Spanish sample. *J*
336 *Clin Child Adolesc Psypchol* 2011; **40**(3): 424-433.

337

- 338 6. Jansen A, Vanreyten A, van Balveren T, Roefs A, Nederkoorn C, Havermans R.
339 Negative affect and cue-induced overeating in non-eating disordered obesity.
340 *Appetite* 2009; **51**(3): 56-562.
341
- 342 7. Webber L, Hill C, Saxton J, Van Jaarsveld C, Wardle J. Eating behaviour and
343 weight in children. *Int J Obes (Lond)* 2009; **33**(1): 21-28.
344
- 345 8. Reise S. The rediscovery of bifactor measurement models. *Multivar Behav Res*
346 2012; **47**(5): 667-696.
347
- 348 9. Reise S, Moore T, Haviland M. Bifactor models and rotations: exploring the
349 extent to which multidimensional data yield univocal scale scores. *J Pers Assess*
350 2010; **92**(6): 544-559.
351
- 352 10. Reise S, Morizot J, Hays R. The role of the bifactor model in resolving
353 dimensionality issues in health outcomes measures. *Qual Life Res* 2007;
354 **16**(Suppl 1): 19-31.
355
- 356 11. Rodriguez A, Reise S, Haviland M. Applying bifactor statistical indices in the
357 evaluation of psychological measures. *J Pers Assess* 2016; **98**(3): 223-237.
358
- 359 12. Boutelle K, Braden A, Douglas J, Rhee K, Strong D, Rock C *et al.* Design of the
360 FRESH study: A randomized controlled trial of a parent-only and parent-child

361 family-based treatment for childhood obesity. *Contemp Clin Trials* 2015; **45**(Pt B):
362 364-370.

363

364 13. Boutelle K, Rhee K, Liang J, Braden A, Douglas J, Strong D *et al.* Effect of
365 attendance of the child on body weight, energy intake, and physical activity in
366 childhood obesity treatment: A randomized clinical trial. *JAMA Pediatr* 2017;
367 **171**(7): 622-628.

368

369 14. Fairburn C, Cooper Z, O'Connor M. Eating disorder examination. In : Fairburn
370 CG, editor. *Cognitive Behavior Therapy and Eating Disorders*, 16th edition. New
371 York: Guilford Press, pages 265-308. 2008.

372

373 15. Stojek M, Tanofsky-Kraff M, Shomaker L, Kelly N, Thompson K, Mehari R *et al.*
374 Associations of adolescent emotional and loss of control eating with 1-year
375 changes in disordered eating, weight, and adiposity. *Int J Eat Disord* 2017; **50**(5):
376 551-560.

377

378 16. Achenbach T. *Manual for Child Behavior Checklist/4-18 and 1991 Profile.*
379 *University of Vermont, Department of Psychiatry.* Burlington VT, ISBN
380 0938565087, 9780938565086, 1991.

381

- 382 17. Dutra L, Campbell L, Western D. Quantifying clinical judgment in the assessment
383 of adolescent psychopathology: Reliability, validity, and factor structure of the
384 Child Behavior Checklist for clinician report. *J Clin Psychol* 2004; **60**(1): 65-85.
385
- 386 18. Team R. R: A language and environment for statistical computing. R Foundation
387 for Statistical Computing, Vienna, Austria, 2013,. 2014.
388
- 389 19. Statistic I. SPSS 23.0 for windows. Chicago, IL. 2015.
390
- 391 20. Ruscio J, Roche B. Determining the number of factors to retain in an exploratory
392 factor analysis using comparison data of known factorial structure. *Psychol*
393 *Assess* 2012; **24**(282-292).
394
- 395 21. Kaiser H. The application of electronic computers to factor analysis. *Educ &*
396 *Psychol Measurement* 1960; **20**: 141-151.
397
- 398 22. Velicer W. Determining the number of components from the matrix of partial
399 correlations. *Psychometrika* 1976; **41**(3): 321-327.
400
- 401 23. Ledesma R, Valero-Mora P. Determining the number of factors to retain in EFA:
402 An easy-to-use computer program for carrying out parallel analysis *Practical*
403 *Assess Res & Eval* 2007; **12**(2): 1-11.
404

- 405 24. Raiche G, Walls T, Jagis D, Riopel M, Blais J-G. Non-graphical solutions for
406 Cattell's scree test. *Methodology* 2013; **Methodology**(9).
407
- 408 25. Revelle W, Rocklin T. Very simple structure: An alternative procedure for
409 estimating the optimal number of interpretable factors. *Multivar Behav Res* 1979;
410 **14**(4): 403-414.
411
- 412 26. Hancock G, Mueller R. Rethinking construct reliability within latent variable
413 systems. Structural equation modeling: Present and future. *Scientific Software*
414 *International* 2001: 195-216.
415
- 416 27. Bonifay W, Reise S, Scheines R, Meijer R. When are multidimensional data
417 unidimensional enough for structural equation modeling? An evaluation of the
418 DETECT Multidimensionality Index. *Structural Equation Modeling: A*
419 *Multidisciplinary Journal* 2015; **22**(4): 504-516.
420
- 421 28. Cortina j. What is coefficient alpha? An examination of theory and applications. *J*
422 *Applied Psychology* 1993; **78**(1): 98-104.
423
- 424 29. McDonald R. Test theory: A unified treatment: Psychology Press. 2013.
425

- 426 30. Bentler P. Covariance structure models for maximal reliability of unit-weighted
427 composites. In Handbook of Latent Variable and Related Models, S-Y Lee (ed).
428 *North Holland* 2007: 1-19.
- 429
- 430 31. Bentler P. Alpha, dimension-free, and model-based internal consistency reliability.
431 *Psychometrika* 2009; **74**(1): 137-143.
- 432
- 433 32. Kelley K, Pornprasertmanit S. Confidence intervals for population reliability
434 coefficients: Evaluation of methods, recommendations, and software for
435 composite measures. *Psychol Methods* 2016; **21**(1): 69-92.
- 436
- 437 33. DeVellis R. Scale Development: Theory and Applications Second Edition
438 (Applied Social Research Methods). *Sage Publications, Inc.* 2003.
- 439
- 440 34. Lindquist K, Barrett L. Emotional complexity. Chapter in M. Lewis, J. M. Haviland-
441 Jones, & L.F. Barrett (Eds.), Handbook of emotions, 3rd edition. New York:
442 Guilford. 2008: 513-430.
- 443
- 444 35. Flora D, Curran P. An empirical evaluation of alternative methods of estimation
445 for confirmatory factor analysis with ordinal data. *Psychol Methods* 2004; **9**(4):
446 466-491.
- 447

448 36. Crutzen R, Peters G. Scale quality: alpha is an inadequate estimate and factor-
449 analytic evidence is needed first of all. *Health Psychol Rev* 2017; **11**(3): 242-247.

450

451 37. McNeish D. Thanks coefficient alpha, we'll take it from here. *Psychol Methods*
452 2017; **[Epub ahead of print]**.

453

454

Figure 1. Screen plot of indices for the optimal number of factors to be retained




Table 1. Sample characteristics by high or low in emotional eating (EE) scale

Variable names	Total		High EE		Low EE	
Total EE	14.87	(15.44)	26.29	(14.38)	3.47	(3.07)
Gender (male)	50	(33.33)	20	(26.00)	30	(40.00)
Hispanic *	47	(32.00)	17	(24.30)	30	(40.00)
BMI z-score	2.00	(0.34)	1.99	(0.35)	2.01	(0.33)
Loss of control eating (%) *	43	(29.30)	29	(40.80)	13	(17.60)
CBCL						
Internalizing percentile	42.54	(29.19)	44.36	(28.08)	40.98	(30.57)
Externalizing percentile	34.80	(27.53)	38.18	(27.97)	31.03	(26.68)
Total percentile	40.68	(28.64)	43.06	(28.01)	37.94	(29.17)

Mean (SD) or N (%) were reported; t-statistics were used; * <0.05; EE- emotional eating; CBCL – child behavioral check list

Table 2. Correlation coefficient between percentile of internalizing, externalizing, and total behavior problems and sum of total EES-C and extracted factor structures

	Internalizing	Externalizing	Total
EES-C Total	-.03	.02	.00
EES-C TFS-F1	-.03	.02	.02
EES-C TFS-F2	-.05	.03	-.02
EES-C TFS-F3	-.08	-.06	-.08
EES-C FFS-F1	-.04	.05	-.01
EES-C FFS-F2	-.06	.04	.01
EES-C FFS-F3	-.02	.04	.04
EES-C FFS-F4	-.08	-.06	-.08
EES-C FFS-F5	-.01	.06	.06

No factor structures were significant at the .05 level

Table 3. Standardized bi-factor loadings and indices from three-factor solution (TFS)

	GF	TFS-F1	TFS-F2	TFS-F3
1 Resentful	0.70			
2 Discouraged	0.66	0.31		
3 Shaky	0.74		0.24	
4 Worn out	0.57			
5 Not doing enough	0.70	0.24		
6 Excited	0.72		0.35	
7 Disobedient	0.75			
8 Down	0.64	0.36		
9 Stressed out	0.70			
10 Sad	0.66	0.30		
11 Uneasy	0.73		0.28	
12 Irritated	0.74		0.28	
13 Jealous	0.71			
14 Worried	0.65	0.36		
15 Frustrated	0.77			
16 Lonely	0.61	0.34		
17 Furious	0.64			0.79
18 On edge	0.76	0.23		
19 Confused	0.71	0.25		
20 Nervous	0.67		0.21	
21 Angry	0.79			0.28
22 Guilty	0.58	0.42		
23 Bored	0.62		0.20	
24 Helpless	0.73	0.25		
25 Upset	0.72	0.38		
Indices				
Eigenvalue	11.98	1.31	0.61	1.01
Coefficient α	0.96			
Coefficient ω total	0.97	0.94	0.91	0.89
ω hierarchical and subscale	0.88	0.16	0.11	0.33
Reliable variance from ω	90.82	17.55	11.75	37.30
Explained common variance	0.80			
Percent uncontaminated corr	0.58			
Scalability (H)	0.94	0.55	0.31	0.63

GF= general factor; TFS-F1: Depression; TFS-F2: Anxiety; TFS-F3: Angry

Table 4. Standardized bifactor loadings and indices from five-factor solution (FFS)

	GF	FFS-F1	FFS-F2	FFS-F3	FFS-F4	FFS-F5
1 Resentful	.69	.20				
2 Discouraged	.67					.22
3 Shaky	.73	.27				
4 Worn out	.59		.24			
5 Not doing enough	.70					.23
6 Excited	.71	.37				
7 Disobedient	.76					
8 Down	.64			.44		
9 Stressed out	.70			.40		
10 Sad	.67			.31		
11 Uneasy	.71	.32				
12 Irritated	.75	.28				
13 Jealous	.70					
14 Worried	.69		.32			
15 Frustrated	.77			.25		
16 Lonely	.61					.48
17 Furious	.62				.81	
18 On edge	.78		.25			
19 Confused	.73		.26			
20 Nervous	.67	.22				
21 Angry	.80				.25	
22 Guilty	.62		.46			
23 Bored	.61	.22				
24 Helpless	.73					.32
25 Upset	.74			.37		
Indices						
Eigenvalue	12.17	.70	.68	.76	1.05	.74
Coefficient α	.96					
Coefficient ω total	.97	.92	.87	.91	1.04	.84
ω hierarchical and subscale	.92	.11	.14	.18	.33	.14
Reliable variance from ω	95.25	12.10	16.66	19.78	31.82	17.50
Explained common variance	.80					
Percent uncontaminated corr	.79					
Scalability (H)	.95	.36	.36	.43	.66	.34

GF= general factor; FFS-F1: Anxiety; FFS-F2: Guilty; FFS-F3: Down; FFS-F4: Angry; FFS-F5: Loneliness