

Lawrence Berkeley National Laboratory

Recent Work

Title

THE ELASTIC SCATTERING OF PROTONS FROM HELIUM 4

Permalink

<https://escholarship.org/uc/item/2gj5v9kv>

Author

Cork, Bruce.

Publication Date

1952-02-11

UNIVERSITY OF CALIFORNIA - BERKELEY

UCRL- 1673

UNCLASSIFIED

TWO-WEEK LOAN COPY

*This is a Library Circulating Copy
which may be borrowed for two weeks.
For a personal retention copy, call
Tech. Info. Division, Ext. 5545*

RADIATION LABORATORY

DISCLAIMER

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

UCRL-1673
Unclassified-Physics Distribution

UNCLASSIFIED

UNIVERSITY OF CALIFORNIA

Radiation Laboratory

Contract No. W-7405-eng-48

THE ELASTIC SCATTERING OF PROTONS FROM HELIUM 4

Bruce Cork

February 11, 1952

Berkeley, California

THE ELASTIC SCATTERING OF PROTONS FROM HELIUM 4

Bruce Cork

Radiation Laboratory, Department of Physics
University of California, Berkeley, California

February 11, 1952

The absolute differential cross section has been measured for 31.6 Mev protons scattered elastically from helium. Seven increments of angles from 15° to 51° in the laboratory system of coordinates have been measured simultaneously using proportional counters.

The observed nuclear scattering is approximately three times the calculated Rutherford scattering for a scattering angle of 17° in the center of mass system and approximately 100 times the calculated Rutherford scattering for an angle of 62° in the center of mass system. Also, the differential cross section for a center of mass angle of 55° has been observed to be 59.1 millibarns/steradian for incident protons of 19.5 Mev in the laboratory system.

INTRODUCTION

The elastic cross section for scattering of protons from He^4 has been measured at energies up to 9.5 Mev^{1,2,3,4,5} and a phase shift analysis made for low energy protons.⁶ The Berkeley proton linear accelerator has allowed the region of 32 Mev and below to be investigated. The apparatus which was used for proton-proton scattering⁷ was used

without modification to obtain data for elastic scattering of protons from helium 4 in the forward directions. An independent experiment giving the elastic and inelastic cross sections will also be reported in another paper.^{8,9}

METHOD

The 31.6 Mev proton beam from the linear accelerator was deflected 12.6° in an analyzing magnet and a collimator 6 meters long gave a beam of 1 cm diameter, having a divergence of ± 0.001 radius. The energy was calculated by measuring the deflection of the beam and the magnitude of the magnetic field along the trajectory. The beam entered the scattering chamber through a double 0.0002 in. nylon foil and was scattered by helium at a pressure of slightly greater than one atmosphere. The scattered protons could be detected in either of nine proportional counters operated simultaneously. An additional counter detected only background neutrons and x-rays.

The fraction of the beam which was not scattered continued on through a 0.001 in. thick duraluminum foil, and on into the faraday cup, arranged as a charge integrator.

PROCEDURE

The scattering chamber was evacuated to a pressure of less than 10^{-5} mm of Hg and observed to be vacuum tight. Helium of greater than 99.5 percent purity (Grade A), obtained from the Mathieson Company,

was allowed to enter the scattering chamber through a liquid nitrogen trap made of stainless steel. The impurities before trapping were reported to be mostly hydrogen.

It was possible to measure the amount of hydrogen impurity by detecting the 90° coincidence scattered protons in the scattering chamber, (see below).

The helium was admitted to the scattering chamber at a constant rate of approximately one liter per minute and allowed to bubble out through an oil lock column of Litton oil 5 cm high. The number of scattering nuclei was determined by measuring the height of this column, the barometric pressure, and the temperature of the gas in the scattering chamber.

The position of the beam and the amount of multiple scattering of the beam were detected by inserting a photographic emulsion at the charge integrator. The beam was observed to have a mean diameter of 2.4 cm at the charge integrator. The charge integrator had an aperture of 6 cm, thus the amount of the beam lost was negligible.

The scattered protons were detected by the same proportional counters which were used for proton-proton scattering.⁷ The counter plateaus were determined by adjusting the gas multiplication and amplifier gain, making a run, then a background run with the shutter closed, and then repeating the run with a higher value of gas multiplication. The number of protons plus background was recorded for each of the ten counters. Also, the number of 90° coincidence counts was measured in

the 45° - 45° counters, and in the 51° - 39° counters. This was a measure of the hydrogen contamination in the helium, plus accidental coincidence.

RESULTS

The data for two series of runs are given in Table I. The number of counts for each angle measured is tabulated, corrected for background, counter resolving time, and normalized to NTP. The counting rate was sufficiently low so that the correction for counter resolving time was always less than 1 percent. The "plateau" was such that for the first group of runs, a 50 volt increase in potential of the proportional counter wire indicated a 0.3 percent decrease in the scattering cross section, while in the second group of runs, a 1.1 percent increase in the scattering cross section was indicated. The statistical fluctuation was ± 1.3 percent.

The number of 90° coincidence counts is tabulated for each of two sets of counters. The pulse repetition rate of the linear accelerator was 15 pulses per record, 400 microsecond long pulses. The resolving time of the coincidence circuit was 1.0 microseconds, and from the observed counting rate, the calculated number of accidental coincidence counts is 1.5 (unscaled) per run. From these corrected data and the measured cross section for proton-proton scattering,¹⁰ the amount of hydrogen impurity is determined to be 1.0 ± 0.5 percent. The corrections have been made for each angle assuming 1 percent hydrogen

contamination and no other contamination. The calculated cross section for each angle is given in Table II. The assigned probable errors include:

- a. Collected charge $\pm \frac{1}{2}$ percent.
- b. Mean energy ± 1 percent.
- c. Measurement of temperature and pressure $\pm \frac{1}{2}$ percent.
- d. Slope of plateau ± 1.5 percent.
- e. R.M.S. deviation of counts ± 1.5 percent.
- f. Calculated geometry $\pm 3/4$ percent.
- g. Contamination scattering $\pm \frac{1}{2}$ percent.

The R.M.S. value of these probable errors is ± 2.6 percent.

Fig. 1 is a comparison of the observed cross section with the calculated Rutherford cross section at this energy. A further comparison will be made in a following paper.⁹

Protons of 19.5 Mev incident energy were obtained by adjusting the radio frequency voltage distribution of the linear accelerator cavity.¹⁰ The differential cross section was then measured at a center of mass angle of 55.0 degrees and observed to be 59.1 ± 1.6 millibarns/steradian.

ACKNOWLEDGMENTS

It is a pleasure to acknowledge the help of Professor Luis W. Alvarez who made these experiments possible. Also, the continued patience of the linear accelerator crew, under the supervision of Robert Watt has greatly simplified the process of obtaining data.

TABLE I

1. Actual counts divided by four and normalized to N.T.P. and A collected charge of 306.9×10^{-12} coulombs.
2. Mean energy = 31.6 ± 0.3 Mev.
3. Geometrical scattering length = 5.38 cm.
4. T and B refer to top and bottom halves of a given counter.

MEAN ANGLES, CENTER OF MASS DEGREES P - α SCATTERING

Date										Proton-Proton Scattering	
	62.5°T	62.5°B	55.0°T	55.0°B	47.8°T	47.8°B	40.1°	24.8°	17.1°	102°-78°	90°-90°
2-11-50	352	337	474	469	-	657	1,560	1,554	1,182	1.7	1.1
	365	350	512	474	-	668	1,545	1,490	1,173	1.7	1.9
	386	314	509	479	-	618	1,510	1,520	1,140	1.1	2.1
5-18-51	355	341	496	457	710	634	1,498	1,474	-	1.7	1.2
	350	385	447	480	666	658	1,572	1,530	-	1.9	1.2
	366	343	490	490	654	656	1,546	1,574	-	1.4	2.2
Total	2,174	2,070	2,928	2,849	2,030	3,891	9,231	9,142	3,495	9.5	9.7
Total Both Sectors		4,244		5,777		7,851	9,231	9,142	6,990	9.5	9.7
$\frac{\theta\sigma}{\theta\Omega}$ cm millibarns/ steradian		31.53		45.21		68.7	93.3	145.5	159.9	14.30	14.39
Probable Error		± 0.8		± 1.2		± 1.8	± 2.4	± 3.8	± 4.2	± 0.15	± 0.14

TABLE II

ELASTIC SCATTERING OF PROTONS FROM He⁴ E = 31.6 MEV PROTONS

θ Lab Deg.	θ cm Deg.	Measured $\frac{d\sigma}{d\Omega}$ cm ² $\times 10^{-27}$ cm	Calculated $\frac{d\sigma}{d\Omega}$ cm Rutherford $\times 10^{-27}$ cm ²
13.64	17.07	159.9 \pm 4.2	47.1
19.89	24.80	145.5 \pm 3.8	10.75
26.21	-	-	3.58
32.33	40.13	93.3 \pm 2.4	1.65
38.82	47.80	68.7 \pm 1.8	0.847
44.82	55.00	45.2 \pm 1.2	0.503
51.19	62.50	31.5 \pm 0.8	0.315

REFERENCES

1. Chadwick, J., and E. S. Bieler, "The Collisions of Alpha Particles with Hydrogen Nuclei", Phil. Mag. 42, 923 (1921).
2. Heydenburg, N.P., and R. B. Roberts, "Deuteron-Deuteron Proton-Helium, and Deuteron-Helium Scattering", Phys. Rev. 56, 1092 (1939).
3. Heydenburg, N. P., and N. F. Ramsey, "The Scattering of One to Three Mev Protons by Helium", Phys. Rev. 60, 42 (1941).
4. Freier, George, Eugene Lampi, W. Sleator, and J. H. Williams, "Angular Distribution of 1 to 3.5 Mev Protons Scattered by He⁴", Phys. Rev. 75, 1345 (1949).
5. Putnam, T. M., "The Differential Cross Section for the Scattering of 9.5 Mev Protons from Helium". Phys. Rev. Bulletin Vol. 26, No. 8, Pg. 26.
6. Critchfield, C. L., and D. C. Dodder, "Phase Shifts in Proton-Alpha Scattering", Phys. Rev. 76, 602 (1949).
7. B. Cork, L. Johnston and C. Richman, Phys. Rev. 79, 71 (1950).
8. J. Benveniste, and Bruce Cork. Phys. Rev. 83, 894-A (1951).
9. J. Benveniste, and Bruce Cork. To be published.
10. Bruce Cork. Phys. Rev. 80, 321 (1950).




Fig. 1