UC Irvine

UC Irvine Previously Published Works

Title

CORRECTION

Permalink

https://escholarship.org/uc/item/2gm6c280

Journal

EPILEPSY RESEARCH, 3(3)

ISSN

0920-1211

Author

RIBAK, CE

Publication Date

1989

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at https://creativecommons.org/licenses/by/4.0/

Peer reviewed

Corrigendum

Epilepsy Res., 2 (1988) 345–355, Anatomical and behavioral analyses of the inheritance of audiogenic seizures in the progeny of genetically epilepsy-prone and Sprague–Dawley rats, by C.E. Ribak, R.C. Roberts, M.Y. Byun and H.L. Kim.

Page 354, right-hand column, line 15: number of all neurons in the seizing F_2 should read number of small neurons in the seizing F_2 .

Substitute Fig. 2 with the one shown below.




Fig. 2. Histogram showing the percent of animals exhibiting no seizure activity (ARS = 0-1), moderate seizures (ARS = 2-4) or severe seizures (ARS = 5-9) at various ages. The F_1 generation has audiogenic response phenotypes different from the progenitor stains. In addition, scizure intensity continues to vary after 9 weeks of age, whereas the ARSs are stable in the GEPR after this time point. The number of animals (n) tested at each age was: 3 weeks of age, n = 12; 6 weeks of age, n = 18; 9 and 11 weeks of age, n = 25.