

UCLA

UCLA Previously Published Works

Title

Determinants of Medication Adherence to Antihypertensive Medications among a Chinese Population Using Morisky Medication Adherence Scale

Permalink

<https://escholarship.org/uc/item/2j12x3cx>

Journal

PLOS ONE, 8(4)

ISSN

1932-6203

Authors

Lee, Gabrielle KY

Wang, Harry HX

Liu, Kirin QL

et al.

Publication Date

2013

DOI

10.1371/journal.pone.0062775

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Determinants of Medication Adherence to Antihypertensive Medications among a Chinese Population Using Morisky Medication Adherence Scale

Gabrielle K. Y. Lee¹, Harry H. X. Wang¹, Kirin Q. L. Liu¹, Yu Cheung², Donald E. Morisky³, Martin C. S. Wong^{1*}

1 School of Public Health and Primary Care, The Chinese University of Hong Kong, Hong Kong SAR, China, **2** New Territories East Cluster, Hospital Authority, Hong Kong SAR, China, **3** Department of Community Health Sciences, UCLA Fielding School of Public Health, United States of America

Abstract

Background and Objectives: Poor adherence to medications is one of the major public health challenges. Only one-third of the population reported successful control of blood pressure, mostly caused by poor drug adherence. However, there are relatively few reports studying the adherence levels and their associated factors among Chinese patients. This study aimed to study the adherence profiles and the factors associated with antihypertensive drug adherence among Chinese patients.

Methods: A cross-sectional study was conducted in an outpatient clinic located in the New Territories Region of Hong Kong. Adult patients who were currently taking at least one antihypertensive drug were invited to complete a self-administered questionnaire, consisting of basic socio-demographic profile, self-perceived health status, and self-reported medication adherence. The outcome measure was the Morisky Medication Adherence Scale (MMAS-8). Good adherence was defined as MMAS scores greater than 6 points (out of a total score of 8 points).

Results: From 1114 patients, 725 (65.1%) had good adherence to antihypertensive agents. Binary logistic regression analysis was conducted. Younger age, shorter duration of antihypertensive agents used, job status being employed, and poor or very poor self-perceived health status were negatively associated with drug adherence.

Conclusion: This study reported a high proportion of poor medication adherence among hypertensive subjects. Patients with factors associated with poor adherence should be more closely monitored to optimize their drug taking behavior.

Citation: Lee GKY, Wang HHX, Liu KQL, Cheung Y, Morisky DE, et al. (2013) Determinants of Medication Adherence to Antihypertensive Medications among a Chinese Population Using Morisky Medication Adherence Scale. PLoS ONE 8(4): e62775. doi:10.1371/journal.pone.0062775

Editor: D. William Cameron, University of Ottawa, Canada

Received: August 21, 2012; **Accepted:** March 26, 2013; **Published:** April 25, 2013

Copyright: © 2013 Lee et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Funding: The funding source is the Direct Grant of the Research Committee, The Chinese University of Hong Kong. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Competing Interests: The authors have declared that no competing interests exist.

* E-mail: wong_martin@cuhk.edu.hk

Introduction

Poor adherence to medications is a major public health challenge. Adherence to medication has been defined as the extent to which patients' behaviors coincide with health care providers' recommendations for health and medical advice [1]. It can be defined as the extent to which a patient's behavior, with respect to taking medication, corresponds with agreed recommendations from a healthcare provider [2]. Hypertension is one of the most common non-communicable diseases globally. More than 26% of the adult populations worldwide have been diagnosed as having hypertension, and the prevalence of hypertension increases with age [3]. Globally, it is also one of the major causes of premature death, and 7.1 million of people die from hypertension related diseases annually and the problem is still growing [4].

Successful control of blood pressure is of paramount importance in the reduction of morbidity and mortality rates [5], and many studies have demonstrated the impact of antihypertensive agents on improving clinical outcomes [6–8]. However, the effectiveness

of antihypertensive agents must be achieved by optimal adherence to prescribed medications according to healthcare providers' instructions [9].

Although the control of blood pressure has improved considerably, poor adherence with medication treatment remains a major problem among hypertensive patients, and has been identified as one of the main causes of failure in achieving blood pressure control [10]. Only 29% of hypertensive patients in the United States achieved good control, and even worse rates have been reported in Canada and European countries [11]. It is estimated that the overall adherence rates of medications were approximately 50% [12]. Among hypertensive patients who have poor blood pressure control; poor drug adherence is one of the causes, and accounts for increasingly significant and substantial public health burden. Of all avoidable hospital admissions in the United States, 33 to 69 percent are due to poor medication adherence, with expenses of approximately \$100 billion a year [13,14]. Poor adherence has been attributed to unnecessary over-prescription of drugs, substantial worsening of diseases, avoidable increases in

hospital admission rates, longer hospital stays, leading to a significant medical burden [15,16]. It is a crucial public health agenda to improve adherence with antihypertensive medications for improvements of medication-taking behavior.

Barriers to drug adherence consist of multiple factors that include complex medication regimens, dosing frequency, behavioral factors and side effects of treatment [16]. The most typical barriers to drug adherence are under the patient's control, including patient's knowledge and attitudes towards medications. Therefore, attention to these barriers is a necessary and important step to improve adherence [16]. Various methods have been developed to measure medication adherence, and they can be grouped into three categories: subjective (self-report), direct (serum or urine drug-level), and indirect (pharmacy database records, pharmacy refill rates, or pill counts) [10,17]. Each method has its strengths and drawbacks; and subjective and indirect methods are more frequently used in adherence-related studies. Self-reported measures are a relatively simple and inexpensive method, and it could also include information on social, situational and behavioral factors that affect adherence [10,17].

Antihypertensive drug adherence has been extensively evaluated in many developed countries in the West using different methods of medication adherence measurement [18–20]; however there is a scarcity of studies evaluating the adherence levels of antihypertensive pharmacotherapies among Chinese population.

This study aims to assess the drug adherence profiles of Chinese patients prescribed antihypertensive agents, and to examine the factors associated with antihypertensive drug adherence among Chinese patients. Self-reported survey as a tool to measure drug adherence was used in this study to supplement the findings of previous research reports. As this study evaluated additional factors including education level, and self-perceived health status which have not been explored among Chinese patients in previous studies [21–23], the findings of this study could provide more comprehensive information for physicians and health care policy makers with targets to improve medication adherence.

Methods

Ethics Statement

The study was approved by Joint Chinese University of Hong Kong - New Territories East Cluster Clinical Research Ethics Committee (CREC) of Hong Kong. The study was conducted within the researcher's country of residence. All participants provided their written informed consent to participate in this study.

Setting and Subjects

This study was conducted in one of the out-patient clinics in the New Territories Region from 01 February 2012 to 30 April 2012. This clinic is a public, primary care clinic run by trained family physicians. There are around 500 patient visits daily, excluding evening sessions. All adult patients aged 18 years or older; who are taking at least one long-term antihypertensive drug; and are able to communicate and understand Cantonese were invited to participate. Each potential participant was verbally asked about the nature of their chronic medication and verified by their hand-held records to ascertain study eligibility.

Study Design and Protocol

This study is a cross-sectional survey and the data were collected primarily by self-administered questionnaires. All eligible patients were invited to complete a self-administered questionnaire consisting of basic socio-demographic profile, self-perceived health

status, and self-reported medication adherence. University students who have attended a standardized workshop on study protocol and questionnaire administration consecutively screened for eligible participants and invited them to complete a survey after obtaining patient consent. For illiterate patients, the student researchers read the question items word-by-word as exactly printed on the questionnaire. Each participant had their blood pressure and body mass index (BMI) measured after survey completion. The standardized workshops aim to enhance inter-rater reliability.

Demographic and Clinical Information of the Study Participants

Participants' age, gender, level of education, marital status, number of antihypertensive drug taking, duration of taking antihypertensive drugs, educational levels, occupation, marital status and self-perceived health status were obtained through a self-reported survey. The BMI was measured by a calibrated stadiometer and weight scale for the study participants on light clothing without shoes. BMI was further classified by the following groups: underweight ($<18 \text{ kg/m}^2$); normal ($18\text{--}23 \text{ kg/m}^2$); overweight or obesity ($>23 \text{ kg/m}^2$). Self-perceived health status consists of the following categories: excellent and good, not good and not bad, poor and very poor. It was evaluated by a questionnaire item designed based on the Health Belief Model [24] using a 5-point Likert scale.

Assessment of Medication Adherence

Self-reported medication adherence was measured by the eight-item Morisky Medication Adherence Scale (MMAS-8). The MMAS-8 has been proven reliable ($\alpha = 0.83$) for assessment of adherence in patients with hypertension, and is significantly associated with blood pressure control [25–27]. Using a cut-off of 6, its sensitivity or identifying low vs. higher adherers was estimated to be 93%. and the specificity was 53% [26]. The MMAS-8 has been demonstrated to have good concurrent and predictive validity and might function as a screening tool in outpatient settings with different patient groups [26]. MMAS-8 scores can range from zero to eight in integers (see Appendix S1). The precise scoring criteria can be obtained from the developer/owner. The advantages of this instrument over other methods of measurement include its simplicity, quick administration and low-cost. A Chinese version of the MMAS-8 [28] was used and adapted specifically for antihypertensive agents in this study. The survey was in Cantonese, tailored-made for use in the Hong Kong population.

Outcomes Variables and Covariates

The outcome variable in this study is the measure of drug adherence as assessed by self-reported Morisky scores [26,27,29], transformed into a dichotomous variable with a score greater than 6 considered as "good adherence". The independent variables that were controlled for include age, gender, the total number of antihypertensive drugs used irrespective of medication class, duration of antihypertensive drug use, educational levels, marital status, occupation, monthly household income, self-perceived health status, systolic blood pressure and diastolic blood pressure. Blood pressure was measured using standardized methodology based on the 2007 guidelines for the management of arterial hypertension [30]. All patients had their BP measured twice, each separated 10 minutes apart. The average of two BP values was used in the analysis.

Statistical Analysis

The PASW Statistics version 18.0 (SPSS) was used for all data analysis. In the univariate analysis, all variables listed above were consecutively tested for significant associations with “good adherence” as the outcome variable. Chi-square tests and Student’s t-tests were used for categorical and continuous variables, respectively. All variables were unconditionally entered into a binary logistic regression model for adjustment. The absence of multicollinearity was ascertained before the regression modeling. As part of quality control, each data entered into the computer were double checked for their accuracy by random assessment of the survey responses and the data entered. All p values ≤ 0.05 were regarded as statistically significant.

Results

A total of 1,154 consecutive patients completed and returned the questionnaires. Of those, 24 (2.1%) reported not taking antihypertensive agents and 7 (0.6%) were missing, and they were excluded from the analysis. There were 1,114 completed surveys included in the analysis. The basic characteristics of these patients were shown in Table 1. Among the eligible patients, the majority of patients (60.1%) reported taking one antihypertensive agent and the mean number of antihypertensive agents used was 1.6 (SD 1.0). The mean year of antihypertensive used was 7.7 years (SD 6.9). The mean systolic blood pressure was 130.6 mmHg (± 16.9) and the mean diastolic blood pressure was 74.7 mmHg (± 9.3). The mean score of the MMAS-8 was 6.7 (± 1.4), ranging from 0 to 8. A total of 65.1% of patients obtained a score of greater than 6 and their adherence was categorized as “good”, while 32.6% of patients obtained a score of 6 or below and adherence levels were classified as “poor”.

The characteristics of patients who had good adherence to antihypertensive agents and patients who had poor adherence were mostly similar (Table 2). When compared with patients who had good adherence to antihypertensive agents, more patients who had poor adherence to antihypertensive agents were younger (39.2% vs 27.0% were aged <60), had shorter duration of antihypertensive agents used (45.2% vs 36.7% were first users of antihypertensive agents within the past 5 years), more were employed (31.9% vs 22.2%), and reported a self-perceived health status of “poor or very poor” (18.9% vs 15.2%).

All variables, which showed significant correlations with the adherence score, were entered into a multivariate logistic regression model. After adjustment of potential confounders, good adherence to antihypertensive agents was more common among those with advanced age, patients who have used antihypertensive agents for more than ten years, and those who were unemployed or retired (Table 3). As a sensitivity analysis, unemployed patients were separated from the retired and the identical logistic regression model was repeated. The associated factors found were similar to that in the original regression analysis (data not shown). In addition, patients who reported a self-perceived health status of “poor or very poor” were more likely to report poor adherence to antihypertensive agents (Table 3).

Discussion

The purpose of this study was to study the adherence profiles of Chinese patients prescribed antihypertensive agents and the factors associated with drug adherence using a self-administered questionnaire. The overall rate of good adherence to antihypertensive agents (65.1%) was not as high as a previous study among Chinese population (85.5%) [10], and was slightly lower than the

adherence rate of 71.6% among patients in Urban health Clinic Setting using the MMAS-8 in the U.S [31]. Other studies in western population reported rates ranging from 53% to 91% [32,33]. The rate of adherence may differ by many factors, including types of population, study design, method of measurement, to quote a few. It has been reported that cultural factors might influence antihypertensive adherence which could explain different levels of adherence among different populations. These include cultural health perception of hypertension, health perceptions of Western medications, self-care behavior and social support [34].

Several studies have also evaluated the factors associated with antihypertensive drug adherence [35–37]. Age, gender, the number of antihypertensive agents, and socioeconomic status were found to be associated with drug adherence. One of the studies conducted in a Chinese population [38] has found that many of the factors associated with antihypertensive drug adherence among Chinese patients were similar to those identified by studies conducted in Western populations. Among Chinese patients, there were previous studies which utilized Medication Possession Ratio (MPR) as a measure of adherence from retrospective dataset analysis [30,38–42]. These reports were different from this study which prospectively evaluated medication adherence using a self-reported instrument. We are of the view that both types of studies could be interpreted in different perspectives to inform the current situation of medication non-adherence in Chinese patients.

The association between age and drug adherence is complex and the conclusion is currently mixed in the literature. In this study, patients with older age were found to be more adherent than patients with younger age. This finding was compatible with the majority of studies that advanced age was reported to be associated with good adherence to antihypertensive agents [10,17,31,43,44]. A possible explanation is the greater number of comorbidities among elderly patients, who might perceive themselves as sicker and hence adhere better to antihypertensive prescriptions [34]. But conversely, some researchers reported the opposite correlation where poor adherence was associated with increasing age [36,45]. Other variables like individual’s cognitive level, physical mobility, and self-care abilities may affect the relationship between age and adherence.

In this study, patients with longer duration of antihypertensive agents used (over 10 years) reported better adherence than patients with shorter duration (5 years or less). This finding was consistent with a previous study using the MMAS-8 where poor antihypertensive drug adherence was more commonly found among those first diagnosed with hypertension for less than 10 years [31]. One explanation for this could be that patients having longer duration of taking antihypertensive agents could have gained more experience with hypertension, established a better physician-patient relationship and had greater faith on physicians’ advice. In addition, they might become more knowledgeable about their own health condition and the appropriate management of disease control [46]. Turning to occupation, patients who were unemployed or retired were more likely to be compliant. Few studies have reported the relationship between occupation and adherence. Some studies have shown that unemployed patients tended to have poorer adherence [47]; and the lack of health care coverage is a hidden issue that has emerged as important. Health care systems and practices of different countries may affect the result [48]. It is possible that the positive association between unemployed or retired statuses and good adherence to antihypertensive agents was mediated by having low-cost subsidy, and that medical service was relatively accessible in Hong Kong.

Table 1. Patient characteristics.

		Total study population (N = 1,114)	
		N	%
Gender	Male	464	41.7
	Female	648	58.3
Age	Mean (SD)		65.7 (11.1)
	<60	341	31.0
	60–65	242	22.0
	66–69	110	10.1
	≥70	406	36.9
No. of antihypertensive drugs used	Mean (SD)		1.6 (1.0)
	1	662	60.1
	2	325	29.5
	≥3	114	10.4
Duration of antihypertensive drugs used (year)	Mean (SD)		7.7 (6.9)
	<5 years	531	47.9
	5–10 years	336	30.3
	>10 years	241	21.8
Body Mass Index*	Underweight	26	3.2
	Normal Weight	210	25.9
	Overweight or Obesity	583	70.9
Marital status	Married	838	76.1
	Single/Divorced/Widowed/Cohabited	263	23.9
Education level	Primary or below	597	54.0
	Secondary	440	39.8
	Tertiary or above	68	6.2
Occupation	Employed	282	25.8
	Unemployed/Retired	531	48.5
	Housewife	281	25.7
Monthly Household income	0–10000	501	57.0
	10001–20000	222	25.3
	20001–30000	89	10.1
	30001+	67	7.6
Current Health Status	Excellent or Good	269	24.5
	Not Good and Not Bad	645	58.9
	Poor or Very Poor	182	16.6
Systolic blood pressure		1,081	130.6/16.9 (Mean/SD)
Diastolic blood pressure		1,080	74.7/9.3 (Mean/SD)

*Body Mass Index is calculated as weight in kg/(height in meters)².
doi:10.1371/journal.pone.0062775.t001

In addition, patients who reported a self-perceived health status of “poor or very poor” were less likely to be adherent to antihypertensive agents when compared with those who reported a self-perceived health status of “excellent or good”. Patients’ perception and attitudes towards health have been recognized as one of the barriers to drug adherence [16]. It is therefore possible that patients reported a self-perceived health status of “poor or very poor” have a relatively poorer self-control and negative beliefs about hypertension, limiting their adherence to antihypertensive medication regimens.

To our knowledge, this is the first study that evaluated the adherence profile of antihypertensive agents among Chinese

patients using the measuring tool of the MMAS-8. Self-reported survey as a tool to measure drug adherence used in this study supplemented the findings of previous studies. As additional factors including education level, and self-perceived health status where previous studies have not explored in Chinese population were evaluated in this study, the findings of this study provide clear implications to physicians and health care policy makers. It is obvious that patients with the identified associated factors should receive more intensive monitoring for drug adherence. Effective strategies to improve drug adherence can be achieved by reinforcement of regular monitoring of blood pressure at home, educational programmes involving family member, consultations

Table 2. Patient characteristics according to antihypertensive drug adherence.

		Good adherence		Poor adherence		<i>p</i>
		n	%	n	%	
Total Study Population		725	65.1	363	32.6	
Age	Mean (SD)	718	67.4 (10.9)	355	64.8 (11.2)	<0.001
	<60	194	27	139	39.2	
	60–65	149	20.8	85	23.9	
	66–69	80	11.1	27	7.6	
	≥70	295	41.4	104	29.3	
Gender		723		363		0.073
	Male	314	43.3	137	37.7	
	Female	409	56.4	226	62.3	
No. of antihypertensive used		716	1.5 (0.8)	359	1.6 (1.1)	0.813
	1	427	59.6	221	61.6	
	2	215	30	104	29	
	≥3	74	10.3	34	9.5	
Duration of antihypertensive used (year)		722	8.0 (6.8)	361	7.6 (6.9)	0.006
	<5 years	265	36.7	163	45.2	
	5–10 years	283	39.2	137	37.7	
	>10 years	174	24.1	61	16.8	
Body Mass Index (kg/m²)*		526		265		0.989
	Underweight	17	3.2	9	3.4	
	Normal Weight	133	25.3	70	26.4	
	Overweight or Obesity	376	71.5	186	70.2	
Education level		719		360		0.246
	Primary or below	402	55.9	183	50.8	
	Secondary	272	37.8	155	43.1	
	Tertiary or above	45	6.3	22	6.1	
Marital status		716		360		0.961
	Married	544	76.0	274	76.1	
	Single/Divorced/Widowed/ Cohabited	172	24.0	86	23.9	
Occupation		715		357		<0.001
	Employed	159	22.2	114	31.9	
	Unemployed/retired	386	54	137	38.4	
	Housewife	170	23.8	106	29.7	
Monthly Household income (\$HK)		571		291		0.472
	0–10000	335	58.7	155	53.3	
	100001–20000	138	24.2	80	27.5	
	20001–30000	54	9.5	33	11.3	
	30000+	44	7.7	23	7.9	
Current Health Status		719		355		0.069
	Excellent or Good	189	26.3	73	20.6	
	Not Good and not Bad	421	58.6	215	60.6	
	Poor or Very Poor	109	15.2	67	18.9	
Systolic Blood Pressure	Mean (SD)	724	130.8 (16.36)	358	130.5 (17.11)	0.704
Diastolic Blood Pressure	Mean (SD)	724	73.5 (9.04)	357	75.3 (9.43)	0.853

*Body Mass Index is calculated as weight in kg/(height in metres)².
doi:10.1371/journal.pone.0062775.t002

Table 3. Factors associated with good adherence to antihypertensive drugs.

	No.	Adjusted OR (95% C.I.)	p
Age (years)	1,042	1.016 (1.001–1.032)	0.037
Duration of taking antihypertensive drugs (years)			0.022
<5 years	492	1.000	0.065
5–10 years	319	1.334 (0.982–1.810)	0.011
>10 years	231	1.598 (1.115–2.291)	
Occupation			
Employed (Full-time/Part-time)	266	1.000	0.041
Unemployed/Retired	513	1.496 (1.017–2.200)	0.041
Housewife	263	1.012 (0.696–1.473)	0.949
Self-perceived health status			
Excellent/Good	256	1.000	1.126
Not good and Not bad	619	0.789 (0.570–1.092)	0.153
Poor/Very poor	167	0.647 (0.422–0.991)	0.045

The following variables were entered into a binary logistic regression model: age, gender, number of antihypertensive agents used, duration of antihypertensive agents used, body mass index, educational level, marital status, occupation, monthly household income, self-perceived health status, systolic blood pressure and diastolic blood pressure.

doi:10.1371/journal.pone.0062775.t003

by community pharmacists, and sending reminders of clinical appointments and regular drug taking [48].

Several limitations of our study should be addressed here. First, our sample was taken in a single out-patient clinic in the New Territories region, and patients' characteristics may differ from other districts, hence limiting its representativeness and it may not be generalizable to other health care settings. Secondly, we have excluded those hypertensive patients who were unable to communicate and understand Cantonese and this could overestimate the adherence level by including patients with higher educational levels than the general public. Also, research methodologies involving self-reported measures depend largely on individuals' memory, and recall bias may exist. In addition, many other factors such as barriers to medication taking, patients' knowledge and attitudes towards long-term drug use which could influence adherence were unable to be controlled. Lastly, the absence of association between the Morisky scores and the blood pressure levels is unexpected. One explanation is that we have not excluded patients having white-coat hypertension despite the standard methodology to measure BP. Further studies are needed to evaluate the relationship between this scale and BP levels among Chinese hypertensive patients.

Conclusions

In conclusion, some factors associated with antihypertensive drug adherence among Chinese patients using the MMAS-8 were novel findings as compared with studies conducted in Chinese patients using Medication possession ratio (MPR) [30,38–42] and other studies conducted in Western population. Since studies on

drug adherence to antihypertensive agents among Chinese population is scarce, further investigation is required to evaluate the applicability of MMAS as a tool of drug adherence measurement among Chinese patients. The factors identified in this study denote those individuals at risk for poor antihypertensive drug adherence, and future research studies should identify the reasons for their non-adherence. More drug adherence-enhancing strategies should be focused on these groups of patients, and the most effective models of which should be evaluated in separate patient groups.

Supporting Information

Appendix S1 The Morisky Medication Adherence Scale (MMAS-8).

(DOC)

Acknowledgments

Permission to use the Morisky Medication Adherence Scale was granted by Dr. Donald E. Morisky (University of California, Los Angeles) and we are grateful for his permission to reprint the questionnaire items of this scale in the manuscript.

Author Contributions

Conceived and designed the experiments: MCSW. Performed the experiments: MCSW GKYL HHXW KQLL. Analyzed the data: MCSW GKYL HHXW KQLL DEM. Contributed reagents/materials/analysis tools: YC DEM. Wrote the paper: MCSW GKYL DEM.

References

- Haynes RB, Sackett DL (1976) Compliance with therapeutic regimens: Johns Hopkins University Press.
- Sabate E, World Health Organization, WHO Adherence to Long Term Therapies Project, Global Adherence Interdisciplinary Network (2003) Adherence to long-term therapies : evidence for action. Geneva: World Health Organization. xv, 198 p.
- Kearney PM, Whelton M, Reynolds K, Muntner P, Whelton PK, et al. (2004) Global burden of hypertension: analysis of worldwide data. *Lancet* 365: 217–223.
- Organization WHO (2002) The World Health Report 2002: Reducing Risks, Promoting Healthy Life: Stylus Pub Llc.
- Lenfant C, Chobanian AV, Jones DW, Roccella EJ (2003) Seventh Report of the Joint National Committee on the Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC 7) - Resetting the hypertension sails (Reprinted from Hypertension, vol 41, pg 1178–1179, 2003). *Circulation* 107: 2993–2994.
- Turnbull F, Neal B, Algert C, Chalmers J, Woodward M, et al. (2003) Effects of different blood-pressure-lowering regimens on major cardiovascular events:

- results of prospectively-designed overviews of randomised trials. *Lancet* 362: 1527–1535.
7. Cushman WC, Ford CE, Cutler JA, Margolis KL, Davis BR, et al. (2002) Success and predictors of blood pressure control in diverse North American settings: the antihypertensive and lipid-lowering treatment to prevent heart attack trial (ALLHAT). *J Clin Hypertens (Greenwich)* 4: 393–404.
 8. Black HR, Elliott WJ, Neaton JD, Grandits G, Grambsch P, et al. (2001) Baseline Characteristics and Early Blood Pressure Control in the CONVINC Trial. *Hypertension* 37: 12–18.
 9. Sabaté E, Organization ESWH (2003) Adherence to Long-Term Therapies: Evidence for Action: Stylus Pub Llc.
 10. Burnier M (2006) Medication adherence and persistence as the cornerstone of effective anti hypertensive therapy. *American Journal of Hypertension* 19: 1190–1196.
 11. Wolf-Maier K, Cooper RS, Kramer H, Banegas JR, Giampaoli S, et al. (2004) Hypertension treatment and control in five European Countries, Canada, and the United States. *Hypertension* 43: 10–17.
 12. Haynes RB, McDonald HP, Garg AX (2002) Helping patients follow prescribed treatment - Clinical applications. *Jama-Journal of the American Medical Association* 288: 2880–2883.
 13. McDonnell PJ, Jacobs MR (2002) Hospital admissions resulting from preventable adverse drug reactions. *Annals of Pharmacotherapy* 36: 1331–1336.
 14. Levy G, Zamacona MK, Jusko WJ (2000) Developing compliance instructions for drug labeling. *Clinical Pharmacology & Therapeutics* 68: 586–591.
 15. Ogden LG, He JA, Lydick E, Whelton PK (2000) Long-term absolute benefit of lowering blood pressure in hypertensive patients according to the JNC VI risk stratification. *Hypertension* 35: 539–543.
 16. Osterberg L, Blaschke T (2005) Drug therapy: Adherence to medication. *New England Journal of Medicine* 353: 487–497.
 17. Krousel-Wood M, Thomas S, Muntner P, Morisky D (2004) Medication adherence: a key factor in achieving blood pressure control and good clinical outcomes in hypertensive patients. *Current Opinion in Cardiology* 19: 357–362.
 18. Friedman O, McAlister FA, Yun LS, Campbell NRC, Tu K, et al. (2010) Antihypertensive Drug Persistence and Compliance Among Newly Treated Elderly Hypertensives in Ontario. *American Journal of Medicine* 123: 173–181.
 19. Esposti LD, Saragoni S, Benemei S, Batacchi P, Geppetti P, et al. (2011) Adherence to antihypertensive medications and health outcomes among newly treated hypertensive patients. *ClinicoEconomics and outcomes research : CEOR* 3: 47–54.
 20. Inkster ME, Donnan PT, MacDonald TM, Sullivan FM, Fahey T (2006) Adherence to antihypertensive medication and association with patient and practice factors. *Journal of Human Hypertension* 20: 295–297.
 21. Wang PS, Bohn RL, Knight E, Glynn RJ, Mogun H, et al. (2002) Noncompliance with antihypertensive medications: the impact of depressive symptoms and psychosocial factors. *Journal of General Internal Medicine* 17: 504–511.
 22. Yiannakopoulou E, Papadopulos JS, Cokkinos DV, Mountokalakis TD (2005) Adherence to antihypertensive treatment: a critical factor for blood pressure control. *Eur J Cardiovasc Prev Rehabil* 12: 243–249.
 23. Ho PM, Bryson CL, Rumsfeld JS (2009) Medication Adherence Its Importance in Cardiovascular Outcomes. *Circulation* 119: 3028–3035.
 24. Champion VL (1984) Instrument development for health belief model constructs. *ANS Adv Nurs Sci* 6: 73–85.
 25. Krousel-Wood MA, Muntner P, Islam T, Morisky DE, Webber LS (2009) Barriers to and Determinants of Medication Adherence in Hypertension Management: Perspective of the Cohort Study of Medication Adherence Among Older Adults. *Medical Clinics of North America* 93: 753–+.
 26. Morisky DE, Ang A, Krousel-Wood M, Ward HJ (2008) Predictive Validity of a Medication Adherence Measure in an Outpatient Setting. *Journal of Clinical Hypertension* 10: 348–354.
 27. Krousel-Wood M, Islam T, Webber LS, Re RN, Morisky DE, et al. (2009) New medication adherence scale versus pharmacy fill rates in seniors with hypertension. *American Journal of Managed Care* 15: 59–66.
 28. Zhang SY, Bai LQ, Tan HZ, Yang HL (2010) Application of the morisky medication adherence scale on pulmonary tuberculosis patients. *Chinese Journal of Antituberculosis* 32: 527–530.
 29. Morisky DE, DiMatteo MR (2011) Improving the measurement of self-reported medication nonadherence: Final response. *Journal of Clinical Epidemiology* 64: 262–263.
 30. Mancia G, De Backer G, Dominiczak A, Cifkova R, Fagard R, et al. (2007) 2007 Guidelines for the management of arterial hypertension: The Task Force for the Management of Arterial Hypertension of the European Society of Hypertension (ESH) and of the European Society of Cardiology (ESC). *Eur Heart J* 28: 1462–1536.
 31. Hyre AD, Krousel-Wood MA, Muntner P, Kawasaki L, DeSalvo KB (2007) Prevalence and predictors of poor antihypertensive medication adherence in an urban health clinic setting. *J Clin Hypertens (Greenwich)* 9: 179–186.
 32. Wetzels GEC, Nelemans P, Schouten JS, Prins MH (2004) Facts and fiction of poor compliance as a cause of inadequate blood pressure control: a systematic review. *Journal of Hypertension* 22: 1849–1855.
 33. Gerth WC (2002) Compliance and persistence with newer antihypertensive agents. *Current Hypertension Reports* 4: 424–433.
 34. Billups SJ, Malone DC, Carter BL (2000) The relationship between drug therapy noncompliance and patient characteristics, health-related quality of life, and health care costs. *Pharmacotherapy* 20: 941–949.
 35. Morris AB, Li J, Kroenke K, Bruner-England TE, Young JM, et al. (2006) Factors associated with drug adherence and blood pressure control in patients with hypertension. *Pharmacotherapy* 26: 483–492.
 36. Lam P, Lum C, Leung M (2007) Drug non-adherence and associated risk factors among Chinese geriatric patients in Hong Kong. *Hong Kong Medical Journal* 13: 284.
 37. Elliott WJ, Plauschnat CA, Skrepnek GH, Gause D (2007) Persistence, adherence, and risk of discontinuation associated with commonly prescribed antihypertensive drug monotherapies. *The Journal of the American Board of Family Medicine* 20: 72–80.
 38. Wong MC, Jiang JY, Griffiths SM (2010) Factors associated with antihypertensive drug compliance in 83,884 Chinese patients: a cohort study. *J Epidemiol Community Health* 64: 895–901.
 39. Wong MCS, Jiang JY, Griffiths SM (2011) Factors associated with compliance to thiazide diuretics among 8551 Chinese patients. *Journal of Clinical Pharmacy and Therapeutics* 36: 179–186.
 40. Wong MCS, Jiang JY, Griffiths SM (2009) Short-Term Adherence to beta-Blocker Therapy Among Ethnic Chinese Patients With Hypertension: A Cohort Study. *Clinical Therapeutics* 31: 2170–2177.
 41. Wong MCS, Jiang JY, Griffiths SM (2010) Antihypertensive Drug Adherence Among 6408 Chinese Patients on Angiotensin-Converting Enzyme Inhibitors in Hong Kong: A Cohort Study. *Journal of Clinical Pharmacology* 50: 598–605.
 42. Wong MCS, Jiang JY, Griffiths SM (2009) Factors Associated With Compliance, Discontinuation and Switching of Calcium Channel Blockers in 20,156 Chinese Patients. *American Journal of Hypertension* 22: 904–910.
 43. Degli Esposti E, Sturani A, Di Martino M, Falasca P, Novi MV, et al. (2002) Long-term persistence with antihypertensive drugs in new patients. *Journal of Human Hypertension* 16: 439–444.
 44. Carter BL (2007) Adherence, Quality of Life, Cost Effectiveness, and the Role of the Pharmacist. In: Lip GYH, Hall JE, editors. *Comprehensive hypertension*. Philadelphia: Mosby Elsevier. 1119–1127.
 45. Jackevicius CA, Mamdami M, Tu JV (2002) Adherence with statin therapy in elderly patients with and without acute coronary syndromes. *Jama-Journal of the American Medical Association* 288: 462–467.
 46. Svensson S, Kjellgren KI, Ahlner J, Saljo R (2000) Reasons for adherence with antihypertensive medication. *International Journal of Cardiology* 76: 157–163.
 47. Zyczynski TM, Coyne KS (2000) Hypertension and current issues in compliance and patient outcomes. *Current Hypertension Reports* 2: 510–514.
 48. Jin J, Sklar GE, Min Sen Oh V, Chuen Li S (2008) Factors affecting therapeutic compliance: A review from the patient's perspective. *Ther Clin Risk Manag* 4: 269–286.