

UCLA

Proceedings of UCLA Health

Title

Severe Vitamin B12 Deficiency and Pancytopenia

Permalink

<https://escholarship.org/uc/item/2nt220cr>

Journal

Proceedings of UCLA Health, 26(1)

Authors

Hsu, Sherwin

An, Eric

Publication Date

2023-01-13

CLINICAL VIGNETTE

Severe Vitamin B₁₂ Deficiency and Pancytopenia

Sherwin Hsu, MD and Eric An

A 59-year-old male presented with 2 months of progressively worsening fatigue, poor oral intake, nausea, vomiting, dizziness, and unintentional weight loss of 20-lbs. He had recently been diagnosed with severe anemia (hemoglobin 6.1 grams/dL) 2 weeks prior and was transfused 2 units of red blood cells at another hospital. He denied any hematemesis, hematochezia, melena, hematuria, or prior history of anemia. He had no other known medical problems and was taking no medications and not using any recreational drugs or alcohol.

Vital signs and physical examination on presentation were all within normal limits. Initial lab findings were significant for pancytopenia with a hemoglobin of 5.6 grams/dL, mean corpuscular volume of 93.7 fL, white blood cell count of 3200 cells/mm³ (absolute neutrophil count of 1700 cells/mm³), and platelets of 77000 cells/mm³. He was transfused red blood cells with an appropriate response in his hemoglobin. Unfortunately, iron studies were not obtained prior to transfusion. Further evaluation of his pancytopenia included an international normalized ratio and prothrombin time that were within normal limits. However, he had a low haptoglobin (<15 milligrams/dL), elevated bilirubin (1.5 milligrams/dL), and elevated lactate dehydrogenase (>2900 Units/L) which were concerning for a hemolytic process. In addition, his calculated reticulocyte index was 0.55 which was consistent with a hypoproliferative process.

During his hospitalization, the patient had a CT abdomen pelvis and right upper quadrant ultrasound performed due to his nausea, vomiting, and weight loss. Imaging studies were only remarkable for borderline hepatomegaly with hepatic steatosis. When his vitamin B₁₂ level returned severely low (<146 picograms/mL) a diagnosis of pancytopenia and intramedullary hemolysis secondary to vitamin B₁₂ deficiency was made as folate level was within normal limits). He was initially started on intramuscular vitamin B₁₂ injections due to concerns about impaired oral absorption since he presented with gastrointestinal complaints. Eventually a diagnosis of pernicious anemia was confirmed after both his intrinsic factor and parietal cell antibodies returned positive. After nearly a week of intramuscular vitamin B₁₂ injections, his reticulocyte index significantly increased to 8.68. An esophagogastroduodenoscopy was scheduled as an outpatient for further evaluation of his pernicious anemia.

Discussion

Severe vitamin B₁₂ deficiency can cause extensive hematologic changes, at times mimicking features of microangiopathic hemolytic anemia, myelodysplastic syndrome, or acute leukemia.¹⁻⁴ Low B₁₂ or folate leads to impaired DNA synthesis,⁵ thus affecting all hematopoietic cell lines. Despite the association between pancytopenia and B₁₂ deficiency, only about 2.5% of patients with B₁₂ deficiency have symptomatic pancytopenia.¹ Severe B₁₂ deficiency more classically presents as megaloblastic anemia due to delayed maturation of the nucleus relative to cytoplasm of precursor RBCs. This nuclear-cytoplasmic asynchrony leads to hypersegmentation of granulocytes and intramedullary hemolysis due to ineffective erythropoiesis.^{5,6} Severe B₁₂ deficiency can also lead to increased hypercellular, dysplastic bone marrow,^{7,8} which may result in thrombocytopenia as well as neutropenia due to intramedullary destruction of myeloid cells.⁹

More typical manifestations of B₁₂ deficiency include neurological symptoms such as general weakness, unsteady gait, poor proprioception, and vibratory sensory changes.⁵ Deficiency of B₁₂, a cofactor for only two enzymes, methionine synthase and L-methylmalonyl-coenzyme A mutase, leads to accumulation of methylmalonyl CoA which inhibits fatty acid synthesis and impairs myelin synthesis. The resulting “subacute combined degeneration” involves demyelination in the dorsal and lateral columns of the spinal cord, cranial and peripheral nerves, and cortical white matter.⁵ Neurological manifestations associated with B₁₂ deficiency are helpful diagnostically but may not always occur. In some cases, the severity of megaloblastic anemia was found to be inversely correlated with the degree of neurologic dysfunction.¹⁰ Other less common conditions associated with B₁₂ deficiency include atrophic glossitis, malabsorption, and thrombosis.^{5,11}

The most common cause of severe vitamin B₁₂ deficiency is pernicious anemia which results from autoimmune gastritis destruction of gastric parietal cells.¹² Without parietal cells, there is no production of intrinsic factor to bind ingested B₁₂. A positive test for anti-intrinsic factor or anti-parietal cell antibodies confirms the diagnosis of pernicious anemia with 100% specificity. However, there is a high possibility of false negative results given its sensitivity of only 50%.⁵ Conditions causing severe malabsorption including ileal resection, gastric bypass surgery, gastrectomy, inflammatory bowel disease, and celiac disease can also lead to B₁₂ deficiency.⁵ To diagnose

malabsorption as the primary cause, B₁₂ levels would need to remain low after consumption of sufficient amounts of B₁₂.⁵ In addition, some drugs are known to cause B₁₂ deficiency, including metformin and proton pump inhibitors.⁵ Because B₁₂ is found only in animal products, a vegan diet or diet low in meat and dairy products can also lead to B₁₂ deficiency.

Treatment of vitamin B₁₂ deficiency involves parenteral or oral replacement. Most adults with B₁₂ deficiency present asymptotically and can take weekly oral vitamin repletion for a few weeks.¹³ Adults with severe B₁₂ deficiency including symptomatic anemia or neurologic findings are treated more urgently with 1000 micrograms via intramuscular injection at least several times per week for 1-2 weeks, then weekly until clear improvement is shown.^{5,13} In rare cases of severe anemia with hemodynamic instability, blood transfusion may be considered.⁵ There is usually rapid hematologic response to parenteral treatment—increase in reticulocyte count in 1 week and resolution of megaloblastic anemia in 6-8 weeks.⁵ Although neurologic symptoms may worsen transiently during treatment, these symptoms typically subside over weeks to months.¹⁴ In addition to lifelong treatment of vitamin B₁₂ deficiency for individuals with pernicious anemia,¹⁵ such individuals may also require additional gastrointestinal evaluations. Pernicious anemia is associated with increased gastrointestinal malignancies and carcinoid tumors for which endoscopic evaluation may be appropriate.¹⁶

REFERENCES

1. **Andrès E, Affenberger S, Zimmer J, Vinzio S, Grosu D, Pistol G, Maloisel F, Weitten T, Kaltenbach G, Blicklé JF.** Current hematological findings in cobalamin deficiency. A study of 201 consecutive patients with documented cobalamin deficiency. *Clin Lab Haematol.* 2006 Feb;28(1):50-6. doi: 10.1111/j.1365-2257.2006.00755.x. PMID: 16430460.
2. **Halfdanarson TR, Walker JA, Litzow MR, Hanson CA.** Severe vitamin B12 deficiency resulting in pancytopenia, splenomegaly and leukoerythroblastosis. *Eur J Haematol.* 2008 May;80(5):448-51. doi: 10.1111/j.1600-0609.2008.01043.x. Epub 2008 Jan 23. PMID: 18221385.
3. **Belen B, Hismi BO, Kocak U.** Severe vitamin B12 deficiency with pancytopenia, hepatosplenomegaly and leukoerythroblastosis in two Syrian refugee infants: a challenge to differentiate from acute leukaemia. *BMJ Case Rep.* 2014 Mar 5;2014:bcr2014203742. doi: 10.1136/bcr-2014-203742. PMID: 24599434; PMCID: PMC3948160.
4. **Konda M, Godbole A, Pandey S, Sasapu A.** Vitamin B12 deficiency mimicking acute leukemia. *Proc (Bayl Univ Med Cent).* 2019 Jul 30;32(4):589-592. doi: 10.1080/08998280.2019.1641045. PMID: 31656431; PMCID: PMC6793970.
5. **Stabler SP.** Clinical practice. Vitamin B12 deficiency. *N Engl J Med.* 2013 Jan 10;368(2):149-60. doi: 10.1056/NEJMcp1113996. PMID: 23301732.
6. **Noël N, Maigné G, Tertian G, Anguel N, Monnet X, Michot JM, Goujard C, Lambotte O.** Hemolysis and schistocytosis in the emergency department: consider pseudothrombotic microangiopathy related to vitamin B12 deficiency. *QJM.* 2013 Nov;106(11):1017-22. doi: 10.1093/qjmed/hct142. Epub 2013 Jul 10. PMID: 23842487.
7. **Parmentier S, Meinel J, Oelschlaegel U, Mohr B, Ehninger G, Schaich M, Platzbecker U.** Severe pernicious anemia with distinct cytogenetic and flow cytometric aberrations mimicking myelodysplastic syndrome. *Ann Hematol.* 2012 Dec;91(12):1979-81. doi: 10.1007/s00277-012-1488-0. Epub 2012 May 11. PMID: 22576312.
8. **Mishra VA, Harbada R, Sharma A.** Vitamin B12 and vitamin d deficiencies: an unusual cause of Fever, severe hemolytic anemia and thrombocytopenia. *J Family Med Prim Care.* 2015 Jan-Mar;4(1):145-8. doi: 10.4103/2249-4863.152276. PMID: 25811010; PMCID: PMC4366991.
9. **Boxer L, Dale DC.** Neutropenia: causes and consequences. *Semin Hematol.* 2002 Apr;39(2):75-81. doi: 10.1053/shem.2002.31911. PMID: 11957188.
10. **McCaddon A.** Vitamin B12 in neurology and ageing: clinical and genetic aspects. *Biochimie.* 2013 May;95(5):1066-76. doi: 10.1016/j.biochi.2012.11.017. Epub 2012 Dec 7. PMID: 23228515.
11. **Graells J, Ojeda RM, Muniesa C, Gonzalez J, Saavedra J.** Glossitis with linear lesions: an early sign of vitamin B12 deficiency. *J Am Acad Dermatol.* 2009 Mar;60(3):498-500. doi: 10.1016/j.jaad.2008.09.011. PMID: 19231648.
12. **Bizzaro N, Antico A.** Diagnosis and classification of pernicious anemia. *Autoimmun Rev.* 2014 Apr-May;13(4-5):565-8. doi: 10.1016/j.autrev.2014.01.042. Epub 2014 Jan 11. PMID: 24424200.
13. **Devalia V, Hamilton MS, Molloy AM; British Committee for Standards in Haematology.** Guidelines for the diagnosis and treatment of cobalamin and folate disorders. *Br J Haematol.* 2014 Aug;166(4):496-513. doi: 10.1111/bjh.12959. Epub 2014 Jun 18. PMID: 24942828.
14. **Healton EB, Savage DG, Brust JC, Garrett TJ, Lindenbaum J.** Neurologic aspects of cobalamin deficiency. *Medicine (Baltimore).* 1991 Jul;70(4):229-45. doi: 10.1097/00005792-199107000-00001. PMID: 1648656.
15. **Annibale B, Lahner E, Fave GD.** Diagnosis and management of pernicious anemia. *Curr Gastroenterol Rep.* 2011 Dec;13(6):518-24. doi: 10.1007/s11894-011-0225-5. PMID: 21947876.
16. **Shah SC, Piazuolo MB, Kuipers EJ, Li D.** AGA Clinical Practice Update on the Diagnosis and Management of Atrophic Gastritis: Expert Review. *Gastroenterology.* 2021 Oct;161(4):1325-1332.e7. doi: 10.1053/j.gastro.2021.06.078. Epub 2021 Aug 26. PMID: 34454714; PMCID: PMC8740554.