

UCSF

UC San Francisco Previously Published Works

Title

The Inclusion of Women in Global Oncology Drug Trials Over the Past 20 Years

Permalink

<https://escholarship.org/uc/item/2pb3b5w5>

Journal

JAMA Oncology, 7(10)

ISSN

2374-2437

Authors

Jenei, Kristina
Meyers, Daniel E
Prasad, Vinay

Publication Date

2021-10-01

DOI

10.1001/jamaoncol.2021.3686

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

Letters

RESEARCH LETTER

The Inclusion of Women in Global Oncology Drug Trials Over the Past 20 Years

Thirty years have passed since the enactment of the National Institutes of Health (NIH) Revitalization Act, which encouraged NIH-funded investigators to include adequate numbers of women in clinical studies.¹ Since then, there have been important steps taken to ensure better representation of women and racial and ethnic minority groups in biomedical trials.² However, lack of representation remains problematic in oncology. Previous research suggests that women represent approximately 30% to 40% of participants in trials leading to drug approvals in the US.^{3,4}

In 2010, the NIH Office of Research on Women's Health⁵ set forth a vision to advance the understanding of sex-specific disease differences by 2020. As clinical trials become international in scope, we sought to evaluate the global movement toward this vision in oncology. In this cohort study, we reviewed enrollment patterns of completed cancer drug trials over the past 20 years to compare sex-specific trial participation to current cancer incidence rates. Using data from the International Agency for Research on Cancer (IARC),⁶ we

identified 6 common solid tumor types for women (lung, colon, thyroid, melanoma, kidney, and pancreas).

Methods | We conducted a systematic search of cancer drug trials registered on ClinicalTrials.gov between 2000 and 2020 for 6 common cancers for women (excluding breast and uterus). The investigation of publicly available trial protocols was exempt from institutional review board approval. We included completed drug trials with results for adults (≥ 18 years). Behavioral, device, procedure, and radiotherapy trials were excluded (eFigure in the Supplement). Pearson χ^2 test of independence was used to evaluate the association between female and male enrollment and phase (1, 2, 3), tumor type, and funding source. *P* values used 2-tailed tests, with *P* < .05 being significant. We explored changes in sex-specific enrollment patterns over 2 decades by stratifying trials as 2000 to 2010 and 2011 to 2020. Cancer incidence rates were calculated by dividing the number of new male and female cases by the total number of new cases per tumor type (2020).⁶ Data analysis was performed using R statistical software, version 1.1.463 (R Foundation).

Results | We identified 505 oncology clinical trials that met the eligibility criteria between 2000 and 2020. Of the total 182 416 participants, 73 103 (40%) were women, while 109 313 (60%) were


Table. Comparison Between Sex-Specific Enrollment and Clinical Trial Characteristics

Characteristic	Sex, No. (%)		P value ^a
	Female	Male	
Total enrolled	73 103 (40)	109 313 (60)	<.001
Trial phase			
1	3034 (48)	3322 (52)	.001
2	18 838 (43)	24 508 (57)	<.001
3	40 139 (38)	66 611 (62)	<.001
Year			
2000-2010	23 350 (40)	34 745 (60)	<.001
2011-2020	49 753 (42)	68 022 (58)	
Tumor type			
Lung	40 829 (41)	57 979 (59)	<.001
Colon	7600 (33)	15 266 (67)	<.001
Thyroid	904 (51)	875 (49)	.50
Melanoma	11 317 (42)	15 529 (58)	<.001
Kidney	6586 (33)	13 127 (67)	<.001
Pancreas	5867 (47)	6537 (53)	<.001
Sites			
US	49 911 (40)	75 755 (60)	<.001
Canada	29 603 (39)	45 372 (61)	<.001
China	23 456 (41)	33 645 (56)	<.001
United Kingdom	28 472 (39)	44 478 (61)	<.001
Australia	28 505 (39)	44 332 (61)	<.001
Funding (US)			
Industry	41 391 (41)	60 473 (59)	<.001
NIH	6828 (48)	7285 (52)	

Abbreviation: NIH, National Institutes of Health.

^a *P* values from Pearson χ^2 test of independence.

Figure. Composition of Trial Enrollment and Incidence by Sex per Tumor Type



^a New cases by sex per International Agency for Research on Cancer 2020 data.⁶

were men. We observed significant differences between the enrollment of sexes across all trial phases and tumor types except thyroid (Table). Drug trials for colon and kidney cancer enrolled the least number of women, with 33% participation rates (Figure).

Comparing 2000 to 2010 with 2011 to 2020, we observed a marginal increase in female participation, from 40% to 42%, respectively ($P < .001$). We evaluated the association between industry and NIH-sponsored studies in the US ($n = 358$) and found that NIH-funded trials enrolled a higher proportion of women (48%) compared with industry trials (41%) ($P < .001$).

Discussion | It is essential that women be enrolled in clinical trials in numbers that, at least, mirror the distribution of the disease in the population so that potential biological differences can be understood. Our analysis, although limited to 6 tumor types, suggests that sex differences persist in clinical trials. In 2020, women represented 77% of newly diagnosed thyroid cancer cases worldwide⁶ yet comprised only 51% of participants in trials investigating thyroid drugs (Figure). Similarly, women represented 48% of global colon cancer cases⁶ yet accounted for only 33% of trial participants for colon cancer therapeutics (Figure).

Higher enrollment of women in NIH-funded studies (48%) compared with industry studies (41%) warrants further exploration, but the stagnant rates of women in trials suggest that regulatory initiatives² over the past 2 decades may be insufficient. Our analysis, although limited to sponsor- and manufacturer-disclosed information, demonstrates that persistent inequities remain in the recruitment of female participants in trials investigating new therapeutics for certain tumor types in oncology.

Kristina Jenei, BSN, MSc
Daniel E. Meyers, MD, MSc
Vinay Prasad, MD, MPH

Author Affiliations: School of Population and Public Health, The University of British Columbia, Vancouver, British Columbia, Canada (Jenei); Department of Medicine, University of Calgary, Calgary, Alberta, Canada (Meyers); Department of Epidemiology and Biostatistics, University of California, San Francisco (Prasad).

Accepted for Publication: June 21, 2021.

Published Online: August 26, 2021. doi:10.1001/jamaoncol.2021.3686

Corresponding Author: Kristina Jenei, BSN, MSc, School of Population and Public Health, The University of British Columbia, 2206 East Mall, Vancouver, BC V6T 1Z3, Canada (kjenei@mail.ubc.ca).

Author Contributions: Ms Jenei had full access to all of the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Concept and design: All authors.

Acquisition, analysis, or interpretation of data: Jenei, Meyers.

Drafting of the manuscript: Jenei.

Critical revision of the manuscript for important intellectual content: All authors.

Statistical analysis: Jenei.

Administrative, technical, or material support: Meyers.

Supervision: Prasad.

Conflict of Interest Disclosures: Dr Prasad reported receiving research funding from Arnold Ventures during the conduct of the study; and royalties from Johns Hopkins Press, Medscape, and MedPage; consulting fees from UnitedHealthcare; speaking fees from New Century Health and Evicore; and personal fees from Patreon Plenary Session (podcast has Patreon backers) outside the submitted work. No other disclosures were reported.

1. NIH Revitalization Act of 1993 Public Law 103-43. In: Mastroianni AC, Faden R, Federman D, eds. *Women and Health Research: Ethical and Legal Issues of Including Women in Clinical Studies: Volume I*. National Academies Press; 1994:appendix B. Accessed May 5, 2021. <https://www.ncbi.nlm.nih.gov/books/NBK236531/>
2. Mazure CM, Jones DP. Twenty years and still counting: including women as participants and studying sex and gender in biomedical research. *BMC Womens Health*. 2015;15(1):94. doi:10.1186/s12905-015-0251-9
3. Murthy VH, Krumholz HM, Gross CP. Participation in cancer clinical trials: race-, sex-, and age-based disparities. *JAMA*. 2004;291(22):2720-2726. doi:10.1001/jama.291.22.2720
4. Duma N, Vera Aguilera J, Paludo J, et al. Representation of minorities and women in oncology clinical trials: review of the past 14 years. *J Oncol Pract*. 2018;14(1):e1-e10. doi:10.1200/JOP.2017.025288
5. National Institutes of Health Office of Research on Women's Health. Strategic plan. moving into the future with new dimensions and strategies: a vision for 2020 for women's health research. Accessed July 22, 2021. https://orwh.od.nih.gov/sites/orwh/files/docs/ORWH_StrategicPlan2020_Vol1.pdf
6. Global Cancer Observatory. Accessed June 16, 2021. <https://gco.iarc.fr/>

Use of an Analytics and Electronic Health Record-Based Approach for Targeted COVID-19 Vaccine Outreach to Marginalized Populations

Equity in vaccine outreach and delivery has been prioritized given the disproportionate harms of the COVID-19 pandemic on communities of color and those with lower socioeco-