

UCSF

UC San Francisco Previously Published Works

Title

Depression Among Older Adults with Diabetes Mellitus

Permalink

<https://escholarship.org/uc/item/2qj0w393>

Journal

Clinics in Geriatric Medicine, 31(1)

ISSN

0749-0690

Authors

Park, Mijung
Reynolds, Charles F

Publication Date

2015-02-01

DOI

10.1016/j.cger.2014.08.022

Peer reviewed



Published in final edited form as:

Clin Geriatr Med. 2015 February ; 31(1): 117–137. doi:10.1016/j.cger.2014.08.022.

Depression among older adults with diabetes mellitus

Mijung Park, PhD MPH RN [Assistant Professor] and

Department of Health and Community Systems, University of Pittsburgh, School of Nursing, 3500 Victoria Street, 421 Victoria building, Pittsburgh, PA 15213

Charles F. Reynolds III, MD [UPMC Endowed Professor of Geriatric Psychiatry Director]

NIMH Center of Excellence in Late Life Depression Prevention and Treatment, Hartford Center of Excellence in Geriatric Psychiatry, and Aging Institute of UPMC Senior Services and the University of Pittsburgh, 3811 O'Hara Street, Pittsburgh, PA 15213-2582, Telephone: 412-246-5991, ReynoldsCF@upmc.edu

Synopsis

Depression is among the leading causes of decreased disability-adjusted life years in the world¹ and a serious public health problem.² Older adults with DM experience greater risk for comorbid depression compared to those who do not have DM.³ Having DM increases the risk of subsequent development or recurrence of depression. Conversely, history of depression increases the risk for new onset DM.⁴ As an unwanted co-traveler of DM, undetected, untreated or undertreated depression impinges an individual's ability to manage their DM successfully, hindering their adherence to treatment regime.⁵ It also undermines the effectiveness of provider-patient communication and decays therapeutic relationships. Thus, in the context of caring for older adults with DM, comorbid depression presents special challenges and opportunities for clinicians. Moreover, recent studies have suggested that co-occurring depression and DM may accelerate cognitive decline, highlighting the importance of treating depression and DM. Several treatment modalities are available, which can be used to treat and manage depression in primary care settings: pharmaceutical, brief psychotherapeutic, behavioral and life style interventions, and combination therapies. An evidence-based health care delivery model is also available for treating depression in primary care settings. In this article, we summarize the clinical presentation of late-life depression, potential mechanisms of comorbidity of depression and DM, importance of depression in the successful management of DM, and available best practice models for depression treatment.

Keywords

Diabetes; Depression; Mood disorders; Aging; Collaborative Care

© 2014 Elsevier Inc. All rights reserved.

Mijung Park, PhD MPH RN (corresponding author): Telephone: 412-624-9647, parkm@pitt.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Introduction

Diabetes mellitus (DM) is one of the most common chronic conditions among older adults. About 26.9%, or 10.9 million U.S. residents aged 65 years and older had diabetes in 2010.⁶ Depressive disorders are serious chronic diseases that increase morbidity and mortality,⁷ erode quality of life,⁸ and increase medical expenditure. Depression and DM often co-occur. Data from a range of settings suggest that the prognosis of both DM and depression—in terms of severity of disease, complications, treatment resistance and mortality—is worse for either disease when they are co-morbid than when they occur separately. Comorbid depression in patients with DM is strongly associated with increased burdens of DM symptoms⁹, poor self-management and treatment adherence¹⁰, increase in health care services utilization and medical expenditures¹¹ and an increased risk of DM complications.¹² DM complications such as myocardial infarction, amputation or loss of vision can in turn precipitate or worsen depressive episodes. Yet, few studies have extensively examined the associations between depression and DM in the older adults populations. Also, recent studies have found that combination of DM and depression may increase the risk for dementia, suggesting increased brain toxicity.¹³ The purpose of this review is to summarize the clinical presentation of late-life depression among older adults with DM, potential mechanisms of comorbidity of depression and DM, importance of depression in the successful management of DM, and available best practice models for depression treatment.

Disease description

Depression is a mood disorder that causes a persistent feeling of sadness and loss of interest. To be diagnosed with major depression, a patient must have depressed mood or anhedonia and at least 5 of the following 9 symptoms nearly every day for at least 2 weeks:

- Depressed mood
- Marked diminished interest/pleasure
- Sleep disturbance (increased or decreased sleep)
- Appetite disturbance (increased or decreased appetite; typically with weight change)
- Fatigue/loss of energy
- Diminished concentration or indecisiveness
- Feelings of worthlessness or excessive or inappropriate guilt
- Psychomotor retardation or agitation (a change in mental and physical speed perceived by other people)
- Recurrent thoughts of death or suicide (not just fear of dying)

They must also experience functional impairment related to these depressive symptoms.

Older adults, however, do not always present with the typical symptoms of depression. In particular, depressed or sad mood may be less evident or not even present. In these cases,

anhedonia may be better indicator for depression.^{14,15} Depressed older adults may experience sleep disturbances (sleeping too much or too little) or changes in appetite (eating too much or too little). Decrease in self-efficacy, motivation, and ability to participate in self-care may also indicate underlying depressive symptoms. Signs of such symptoms can be subtle. Older adults may reply with “I don’t know” to simple questions, decline to participate in physical, speech or occupational therapy, and feeling negative or hopeless about treatments offered. Some may easily give up tasks during these therapies. Older adult who experience loss of self-worth or sense of loneliness due to depression, may complain “nobody needs me,” or “I feel I am just in everyone’s way.” The symptoms of late-life depression are often attributed to normal aging, grief, physical illness or dementia and providers and patients miss important opportunities to initiate treatment for what is an treatable health problem.¹⁶

The majority of older adults with clinically significant depressive symptoms do not meet standard diagnostic criteria for major depression or dysthymic disorder.^{17,18} Patients in this group fall short of meeting diagnostic criteria for major depression because of fewer or limited duration of depression symptoms. Nonetheless, studies suggest that these patients carry similar disease burden: poorer health outcomes, functional impairment, and higher health utilization and treatment costs.¹⁹⁻²¹ Moreover, these patients are at very high risk for subsequently developing major depression and may also develop suicidal ideation.²²⁻²⁴

Although depression can be successfully treated,²⁵ many older adult suffer from a chronic or recurrent depression.²⁶ In a prospective study about courses of major depression, approximately 85% of individuals who recover from depression experience recurrence within 15 years.²⁷ Findings meta-analyses of chronic depression in primary care and community sample suggests that about one in three depressed older adults experience chronic and persist course.^{28,29} Depressed patients with DM are at greater risk for a chronic course of depression or less complete recovery.³⁰ Such chronicity, in turn, makes it more difficult for older adults and their family caregivers to optimally self-manage DM.

In the older adults with DM, depressive symptoms may be overlooked because they are assumed to be due to concurrent DM and other medical illnesses. Many of the symptoms of depression such as lower energy, fatigue, loss of appetite, and sleep disturbance are also associated with DM. Thus differentiating stress related to DM self-management and depression can be challenging.³¹ Somatic complaints may suggest presence of depression, especially if they are out of proportion to underlying physical disorders.³² Only 25-30% of primary care patients present with purely affective or cognitive symptoms of depression.³³

Risk factors

Risk factors for developing depression after age 65 are similar to those in younger individuals and include being female, unmarried, low socioeconomic status, having chronic physical illness, social isolation, a history of depression, and a family history. The risk of major depression increases up to three-fold if a first-degree relative has the illness.³⁴ Additional risk factors that are particularly important in older adults include loss and grief,³⁵ social isolation or limited social support,³⁶ high degrees of family conflict,^{37,38} and care-

taking responsibilities.³⁹ Other risk factors that increase the likelihood of depression in the medically ill elderly include presence of cognitive impairment, age greater than 75, active alcohol abuse, and lower educational attainment.⁴⁰⁻⁴⁹ Table 1 summarized the risk factors for depression in older adults.

Protective factors

Strong and supportive social context has been identified as a protective factor. A compelling body of evidence has shown that social support decreases the risks for depression^{50,51} and for depression relapse,⁵² increases adherence to depression treatment,^{53,54} and improves treatment outcomes. Social support may have positive effects on psychological wellbeing independent of whether or not individuals are exposed to stress.^{55,56} Social support may also promote wellbeing through modulation of neuro-endocrine response to stress.⁵⁷

Social activities, such as volunteering suggested to have positive effect on depression outcomes in older adults.⁵⁸ Studies have also suggested that religion and spirituality may play an important part in many older adults' lives and social connectedness and support are an important part of organized religion.⁵⁹ Religion may allow older adults to experience life as meaningful despite losses and challenges and, thereby, reduce the risk of depression. It is also possible that the positive effect of religion on mental health is mediated by the social connectedness and the social support derived from taking part in religious activities.

Prevalence/Incidence

Depression is one of the most common mental disorders in late-life.⁶⁰ About one in four U.S. resident projected to experience major depression by age 75.⁶¹ In community settings, about 5% of adults aged 65 and older meet research diagnostic criteria of major depression,^{62,63} with rates of subsyndromal, clinically significant depression estimated at 8% -16%.⁶⁴ The rates of geriatric depression increase to 12-30% in institutional settings, and up to 50% for residents in long-term care facilities.^{65,66} Approximately 5-10% of older adults seen in primary care settings have clinically significant depression.⁶⁷

Variations in prevalence in late-life depression across racial/ethnic group are rarely examined and existing literature present mixed findings. However, given that differences in diabetes prevalence exist by racial/ethnic group, these may be important to appreciate. Cross-sectional and longitudinal epidemiological study showed that Non-Hispanic White and Latino older adults have higher rates of depression.⁶⁸⁻⁷⁰ A longitudinal population study of community-dwelling Hispanic older adults showed approximately 9% of sample met the criteria for lifetime diagnosis of major depression, while 24% of sample reported minor depression.⁷⁰ The prevalence of depression in African American is generally lower than their White counterparts.⁷¹

Minority older adults are less likely to be diagnosed with or treated for depression than their white counterparts.^{72,73} These health service disparities in minority populations become increasingly complicated when considering cultural beliefs and attitudes towards depression care. Culture influences how individuals experience and express depression.^{74,75} Minority patients from certain ethnic groups may express their depression somatically than

psychologically.^{33,76} Such somatic presentations may reduce the recognition of depression by primary care providers or leads to the perception of a patient as ‘difficult.’⁷⁷ Some minorities may also have less faith in the biological etiology of depression, be more skeptical about antidepressant use, and show stronger preferences for counseling than their White counterparts.^{78,79} When pharmaceutical treatment is the only available option, minority older adults may be less likely to engage in treatment and more likely to be non-adherent. The present primary care systems that focus primarily on pharmacological treatment without considering the unique barriers faced by ethnic and racial minority populations may not affect the pattern of disparities observed.⁸⁰

Scant information on the cross-national prevalence of late-life depression is available. A report from a World Health Organization concluded that older adults in developed countries had relatively low average depression rates of 2.6% while those in developing countries had an average rate almost three times higher (7.5%).⁸¹ The rapid increase in diabetes across the world, particularly in developing countries,^{82,83} indicate that these international discrepancies are critical to understand.

Among older adults with DM, depression is highly prevalent.⁸⁴ Up to 30% of individuals with DM have a significant number of depressive symptoms and 12 to 18% meet diagnostic criteria for major depression.^{3,85} Patients with DM experience significantly higher rates of depression compared with their age- and gender-matched counterparts.⁸⁴ A meta-analysis of ten studies showed that the prevalence of depression was significantly higher in patients with DM compared with those without (17.6 vs. 9.8%).⁸⁴ The prevalence of depression was higher in females with diabetes (23.8%) compared with males (12.8%); however, the odds ratio for depression in patients with Type 2 DM compared with those without was higher in males (OR = 1.9, 95% CI 1.7-2.1) than females (OR = 1.3, 95% CI 1.2-1.4).

Depression and mortality in individuals with DM

Based on several meta-analyses, depression is associated 1.5- to 2.6-fold increase of mortality among individuals with in DM.^{7,86,87} Few studies have examined if treating depression may decrease mortality among individuals with depression. Data from a large clinical trial of collaborative late-life depression treatment program in primary care setting (PROSPECT) showed that evidence-based treatment of depression can reduce mortality rates among those with DM (adjusted hazard ratio 0.49, 95% CI: 0.24-0.98).⁸⁸ Another study of all cause mortality with same sample also found that patients with major depression in intervention practice were 24% less likely to have died, compared to depressed older adults in usual primary care practices.⁸⁹

Underlying mechanisms of comorbidity of depression and DM

Katon⁹⁰ proposed a complex bidirectional relationship between depression and type 2 DM (Figure 1). Depression early in adult life is a risk factor for subsequent development of DM.⁹¹ The increased risk of DM in patients with depression has been hypothesized to be the result of maladaptive health risk behaviors associated with depression such as smoking, obesity and lack of physical exercise⁹⁰ as well as psychobiologic factors such as increased cortisol levels, increased inflammatory factors⁹² and insulin resistance.⁹³

On the other hand, DM may increase risk of depression or worsen the depression symptoms due to increased symptom burden, DM complications causing functional impairment and decreased quality of life, as well as vascular brain changes secondary to DM. Comorbid depression has been found to impair the ability to perform self-care activities necessary to control DM by affecting memory, energy level, and executive function.⁹⁴⁻⁹⁶ Lack of self-care and the psychobiologic changes associated with comorbid depression may explain why individuals with comorbid depression experience increased risk of macro- and micro-vascular complications¹² and dementia.¹³

While DM and depression independently associated with memory and cognitive impairment,⁹⁷⁻⁹⁹ recent clinical epidemiological studies found that concurrent DM and depression may have synergistic negative effect on brain health, posing greater risk for developing dementia and Alzheimer's disease.¹³ There are several biological mechanisms that can explain such a toxic effect of the DM and depression combination on brain health. First, DM and depression are risk factors for cardiovascular and cerebrovascular diseases (e.g., vascular dementia and Alzheimer's disease). Concurrent DM and depression may increase risk for cardiovascular and cerebrovascular events in additive fashion.¹³ Secondly, depression is associated with dysregulation of the hypothalamic-pituitary axis,^{100,101} which increases glucocorticoid production, and impairs negative feedback. Hypercortisolemia is associated with metabolic syndrome—a risk factor for vascular dementia and Alzheimer's disease.¹⁰²⁻¹⁰⁴ Finally, chronic or recurrent depression is associated with hippocampal atrophy.^{105,106}

The temporal relationship between depression and DM has not been fully established. Existing evidence suggests that the association between history of depression and subsequent development of DM is stronger than that between history DM and subsequent development of depression.¹⁰⁷ Several prospective epidemiological studies have found that having depression increased risk for developing DM subsequent years by 1.62-to 2.52-fold: Demakakos and colleagues¹⁰⁸ analyzed data of 6,111 individuals who reported not having doctor-diagnosed DM at baseline in 2002-2003 and concluded that, after adjusting for several demographic and clinical characteristics, individuals who had greater depressive symptomatology at baseline experienced 1.62-fold increased risk for developing DM in the subsequent 45.8 months. In another longitudinal study of community-residing adult population by Camethon et al, after adjusting for age, race, and gender, the adjusted risk of developing DM in subsequent years are 2.52-fold higher among those with greater depressive symptoms than those with low depressive symptoms.¹⁰⁹ A prospective epidemiological study of 1,715 residents of Baltimore Catchment area showed that having major depressive disorder was associated with more than 2-fold increase in risk (adjusted odds ratio: 2.23; 95% CI 0.90-5.55) for developing DM in the subsequent 13 years.¹¹⁰ However, this was not statistically significant.

Having DM increases the risk for developing depression subsequently. A meta-analysis of 11 epidemiological studies concluded that individuals who had Type 2 DM without depression at baseline experienced 1.24-fold increased risk (95% CI 1.09-1.40) for subsequently developing or recurring depression in subsequent years.¹¹¹ Data from the study of 5,201 multi-racial participants of Women's Health Across the Nation (SWAN) concluded

that having DM increase the odds for developing DM in the subsequent 2 years by 2.8-fold (95% CI: 1.2-6.4) among depressed African American Women.⁹¹ Therefore, older adults with diabetes should be considered at higher risk for future depression.

Prevention and early treatment

Prevention and early treatment of depression matter to providers who treat individuals with DM for several reasons. First, as mentioned earlier, having depression may increase the risk for subsequently developing DM. Thus, prevention and early treatment of depression may lower the incidence of DM. Second, having comorbid depression decreases individuals' ability to successfully self-manage DM and increases likelihood for poor DM outcomes. Thirdly, depression is treatable and several treatment approaches are developed for primary care settings. In a recent clinical trial with 247 older adults¹¹² suggests that both problem solving therapy for primary care (PST-PC) and coaching in health dietary practices could effectively reduce the incidence of major depression in older adults, including in older black adults with high BMI. Despite the fact that black participants in the trial carried a greater risk for depression than did their white counterparts – including obesity –the interventions were equally effective in reducing the incidence of major depression from an expected rate of 20-25% over 2 years to only 8-9%, in both black older adults and white older adults. Thus, given the challenges of treating prevalent depression in people living with both depression and DM mellitus, especially in minority populations –where stigma represents an important barrier, it seems particularly important to develop depression prevention strategies that utilize, variously, active coping strategies and life style interventions that promote health and protective factors.

The U.S. Preventive Services Task Force found at least fair evidence that screening adults for depression improves health outcomes and that benefits outweigh harms (B rating), and recommends screening for depression if practices have systems in place to assure accurate diagnosis, effective treatment, and follow-up.¹¹³ Based on these recommendations, the Centers for Medicare & Medicaid Services recently determined to cover annual screening for depression for Medicare beneficiaries in primary care settings with staff-assisted depression care supports.

Treatment strategies

In managing DM, clinicians use several clinical data to establish a baseline and treatment goals – e.g. fasting serum glucose level and HgA1C. The same stepped, measurement-based treatment principles can be applied to treating depression: establishing a baseline; setting clinical and functional goal for depression; and assessing patient's progress accordingly. Figure 2 illustrates one of widely used clinical guidelines for depression treatment.

Various treatment options are available for late-life depression: antidepressant medications, psychotherapy, or a combination. Other options include exercise programs or other life-style modifying interventions and electroconvulsive therapy (ECT). Several meta-analyses showed that combination therapy is more efficacious than monotherapeutic approaches in treating and in preventing relapse.¹¹⁴⁻¹¹⁷ In a recent meta-analysis of 14 RCTs of intervention for depression among patient with diabetes, all treatments have shown to be

effective in reduction of depressive symptoms ($g=-0.51$, 95% CI: $-0.63, -0.39$).¹¹⁸ The pooled effect size was large in psychotherapeutic interventions combined with diabetes self-management education ($g= -0.58$, 95% CI $-0.77, -0.39$); and pharmacological treatment showed moderate pooled effect size ($g=-0.47$; 95% CI -0.67 to -0.27).¹¹⁸ Unlike major depression, subsyndromal depressive conditions have a relatively small evidence base regarding treatments; existing data suggest that available therapies have modest effect sizes when compared to usual care or placebo.^{16,119,120}

Targeting interventions for patients with minor and subsyndromal depression may prove useful as both primary and secondary prevention strategies and clinicians should watch such patients carefully because of the high risk of worsening depression, especially if patients have experienced prior episodes of major depression. Psychosocial treatments may be more helpful than medications for older adults with less severe forms of depression.¹²¹

Treatment delivery model for comorbid depression and DM

To date, more than 40 randomized controlled trials have established a robust evidence base for an approach called 'collaborative care models (also called integrated care models).¹²²⁻¹²⁵ Recently, several trials of collaborative care model have documented the effectiveness of treating comorbid depression and DM^{125,126} (e.g., PATHWAY¹²⁶) and comorbid depression, and DM and/or coronary heart disease (TEAMcare¹²⁷). In such programs, a depression care manager (usually a nurse or clinical social worker) supports medication management prescribed by PCPs through patient education, close and proactive follow-up, and brief, evidence-based psychosocial treatments such as behavioral activation or problem solving treatment in primary care. The care manager may also facilitate referrals to additional services as needed. A psychiatric consultant regularly (usually weekly) reviews all patients in the care manager's caseload who are not improving as expected and provides focused treatment recommendations to the patient's PCP.^{122,128-130}

Tools for assessing and tracking depression

Several tools for assessing depression severity are available and can be easily administered by office staff or physicians during a clinic care visit. The most frequently used tool is the 9-item Patient Health Questionnaire (PHQ-9)¹³¹, which systematically explores the 9 DSM-IV¹³² symptoms of major depression. A score of 10 or greater on the PHQ-9 indicates an elevated level of depressive symptomatology and increased risk of clinically significant depression. PHQ-9 have good sensitivity and specificity (both about 88%) for detecting depressive disorders.^{133,131}

Pharmacological management

Approximately 80% of antidepressant prescribed in the U.S. is prescribed in primary care settings. More than 20 antidepressant medications have been approved by the FDA for treatment of depression in older adults. Newer generations of antidepressants such as SSRIs and SNRIs have more tolerable side effect profile compared to tricyclic antidepressants (TCAs).^{134,135} Comparing second generation antidepressants, studies found no evidence that specific medications are more effective than others.¹³⁶ Thus, physicians may discuss the choice of medication with patients and their family. Because of the changes in

pharmacokinetics (e.g., decreased renal or hepatic clearance), older adults may require lower doses of medications than their younger counterparts. Furthermore, considering high rates of multimorbidity and polypharmacy among older adults, drug-drug interactions are particular concern in this population. On the other hand, medication doses should be titrated upwards to full adult doses in patients who experience partial responses without substantial side effects. Table 2 summarizes common antidepressant medications.

Psychological management

Older adults patients with depression tend to prefer psychotherapy over medications.¹³⁷ Several psychotherapeutic modalities are effective for late-life depression¹³⁸⁻¹⁴⁰: cognitive behavioral therapy (CBT),^{121,141} interpersonal therapy (IPT),¹⁴² problem solving treatment (PST),¹⁴³ and Behavioral Activation and Pleasant Activity Scheduling (BA).¹⁴⁴⁻¹⁴⁶ A recent meta-analysis of 44 studies comparing psychotherapies and control conditions,¹⁴⁰ concluded that the overall effect size of psychotherapy over control group were $g = 0.64$ (95% CI: 0.47-0.80). Although such treatments can be effectively provided to older adults in primary care,^{147,148} they are not widely available in most primary care settings today and we encourage clinicians to develop relationships with mental health specialists who can offer such treatments or to train a staff member in the clinic to provide evidence-based brief psychotherapies. Table 3 summarizes brief- and evidence-based psychotherapeutic approaches that can be easily administered in the primary care context and by nurses, social workers, or other licensed psychotherapists.

Electroconvulsive Treatment (ECT)

ECT is an important and viable treatment option for severely depressed older adults. ECT should be strongly considered for patients who have severe, persistent depression that does not respond to several trials of antidepressant medications or psychotherapy and/or puts patient at high risk of harm (e.g. severe weight loss, malnutrition, refusal of food, suicidal ideation). Poor tolerance or limited response to medications and a history of successful treatment with ECT are also indications for ECT. Rates of ECT use in depressed adults vary substantially.^{149,150} Older adults are more likely to receive ECT than younger adults, African Americans are less likely to receive ECT than Whites and individuals with poor health insurance or living in rural areas are also less likely to receive ECT.¹⁴⁹

Summary/Discussion

DM and depression is frequently comorbid in older adult patients. Considering the negative synergistic effects on range of health outcomes resulted by comorbid depression and DM, prevention and early treatment of comorbid depression are a critical component of successful health management in older adult patients with DM.

References

1. Murray CJ, Lopez AD. Measuring the global burden of disease. *New England Journal of Medicine*. 2013; 369(5):448–457. [PubMed: 23902484]

2. Park, M.; Unützer, J. Public Health Burden of Late-Life Mood Disorders. In: Levretsky, H.; Sajatovic, M.; Reynolds, C., editors. *Late-life Mood Disorders*. Oxford University Press; NY: 2013. p. 42-60.
3. Anderson RJ, Freedland KE, Clouse RE, Lustman PJ. The prevalence of comorbid depression in adults with diabetes: a meta-analysis. *Diabetes Care*. Jun; 2001 24(6):1069–1078. [PubMed: 11375373]
4. Palinkas L, Barrett-Connor E, Wingard D. Type 2 Diabetes and Depressive Symptoms in Older Adults: a Population-based Study. *Diabetic medicine*. 1991; 8(6):532–539. [PubMed: 1832352]
5. Kramer MK, Kriska AM, Venditti EM, et al. Translating the Diabetes Prevention Program: a comprehensive model for prevention training and program delivery. *American journal of preventive medicine*. Dec; 2009 37(6):505–511. [PubMed: 19944916]
6. Centers for Disease Control Prevention. National diabetes fact sheet: national estimates and general information on diabetes and prediabetes in the United States, 2011. US Department of Health and Human Services, Centers for Disease Control and Prevention; Atlanta, GA: 2011. p. 201
7. Park M, Katon WJ, Wolf FM. Depression and risk of mortality in individuals with diabetes: a meta-analysis and systematic review. *General hospital psychiatry*. 2013; 35(3):217–225. [PubMed: 23415577]
8. Cuijpers P, Beekman AT, Reynolds CF 3rd. Preventing depression: a global priority. *JAMA: the journal of the American Medical Association*. Mar 14; 2012 307(10):1033–1034.
9. Ludman EJ, Katon W, Russo J, et al. Depression and diabetes symptom burden. *General hospital psychiatry*. 2004; 26(6):430–436. [PubMed: 15567208]
10. Gonzalez JS, Peyrot M, McCarl LA, et al. Depression and diabetes treatment nonadherence: a meta-analysis. *Diabetes Care*. Dec; 2008 31(12):2398–2403. [PubMed: 19033420]
11. Egede LE, Zheng D, Simpson K. Comorbid Depression is Associated With Increased Health Care Use and Expenditures in Individuals With Diabetes. *Diabetes Care*. Mar; 2002 25(3):464–470. 2002. [PubMed: 11874931]
12. de Groot M, Anderson R, Freedland KE, Clouse RE, Lustman PJ. Association of depression and diabetes complications: a meta-analysis. *Psychosom Med*. Jul-Aug;2001 63(4):619–630. [PubMed: 11485116]
13. Katon W, Lyles CR, Parker MM, Karter AJ, Huang ES, Whitmer RA. Association of depression with increased risk of dementia in patients with type 2 diabetes: the Diabetes and Aging Study. *Arch Gen Psychiatry*. Apr; 2012 69(4):410–417. [PubMed: 22147809]
14. Blazer DG. Psychiatry and the Oldest Old. *Am J Psychiatry*. Dec 1; 2000 157(12):1915–1924. 2000. [PubMed: 11097951]
15. Evans DL, Charney DS, Lewis L, et al. Mood disorders in the medically ill: scientific review and recommendations. *Biol Psychiatry*. Aug 1; 2005 58(3):175–189. [PubMed: 16084838]
16. Unutzer J. Diagnosis and treatment of older adults with depression in primary care. *Biol Psychiatry*. Aug 1; 2002 52(3):285–292. [PubMed: 12182933]
17. Lyness JM, Kim J, Tang W, et al. The Clinical Significance of Subsyndromal Depression in Older Primary Care Patients. *American Journal of Geriatric Psych*. 2007; 15(3):214–223. 210.1097/1001.JGP.0000235763.0000250230.0000235783.
18. Meeks TW, Vahia IV, Lavretsky H, Kulkarni G, Jeste DV. A tune in “a minor” can “b major”: A review of epidemiology, illness course, and public health implications of subthreshold depression in older adults. *Journal of Affective Disorders*. 2011; 129(1-3):126–142. [PubMed: 20926139]
19. Judd LL, Akiskal HS. The Clinical and Public Health Relevance of Current Research on Subthreshold Depressive Symptoms to Elderly Patients. *American Journal of Geriatric Psych*. 2002; 10(3):233–238.
20. Lavretsky H, Kumar A. Clinically Significant Non-Major Depression: Old Concepts, New Insights. *American Journal of Geriatric Psych*. 2002; 10(3):239–255.
21. Lyness J, King D, Cox C, Yoediono Z, Caine E. The importance of subsyndromal depression in older primary care patients: prevalence and associated functional disability. *J Am Geriatr Soc*. 1999; 47(6):647–652. [PubMed: 10366161]
22. Remick RA. Diagnosis and management of depression in primary care: a clinical update and review. *CMAJ*. Nov 26; 2002 167(11):1253–1260. 2002. [PubMed: 12451082]

23. Lyness JM, Yu Q, Tang W, Tu X, Conwell Y. Risks for Depression Onset in Primary Care Elderly Patients: Potential Targets for Preventive Interventions. *Am J Psychiatry*. Dec 1; 2009 166(12): 1375–1383. 2009. [PubMed: 19833788]
24. Grabovich A, Lu N, Tang W, Tu X, Lyness JM. Outcomes of subsyndromal depression in older primary care patients. *Am J Geriatr Psychiatry*. Mar; 2010 18(3):227–235. [PubMed: 20173424]
25. Unutzer J. Clinical practice. Late-life depression. *N Engl J Med*. Nov 29; 2007 357(22):2269–2276. [PubMed: 18046030]
26. Klein DN, Shankman SA, Rose S. Ten-year prospective follow-up study of the naturalistic course of dysthymic disorder and double depression. *Am J Psychiatry*. May; 2006 163(5):872–880. [PubMed: 16648329]
27. Mueller TI, Leon AC, Keller MB, et al. Recurrence after recovery from major depressive disorder during 15 years of observational follow-up. *Am J Psychiatry*. 1999; 156(7):1000–1006. [PubMed: 10401442]
28. Licht-Strunk E, van der Windt DLA, van Marwijk HW, de Haan M, Beekman AT. The prognosis of depression in older patients in general practice and the community. A systematic review. *Family Practice*. 2007; 24(2):168–180. [PubMed: 17237495]
29. Cole MG, Bellavance F, Mansour A. Prognosis of depression in elderly community and primary care populations: a systematic review and meta-analysis. *Am J Psychiatry*. Aug; 1999 156(8): 1182–1189. [PubMed: 10450258]
30. Iosifescu DV, Nierenberg AA, Alpert JE, et al. Comorbid Medical Illness and Relapse of Major Depressive Disorder in the Continuation Phase of Treatment. *Psychosomatics*. 2004; 45(5):419–425. [PubMed: 15345787]
31. Fisher L, Skaff MM, Mullan JT, et al. Clinical depression versus distress among patients with type 2 diabetes not just a question of semantics. *Diabetes Care*. 2007; 30(3):542–548. [PubMed: 17327318]
32. Drayer RA, Mulsant BH, Lenze EJ, et al. Somatic symptoms of depression in elderly patients with medical comorbidities. *International Journal of Geriatric Psychiatry*. 2005; 20(10):973–982. [PubMed: 16163749]
33. Kirmayer LJ, Young A. Culture and somatization: clinical, epidemiological, and ethnographic perspectives. *Psychosom Med*. Jul 1; 1998 60(4):420–430. 1998. [PubMed: 9710287]
34. Sadovnick AD, Remick RA, Lam R, et al. Mood disorder service genetic database: Morbidity risks for mood disorders in 3,942 first-degree relatives of 671 index cases with single depression, recurrent depression, bipolar I, or bipolar II. *American Journal of Medical Genetics*. 1994; 54(2): 132–140. [PubMed: 8074163]
35. Stroebe M, Schut H, Stroebe W. Health outcomes of bereavement. *The Lancet*. Dec 14; 2007 370(9603):1960–1973. 2007.
36. Pinquart M, Sorensen S. Influences of socioeconomic status, social network, and competence on subjective well-being in later life: a meta-analysis. *Psychol Aging*. Jun; 2000 15(2):187–224. [PubMed: 10879576]
37. Park M, Unutzer J, Grembowski D. Ethnic and Gender Variations in The Associations between Family Cohesion, Family Conflict, and Depression in Older Asian and Latino Adults. *Journal of Immigrant and Minority Health*. 2013:1–8. [e-pub ahead of time]. [PubMed: 22307545]
38. Park M, Unutzer J. Hundred forty eight more days with depression: the association between marital conflict and depression-free days. *Int J Geriatr Psychiatry*. Apr 4.2014 n/a-n/a.
39. Thompson A, Fan MY, Unutzer J, Katon W. One extra month of depression: the effects of caregiving on depression outcomes in the IMPACT trial. *Int J Geriatr Psychiatry*. May; 2008 23(5):511–516. [PubMed: 17944005]
40. Abe-Kim J, Takeuchi DT, Hong S, et al. Use of mental health-related services among immigrant and US-born Asian Americans: results from the National Latino and Asian American Study. *Am J Public Health*. Jan; 2007 97(1):91–98. [PubMed: 17138905]
41. Abe-Kim JS, Takeuchi DT. Cultural Competence and Quality of Care: Issues for Mental Health Service Delivery in Managed Care. *Clinical Psychology: Science and Practice*. 1996; 3(4):273–295.

42. Aber JL, Bennett NG, Conley DC, Li J. The effects of poverty on child health and development. *Annu Rev Public Health*. 1997; 18:463–483. [PubMed: 9143727]
43. Abueg, FR.; Chun, KM. Traumatized Asian and Asian American populations. In: Marsella, A.; Friedman, M.; Gerrity, E.; Scurfield, R., editors. *Ethnocultural approaches to understanding posttraumatic stress disorder: Issues, research, and clinical applications*. American Psychological Association; Washington D.C.: 1996.
44. Abueg, FR.; Chun, KM. Traumatization stress among Asians and Asian Americans. In: Marsella, AJEF.; Matthew, J., et al., editors. *Ethnocultural aspects of posttraumatic stress disorder: Issues, research, and clinical applications*. American Psychological Association; Washington, DC: 1996. p. 285-299.
45. Ackerman DL, Unutzer J, Greenland S, Gitlin M. Inpatient treatment of depression and associated hospital charges. *Pharmacoepidemiol Drug Saf*. Apr-May;2002 11(3):219–227. [PubMed: 12051121]
46. Adler N, Boyce W, Chesney M, Folkman S, Syme S. Socioeconomic inequalities in health: no easy solution. *JAMA: the journal of the American Medical Association*. 1993; 269:3140.
47. Adler NE, Newman K. Socioeconomic disparities in health: pathways and policies. *Health Aff (Millwood)*. Mar-Apr;2002 21(2):60–76. [PubMed: 11900187]
48. Adler NE, Newman K. Socioeconomic Disparities In Health: Pathways And Policies. *Health Aff*. Mar 1; 2002 21(2):60–76. 2002.
49. Park, M. *The Variations in Associations between Family Contexts and Late-Life Depression Outcomes [Thesis]*: Health Services. University of Washington; 2013.
50. Martire LM, Schulz R. Involving Family in Psychosocial Interventions for Chronic Illness. *current directions in psychological science*. 2007; 16(2):90–94.
51. Lee MS, Crittenden KS, Yu E. Social support and depression among elderly Korean immigrants in the United States. *International Journal of Aging and Human Development*. 1996; 42(4):313–327. [PubMed: 8835613]
52. George LK, Blazer DG, Hughes DC, Fowler N. Social support and the outcome of major depression. *Br J Psychiatry*. Apr; 1989 154(4):478–485. [PubMed: 2590779]
53. Smith F, Francis S-A, Gray N, Denham M, Graffy J. A multi-centre survey among informal carers who manage medication for older care recipients: problems experienced and development of services. *Health & Social Care in the Community*. 2003; 11(2):138–145. [PubMed: 14629216]
54. Voils CI, Steffens DC, Flint EP, Bosworth HB. Social Support and Locus of Control as Predictors of Adherence to Antidepressant Medication in an Elderly Population. *American Journal of Geriatric Psych*. 2005; 13(2):157–165.
55. Agency for Healthcare Research and Quality. 2004 National Healthcare Disparities Report. Rockville, MD2004:
56. Agency for Healthcare Research and Quality. National Healthcare Disparities Report. Rockville, MD2007: 2007.
57. Kawachi I, Berkman L. Social ties and mental health. *Journal of Urban Health*. 2001; 78(3):458–467. [PubMed: 11564849]
58. Hong S-I, Hasche L, Bowland S. Structural Relationships Between Social Activities and Longitudinal Trajectories of Depression Among Older Adults. *The Gerontologist*. Feb 1; 2009 49(1):1–11. 2009. [PubMed: 19362999]
59. Unutzer J, Katon W, Sullivan M, Miranda J. Treating depressed older adults in primary care: narrowing the gap between efficacy and effectiveness. *Milbank Q*. 1999; 77(2):225–256. 174. [PubMed: 10392163]
60. Centers for Disease Control and Prevention and National Association of Chronic Disease Directors. *The State of Mental Health and Aging in America Issue Brief 2: Addressing Depression in Older Adults: Selected Evidence-Based Programs*. National Association of Chronic Disease Directors. , editor. Atlanta, GA: 2009.
61. Kessler RC, Berglund P, Demler O, Jin R, Merikangas KR, Walters EE. Lifetime Prevalence and Age-of-Onset Distributions of DSM-IV Disorders in the National Comorbidity Survey Replication. *Arch Gen Psychiatry*. Jun 1; 2005 62(6):593–602. 2005. [PubMed: 15939837]

62. Mojtabai R, Olfson M. Major depression in community-dwelling middle-aged and older adults: prevalence and 2- and 4-year follow-up symptoms. *Psychological Medicine*. 2004; 34(04):623–634. [PubMed: 15099417]
63. Byers AL, Yaffe K, Covinsky KE, Friedman MB, Bruce ML. High occurrence of mood and anxiety disorders among older adults: The National Comorbidity Survey Replication. *Arch Gen Psychiatry*. May; 2010 67(5):489–496. [PubMed: 20439830]
64. Blazer DG. Depression in Late Life: Review and Commentary. *Focus*. Jan 1; 2009 7(1):118–136. 2009.
65. Teresi J, Abrams R, Holmes D, Ramirez M, Eimicke J. Prevalence of depression and depression recognition in nursing homes. *Social Psychiatry and Psychiatric Epidemiology*. 2001; 36(12):613–620. [PubMed: 11838834]
66. Hoover DR, Siegel M, Lucas J, et al. Depression in the first year of stay for elderly long-term nursing home residents in the U.S.A. *International Psychogeriatrics*. 2010:1–11. First View.
67. Lyness JM, Caine ED, King DA, Cox C, Yoediono Z. Psychiatric Disorders in Older Primary Care Patients. *Journal of General Internal Medicine*. 1999; 14(4):249–254. [PubMed: 10203638]
68. Kuo BCH, Chong V, Joseph J. Depression and Its Psychosocial Correlates Among Older Asian Immigrants in North America. *Journal of Aging and Health*. 2008; 20(6):615–652. [PubMed: 18768704]
69. Simpson S, Krishnan L, Kunik M, Ruiz P. Racial Disparities in Diagnosis and Treatment of Depression: A Literature Review. *Psychiatric Quarterly*. 2007; 78(1):3–14. [PubMed: 17102936]
70. Black SA, Markides KS, Ray LA. Depression predicts increased incidence of adverse health outcomes in older Mexican Americans with type 2 diabetes. *Diabetes Care*. Oct; 2003 26(10):2822–2828. [PubMed: 14514586]
71. Steffens DC, Fisher GG, Langa KM, Potter GG, Plassman BL. Prevalence of depression among older Americans: the Aging, Demographics and Memory Study. *Int Psychogeriatr*. Oct; 2009 21(5):879–888. [PubMed: 19519984]
72. Strothers HS, Rust G, Minor P, Fresh E, Druss B, Satcher D. Disparities in Antidepressant Treatment in Medicaid Elderly Diagnosed with Depression. *Journal of the American Geriatrics Society*. 2005; 53(3):456–461. [PubMed: 15743289]
73. Crystal S, Sambamoorthi U, Walkup JT, Akincigil A. Diagnosis and treatment of depression in the elderly medicare population: predictors, disparities, and trends. *Journal of the American Geriatrics Society*. Dec; 2003 51(12):1718–1728. [PubMed: 14687349]
74. Kleinman, A. *Patients and healers in the context of culture: an exploration of the borderland between anthropology, medicine, and psychiatry*. University of California Press; Berkeley and Los Angeles, California: 1981.
75. Kleinman A. Culture and Depression. *N Engl J Med*. Sep 2; 2004 351(10):951–953. 2004. [PubMed: 15342799]
76. Pang KYC. Symptom expression and somatization among elderly Korean immigrants. *Journal of Clinical Geropsychology*. 2000; 6(3):199–212.
77. Jackson JL, Kroenke K. Difficult Patient Encounters in the Ambulatory Clinic: Clinical Predictors and Outcomes. *Arch Intern Med*. May 24; 1999 159(10):1069–1075. 1999. [PubMed: 10335683]
78. Cooper LA, Gonzales JJ, Gallo JJ, et al. The Acceptability of Treatment for Depression among African-American, Hispanic, and White Primary Care Patients. *Medical care*. 2003; 41(4):479–489. [PubMed: 12665712]
79. Givens JL, Houston TK, Van Voorhees BW, Ford DE, Cooper LA. Ethnicity and preferences for depression treatment. *General hospital psychiatry*. May-Jun;2007 29(3):182–191. [PubMed: 17484934]
80. Alegria M, Chatterji P, Wells K, et al. Disparity in depression treatment among racial and ethnic minority populations in the United States. *Psychiatr Serv*. Nov; 2008 59(11):1264–1272. [PubMed: 18971402]
81. Kessler RC, Birnbaum HG, Shahly V, et al. Age differences in the prevalence and co-morbidity of DSM-IV major depressive episodes: results from the WHO World Mental Health Survey Initiative. *Depression and Anxiety*. 2010; 27(4):351–364. [PubMed: 20037917]

82. Chen L, Magliano DJ, Zimmet PZ. The worldwide epidemiology of type 2 diabetes mellitus--present and future perspectives. *Nat Rev Endocrinol.* Apr; 2012 8(4):228–236. [PubMed: 22064493]
83. Wild S, Roglic G, Green A, Sicree R, King H. Global prevalence of diabetes: estimates for the year 2000 and projections for 2030. *Diabetes Care.* May; 2004 27(5):1047–1053. [PubMed: 15111519]
84. Ali S, Stone MA, Peters JL, Davies MJ, Khunti K. The prevalence of co-morbid depression in adults with Type 2 diabetes: a systematic review and meta-analysis. *Diabet Med.* Nov; 2006 23(11):1165–1173. [PubMed: 17054590]
85. Li C, Ford E, Strine T, Mokdad A. Prevalence of depression among U.S. adults with diabetes: findings from the 2006 Behavioral Risk Factor Surveillance System. *Diabetes Care.* 2008; 31:105. [PubMed: 17934145]
86. Hofmann M, Köhler B, Leichsenring F, Kruse J. Depression as a risk factor for mortality in individuals with diabetes: a meta-analysis of prospective studies. *PloS one.* 2013; 8(11):e79809. [PubMed: 24278183]
87. van Dooren FE, Nefs G, Schram MT, Verhey FR, Denollet J, Pouwer F. Depression and risk of mortality in people with diabetes mellitus: a systematic review and meta-analysis. *PloS one.* 2013; 8(3):e57058. [PubMed: 23472075]
88. Bogner HR, Morales KH, Post EP, Bruce ML. Diabetes, depression, and death: a randomized controlled trial of a depression treatment program for older adults based in primary care (PROSPECT). *Diabetes Care.* 2007; 30(12):3005–3010. [PubMed: 17717284]
89. Gallo JJ, Morales KH, Bogner HR, et al. Long term effect of depression care management on mortality in older adults: follow-up of cluster randomized clinical trial in primary care. *BMJ.* 2013; 346:f2570. [PubMed: 23738992]
90. Katon WJ. Clinical and health services relationships between major depression, depressive symptoms, and general medical illness. *Biological Psychiatry.* 2003; 54(3):216–226. [PubMed: 12893098]
91. Golden S, Lazo M, Carnethon M, Bertoni A, Schreiner P. Examining a bidirectional association between depressive symptoms and diabetes. *JAMA: the journal of the American Medical Association.* 2008; 299:2751.
92. Stuart MJ, Baune BT. Depression and type 2 diabetes: Inflammatory mechanisms of a psychoneuroendocrine co-morbidity. *Neuroscience & Biobehavioral Reviews.* 2012; 36(1): 658–676. 1//. [PubMed: 22020230]
93. Silva N, Atlantis E, Ismail K. A review of the association between depression and insulin resistance: pitfalls of secondary analyses or a promising new approach to prevention of type 2 diabetes? *Curr Psychiatry Rep.* Feb; 2012 14(1):8–14. [PubMed: 22094982]
94. Kuo H-K. Cognitive dysfunction and depression in older adults with diabetes. *Geriatric Diabetes.* 2007:63–84.
95. Ciechanowski PS, Katon WJ, Russo JE, Hirsch IB. The relationship of depressive symptoms to symptom reporting, self-care and glucose control in diabetes. *General hospital psychiatry.* Jul-Aug;2003 25(4):246–252. [PubMed: 12850656]
96. Ciechanowski PS, Katon WJ, Russo JE. Depression and diabetes: impact of depressive symptoms on adherence, function, and costs. *Arch Intern Med.* Nov 27; 2000 160(21):3278–3285. [PubMed: 11088090]
97. McDermott LM, Ebmeier KP. A meta-analysis of depression severity and cognitive function. *J Affect Disord.* Dec; 2009 119(1-3):1–8. [PubMed: 19428120]
98. Byers AL, Yaffe K. Depression and risk of developing dementia. *Nat Rev Neurol.* Jun; 2011 7(6): 323–331. [PubMed: 21537355]
99. Rapp MA, Schnaider-Beeri M, Wysocki M, et al. Cognitive decline in patients with dementia as a function of depression. *The American Journal of Geriatric Psychiatry.* 2011; 19(4):357–363. [PubMed: 20808140]
100. Stetler C, Miller GE. Depression and hypothalamic-pituitary-adrenal activation: a quantitative summary of four decades of research. *Psychosom Med.* Feb-Mar;2011 73(2):114–126. [PubMed: 21257974]

101. Lok A, Mocking RJ, Ruhe HG, et al. Longitudinal hypothalamic-pituitary-adrenal axis trait and state effects in recurrent depression. *Psychoneuroendocrinology*. Jul; 2012 37(7):892–902. [PubMed: 22094110]
102. Vanhanen M, Koivisto K, Moilanen L, et al. Association of metabolic syndrome with Alzheimer disease A population-based study. *Neurology*. 2006; 67(5):843–847. [PubMed: 16966548]
103. Craft S. The role of metabolic disorders in Alzheimer disease and vascular dementia: two roads converged. *Arch Neurol*. Mar; 2009 66(3):300–305. [PubMed: 19273747]
104. Notarianni E. Hypercortisolemia and glucocorticoid receptor-signaling insufficiency in Alzheimer's disease initiation and development. *Curr Alzheimer Res*. Sep; 2013 10(7):714–731. [PubMed: 23906001]
105. Cole J, Costafreda SG, McGuffin P, Fu CH. Hippocampal atrophy in first episode depression: a meta-analysis of magnetic resonance imaging studies. *J Affect Disord*. Nov; 2011 134(1-3):483–487. [PubMed: 21745692]
106. Taylor WD, McQuoid DR, Payne ME, Zannas AS, Macfall JR, Steffens DC. Hippocampus Atrophy and the Longitudinal Course of Late-life Depression. *Am J Geriatr Psychiatry*. Nov 22.2013
107. Mezuk B, Eaton WW, Albrecht S, Golden SH. Depression and type 2 diabetes over the lifespan: a meta-analysis. *Diabetes Care*. Dec; 2008 31(12):2383–2390. [PubMed: 19033418]
108. Demakakos P, Pierce MB, Hardy R. Depressive Symptoms and Risk of Type 2 Diabetes in a National Sample of Middle-Aged and Older Adults. *Diabetes Care*. Apr 1; 2010 33(4):792–797. 2010. [PubMed: 20086253]
109. Carnethon MR, Kinder LS, Fair JM, Stafford RS, Fortmann SP. Symptoms of depression as a risk factor for incident diabetes: findings from the National Health and Nutrition Examination Epidemiologic Follow-up Study, 1971-1992. *American journal of epidemiology*. 2003; 158(5): 416–423. [PubMed: 12936896]
110. Eaton W, Pratt L, Armenian H, Ford D, Gallo J. Depression and risk for onset of type II diabetes: a prospective populationbased study. *Diabetes Care*. 1996; 19:1097–1102. [PubMed: 8886555]
111. Nouwen A, Winkley K, Twisk J, et al. Type 2 diabetes mellitus as a risk factor for the onset of depression: a systematic review and meta-analysis. *Diabetologia*. 2010; 53(12):2480–2486. [PubMed: 20711716]
112. Reynolds CF, Thomas SB, Morse JQ, et al. Early Intervention to Preempt Major Depression Among Older Black and White Adults. *Psychiatric Services*. 2014
113. U.S. Preventive Services Task Force. [Accessed October 30, 2011] Screening for Depression: Recommendations and Rationale. 2002. <http://www.uspreventiveservicestaskforce.org/3rduspstf/depression/depressrr.htm>
114. Cuijpers P, van Straten A, Hollon SD, Andersson G. The contribution of active medication to combined treatments of psychotherapy and pharmacotherapy for adult depression: a meta-analysis. *Acta Psychiatr Scand*. Jun; 2010 121(6):415–423. [PubMed: 19922522]
115. Kohler S, Hoffmann S, Unger T, Steinacher B, Dierstein N, Fydrich T. Effectiveness of cognitive-behavioural therapy plus pharmacotherapy in inpatient treatment of depressive disorders. *Clin Psychol Psychother*. Mar-Apr;2013 20(2):97–106. [PubMed: 22095701]
116. Rocha FL, Fuzikawa C, Riera R, Hara C. Combination of antidepressants in the treatment of major depressive disorder: a systematic review and meta-analysis. *J Clin Psychopharmacol*. Apr; 2012 32(2):278–281. [PubMed: 22367652]
117. Reynolds CF, Frank E, Perel JM, et al. Nortriptyline and Interpersonal Psychotherapy as Maintenance Therapies for Recurrent Major Depression. *JAMA: The Journal of the American Medical Association*. 1999; 281(1):39–45.
118. van der Feltz-Cornelis CM, Nuyen J, Stoop C, et al. Effect of interventions for major depressive disorder and significant depressive symptoms in patients with diabetes mellitus: a systematic review and meta-analysis. *General hospital psychiatry*. 2010; 32(4):380–395. [PubMed: 20633742]
119. Bruce ML, Ten Have TR, Reynolds CF III, et al. Reducing Suicidal Ideation and Depressive Symptoms in Depressed Older Primary Care Patients: A Randomized Controlled Trial. *JAMA: the journal of the American Medical Association*. 2004; 291(9):1081–1091.

120. Oxman TE, Sengupta A. Treatment of Minor Depression. *American Journal of Geriatric Psych*. 2002; 10(3):256–264.
121. Pinquart M, Duberstein PR, Lyness JM. Treatments for Later-Life Depressive Conditions: A Meta-Analytic Comparison of Pharmacotherapy and Psychotherapy. *Am J Psychiatry*. Sep 1; 2006 163(9):1493–1501. 2006. [PubMed: 16946172]
122. Gilbody S, Bower P, Fletcher J, Richards D, Sutton AJ. Collaborative care for depression: A cumulative meta-analysis and review of longer-term outcomes. *Arch Intern Med*. 2006; 166:2314–2321. [PubMed: 17130383]
123. Simon G. Collaborative care for depression. *BMJ*. Feb 4; 2006 332(7536):249–250. [PubMed: 16455698]
124. Simon G. Collaborative care for mood disorders. *Curr Opin Psychiatry*. Jan; 2009 22(1):37–41. [PubMed: 19122533]
125. Bogner HR, Morales KH, de Vries HF, Cappola AR. Integrated management of type 2 diabetes mellitus and depression treatment to improve medication adherence: a randomized controlled trial. *The Annals of Family Medicine*. 2012; 10(1):15–22.
126. Katon W, Von Korff M, Lin E, et al. The pathways study: A randomized trial of collaborative care in patients with diabetes and depression. *Archives of General Psychiatry*. 2004; 61(10):1042–1049. [PubMed: 15466678]
127. Katon W, Lin E, Von Korff M, et al. Integrating depression and chronic disease care among patients with diabetes and/or coronary heart disease: The design of the TEAMcare study. *Contemporary Clinical Trials*. 2010; 31(4):312–322. [PubMed: 20350619]
128. Katon W, Unutzer J. Collaborative Care Models for Depression: Time to Move From Evidence to Practice. *Arch Intern Med*. Nov 27; 2006 166(21):2304–2306. 2006. [PubMed: 17130381]
129. Katon W, Unutzer J, Wells K, Jones L. Collaborative depression care: history, evolution and ways to enhance dissemination and sustainability. *Gen Hosp Psychiatry*. Sep-Oct; 2010 32(5): 456–464. [PubMed: 20851265]
130. Katon W, Von Korff M, Lin E, Simon G. Rethinking practitioner roles in chronic illness: the specialist, primary care physician, and the practice nurse. *General Hospital Psychiatry*. May-Jun; 2001 23(3):138–144. [PubMed: 11427246]
131. Kroenke K, Spitzer RL, Williams JBW, Lin E, et al. The Patient Health Questionnaire Somatic, Anxiety, and Depressive Symptom Scales: a systematic review. *General hospital psychiatry*. 2010; 32(4):345–359. [PubMed: 20633738]
132. American Psychiatric Association. *Diagnostic and Statistical Manual of Mental Disorders: DSM-IV-TR*. 4th ed.. Washington, DC 2000:
133. Kroenke K, Spitzer RL, Williams JBW. The Patient Health Questionnaire-2: Validity of a Two-Item Depression Screener. *Medical care*. 2003; 41(11):1284–1292. [PubMed: 14583691]
134. Kok RM, Heeren TJ, Nolen WA. Continuing Treatment of Depression in the Elderly: A Systematic Review and Meta-Analysis of Double-Blinded Randomized Controlled Trials With Antidepressants. *American Journal of Geriatric Psych*. 2011; 19(3):249–255. 210.1097/JGP.1090b1013e3181ec8085.
135. Nelson JC, Delucchi K, Schneider L. Efficacy of Second Generation Antidepressants in Late-Life Depression: A Meta-Analysis of the Evidence. *American Journal of Geriatric Psychiatry*. 2008; 16(7):558–567. [PubMed: 18591576]
136. Gartlehner G, Gaynes BN, Hansen RA, et al. Comparative benefits and harms of second-generation antidepressants: background paper for the American College of Physicians. *Ann Intern Med*. Nov 18; 2008 149(10):734–750. [PubMed: 19017592]
137. Gum AM, Arean PA, Hunkeler E, et al. Depression treatment preferences in older primary care patients. *Gerontologist*. Feb; 2006 46(1):14–22. [PubMed: 16452280]
138. Cuijpers P, van Straten A, Smit F. Psychological treatment of late-life depression: a meta-analysis of randomized controlled trials. *Int J Geriatr Psychiatry*. Dec; 2006 21(12):1139–1149. [PubMed: 16955421]
139. Pinquart M, Duberstein PR, Lyness JM. Effects of psychotherapy and other behavioral interventions on clinically depressed older adults: A meta-analysis. *Aging and Mental Health*. 2007; 11(6):645–657. [PubMed: 18074252]

140. Cuijpers P, Karyotaki E, Pot AM, Park M, Reynolds CF III. Managing depression in older age: psychological interventions. *Maturitas*. 2014 [Epub ahead of time].
141. Wilson KC, Mottram PG, Vassilas CA. Psychotherapeutic treatments for older depressed people. *Cochrane Database Syst Rev*. 2008; 1(1):CD004853. [PubMed: 18254062]
142. de Mello MF, de Jesus Mari J, Bacaltchuk J, Verdelli H, Neugebauer R. A systematic review of research findings on the efficacy of interpersonal therapy for depressive disorders. *Eur Arch Psychiatry Clin Neurosci*. Apr; 2005 255(2):75–82. [PubMed: 15812600]
143. Bell AC, D’Zurilla TJ. Problem-solving therapy for depression: a meta-analysis. *Clin Psychol Rev*. Jun; 2009 29(4):348–353. [PubMed: 19299058]
144. Cuijpers P, van Straten A, Warmerdam L. Behavioral activation treatments of depression: a meta-analysis. *Clin Psychol Rev*. Apr; 2007 27(3):318–326. [PubMed: 17184887]
145. Mazzucchelli T, Kane R, Rees C. Behavioral Activation Treatments for Depression in Adults: A Meta-analysis and Review. *Clinical Psychology: Science and Practice*. 2009; 16(4):383–411.
146. Ekers D, Richards D, Gilbody S. A meta-analysis of randomized trials of behavioural treatment of depression. *Psychol Med*. May; 2008 38(5):611–623. [PubMed: 17903337]
147. Arean P, Hegel M, Vannoy S, Fan MY, Unutzer J. Effectiveness of problem-solving therapy for older, primary care patients with depression: results from the IMPACT project. *Gerontologist*. Jun; 2008 48(3):311–323. [PubMed: 18591356]
148. Schulberg HC, Post EP, Raue PJ, Have TT, Miller M, Bruce ML. Treating late-life depression with interpersonal psychotherapy in the primary care sector. *International Journal of Geriatric Psychiatry*. 2007; 22(2):106–114. [PubMed: 17096458]
149. Olfson M, Marcus S, Sackeim HA, Thompson J, Pincus HA. Use of ECT for the Inpatient Treatment of Recurrent Major Depression. *Am J Psychiatry*. 1998; 155:22–29. [PubMed: 9433334]
150. Case BG, Bertollo DN, Laska EM, Siegel CE, Wanderling JA, Olfson M. Racial differences in the availability and use of electroconvulsive therapy for recurrent major depression. *Journal of Affective Disorders*. 2012; 136(3):359–365. [PubMed: 22169249]
151. Association, AP. Diagnostic and statistical manual of mental disorders. 5th ed.. American Psychiatric Association; Washington, DC: 2013.
152. Cole MG, Dendukuri N. Risk factors for depression among elderly community subjects: a systematic review and meta-analysis. *Am.J.Psychiatry*. 2003; 160(6):1147–1156. [PubMed: 12777274]
153. Snowdon J. Is depression more prevalent in old age? *Australian and New Zealand Journal of Psychiatry*. 2001; 35(6):782–787. [PubMed: 11990889]
154. Sonnenberg CM, Beekman AT, Deeg DJ, van Tilburg W. Sex differences in late-life depression. *Acta Psychiatr Scand*. Apr; 2000 101(4):286–292. [PubMed: 10782548]
155. Zisook S, Shuchter SR. Depression through the first year after the death of a spouse. *Am J Psychiatry*. Oct; 1991 148(10):1346–1352. [PubMed: 1897615]
156. Turvey CL, Carney C, Arndt S, Wallace RB, Herzog R. Conjugal loss and syndromal depression in a sample of elders aged 70 years or older. *Am J Psychiatry*. Oct; 1999 156(10):1596–1601. [PubMed: 10518172]
157. Prince MJ, Harwood RH, Thomas A, Mann AH, The Gospel Oak Project VII. A prospective population-based cohort study of the effects of disablement and social milieu on the onset and maintenance of late-life depression. *Psychol Med*. Mar; 1998 28(2):337–350. [PubMed: 9572091]
158. Cacioppo JT, Hughes ME, Waite LJ, Hawkley LC, Thisted RA. Loneliness as a specific risk factor for depressive symptoms: cross-sectional and longitudinal analyses. *Psychology and aging*. 2006; 21(1):140. [PubMed: 16594799]
159. Gould RL, Coulson MC, Howard RJ. Cognitive behavioral therapy for depression in older people: a meta-analysis and meta-regression of randomized controlled trials. *Journal of the American Geriatrics Society*. Oct; 2012 60(10):1817–1830. [PubMed: 23003115]
160. Cuijpers P, van Straten A, Warmerdam L. Behavioral activation treatments of depression: A meta-analysis. *Clinical Psychology Review*. 2007; 27(3):318–326. [PubMed: 17184887]

161. Ekers D, Webster L, Van Straten A, Cuijpers P, Richards D, Gilbody S. Behavioural activation for depression: an update of meta-analysis of effectiveness and sub group analysis. *PLoS One*. 2014; 9(6):e100100. [PubMed: 24936656]
162. Samad Z, Brealey S, Gilbody S. The effectiveness of behavioural therapy for the treatment of depression in older adults: a meta-analysis. *Int J Geriatr Psychiatry*. Dec; 2011 26(12):1211–1220. [PubMed: 21308789]
163. Bell AC, D’Zurilla TJ. Problem-solving therapy for depression: A meta-analysis. *Clinical Psychology Review*. 2009; 29(4):348–353. [PubMed: 19299058]
164. Beekman A, van Schaik A, van Marwijk H, et al. Interpersonal Psychotherapy for Elderly Patients in Primary Care. *American Journal of Geriatric Psych*. 2006; 14(9):777–786. 710.1097/1001.JGP.0000199341.0000125431.0000199344b.
165. Bohlmeijer E, Smit F, Cuijpers P. Effects of reminiscence and life review on late-life depression: a meta-analysis. *International Journal of Geriatric Psychiatry*. 2003; 18(12):1088–1094. [PubMed: 14677140]
166. Pinquart M, Forstmeier S. Effects of reminiscence interventions on psychosocial outcomes: a meta-analysis. *Aging mental health*. 2012; 16(5):541–558.

Key Points

1. Depression is highly prevalent in the general population and increases the risk for type 2 DM 2.
2. Comorbid depression and DM is associated negative health outcomes, such as accelerated cognitive decline and increased mortality.
3. Depression impinges the patient and the family caregiver's ability to effectively manage DM, decreases adherence to treatment, and undermines successful physician-patient relationship
4. Effective models for treating comorbid depression and DM exists and some components of these models are implementable in individual clinics.




Figure 1. A conceptual model of interaction between major depression and medical illness
 Reproduced from Katon, Wayne J. "Clinical and Health Services Relationships between Major Depression, Depressive Symptoms, and General Medical Illness." *Biological Psychiatry* 54, no. 3 (2003): 216-26.




Figure 2.
Algorithm for Screening and Treating Depression in Primary Care Setting

Table 1

Risk Factors for Depression in Older Adults

Authors, year	Sample	Finding
Age		
Cole et al (2003) ¹⁵²	Age 50	Meta-analysis of epidemiological studies. Pooled OR for depression associated with age: 1.2 (95% CI: 0.9-1.7)
Snowden (2008) ¹⁵³	Mixed age	Comprehensive review of epidemiological studies: Depression is as common in older age as in earlier life.
Being Female		
Cole et al (2003) ¹⁵²	Age 50	Meta-analysis of epidemiological studies. Pooled OR for depression associated with being female: 1.4 (95% CI: 1.2-1.8)
Sonnenberg et al (2001) ¹⁵⁴	55 Age 85	A random, age and sex-stratified community sample of 3056 older Dutch people; Prevalence of depression in women was almost twice as high as in men.
Grief and Loss		
Zisook et al (1991) ¹⁵⁵	Mixed age	Major depression and anxiety disorders are common within the first year of the spouse's death: 29-58% meet criteria for major depression at one month, 24-30% at two months, and 25% at three months.
Cole et al (2003) ¹⁵²	Age 50	Meta-analysis of epidemiological studies. Pooled OR for depression associated with recent bereavement: 3.3 (95% CI: 1.7-4.9)
Turkey et al (1999) ¹⁵⁶	Age 70	The rate of syndromal depression in the newly bereaved was nearly nine times as high as the rate for married individuals, and the rate of depressive symptoms was nearly four times as high.
Social Isolation		
Cole et al (2003) ¹⁵²	Age 50	Meta-analysis of epidemiological studies. Pooled OR for depression associated with living alone: 1.7 (95% CI: 0.6-4.7)
Prince et al (1998) ¹⁵⁷	Age 65	Prospective epidemiological study. Lack of contact with friends was a direct risk factor but also modified the association between handicap and depression.
Cacioppo et al (2006) ¹⁵⁸	Age 55	There is a reciprocal relationship between loneliness depressive symptomatology and over time
Cognitive impairment		
Cole et al (2003) ¹⁵²	Age 50	Meta-analysis of epidemiological studies. Pooled OR for depression associated with recent bereavement: 2.1 (95% CI: 0.6-8.6)

Table 2

Commonly used antidepressant for late-life depression

Medication	Starting dose	Common Therapeutic Doses	Half-life (hour)	Common side effects	Comments
SSRI (Selective serotonin-reuptake inhibitors):					
Fluoxetine	5 mg	10 – 40 mg once daily	70-80	Nausea, dyspepsia, anorexia, tremors, anxiety, insomnia, sexual dysfunction, jitteriness, hyponatremia	Risk of serotonin syndrome if combined with certain drugs.
Sertraline	12.5 mg	50 – 200 mg once daily	25-30	Loose stools, diarrhea	Very long acting.
Citalopram	10 mg	20 – 40 mg once daily	40-50		
Escitalopram	2.5 mg	10– 30 mg once daily	40-50		
Paroxetine	10 mg	20 – 50 mg once daily	10-20	Dry mouth, drowsiness, fatigue, weight gain,	More anticholinergic side effects. High risk of discontinuation syndrome if drug stopped abruptly.
SNRIs (Serotonin-norepinephrine reuptake inhibitors)					
Venlafaxine XR	37.5 mg	75-225g once daily	5 – 9	Nausea, drowsiness, fatigue, weight gain, hyponatremia, diastolic hypertension at higher doses	Risk of serotonin syndrome if combined with certain drugs. High risk of discontinuation syndrome if medication stopped abruptly.
Duloxetine	20 mg	20-60mg once daily	8 – 17		
Other newer antidepressants					
Mirtazapine	15mg	15-45mg at bedtime	20 -40	Sedation, increased appetite / weight gain	No sexual side effects.
Bupropion SR	100mg	100-150 mg twice daily	15		No sexual side effects. Contraindicated in patients with seizures. Not recommended for patients with comorbid anxiety.
High risk in overdose: 10 days of typical daily dose may result in a fatal cardiac arrhythmia. Get baseline ECG and follow-up ECG at steady state or if new cardiac symptoms occur.					
TCA's (Tricyclic antidepressants)					
Nortriptyline	10mg	50-125mg every night	18-56	Fatigue	Therapeutic blood levels range from 50 to 150 ng/mL
Desipramine	25mg	100-200mg once daily	12-28	Insomnia, agitation	

Table 3
Efficacious non-pharmacologic treatments for late-life depression

Treatment type	Studies	Level of evidence	Comments
Cognitive Behavioral Therapy (CBT)	Gould et al 2012 ¹⁵⁹	Meta-analysis of RCTs	CBT significantly more efficacious at reducing depressive symptoms Combination with pharmacotherapy more efficacious than CBT alone
	Cuijpers et al 2014 ¹⁴⁰	Meta-analysis of RCTs	Older adults populations studies
	Cuijpers et al 2007 ¹⁶⁰	Meta-analysis of RCTs	Mixed population studies Effect is weaker in older adults
Behavior activation (BA)	Ekers et al, 2014 ¹⁶¹	Meta-analysis of RCTs	Mixed population studies
	Samad et al ¹⁶²	Meta-analysis of RCTs	Older adults populations studies
Problem solving therapy (PST)	Bell et al 2009 ¹⁶³	Meta-analysis of RCTs	Mixed population studies
Interpersonal psychotherapy (IPT)	Beekman et al 2006 ¹⁶⁴	Pragmatic RCT	Older adults populations studies
	Bohlmeijer et al 2003 ¹⁶⁵	Meta-analysis of RCTs	Older adults populations studies
Reminiscence (life review)	Pinquart & Forstmeier ¹⁶⁶	Meta-analysis of RCTs	Efficacious for broad range of outcomes, and therapeutic as well as preventive effects are similar to those observed in other frequently used interventions.