

# Lawrence Berkeley National Laboratory

## LBL Publications

### Title

CymA and Exogenous Flavins Improve Extracellular Electron Transfer and Couple It to Cell Growth in Mtr-Expressing *Escherichia coli*

### Permalink

<https://escholarship.org/uc/item/2qm6f21z>

### Journal

ACS Synthetic Biology, 5(7)

### ISSN

2161-5063

### Authors

Jensen, Heather M  
TerAvest, Michaela A  
Kokish, Mark G  
et al.

### Publication Date

2016-07-15

### DOI

10.1021/acssynbio.5b00279

Peer reviewed

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23

**CymA and exogenous flavins improve extracellular electron transfer  
and couple it to cell growth in Mtr-expressing *Escherichia coli***

Heather M. Jensen<sup>1,3,#a</sup> , Michaela A. TerAvest<sup>5,#b</sup> , Mark G. Kokish<sup>2,3,#c</sup> , Caroline M. Ajo-Franklin<sup>1,2,4\*</sup>

<sup>1</sup>Physical Biosciences Division, Lawrence Berkeley National Laboratory, Berkeley, California, United States of America.

<sup>2</sup>Materials Sciences Division, Lawrence Berkeley National Laboratory, Berkeley, California, United States of America.

<sup>3</sup>Department of Chemistry, University of California, Berkeley, California, United States of America.

<sup>4</sup>Synthetic Biology Institute, Berkeley, California, United States of America.

<sup>5</sup>California Institute for Quantitative Biosciences, University of California, Berkeley, California, United States of America

<sup>#a</sup>Current Institution: Joint BioEnergy Institute, Emeryville, California, United States of America.

<sup>#b</sup>Current Institution: Michigan State University, East Lansing, Michigan, United States of America.

<sup>#c</sup>Current Institution: Northwestern University, Evanston, Illinois, United States of America.

\*Corresponding author:

24 E-mail: cajo-franklin@lbl.gov (CMAF)

25

26 **Abstract**

27 Introducing extracellular electron transfer pathways into heterologous organisms offers the  
28 opportunity to explore fundamental biogeochemical processes and to biologically alter redox  
29 states of exogenous metals for various applications. While expression of the MtrCAB electron  
30 nanoconduit from *Shewanella oneidensis* MR-1 permits extracellular electron transfer in  
31 *Escherichia coli*, the low electron flux and absence of growth in these cells limits their  
32 practicality for such applications. Here we investigate how the rate of electron transfer to  
33 extracellular Fe(III) and cell survival in engineered *E. coli* are affected by mimicking different  
34 features of the *S. oneidensis* pathway: the number of electron nanoconduits, the link between the  
35 quinol pool and MtrA, and the presence of flavin-dependent electron transfer. While increasing  
36 the number of pathways does not significantly improve the extracellular electron transfer rate or  
37 cell survival, using the native inner membrane component, CymA, significantly improves the  
38 reduction rate of extracellular acceptors and increases cell viability. Strikingly, introducing both  
39 CymA and riboflavin to Mtr-expressing *E. coli* also allowed these cells to couple metal reduction  
40 to growth, which is the first time an increase in biomass of an engineered *E. coli* has been  
41 observed under  $\text{Fe}_2\text{O}_3$  (*s*) reducing conditions. Overall, this work provides engineered *E. coli*  
42 strains for modulating extracellular metal reduction and elucidates critical factors for engineering  
43 extracellular electron transfer in heterologous organisms.

44

45

46 **Keywords:** synthetic biology, dissimilatory metal-reducing bacteria, bioelectrochemical systems,  
47 energy conservation, multi-heme cytochrome *c*

## 48 **Introduction**

49           Some metal-reducing bacteria, such as those from the *Geobacter* and *Shewanella* genera,  
50 have extracellular electron transfer pathways that can route electrons across the cell membrane to  
51 alter the redox state of exogenous metals. This extracellular metal reduction by microorganisms  
52 plays a key role in microbial-driven mineral transformations (1) and can be used to drive  
53 nanoparticle synthesis under mild conditions (2, 3) or alter the phase of metals for  
54 bioremediation (4) or biomining (3). The ability to modulate extracellular electron transfer then  
55 offers the opportunity to dissect and control these processes. For example, systematically varying  
56 the rate of extracellular electron transfer should allow synthetic control over the number and size  
57 of nanoparticles and should elucidate the role of redox kinetics on mineral transformation.  
58 However, the sparse availability of genetic tools in metal-reducing bacteria and the multiple  
59 overlapping pathways for extracellular electron transfer make it challenging to modulate electron  
60 transfer and/or introduce other functions of interest. In response to this challenge, we (5-7) and  
61 others (8-10) have taken the complementary approach of engineering portions of extracellular  
62 electron transfer pathways into the well-studied industrial microbe *Escherichia coli* (11). While  
63 this work has shown that the Mtr pathway can route electrons to extracellular metal oxides (6),  
64 the low electron flux and the inability of cells to maintain biomass using solid minerals as the  
65 terminal electron acceptor has hampered the use of these strains for applications such as  
66 nanoparticle synthesis and bioremediation (7).

67           Since the extracellular electron transfer pathways of *Shewanella oneidensis* MR-1 have  
68 been extensively studied at the molecular level (12, 13), they have provided the basis for these  
69 engineering efforts (**Figure 1A**). When *S. oneidensis* is grown under metal-respiring conditions,  
70 reducing equivalents from oxidation of electron donors are directed to the menaquinol pool and

71 then to the inner membrane tetraheme cytochrome *c* CymA (14). When extracellular metals or  
72 electrodes are present, electrons from CymA are passed to the periplasmic decaheme cytochrome  
73 *c* MtrA. The re-oxidation of CymA completes the Q-cycle, increasing the proton motive force  
74 (15) and allowing the cell to conserve energy. Notably, it is debated whether CymA directly  
75 reduces MtrA (9, 16) or whether FccA, STC (also known as CctA), or another redox protein,  
76 serve as an intermediate to pass electrons from CymA to MtrA (17-19). MtrA reduces MtrC, a  
77 decaheme cytochrome *c* located on the extracellular face of the outer membrane, as part of the  
78 MtrCAB complex. This complex is proposed to form a porin cytochrome complex (12) that  
79 spans the outer membrane and allows MtrA to directly contact MtrC (20). There are  
80 approximately 70,000 MtrC proteins per *S. oneidensis* MR-1 cell (21), and these proteins can  
81 either directly reduce a mineral or electrode surface or use a flavin-dependent process to  
82 indirectly reduce the solid electron acceptor. While the mechanism underpinning flavin-mediated  
83 electron transfer is still being elucidated (22-24), it is clear that ~80% of the electron transfer  
84 occurs through a flavin-dependent process and ~20% of the electron transfer occurs directly (25-  
85 27). Electron transfer through MtrCAB does not affect the proton motive force; however, it  
86 indirectly contributes to energy conservation by re-oxidizing CymA.

87 *E. coli* strains expressing combinations of MtrA and CymA can reduce soluble Fe(III),  
88 but not extracellular Fe(III) oxides (8, 9). In contrast, expression of the MtrCAB electron  
89 nanoconduit in *E. coli* confers the ability to reduce Fe<sub>2</sub>O<sub>3</sub> (s) (6). However, this strain reduces  
90 Fe(III) several orders of magnitude more slowly than *S. oneidensis* MR-1. Likely as a result of  
91 this slow reduction rate, this *E. coli* strain is unable to conserve energy and its biomass sharply  
92 decreases, which poses a particular challenge for further applications. Introduction of  
93 extracellular electron transfer pathways into another industrial microbe, *Pseudomonas putida*,

94 has not translated into cell growth (28-30), indicating this may be a common challenge. Our  
95 previous work identified re-reduction of MtrA as a rate-limiting step in Fe(III) reduction (6).  
96 This strain also lacks CymA, extracellular flavins, and has fewer Mtr electron conduits (30,000  
97 (5) vs. 70,000) (**Figure 1A**). Here we systematically test how co-expression of CymA, changing  
98 the number of electron conduits per cell, and introduction of flavins affect the flux of electrons to  
99 extracellular iron and *E. coli* survival. We found that co-expression of CymA with the Mtr  
100 nanoconduit and introduction of exogenous flavins significantly increased the electron transfer  
101 rate over the original strains. We also found that *E. coli* strains expressing CymA coupled these  
102 improvements in extracellular reduction to enhanced cellular viability. Importantly, the  
103 combination of CymA and riboflavin allowed Mtr *E. coli* cells to couple Fe(III) reduction to  
104 growth under Fe<sub>2</sub>O<sub>3</sub>(s)-reducing conditions.

105

## 106 **Results and Discussion**

### 107 **The *cymA*, *cymAmtrA*, and *cymAmtr* strains express the full-length and redox active** 108 **cytochromes *c*.**

109 To systematically explore the effect of CymA on soluble iron and solid iron oxide  
110 reduction, we created three separate plasmids for expression of *cymA*, *cymAmtrA*, and  
111 *cymAmtrCAB* (**Figure S1**); in these plasmids, the RBS sequences were designed to be  
112 approximately the same as the corresponding native RBS strength in *S. oneidensis* (31). The  
113 *cymA*, *cymAmtrA*, and *cymAmtrCAB* plasmids were co-transformed with the cytochrome *c*  
114 maturation (*ccm*) plasmid (32) into *E. coli* to create the *cymA*, *cymAmtrA*, and *cymAmtr* strains,  
115 respectively (**Table 1**). The *mtrA* strain, co-expressing *ccm* and *mtrA*, and the *mtr* strain, which  
116 expresses *ccm* and *mtrCAB*, (**Table 1**), were used as previously described (6).

117 We next characterized the expression and localization of each cyt *c* to validate each  
118 strain. Proteins from whole cell extracts were resolved by SDS-PAGE, and heme *c* containing  
119 proteins were identified by enhanced chemiluminescence (ECL) analysis. These whole cell  
120 extracts show that the MtrA and MtrC proteins in all the strains containing *mtrA* and *mtrC*, i.e.  
121 the *cymAmtrA*, *mtrA*, *mtr*, and *cymAmtr* strains, are full-length (35 kD and 76 kD, respectively;  
122 **Figure 1B**). The *cymA* strain also clearly shows a band corresponding to full length CymA (18  
123 kD, **Figure 1B**). While difficult to discern by eye, densitometry analysis of the ECL  
124 demonstrates the CymA band is also present in the two other strains containing *cymA*, the  
125 *cymAmtrA* and *cymAmtr* strains (**Figure 1C**). ECL analysis of fractionated cells shows CymA,  
126 MtrA, and MtrC in *E. coli* are localized (**Figure S1**) as they are in *S. oneidensis* (33-35). Lastly,  
127 diffuse reflectance spectroscopy of all these strains under oxidizing and reducing conditions  
128 showed characteristic changes in the  $\alpha$ - and  $\beta$ - bands, indicating that the cyt *c* proteins are redox  
129 active. Thus, each strain expresses its respective redox active, full length and correctly localized  
130 cyt *c*.

131 To characterize the concentration range of cyt *c* that could be expressed, we increased the  
132 promoter activity (as defined in (5)) in *mtr* and *cymAmtr* strains via IPTG induction (**SI Table 1**)  
133 and measured the total heme *c* content in these cells using diffuse reflectance as described  
134 previously (5). In brief, diffuse reflectance measures absorbance of visible light with limited  
135 scattering, thus allowing quantitative measurement of the  $\alpha$ -bands in whole cell suspensions  
136 (**Figure S2**). From these spectra, we can determine the concentration of heme *c*, and thus infer  
137 relative abundance of electron conduits in whole cell suspensions. From the 0 to 0.37 promoter  
138 activity, the cyt *c* concentration in the *mtr* and *cymAmtr* strains both increase ~5-fold (**Figure**  
139 **1D**). The maximum heme *c* measured in *mtr* and *cymAmtr* strains are 140 and 106  $\mu$ M heme *c*



140  $\text{OD}_{600\text{nm}}^{-1}$ , respectively (**Figure 1D**), which is about half the total heme *c* content of *S. oneidensis*  
141 when grown under the same conditions ( $228 \mu\text{M heme } c \text{ OD}_{600\text{nm}}^{-1}$ ). ECL analysis shows that the  
142 relative abundance of each cyt *c* within a strain is very similar across different induction levels  
143 (**Figure S3**), indicating that induction boosts the abundance of each cyt *c* rather than a single cyt  
144 *c*. Thus, tuning the promoter strength via induction allows us to vary the number of electron  
145 transfer proteins in the *mtr* and *cymAmtr* strains.

#### 146 **Fe(III) citrate reduction and cell survival are improved by co-expression of CymA.**

147 Previous work by Schuetz *et al.* (9) has shown that co-expression of CymA with MtrA  
148 from genomically-integrated constructs can improve Fe(III)-NTA reduction in *E. coli* over  
149 expression of CymA alone, strongly suggesting that CymA would improve  $\text{Fe}_2\text{O}_3$  reduction in  
150 the *cymAmtr* strain. However, co-expression of CymA alongside other *S. oneidensis* MR-1 cyts *c*  
151 dramatically reduces its expression level in our strains (**Figure 1B-D**), and we have previously  
152 observed that subtle changes in expression levels can profoundly impact extracellular electron  
153 transfer capabilities (5). Thus, we first tested cell suspensions of the *cymA*, *mtrA*, and *cymAmtrA*  
154 strains for the ability to reduce soluble Fe(III) citrate in anaerobic conditions with D,L-lactate as  
155 a carbon source to determine the effect of limited expression of CymA. Since lysed *E. coli* cells  
156 can reduce  $\text{Au}^+$  (36), we also included a homogenized sample of the *mtrA* strain to determine if  
157 cell lysis contributed to Fe(III) reduction (**Figure S4**). The lysed *mtrA* sample reduced at a  
158 similar rate as the *ccm* strain (**Figure S4**), demonstrating cell lysis is not a major contributor to  
159 Fe(III) citrate reduction in this system. While the *mtrA* strain reduced Fe(III) slightly faster than  
160 the *ccm* control, the *cymA* and *cymAmtrA* strains increased the Fe(III) reduction rate over the  
161 *mtrA* strain by a substantial ~3-fold and ~4-fold, respectively (**Figure 2A**).

162 In these iron reduction assays, the *E. coli* strains have only D,L-lactate as a carbon  
163 source, and the only available terminal electron acceptor for respiration is Fe(III) citrate.  
164 Therefore, we hypothesized that aspects which increase Fe(III) reduction would also improve  
165 energy conservation and cell survival. Interestingly, while the cell density of the slowest Fe(III)-  
166 reducing strains, the *ccm* and *mtrA* strains, sharply decreased, the cell density of the fastest  
167 Fe(III)-reducing strains, the *cymA* and *cymAmtrA* strain, stayed the same or slightly increased  
168 (**Figure 2B**). Indeed, a direct comparison between the rate of Fe(III) citrate reduction and the  
169 linear rate of change of the cell density in CymA-expressing strains (**Figure S5**) reveals that the  
170 iron reduction rate and strain fitness are strongly, positively correlated (Pearson correlation  
171 coefficient,  $R = 0.93$ ). This strong positive correlation strongly suggests that increasing the iron  
172 reduction rate in these *E. coli* strains also improves energy conservation under our experimental  
173 conditions. The correlation also suggests that a minimum iron reduction rate ( $\sim 0.4 \text{ mM day}^{-1}$ ,  
174 **Figure S5**) is necessary to maintain or increase cell biomass. Taken together, these observations  
175 show that even low levels of CymA co-expressed with MtrA are sufficient to improve Fe(III)  
176 citrate reduction and strongly suggest that increasing the Fe(III) reduction rate helps maintain  
177 biomass in *E. coli* by increasing energy conservation.

#### 178 **Fe<sub>2</sub>O<sub>3</sub> (s) reduction and cell survival are improved by co-expression of CymA.**

179 We next set out to probe the effects of CymA co-expression on solid Fe<sub>2</sub>O<sub>3</sub> reduction and  
180 cell viability in the *ccm*, *cymAmtr*, and *mtr* strains. In the solid iron oxide reduction assays, the *E.*  
181 *coli* strains have only D,L-lactate as a carbon source, and the only available terminal electron  
182 acceptor for respiration is solid Fe<sub>2</sub>O<sub>3</sub>. To survey a wide set of conditions, we varied the number  
183 of electron conduits with a range of promoter activity (**Figure 1D**, **Table S1**), and we  
184 additionally examined the effect of exogenous riboflavin. The promoter activity is indicated in

185 superscript throughout. Due to the relatively slow reduction of solid Fe<sub>2</sub>O<sub>3</sub> by these engineered  
186 *E. coli* in comparison to *S. oneidensis*, many strains required 21 day incubations before  
187 statistically significant measurements of Fe(II) by ferrozine could be attained. In contrast,  
188 changes in colony forming units (cfu) were apparent over much shorter time scales than changes  
189 in Fe(II) concentration.

190 We first summarize the effects of CymA co-expression with the Mtr electron nanoconduit  
191 in *E. coli*. The *ccm* only strain, which accounts for basal Fe<sub>2</sub>O<sub>3</sub> (*s*) reduction, reduced only 62 ±  
192 19 μM Fe(II) over 21 days (**Figure 3A**). The best reducing *mtr* strain, *mtr*<sup>0.37</sup>, showed a  
193 statistically significant ~2-fold improvement over the *ccm* strain (**Figure 3A**). Interestingly, of  
194 all the strains and induction conditions tested, the uninduced *cymAmtr* strain, *cymAmtr*<sup>0.001</sup>,  
195 reduced the greatest amount of Fe<sub>2</sub>O<sub>3</sub>, ~3.5 fold more Fe<sub>2</sub>O<sub>3</sub> (*s*) than the *ccm* strain. The  
196 *cymAmtr*<sup>0.001</sup> strain reduced statistically more Fe<sub>2</sub>O<sub>3</sub> (*s*) than the *mtr*<sup>0.37</sup> strain, demonstrating that  
197 co-expression of CymA improves the reduction of solid Fe<sub>2</sub>O<sub>3</sub> (**Figure 3A**) as well as soluble  
198 Fe(III) (**Figure 2A**).

199 CymA also significantly enhanced the survival of *cymAmtr* in the iron reduction assay  
200 conditions. The *ccm* strain decreased to 7% of its initial cfu mL<sup>-1</sup> over 4 days (**Figure 3B**),  
201 consistent with the inability of the native *E. coli* strain to conserve energy when Fe<sub>2</sub>O<sub>3</sub> (*s*) is the  
202 sole terminal electron acceptor available. The *mtr*<sup>0.37</sup> strain maintained 27% of the initial cfu mL<sup>-1</sup>  
203 <sup>1</sup>, indicating that *mtr* improved survival somewhat over native *E. coli*. The most robust strain was  
204 the *cymAmtr*<sup>0.001</sup> strain, which increased its cfu mL<sup>-1</sup> 5% above its initial value (**Figure 3B**).  
205 More broadly, comparing the amount of Fe(II) produced to the relative cell density after 21 days  
206 (**Figure S5**), shows there is a strong, positive correlation between the degree of reduction and  
207 cell survival in CymA-containing strains (Pearson correlation coefficient, R = 0.74). Taken

208 together, these observations indicate that these CymA-expressing *E. coli* strains use metal  
209 reduction to maintain biomass under both Fe(III) citrate and Fe<sub>2</sub>O<sub>3</sub>(s) reducing conditions.

210 **The expression of CymA significantly increases extracellular electron transfer by enabling**  
211 **catalytic turnover of Mtr.**

212 Since CymA can reduce MtrA *in vitro*, we hypothesized that CymA co-expression  
213 increased Fe<sub>2</sub>O<sub>3</sub> (s) reduction and cell viability by more rapidly reducing the MtrCAB complex  
214 than native *E. coli* inner membrane quinone dehydrogenases. To directly investigate this, we  
215 turned to bioelectrochemical techniques that can monitor the redox turnover of MtrCAB. Since  
216 these techniques use a carbon felt electrode as the terminal electron acceptor instead of Fe<sub>2</sub>O<sub>3</sub> (s),  
217 we first monitored current production from the *ccm*, *mtr*<sup>0.18</sup>, and *cymAmtr*<sup>0.18</sup> strains in  
218 bioelectrochemical reactors. The working electrode chamber contained cell suspensions in M1  
219 media supplemented with 40 mM D,L-lactate, and the counter electrode chamber contained 50  
220 mM PIPES buffer (pH 7.4), and was separated by a cation exchange membrane. As expected, the  
221 *ccm* strain, which does not express any *S. oneidensis* cyts *c*, produced very low current (**Figure**  
222 **4A**). The *mtr*<sup>0.18</sup> strain did not produce significantly more current than the *ccm* strain, while in  
223 contrast, the *cymAmtr*<sup>0.18</sup> strain produced an average of ~4-fold more current than the *mtr*<sup>0.18</sup>  
224 strain (**Figure 4A**). This increase between these two strains is comparable to the increase in  
225 Fe<sub>2</sub>O<sub>3</sub> (s) reduction, suggesting that the cause of improved extracellular electron transfer routes  
226 in these strains is independent of the identity of the terminal electron acceptor. Next, we  
227 monitored reduction of the Mtr pathway by intracellular reducing equivalents using turnover  
228 cyclic voltammetry, scanning at 2 mV/s. Both the *ccm* and *mtr*<sup>0.18</sup> strains exhibited an  
229 irreversible oxidation peak of unknown origin, but did not show a catalytic wave, i.e. an s-shaped  
230 peak, indicative of re-reduction. This indicates that the catalytic turnover of the Mtr pathway is

231 extremely slow in the absence of CymA. In contrast, the turnover cyclic voltammogram of the  
232 *cymAmtr*<sup>0.18</sup> strain (**Figure 4B**) showed a clear catalytic wave, centered just below 0.0 V<sub>Ag/AgCl</sub>,  
233 which is consistent with previous measurements of the formal potential of the Mtr pathway in *S.*  
234 *oneidensis* MR-1 (37). The presence of this wave indicates that the Mtr pathway can be re-  
235 reduced in the presence of CymA under turnover conditions. This rapid turnover additionally  
236 shows that there are properly assembled electron conduits in the engineered *E. coli*, however we  
237 have not yet determined what fraction of the electron conduits are correctly assembled.

238         Taken together, these data support our initial observation that re-reduction of MtrA is a  
239 rate-limiting step for Fe(III) reduction in *mtrA E. coli* (6) and show that CymA increases the rate  
240 of Fe(III) reduction in both MtrA- and MtrCAB-expressing *E. coli* by increasing the rate of re-  
241 reduction of MtrA. The ~15 nm thick periplasm in *E. coli* (38) is too thick to allow CymA to  
242 directly contact MtrA within an MtrCAB complex, however, we observe a significant amount of  
243 MtrA in the periplasmic fraction (**Figure S6**). If MtrA diffuses in the periplasm with a diffusion  
244 constant similar to comparably sized *E. coli* proteins, i.e. ~2 μm<sup>2</sup>s<sup>-1</sup> (39), it would only require 3  
245 μs on average to diffuse from the inner membrane to the outer membrane. This timescale is 3  
246 orders of magnitude faster than the ms required for electron transfer across the MtrCAB conduit  
247 (20). Coupled with the observation that CymA can reduce MtrA *in vitro* (40), the simplest  
248 explanation of these data is that, in *E. coli*, CymA directly reduces MtrA and MtrA diffuses  
249 across the periplasmic space to shuttle electrons to MtrC.

250 **The number of electron nanoconduits does not significantly affect solid Fe<sub>2</sub>O<sub>3</sub> reduction**  
251 **and cell survival.**

252         Using the data we gathered on Fe(III) reduction for *cymAmtr* and *mtr* strains at different  
253 induction levels, we next probed the relationship between the number of electron conduits in the

254 *mtr* and *cymAmtr* strains and iron reduction. We use [heme *c*] as a measure of the number of  
255 electron conduits because the stoichiometry of cyts *c* is unchanged with different induction levels  
256 (**Figure S3**) in our *E. coli* strains. Pearson correlation analysis shows a strong correlation  
257 between [heme *c*] and [Fe(II)] in both strains, with the *mtr* strain showing a positive correlation  
258 and the *cymAmtr* showing a negative correlation (Pearson correlation coefficients  $R_{mtr} = 0.95$  and  
259  $R_{cymAmtr} = -0.84$ , **Figure S7**). However, the slope of these correlations are fairly small ( $m_{mtr} =$   
260  $0.31 [\text{Fe(II)}] [\text{heme } c]^{-1}$  and  $m_{cymAmtr} = -0.52 [\text{Fe(II)}] [\text{heme } c]^{-1}$ ): for example, the Fe(III)  
261 reduction rate in the *mtr* strain changes only ~1.5-fold when the number of *mtr* electron conduits  
262 changes ~5-fold. This indicates that, for either strain, the number of conduits is not a strong  
263 determinant of the reduction of solid iron oxide.

264 The finding that the number of electron conduits in engineered *E. coli* is not a strong  
265 determinant of  $\text{Fe}_2\text{O}_3$  (*s*) reduction suggests that electron transfer through the Mtr complex is not  
266 a rate-limiting step in iron reduction in these strains. To further test this idea, we estimated what  
267 the rate of iron reduction in *E. coli* would be if transport through the MtrCAB conduit were rate-  
268 limiting. Using our diffuse reflectance data of our best performing *cymAmtr*<sup>0.001</sup> strain (**Figure**  
269 **1D**) and assuming that 90% of the heme *c* is in MtrCAB complexes (**Figure 1B**), we estimate the  
270 number of Mtr electron nanoconduits per *E. coli* cell is 50,000 complexes cell<sup>-1</sup>. This is  
271 comparable to the number of MtrC in *S. oneidensis*, which is ~70,000 cell<sup>-1</sup> (21). Together with  
272 estimates on the maximal rate of electron transfer through the Mtr electron nanoconduit ( $10^3 \text{ e}^- \text{ s}^-$   
273  $^1 \text{ complex}^{-1}$ ) (20), we can approximate the maximal rate of iron reduction per *cymAmtr*<sup>0.001</sup> *E.*  
274 *coli* cell would be  $\sim 8 \times 10^{-17} \text{ M Fe(II) s}^{-1} \text{ cell}^{-1}$  (**Supplemental Calculation 1**). This theoretical  
275 maximum rate is ~2000-fold higher than what we observe ( $\sim 4 \times 10^{-20} \text{ M Fe(II) s}^{-1} \text{ cell}^{-1}$ ). Thus,

276 we conclude that electron transfer through the Mtr conduit is not rate-limiting for extracellular  
277 electron transfer in these strains of engineered *E. coli*.

278 **Flavins couple solid Fe<sub>2</sub>O<sub>3</sub> reduction to transient growth in the *cymAmtr* strain.**

279 Lastly, we considered how addition of 5 μM riboflavin affected iron reduction and cell  
280 survival in the engineered *E. coli* strains. We found that riboflavin increased the total amount of  
281 reduced iron by 2 and 2.5 fold in the *mtr*<sup>0.18</sup> and *cymAmtr*<sup>0.18</sup> strains over those strains without  
282 riboflavin added, respectively (**Figure 5A, Figure S8**). To quantitatively assess the fraction of  
283 electrons that reduce *via* direct contact or flavin-dependent mechanisms, we performed  
284 experiments in M1 medium, a trace medium with no riboflavin or casein added (See  
285 Supplemental Information), with and without supplementation of riboflavin. We assumed that  
286 cultures without riboflavin reduce iron solely by direct contact, while cultures supplemented with  
287 riboflavin reduce iron by both direct contact and through a riboflavin-mediated mechanism. This  
288 allowed us to calculate the percentage of flavin-dependent iron reduction by comparing the iron  
289 reduction in the supplemented and non-supplemented cultures. The resulting percentage of iron  
290 reduced in a flavin-dependent process in the *mtr*<sup>0.18</sup> and *cymAmtr*<sup>0.18</sup> strains was 72% and 77%,  
291 respectively (**Supplemental Calculation 2**). These percentages closely match the ratio observed  
292 in *S. oneidensis* MR-1 (25, 26). This indicates that, like in the native organism, MtrC is able to  
293 utilize solid metal oxides as terminal electron acceptors *via* both direct contact and through  
294 flavin-mediated mechanisms.

295 Interestingly, while riboflavin had similar relative effects on iron reduction in both the  
296 *mtr* and *cymAmtr* strains, it had strikingly different effects on cell growth in these strains. The  
297 addition of riboflavin did not statistically change the survival of the *mtr*<sup>0.18</sup> strain (**Figure S8**).  
298 This observation indicates that riboflavin by itself does not support growth under metal reducing

299 conditions. In the case of the *cymAmtr*<sup>0.18</sup> strain, supplementation with riboflavin allowed the  
300 cells to grow, increasing the cfu mL<sup>-1</sup> 10-fold over the original suspension after 4 days (**Figure**  
301 **5B**). This cfu increase is temporally associated with an initial period of rapid reduction of Fe(III)  
302 in the *cymAmtr*<sup>0.18</sup> strain but not the *ccm* strain (**Figure 5C, Figure S9**). Following this period,  
303 the cfu decrease and the increase in Fe(II) concentration slows (days 4-10). After 10 days, the  
304 cell density drops below its initial value and the Fe(II) concentration remains constant, even  
305 though a large excess of Fe(III) is still present. Given the correlations between iron reduction and  
306 cell viability, we suggest that the initial high rate of Fe(III) reduction conferred by riboflavin and  
307 CymA is enough to allow the *E. coli* to conserve energy and grow. However, as the Fe(III)  
308 reduction rate in this strain slows, perhaps due to a depletion of energy stores and slow oxidation  
309 of lactate, the *E. coli* can no longer conserve energy and the cell density decreases  
310 correspondingly. Eventually, the cell density is so low that there is no detectable change in Fe(II)  
311 concentration. This scenario would be consistent with observations in *E. coli*, *Shewanella sp.*,  
312 and other microbes that the respiratory rate is positively correlated with the growth rate (37, 41).

313 The *cymAmtr*<sup>0.18</sup> *E. coli* strain generates 200 μM Fe(II) over the course of 10 days, which  
314 corresponds to a cell normalized reduction rate of  $\sim 8 \times 10^{-20}$  M Fe(II) s<sup>-1</sup> cell<sup>-1</sup>. Also, this strain  
315 can maintain cell density at or above its initial level over this same period, which marks the first  
316 time an engineered *E. coli* strain has been shown to grow under solid metal-reducing conditions.  
317 These are significant improvements over previously described *E. coli* strains capable of solid  
318 metal reduction. Additionally, this work strongly suggests that further improvements in  
319 extracellular metal reduction will also boost cell growth in these strains.

320 **Opportunities for additional engineering of metal-reducing *E. coli*.**



321 The fastest iron reduction rate in the engineered *E. coli* cultures described here is still ~50  
322 times slower than *S. oneidensis* MR-1 ( $\sim 8 \times 10^{-20}$  M Fe(II) s<sup>-1</sup> cell<sup>-1</sup> vs.  $\sim 4 \times 10^{-18}$  M Fe(II) s<sup>-1</sup> cell<sup>-1</sup>)  
323 (42). We attribute the strain's inability to sustain growth for long periods of time to this slower  
324 rate and suggest further work is still required in strain engineering for demanding applications.  
325 While our cyclic voltammetry data indicates that a significant fraction of the MtrCAB electron  
326 nanoconduits are correctly assembled in the outer membrane of *E. coli*, we cannot rule out that  
327 some fraction of MtrCAB complexes are misassembled and non-functional in *E. coli*. Assessing  
328 and potentially improving this assembly may improve extracellular metal reduction.  
329 Additionally, the difference between our *cymAmtr E. coli* strains and *S. oneidensis* MR-1 may  
330 reflect our still-evolving understanding of electron transfer in *S. oneidensis*. While this work was  
331 in preparation, new data generated by Sturm et al. (19) and Alves et al. (17) strongly suggested  
332 that Stc and/or FccA may be involved in shuttling electrons across the periplasm. Thus, co-  
333 expression of Stc and/or FccA may help extracellular electron transfer in *Mtr E. coli*.

334 The difference between the electron transfer rates in engineered *E. coli* and *S. oneidensis*  
335 MR-1 may also arise from the fact that that efficient extracellular electron transfer relies on  
336 many processes besides transport of electrons from inner membrane to extracellular acceptors  
337 (43). Specifically, extracellular electron transfer also depends on the import and oxidation of an  
338 electron donor, transfer of the electrons *via* intracellular redox carriers to the transmembrane  
339 pathway, and adhesion of the bacterium to a metal oxide. Several of these processes are slow or  
340 disrupted in the *E. coli* background used here. For example, the reaction rates of the L- and D-  
341 lactate dehydrogenases in *E. coli* are ~10 and ~20-fold slower, respectively, than their  
342 counterparts in *S. oneidensis* (44). Additionally, since the *Mtr E. coli* are grown aerobically  
343 before being introduced into anaerobic metal-reducing conditions, these cells may lack the

344 dehydrogenases and quinones needed to efficiently direct reducing equivalents to the Mtr  
345 pathway. Lastly, while *Shewanella sp.* can rapidly attach to Fe<sub>2</sub>O<sub>3</sub> surfaces (45, 46), the  
346 BL21(DE3) derivatives used in this work are disrupted in their ability to initiate attachment to  
347 surfaces because they are non-motile (47). Future work will focus on testing these multiple  
348 possibilities to increase the Fe(III) reduction rate and cell growth rates in Mtr-expressing *E. coli*.

## 349 **Conclusions**

350 This work shows that replicating certain features of the electron transfer pathway of *S.*  
351 *oneidensis* MR-1 are critical for boosting extracellular electron transfer in *E. coli*, while others  
352 are not. Specifically, we found that increasing the number of Mtr complexes did not increase  
353 extracellular electron transfer, but the presence of CymA and riboflavin had a significant impact  
354 on extracellular electron transfer and viability. Additionally, we show that as the rate of metal  
355 reduction in CymA-expressing *E. coli* increases, these strains gradually transition from rapidly  
356 losing biomass to transiently growing under metal-reducing conditions. This knowledge will  
357 allow metal-reduction to be more readily introduced into new heterologous hosts.

358 Despite the relatively modest iron reduction in these newest *E. coli* strains relative to  
359 *Shewanella sp.* or *Geobacter sp.*, the ability of the *cymAmtr E. coli* strain described here to  
360 maintain biomass or, in the presence of riboflavin, grow over limited timescales makes a useful  
361 new tool for both basic and applied studies. Since *E. coli* uses many different electron donors  
362 compared to *Geobacter* or *Shewanella*, it may provide a more versatile tool for bioremediation or  
363 biomining. Also, it is well known that both cell surface structures and metal reduction play a role  
364 in formation and transformation of metal-containing solids such as metallic nanoparticles and  
365 metal-oxide minerals. The ability of our *E. coli* strains to reduce solid and chelated Fe(III) with a  
366 significantly different cell surface chemistry than *Shewanella sp.* or *Geobacter sp.* offers an

367 opportunity to dissect these separate effects (48). Thus, new *E. coli* strains described herein and  
368 those enabled by the design rules described herein, will fuel both basic and applied studies.

369

370 **Methods**

371 **Plasmids and strains.**

372 The primers, plasmids, and strains used in this study are listed in **Tables S2, S3, and**  
373 **Table 1**, respectively. The *ccm* (pEC86), *mtrA* (I5024), and *mtrCAB* (I5023) plasmids were  
374 described previously (6). The *cymA* (I5040), *cymAmtrA* (I5052), and *cymAmtrCAB* (I5049)  
375 plasmids were constructed for this work. In brief, these plasmids were constructed using PCR  
376 amplification of the genes from genomic DNA of *Shewanella oneidensis* MR-1 using Pfx  
377 Platinum polymerase (Invitrogen), digestion of the pSB1ET2 plasmid and PCR fragment(s) with  
378 restriction endonucleases (New England Biolabs), ligation of these fragments with T4 DNA  
379 ligase (Roche) and, in some cases, site directed mutagenesis (QuikChange II, Agilent  
380 Technologies). Detailed descriptions of the assembly of I5040, I5052, and I5049 can be found in  
381 the Supporting Information. After sequence verification, the resulting *cymA* plasmid and  
382 *cymAmtrA* were co-transformed with *ccm* into BL21(DE3) to make the *cymA* and *cymAmtrA*  
383 strains, respectively. In contrast, the *mtrCAB* and *cymAmtrCAB* plasmids, were co-transformed  
384 with pEC86 into the *E. coli* strain C43(DE3) (Lucigen, Middleton, WI) to make the *mtr* and  
385 *cymAmtr* strains, respectively.

386 **Growth conditions and medium composition.**

387 All strains, unless otherwise specified, were grown in 2xYT medium at 30 °C with 50 µg  
388 mL<sup>-1</sup> kanamycin; strains containing the pEC86 plasmid were grown with an additional 30 µg mL<sup>-1</sup>  
389 chloramphenicol. Glycerol stocks were used to inoculate 5 mL 2xYT medium, and cultures  
390 were grown overnight at 37 °C with 250-rpm shaking. Then, 500 µL of overnight cultures were  
391 back-diluted into 50 mL 2xYT medium and grown with 200-rpm shaking for 16 h at 30 °C.

392 When indicated, strains were induced with IPTG at an  $OD_{600nm}$  of 0.5-0.7. IPTG concentration is  
393 displayed here as promoter activity defined in Goldbeck, et al (5).

394 Cell suspensions used for iron reduction assays were resuspended in anaerobic defined  
395 M1 medium supplemented with 40 mM D,L-lactate and 0.2% casamino acids. The composition  
396 of the M1 medium can be found in the Supporting Information.

#### 397 **Subcellular Fractionation.**

398 Periplasmic and membrane preparations were performed as previously described (6).  
399 Membrane samples were solubilized in a solution of 5% (w/v) Triton X-100, 50 mM HEPES pH  
400 7.4, and 200 mM NaCl.

#### 401 **Visible spectra of cytochrome samples by diffuse reflectance.**

402 The concentration of cytochromes in whole cells was determined by diffuse reflectance  
403 as previously described by Goldbeck et al. (5). The reduced samples were chemically reduced  
404 with sodium dithionite crystals (Sigma, St. Louis, MO). The extinction coefficients at 552 nm of  
405  $104 \text{ mM}^{-1} \text{ cm}^{-1}$  (personal communication, Prof. Julea Butt) and  $280 \text{ mM}^{-1} \text{ cm}^{-1}$  (35) were used to  
406 determine the CymA and MtrA concentrations, respectively.

#### 407 **ECL to detect *c*-type cytochromes in whole cell lysates.**

408 Cell pellets from 1.5 mL of culture was resuspended in 0.1 mL Bacterial Protein  
409 Extraction Reagent (B-Per, ThermoScientific, Grand Island, NY). The cells were frozen at -20  
410 °C immediately after growth and then thawed immediately before analysis. Cells were lysed for  
411 30 minutes at room temperature with  $6 \mu\text{g mL}^{-1}$  chicken egg white lysozyme (Sigma),  $1 \mu\text{g mL}^{-1}$   
412 DNAase, 3.9 mM  $\text{MgSO}_4$ , 0.96 mM EDTA, and 0.98 mM phenylmethylsulfonyl fluoride. Total  
413 protein of the cell lysates was determined by BCA Protein Assay Kit (ThermoScientific). Cells  
414 were diluted in 100 mM HEPES, pH 7.4, to normalize to equal total protein concentration.

415 Samples were prepared with NuPAGE 4x Sample Buffer (Bio-Rad) and heated at 95 °C for 5  
416 minutes. A total of 8 µg protein was loaded and run in a 4-20% Tris-HCl polyacrylamide gel  
417 (Bio-Rad) at 200 V for 1 hour. The gel was rinsed twice in water and then equilibrated in cold  
418 Pierce Western Transfer buffer (ThermoScientific) for 15 minutes. The proteins were transferred  
419 to a 0.45 µm nitrocellulose membrane (Bio-Rad, Hercules, CA) in Pierce Western Transfer  
420 buffer at 30 V for 100 minutes. Ponceau S staining was used to confirm uniform transfer across  
421 all lanes.

422 The nitrocellulose membrane was incubated for 5 minutes in 10 mL of Pierce Pico West  
423 Enhanced Chemiluminescence substrate (ThermoScientific), a 1:1 mixture of Pico West  
424 Peroxide Solution and Luminol Enhancer solution. The chemiluminescent signal was detected  
425 using the ChemiDoc™ XRS system. The chemiluminescent signal and molecular weights were  
426 quantified using ImageJ (49).

#### 427 **Assaying soluble Fe(III) citrate reduction and cell density of strains.**

428 Fe(III) reduction assays were performed as previously described (6). Briefly, cultures  
429 grown aerobically were pelleted, washed, and resuspended to an OD<sub>600 nm</sub> of 0.5 in anaerobic M1  
430 medium supplemented with 40 mM D,L-lactate as the sole carbon source. All anaerobic media  
431 and buffers were sparged with nitrogen and anaerobic conditions were maintained in an  
432 anaerobic chamber (Coy Laboratory Products, Grass Lake, MI) with an atmosphere of 2% H<sub>2</sub>  
433 balance N<sub>2</sub>. Fe(III) citrate (Sigma, St. Louis, MO) was added to a final concentration of 10 mM.  
434 The Fe(II) concentration was determined with the ferrozine assay, adapted from Stookey (50).  
435 The concentration of Fe(II) in each culture was subtracted by abiotic iron reduction observed in  
436 sterile medium-only controls at each time point. Additionally, a lysed *mtrA* control was included  
437 to determine the extent of non-metabolic Fe(III) reduction (Supporting Information). Cell density

438 was determined by measuring the optical density at 600 nm and subtracting the scattering of  
439 abiotic media with 10mM Fe(III) citrate. Error bars represent the standard deviation across three  
440 biological replicates. The rate of Fe(III) citrate reduction and linear rate of change in optical  
441 density were calculated by best linear fit for each individual biological replicate.

#### 442 **Bulk Fe<sub>2</sub>O<sub>3</sub> (s) reduction assay.**

443 Fe<sub>2</sub>O<sub>3</sub> (s) reduction assays were performed as previously described (6). Briefly, cultures  
444 grown aerobically were pelleted, washed, and resuspended to an OD<sub>600 nm</sub> of 1.0 in anaerobic M1  
445 medium supplemented with 40 mM D,L-lactate as the carbon source, 6.0 mg mL<sup>-1</sup> Fe<sub>2</sub>O<sub>3</sub> (s)  
446 (Sigma) as the sole terminal electron acceptor, and IPTG, where indicated. Where indicated, the  
447 media was supplemented with 5 μM riboflavin.

448 The colony forming units (cfu) and Fe(II) concentration for each culture was measured  
449 after initial anaerobic inoculation and at time points following. Cfus were determined with  
450 kanamycin selection on LB plates grown aerobically at 37 °C. The Fe(II) concentration was  
451 determined with the ferrozine assay, adapted from Stookey (50). The concentration of Fe(II) in  
452 each culture was subtracted by any abiotic iron reduction observed in sterile media-only controls  
453 at each time point. Error bars represent standard deviation of triplicate cultures.

#### 454 **Cyclic Voltammetry of *mtr* and *cymAmtr* strains.**

455 Cells from 50-mL cultures were pelleted and washed twice with M1 medium  
456 supplemented with 40 mM D,L-lactate. Pellets were resuspended to an OD<sub>600nm</sub> of 0.7 in the  
457 working chamber of 2-chambered bioelectrochemical reactors. The working chamber contained  
458 M1 medium supplemented with 40 mM D,L-lactate as the sole carbon source, the counter  
459 electrode chamber contained 50 mM PIPES buffer (pH 7.4), and the two chambers were  
460 separated by a cation exchange membrane (CMI-7000, Membranes International, Ringwood,

461 NJ). The working electrode was a 25 x 25 mm square piece of graphite felt (GF-S6-06,  
462 Electrolytica, Amherst, NY) and was connected to the potentiostat via a piece of Pt wire. The  
463 counter electrode was a piece of Pt wire and the reference electrode was a pre-made Ag/AgCl  
464 reference (CH Instruments, Bee Cave, TX). During current stabilization, the working chambers  
465 were stirred at ~200 rpm with magnetic stir bars, and stirring was turned off during cyclic  
466 voltammetry analysis. Cyclic voltammograms were recorded in 2 cycles from -0.5 V<sub>Ag/AgCl</sub> to  
467 +0.5 V<sub>Ag/AgCl</sub> at a scan rate of 2 mV/s. Cyclic voltammograms were recorded before and  
468 immediately after inoculation. After initial cyclic voltammetry, the working electrodes were held  
469 at +0.2 V<sub>Ag/AgCl</sub> overnight to allow cells to associate with the electrode surface. Cyclic  
470 voltammograms were recorded again after the overnight period.

471

#### 472 **Supporting Information.**

473 SI Figure 1. Schematic of plasmids and localization of MtrC, MtrA, and CymA.

474 SI Figure 2. Diffuse reflectance spectra show the spectral signatures characteristic of  
475 cytochromes *c* in whole cell suspensions.

476 SI Figure 3. ECL analysis shows that the relative abundance of each cyt *c* within a strain is  
477 similar across different induction levels

478 SI Figure 4. Homogenized MtrA *E. coli* reduce Fe(III) citrate at the same basal rate as *ccm*.

479 SI Figure 5. Fitness is strongly correlated to iron reduction rate.

480 SI Figure 6. A scaled schematic of the gram-negative double membrane and Mtr pathway.

481 SI Figure 7. Correlation analysis of iron reduced against heme *c* concentration.

482 SI Figure 8. Riboflavin mildly improves Fe<sub>2</sub>O<sub>3</sub> reduction, but not survival.



483 SI Figure 9. The *ccm* strain shows no temporal correlation between cell density and Fe<sub>2</sub>O<sub>3</sub>  
484 reduction.

485 SI Table 1. Relative promoter activity as a function of IPTG concentration.

486 SI Table 2. Primers

487 SI Table 3. Plasmids

488 Supporting Calculation 1. Approximation of maximum iron reduction rate.

489 Supporting Calculation 2. Percent Fe<sub>2</sub>O<sub>3</sub> reduced from direct contact or riboflavin-mediated  
490 mechanisms.

491 Supporting Methods.

492

### 493 **Abbreviations**

494 Colony forming units: cfu. Cytochrome *c*: cyt *c*. Cytochrome *c* maturation: ccm. Enhanced  
495 chemiluminescence: ECL. Isopropyl β-D-1-thiogalactopyranoside: IPTG.

496

### 497 **Author Information**

498

### 499 **Authors' contributions**

500 HMJ and CMAF conceived the study. HMJ, MAT, MGK performed the experiments, and  
501 CMAF supervised the work. HMJ analyzed the data. HMJ, MAT, MGK and CMAF drafted the  
502 manuscript, which was read, revised and approved by all authors.

503

### 504 **Acknowledgements**

505 Work at the Molecular Foundry was supported by the Office of Science, Office of Basic Energy  
506 Sciences, of the U.S. Department of Energy under Contract No. DE-AC02-05CH11231. CMAF  
507 and HMJ acknowledge support from the Physical Biosciences Program, Office of Science,  
508 Office of Basic Energy Sciences, of the U.S. Department of Energy under Contract No. DE-  
509 AC02--05CH11231. CMAF and MAT acknowledge support from Office of Naval Research,  
510 Award number N000141310551. This work was supported in part by previous breakthroughs  
511 obtained through the Laboratory Directed Research and Development Program of Lawrence  
512 Berkeley National Laboratory under U.S. Department of Energy Contract No. DE-AC02-  
513 05CH11231.

514

## 515 **References**

- 516 1. Fredrickson, J. K., Zachara, J. M., Kennedy, D. W., Dong, H., Onstott, T. C., Hinman, N.  
517 W., and Li, S.-M. (1998) Biogenic iron mineralization accompanying the dissimilatory  
518 reduction of hydrous ferric oxide by a groundwater bacterium, *Geochim. Cosmochim.*  
519 *Acta* 62, 3239–3257.
- 520 2. Narayanan, K., and Sakthivel, N. (2010) Biological synthesis of metal nanoparticles by  
521 microbes, *Adv. Colloid Interface Sci.* 156, 1-13.
- 522 3. Zhuang, W.-Q., Fitts, J. P., Ajo-Franklin, C. M., Maes, S., Alvarez-Cohen, L., and  
523 Hennebel, T. (2015) Recovery of critical metals using biometallurgy, *Curr. Opin.*  
524 *Biotechnol.* 33, 327-335.
- 525 4. Gadd, G. M. (2010) Metals, minerals and microbes: geomicrobiology and  
526 bioremediation, *Microbiology* 156, 609-643.
- 527 5. Goldbeck, C. P., Jensen, H. M., TerAvest, M. A., Beedle, N., Appling, Y., Hepler, M.,  
528 Cambray, G., Mutalik, V., Angenent, L. T., and Ajo-Franklin, C. M. (2013) Tuning  
529 Promoter Strengths for Improved Synthesis and Function of Electron Conduits in  
530 *Escherichia coli*, *ACS Synth. Biol.* 2, 150-159.
- 531 6. Jensen, H. M., Albers, A. E., Malley, K. R., Londer, Y. Y., Cohen, B. E., Helms, B. A.,  
532 Weigele, P., Groves, J. T., and Ajo-Franklin, C. M. (2010) Engineering of a synthetic  
533 electron conduit in living cells, *Proc. Natl. Acad. Sci. U. S. A.* 107, 19213-19218.
- 534 7. TerAvest, M. A., Zajdel, T. J., and Ajo-Franklin, C. M. (2014) The Mtr Pathway of  
535 *Shewanella oneidensis* MR-1 Couples Substrate Utilization to Current Production in  
536 *Escherichia coli*, *ChemElectroChem* 1, 1874-1879.
- 537 8. Gescher, J. S., Cordova, C. D., and Spormann, A. M. (2008) Dissimilatory iron reduction  
538 in *Escherichia coli*: identification of CymA of *Shewanella oneidensis* and NapC of *E.*  
539 *coli* as ferric reductases, *Mol. Microbiol.* 68, 706-719.

- 540 9. Schuetz, B., Schicklberger, M., Kuermann, J., Spormann, A. M., and Gescher, J. (2009)  
541 Periplasmic Electron Transfer via the c-Type Cytochromes MtrA and FccA of  
542 *Shewanella oneidensis* MR-1, *Appl. Environ. Microbiol.* 75, 7789-7796.
- 543 10. Sturm-Richter, K., Golitsch, F., Sturm, G., Kipf, E., Dittrich, A., Beblawy, S.,  
544 Kerzenmacher, S., and Gescher, J. (2015) Unbalanced fermentation of glycerol in  
545 *Escherichia coli* via heterologous production of an electron transport chain and electrode  
546 interaction in microbial electrochemical cells, *Bioresource Technol* 186, 89-96.
- 547 11. Ajo-Franklin, C. M., and Noy, A. (2015) Crossing Over: Nanostructures that Move  
548 Electrons and Ions across Cellular Membranes, *Adv. Mater.* 27, 5797-5804.
- 549 12. Richardson, D. J., Butt, J. N., Fredrickson, J. K., Zachara, J. M., Shi, L., Edwards, M. J.,  
550 White, G., Baiden, N., Gates, A. J., Marritt, S. J., and Clarke, T. A. (2012) The 'porin-  
551 cytochrome' model for microbe-to-mineral electron transfer, *Mol. Microbiol.* 85, 201-  
552 212.
- 553 13. Shi, L., Rosso, K. M., Clarke, T. A., Richardson, D. J., Zachara, J. M., and Fredrickson,  
554 J. K. (2012) Molecular Underpinnings of Fe(III) Oxide Reduction by *Shewanella*  
555 *oneidensis* MR-1, *Front. Microbiol.* 3, 50.
- 556 14. Marritt, S. J., McMillan, D. G. G., Shi, L., Fredrickson, J. K., Zachara, J. M., Richardson,  
557 D. J., Jeuken, L. J. C., and Butt, J. N. (2012) The roles of CymA in support of the  
558 respiratory flexibility of *Shewanella oneidensis* MR-1, *Biochem. Soc. Trans.* 40, 1217-  
559 1221.
- 560 15. Simon, J., van Spanning, R. J. M., and Richardson, D. J. (2008) The organisation of  
561 proton motive and non-proton motive redox loops in prokaryotic respiratory systems,  
562 *Biochim. Biophys. Acta, Bioenerg.* 1777, 1480-1490.
- 563 16. McMillan, D. G. G., Marritt, S. J., Butt, J. N., and Jeuken, L. J. C. (2012) Menaquinone-7  
564 is specific cofactor in tetraheme quinol dehydrogenase CymA, *J. Biol. Chem.* 287,  
565 14215-14225.
- 566 17. Alves, M. N., Neto, S. E., Alves, A. S., and Fonseca, B. M. (2015) Characterization of  
567 the periplasmic redox network that sustains the versatile anaerobic metabolism of  
568 *Shewanella oneidensis* MR-1, *Front. Microbiol.* 6, 665.
- 569 18. Fonseca, B. M., Paquete, C. M., Neto, S. E., Pacheco, I., Soares, C. M., and Louro, R. O.  
570 (2013) Mind the gap: cytochrome interactions reveal electron pathways across the  
571 periplasm of *Shewanella oneidensis* MR-1, *Biochem. J.* 449, 101-108.
- 572 19. Sturm, G., Richter, K., Doetsch, A., Heide, H., Louro, R. O., and Gescher, J. (2015) A  
573 dynamic periplasmic electron transfer network enables respiratory flexibility beyond a  
574 thermodynamic regulatory regime, *ISME J.* 9, 1802-1815.
- 575 20. White, G. F., Shi, Z., Shi, L., Wang, Z., Dohnalkova, A. C., Marshall, M. J., Fredrickson,  
576 J. K., Zachara, J. M., Butt, J. N., Richardson, D. J., and Clarke, T. A. (2013) Rapid  
577 electron exchange between surface-exposed bacterial cytochromes and Fe(III) minerals,  
578 *Proc. Natl. Acad. Sci. U. S. A.* 110, 6346-6351.
- 579 21. Ross, D. E., Brantley, S. L., and Tien, M. (2009) Kinetic Characterization of OmcA and  
580 MtrC, Terminal Reductases Involved in Respiratory Electron Transfer for Dissimilatory  
581 Iron Reduction in *Shewanella oneidensis* MR-1, *Appl. Environ. Microbiol.* 75, 5218-  
582 5226.
- 583 22. Brutinel, E., D., and Gralnick, J., A. . (2012) Shuttling happens: soluble flavin mediators  
584 of extracellular electron transfer in *Shewanella*, *Appl. Microbiol. Biotechnol.* 93, 41-48.

- 585 23. Edwards, M. J., White, G. F., and Norman, M. (2015) Redox Linked Flavin Sites in  
586 Extracellular Decaheme Proteins Involved in Microbe-Mineral Electron Transfer, *Sci.*  
587 *Rep.* 5, 11677.
- 588 24. Okamoto, A., Nakamura, R., and Nealsen, K. H. (2014) Bound Flavin Model Suggests  
589 Similar Electron Transfer Mechanisms in *Shewanella* and *Geobacter*, *ChemElectroChem*  
590 *1*, 1808-1812.
- 591 25. Jiang, X. C., Hu, J. S., Fitzgerald, L. A., Biffinger, J. C., Xie, P., Ringeisen, B. R., and  
592 Lieber, C. M. (2010) Probing electron transfer mechanisms in *Shewanella oneidensis*  
593 MR-1 using a nanoelectrode platform and single-cell imaging, *Proc. Natl. Acad. Sci. U.*  
594 *S. A.* 107, 16806-16810.
- 595 26. Marsili, E., Baron, D. B., and Shikhare, I. D. (2008) *Shewanella* secretes flavins that  
596 mediate extracellular electron transfer, *Proc. Natl. Acad. Sci. U. S. A.* 105, 3968-3973.
- 597 27. Kotloski, N. J., and Gralnick, J. A. (2013) Flavin electron shuttles dominate extracellular  
598 electron transfer by *Shewanella oneidensis*, *mBio* 4.
- 599 28. Nikel, P. I., and de Lorenzo, V. (2013) Engineering an anaerobic metabolic regime in  
600 *Pseudomonas putida* KT2440 for the anoxic biodegradation of 1, 3-dichloroprop-1-ene,  
601 *Metab. Eng.* 15, 98-112.
- 602 29. Schmitz, S., Nies, S., Wierckx, N., Blank, L. M., and Rosenbaum, M. A. (2015)  
603 Engineering mediator-based electroactivity in the obligate aerobic bacterium  
604 *Pseudomonas putida* KT2440, *Front. Microbiol.* 6, 284.
- 605 30. Steen, A., Ütkür, F., Acuña, B.-d. J. M., and Bunk, B. (2013) Construction and  
606 characterization of nitrate and nitrite respiring *Pseudomonas putida* KT2440 strains for  
607 anoxic biotechnical applications, *J. Biotechnol.* 163, 155-165.
- 608 31. Salis, H. M., Mirsky, E. A., and Voigt, C. A. (2009) Automated design of synthetic  
609 ribosome binding sites to control protein expression, *Nat. Biotechnol.* 27, 946-950.
- 610 32. Arslan, E., Schulz, H., Zufferey, R., Künzler, P., and Thöny-Meyer, L. (1998)  
611 Overproduction of the Bradyrhizobium japonicum c-Type Cytochrome Subunits of the  
612 cbb 3 Oxidase in *Escherichia coli*, *Biochem. Biophys. Res. Commun.* 251, 744-747.
- 613 33. Myers, C. R., and Myers, J. M. (1997) Cloning and sequence of *cymA*, a gene encoding a  
614 tetraheme cytochrome c required for reduction of iron (III), fumarate, and nitrate by  
615 *Shewanella putrefaciens* MR-1, *J. Bacteriol.* 179, 1143-1152.
- 616 34. Myers, J. M., and Myers, C. R. (2002) Genetic complementation of an outer membrane  
617 cytochrome *omcB* mutant of *Shewanella putrefaciens* MR-1 requires *omcB* plus  
618 downstream DNA, *Appl. Environ. Microbiol.* 68, 2781-2793.
- 619 35. Pitts, K. E., Dobbin, P. S., Reyes-Ramirez, F., Thomson, A. J., Richardson, D. J., and  
620 Seward, H. E. (2003) Characterization of the *Shewanella oneidensis* MR-1 Decaheme  
621 Cytochrome MtrA, *J. Biol. Chem.* 278, 27758-27765.
- 622 36. Wang, V. B., Yantara, N., Koh, T. M., Kjelleberg, S., Zhang, Q., Bazan, G. C., Loo, S. C.  
623 J., and Mathews, N. (2014) Uncovering alternate charge transfer mechanisms in  
624 *Escherichia coli* chemically functionalized with conjugated oligoelectrolytes, *Chem.*  
625 *Commun. (Cambridge, U. K.)* 50, 8223-8226.
- 626 37. Carmona-Martinez, A. A., Harnisch, F., Fitzgerald, L. A., Biffinger, J. C., Ringeisen, B.  
627 R., and Schröder, U. (2011) Cyclic voltammetric analysis of the electron transfer of  
628 *Shewanella oneidensis* MR-1 and nanofilament and cytochrome knock-out mutants,  
629 *Bioelectrochemistry* 81, 74-80.

- 630 38. Matias, V. R. F., Al-Amoudi, A., Dubochet, J., and Beveridge, T. J. (2003) Cryo-  
631 transmission electron microscopy of frozen-hydrated sections of *Escherichia coli* and  
632 *Pseudomonas aeruginosa*, *J. Bacteriol.* 185, 6112-6118.
- 633 39. Mullineaux, C. W., Nenner, A., and Ray, N. (2006) Diffusion of green fluorescent  
634 protein in three cell environments in *Escherichia coli*, *J. Bacteriol.* 188, 3442-3448.
- 635 40. Firer-Sherwood, M. A., Bewley, K., D. , Mock, J.-Y., and Elliott, S. J. (2011) Tools for  
636 resolving complexity in the electron transfer networks of multiheme cytochromes c,  
637 *Metalomics* 3, 344-348.
- 638 41. Andersen, K. B., and Meyenburg, K. (1980) Are growth rates of *Escherichia coli* in batch  
639 cultures limited by respiration?, *J. Bacteriol.* 144, 114-123.
- 640 42. Bose, S., Hochella, M. F., Gorby, Y. A., Kennedy, D. W., McCready, D. E., Madden, A.  
641 S., and Lower, B. H. (2009) Bioreduction of hematite nanoparticles by the dissimilatory  
642 iron reducing bacterium *Shewanella oneidensis* MR-1, *Geochim. Cosmochim. Acta* 73,  
643 962-976.
- 644 43. TerAvest, M. A., and Ajo-Franklin, C. M. (2015) Transforming exoelectrogens for  
645 biotechnology using synthetic biology, *Biotechnol. Bioeng.* in press, DOI:  
646 10.1002/bit.25723.
- 647 44. Pinchuk, G. E., Rodionov, D. A., Yang, C., Li, X., Osterman, A. L., Dervyn, E.,  
648 Geydebekht, O. V., Reed, S. B., Romine, M. F., Collart, F. R., Scott, J. H., Fredrickson,  
649 J. K., and Beliaev, A. S. (2009) Genomic reconstruction of *Shewanella oneidensis* MR-1  
650 metabolism reveals a previously uncharacterized machinery for lactate utilization, *Proc.*  
651 *Natl. Acad. Sci. U. S. A.* 106, 2874-2879.
- 652 45. Bonneville, S., Behrends, T., Cappellen, P., Hyacinthe, C., and Röling, W. F. M. (2006)  
653 Reduction of Fe (III) colloids by *Shewanella putrefaciens*: A kinetic model, *Geochim.*  
654 *Cosmochim. Acta* 70, 5842-5854.
- 655 46. Glasauer, S., Langley, S., and Beveridge, T. J. (2001) Sorption of Fe (hydr) oxides to the  
656 surface of *Shewanella putrefaciens*: cell-bound fine-grained minerals are not always  
657 formed de novo, *Appl. Environ. Microbiol.* 67, 5544-5550.
- 658 47. Pratt, L. A., and Kolter, R. (1998) Genetic analysis of *Escherichia coli* biofilm formation:  
659 roles of flagella, motility, chemotaxis and type I pili, *Mol. Microbiol.* 30, 285-293.
- 660 48. De Windt, W., Boon, N., and Van den Bulcke, J. (2006) Biological control of the size and  
661 reactivity of catalytic Pd (0) produced by *Shewanella oneidensis*, *Antonie van*  
662 *Leeuwenhoek* 90, 377-389.
- 663 49. Schneider, C. A., Rasband, W. S., and Eliceiri, K. W. (2012) NIH Image to ImageJ: 25  
664 years of image analysis, *Nat. Methods* 9, 671-675.
- 665 50. Stookey, L. L. (1970) Ferrozine---a new spectrophotometric reagent for iron, *Anal.*  
666 *Chem.* 42, 779-781.

669

670 **Table 1. Strains used in this study.**

Strain	Parental Strain	Plasmid(s)	Gene(s)	Source	Request Name
<i>ccm</i>	BL21(DE3)	pEC86, pSB1ET2	<i>ccmA-H</i>	Jensen 2010	MFe208
<i>mtrA</i>	BL21(DE3)	pEC86, I5024	<i>ccmA-H, mtrA</i>	Jensen 2010	MFe291
<i>cymA</i>	BL21(DE3)	pEC86, I5040	<i>ccmA-H, cymA</i>	this work	MFe431
<i>cymAmtrA</i>	BL21(DE3)	pEC86, I5052	<i>ccmA-H, cymAmtrA</i>	this work	MFe538
<i>ccm</i>	C43(DE3)	pEC86, pSB1ET2	<i>ccmA-H</i>	Goldbeck 2013	MFe408
<i>mtr</i>	C43(DE3)	pEC86, I5023	<i>ccmA-H, mtrCAB</i>	this work	MFe409
<i>cymAmtr</i>	C43(DE3)	pEC86, I5049	<i>ccmA-H, cymAmtrCAB</i>	this work	MFe444

671

672

673 **Figure 1. CymA is co-expressed with MtrCAB to provide an extracellular electron transfer**

674 **pathway that spans both membranes of *E. coli*.** (A) Schematic of the Mtr pathway in *S.*

675 *oneidensis*. Electrons present in menaquinone are routed across the inner membrane by CymA

676 (blue box) and are then sequentially transferred to MtrA and to MtrC through the MtrCAB

677 electron nanoconduits (red box). Electrons can then be passed from MtrC to Fe<sub>2</sub>O<sub>3</sub> either directly

678 or through a flavin-mediated process (green box). (IM = inner membrane. OM = outer

679 membrane.) (B) Heme staining of whole cell lysates of the *cymA*, *cymAmtrA*, and *mtrA* strains

680 show CymA and MtrA are present in the respective strains. Similarly, heme staining of whole

681 cell lysates of the *mtr* and *cymAmtr* strain shows both strains contain MtrA and MtrC. Heme

682 staining was performed via ECL in (B). (C) Densitometry of the *cymAmtrA* (green), *mtr* (red),

683 and *cymAmtr* (blue) containing lanes shown in (B) show that CymA is present only in the

684 *cymAmtr* strain. (D) Heme *c* concentration per cell density for the *mtr* and *cymAmtr* strains as a

685 function of relative promoter activity. Heme *c* concentration was measured by diffused

686 reflectance and relative promoter activity is measured as Goldbeck et al (5).

687

688 **Figure 2. Co-expression of CymA with MtrA improves Fe(III) citrate reduction and**

689 **maintains biomass in *E. coli*.** (A) Fe(II) concentration as a function of time for the *ccm* (open

690 black circles), *mtrA* (closed red circles), *cymA* (half-filled green circles), and *cymAmtrA* (half-

691 filled blue circles) strains. The *cymA* and *cymAmtrA* strains reduce Fe(III) citrate ~3x and ~4x

692 faster than the *mtrA* strain. (B) Change in cell density over time for the *ccm* (open black circles),  
693 *mtrA* (closed red circles), *cymA* (half-filled green circles), and *cymAmtrA* (half-filled blue circles)  
694 strains. Only the *cymAmtrA* strain maintains constant biomass under Fe(III) reducing conditions.  
695

696 **Figure 3. Co-expression of CymA with MtrCAB increases Fe<sub>2</sub>O<sub>3</sub> (s) reduction and**  
697 **maintains biomass in *E. coli*.** (A) The concentration of bulk  $\alpha$ -Fe<sub>2</sub>O<sub>3</sub> reduced to Fe(II) by *ccm*  
698 and the best performing *mtr* and *cymAmtr* strains, showing that *cymAmtr*<sup>0.001</sup> significantly  
699 increased iron reduction over both the *ccm* and the *mtr*<sup>0.37</sup> strains ( $p = 0.002$  and  $0.031$ ,  
700 respectively). This indicates that the co-expression of CymA with Mtr increases the flux of  
701 electrons to iron oxide. (B) Relative cell density of the *ccm*, *mtr*<sup>0.37</sup> and *cymAmtr*<sup>0.001</sup> strains after  
702 4 days under Fe<sub>2</sub>O<sub>3</sub> (s) reducing conditions, showing that only the *cymAmtr* strain maintains  
703 constant biomass. The relative cell density is the cell density after 4 days divided by the initial  
704 cell density. Average values (n=3) are plotted and error bars represent standard deviation.  
705

706 **Figure 4. The expression of CymA significantly increases current production by enabling**  
707 **catalytic turnover of Mtr.** (A) Current production in electrochemical reactors with working  
708 electrodes poised at +0.2 V<sub>Ag/AgCl</sub> increased significantly when *cymA* was co-expressed with *mtr*.  
709 Average values (n=4) are plotted and error bars represent standard deviation. (B) Turnover cyclic  
710 voltammetry (representative scans plotted) at 2 mV/s reveals a catalytic wave only when *cymA* is  
711 co-expressed with *mtr*, indicating that the inner membrane cytochrome is necessary for catalytic  
712 turnover of the outer membrane cytochromes.  
713

714 **Figure 5. Riboflavin improves Fe<sub>2</sub>O<sub>3</sub> (s) reduction and permits growth in the *cymAmtr***  
715 **strain.** (A) Riboflavin significantly enhanced iron reduction by ~2.5x in the *cymAmtr*<sup>0.18</sup> strain ( $p$   
716 = 0.001). Assuming direct contact contributes equally in the cultures with and without the  
717 supplementation of riboflavin, 77% of the iron oxide was reduced by riboflavin. (B) The survival  
718 of the *cymAmtr*<sup>0.18</sup> strain dramatically increased in the presence of riboflavin ( $p = 3 \times 10^{-5}$ ), but not  
719 in the *mtr*<sup>0.18</sup> strain ( $p = 0.13$ ). Average values (n = 3) are plotted and error bars represent  
720 standard deviation. (C) Concentration of Fe(II) produced and colony forming units as a function  
721 of time for the *cymAmtr*<sup>0.18</sup> cultures with riboflavin, showing a that the initial period of rapid

722 Fe(II) production is accompanied by increase in cell density for the *cymAmtr* strain with  
723 riboflavin.

724