

# UC Agriculture & Natural Resources

## California Agriculture

### Title

Blue oak seedling age influences growth and mortality

### Permalink

<https://escholarship.org/uc/item/2rp454q1>

### Journal

California Agriculture, 61(1)

### ISSN

0008-0845

### Authors

Phillips, Ralph L.  
McDougald, Neil K.  
McCreary, Doug D.  
[et al.](#)

### Publication Date

2007

Peer reviewed

# Blue oak seedling age influences growth and mortality

by Ralph L. Phillips, Neil K. McDougald,  
Doug McCreary and Edward R. Atwill

*In some California locations, the natural regeneration of blue oak is limited by the ability of small seedlings to survive long enough to become larger saplings. This study evaluated the growth and survival of different age classes of seedlings. We found that over a 7-year period, older blue oak seedlings had a much higher survival rate than younger seedlings. Under this study's conditions, however, the height of younger seedlings increased while that of older seedlings decreased. These results suggest that once a seedling survives approximately a decade and becomes established, it is much more likely to remain alive compared to newly germinated seedlings. Nonetheless, its height growth rate may be extremely slow.*

Blue oak trees are a valuable economic and aesthetic resource in the Sierra Nevada foothills of California. Ecologically, oak woodlands, including blue oak (*Quercus douglasii*) woodlands, provide some of the richest wildlife habitat in the state (Giusti et al. 2005). However, for nearly a century there have been concerns about the adequacy of natural blue oak regeneration (Jepson 1910). In some locations, natural regeneration may be insufficient to sustain populations (Bartolome et al. 1987; Swiecki et al. 1997a), while in others it may be more than adequate to replace mortality (Wieslander 1935). This conflicting information is cause for concern among landowners, governmental agencies, conservationists and others interested in the long-term sustainability of this species (see page 7).

Standiford et al. (1991) conducted an oak-tree size survey in 1987 and found large numbers of blue oak seedlings less

than 1.0 feet (0.3 meter tall), fewer seedlings in the 4.3-to-10.2 feet (1.3-to-3.1-meters) category, but an adequate stand of mature trees (at least 100 trees per acre). Swiecki et al. (1997b) also reported insufficient numbers of blue oak saplings at 13 of 15 locations sampled throughout the state to replace tree mortality.

Information regarding the mortality of small oak seedlings is limited and not always consistent. Allen-Diaz and Bartolome (1992) reported 50% annual mortality over a 3-year study, while Swiecki et al. (1991) reported that 75.7% of blue oak seedlings survived over a 3-year study. Neither study identified the age of the seedlings.

Livestock grazing, drought and competition for soil moisture from nonnative annual plants have been suggested as factors that influence blue oak seedling survival, but there is limited biological information about which factors are most important (Phillips et al. 1996).



California's oak woodlands are critical, providing rich wildlife habitat, soil stabilization and beautiful scenery, as shown at the UC Hastings Natural History Reservation in Carmel Valley. Oaks are long-lived but also slow-growing trees; regeneration rates for some species may not be adequate to replace natural mortality.

## Age groups and causes of mortality

The first part of this study was initiated in oak woodlands in the southern Sierra Nevada foothills of Kern County, in 1989 (Phillips et al. 1997). At that time, 600 naturally regenerated blue oak seedlings were identified in 13 0.1-acre (0.04-hectare) plots at three locations. In 1993, a technique described by Phillips et al. (1997) was used to estimate the age of the surviving seedlings from this group. The technique involved using regression analysis to develop a model of the relationship between root-crown diameter and the number of growth rings. Root crowns of the seedlings identified in 1989 were measured in 1993 and the regression equation was used to estimate seedling age.

The second part of the study was initiated in 1997. There was a large blue oak acorn crop in fall 1996, and the subsequent winter moisture of late 1996 and early 1997 was adequate to



**Oak seedlings often must compete with dense annual grasses for moisture and nutrients, while rodents and livestock browse on them. A long-term study in Kern County assessed the growth rates and mortality of blue oak seedlings (flagged).**

support what appeared to be above-normal acorn germination. In spring 1997, a large number of newly germinated blue oak seedlings were present in the 13 plots established in 1989. During early summer 1997, after the annual vegetation had dried so that seedlings could be located more easily, each of the original 13 plots was evaluated for the presence of newly sprouted seedlings. In order for seedlings to be classified as newly sprouted, the acorn had to still be attached to the seedling. Each newly sprouted seedling was identified by a number, plot number and year. Seedlings in the old group in these plots had already been individually identified in a similar fashion.

The 1989 seedlings were the first age group (old) and the 1997 seedlings were the second (new). Seedlings from the 1989 and 1997 groups were evaluated annually from 1997 to 2004. These assessments were conducted at the end of the growing season, when the forage had turned brown.

Height measurements were taken in 1997 and 2001 on both the 1997 (new) and 1989 (old) seedling groups, and mortality was assessed annually. The cause of death was categorized as either rodent-caused or non-rodent-caused. To be classified as rodent-caused, there had to be clear evidence that rodents had destroyed the seedling; either rodent burrows present near the seedling, or if

dead seedlings could be pulled up, visible teeth marks on the roots. All other dead seedlings were categorized as non-rodent-caused.

#### Analysis of age groups and height

In 1997, there were 300 seedlings in the old seedling group and 764 seedlings in the new group. The effect of age group and initial height on absolute height change (height in 2001 divided by height in 1997) and the relative height change ( $[\text{height in 2001} \div \text{height in 1997}]$  divided by height in 1997) were analyzed using a generalized least squares regression analysis, with a stratified variance function for each age group due to unequal variance structures (Pinheiro and


**TABLE 1. Generalized least squares regression model for relative effects of age and initial height of blue oak seedlings on the change in height, 1997–2001**

Factor	Regression coefficient	P value*
Model intercept	2.37	< 0.001
Height in 1997 (inches)	-0.72	< 0.001
Age group†		
New seedlings	0.00	—
Old seedlings	-2.05	< 0.001
Height in 1997 × age-group interaction		
Height in 1997 × new‡	0.00	—
Height in 1997 × old	0.58	< 0.001

\* Significance determined by a conditional t-test.

† New = 1997; old = established prior to 1997.

‡ Referent group of seedlings for the regression model.



**Fig. 1. Effect of age and initial height of blue oak seedlings on the difference in height, 1997 to 2001.**



Oak seedlings are often repeatedly browsed by animals such as deer, keeping them small and stunted for decades.



In 1989, the estimated age of this seedling was 11 years.

Bates 2000). The initial height, average age in 1997 and average age in 2001 were analyzed using analysis of variance. The percentage annual death rate was analyzed using a linear mixed-effects regression model (Pinheiro and Bates 2000) after the outcome variable had been transformed using an arcsine transformation (Neter et al. 1990). The percentage annual death loss was calculated as the number of seedlings that were alive the previous year divided into the number of seedlings that died during the following year. The survival and height data reported were from those seedlings not affected by rodents.

### Changes in height

The overall mean height in 1997 was 2.0 inches (5.0 centimeters) for the new group of seedlings and 12.0 inches (30.0 centimeters) for the old (established) group. The 2001 average height was

3.0 inches (7.4 centimeters) for the new group and 11.0 inches (27.4 centimeters) for the old group, with an overall mean change in height of 0.96 inch (2.4 centimeters) and -1.0 inch (-2.5 centimeters), respectively. The generalized least squares regression model showed a significant difference in the change in height between the new and old seedling groups (table 1).

However, this difference was strongly influenced by the initial height of the seedling in 1997 (fig. 1). In other words, regardless of whether the seedling was new or old (established), the taller it was in 1997, the more likely it was to lose height during the 4-year study interval. This was most likely due to the dieback of shoots from lack of moisture. Consistent with this result, Phillips et al. (unpublished data, 2004) observed that shorter seedlings in-

creased in height faster than taller seedlings over a 12-year observation period. In this larger study, 599 seedlings were initially followed from 1998 to 2001, in the same location as the present study.

Similar to the results shown in table 1, the generalized least squares regression model for the percentage change in height between 1997 and 2001 (table 2) showed a significant difference between the old and new seedlings. While the majority of new seedlings gained 50% to 200% in height relative to their 1997 levels, many older seedlings lost 10% to 30% of their height. This difference in the relative percentage change of height was strongly influenced by the initial height of the seedling in 1997.

### Mortality and survival

The initial height of seedlings in the old group that died between 1997 and

TABLE 2. Generalized least squares regression model for relative effects of age and initial height of blue oak seedlings on relative percentage change in height, 1997–2001\*

Factor	Regression coefficient	P value†
Model intercept	2.09	< 0.001
Height in 1997 (inches)	-0.74	< 0.001
Age group‡		
New seedlings	0.00	—
Old seedlings	-2.08	< 0.001
Height in 1997 × age group interaction		
Height in 1997 × new§	0.00	—
Height in 1997 × old	0.73	< 0.001

\* Relative percentage change in height determined as height in 2001 divided by height in 1997.

† Significance determined by a conditional t-test.

‡ New = 1997; old = established prior to 1997.

§ Referent group of seedlings for the regression model.

TABLE 3. Initial height and age of seedlings that lived or died for new and old age groups, 1997–2001

Seedling status	Average initial height inches (cm)	Average age in	
		1997	2001
		years . . . . .	
New seedlings*			
Lived	2.1 (5.2)	< 1.0	5.0
Died	1.9 (4.7)	< 1.0	2.2
Old seedlings			
Lived	11.6 (29.1)†	11.5†	15.0†
Died	6.5 (16.3)†	6.4†	9.4†

\* New = 1997; old = established prior to 1997.

† Means within age groups are different at the 0.05 level.




Fig. 2. Observed survival (%) for old versus new seedlings, 1997 to 2004.




Fig. 3. Average percentage annual death loss for old versus new seedlings from the linear mixed-effects regression model.

2001 was significantly less than the initial height of seedlings that were alive in 2001 (6.5 and 11.6 inches [16.3 versus 29.1 centimeters], respectively) (table 3). Therefore, in the old group shorter seedlings died at a faster rate than taller seedlings. On the other hand, the initial height of

### Blue oak seedlings are especially vulnerable to mortality factors when they are young and small.

living seedlings from the new group in 2001 was not significantly different (2.1 inches [5.2 centimeters]) from that of seedlings that died between 1997 and 2001 (1.9 inches [4.7 centimeters]). Phillips et al. (unpublished data, 2004) found that 2.0-inch (5.0-centimeter) seedlings had less than a 10% probability of surviving, while 16.0-inch (40.0-centimeter) seedlings had a 90% probability. However, this relationship did not hold true for the newer seedlings in this study. Also, in Phillips et al. (unpublished data, 2004), shorter seedlings increased in height at a greater annual rate than did taller seedlings.

The average age of surviving old seedlings in 2001 was 11.5 years, and the average age of seedlings that died between 1997 and 2002 was 6.4 years (table 3). The difference in the age of surviving seedlings versus seedlings that died suggests that once blue oak seedlings survive to a certain age, they

are much more likely to remain alive than are newly germinated seedlings.

After the first year, only 46.0% of the new seedlings survived, as compared to 79.0% for the old group (fig. 2). By the 7th year, only 8.7% of the new seedlings were still alive, compared to 48.2% of the old seedlings. The linear mixed-effects regression model for the percentage annual death loss shows a significant difference in death rate for the two age groups (table 4). New seedlings died at a significantly faster rate than older seedlings. The death rate caused by rodents was not different between the two groups (about 1.5% for both groups).

### Influence of precipitation

Mortality tended to fluctuate from year to year, presumably because of differing environmental conditions (fig. 3). The precipitation for the moisture year 1997-1998 was 27.0 inches (67.4 centimeters), more than double the 40-year average of 12.7 inches (31.8 centimeters). The 1998-1999, 1999-2000 and 2000-2001 annual rainfalls, on the other hand, were closer to the long-term average at 10.6, 12.2 and 9.1 inches (26.5, 30.6 and 22.8 centimeters), respectively. However, there was not a significant correlation between annual death loss and precipitation.

Previous research regarding the relationship between precipitation — or soil moisture — and blue oak regeneration

has not been consistent. For instance, a 15-year study by Phillips et al. (unpublished data, 2004) found a significant, positive correlation between annual death loss and precipitation in older seedlings, while Gordon et al. (1991) reported that both the growth and survival of planted blue oak seedlings were positively related to soil water availability. In this study, the effects of precipitation on seedling growth and survival were unclear due to the shorter duration of the evaluation period and the fact that the study was not replicated in different rainfall zones.

### Protecting seedlings

The two age groups of blue oak seedlings had different survival percentages and rates of height growth during the 7 years of this study. The data indicates that blue oak seedlings are especially vulnerable to mortality factors when they are young and small. Younger seedlings died at a far faster rate than larger seedlings, but those that remained alive tended to grow faster than their older counterparts. Once seedlings had grown for approximately a decade and become established, however, the chances were good that they would remain alive, although many grew extremely slowly or even diminished in height. If enhancing natural oak regeneration is a goal and efforts are made to protect existing oak seedlings to help them grow to the sap-

**TABLE 4. Linear mixed-effects regression model for percentage annual death loss for old versus new blue oak seedlings\***


Factors	Regression coefficient	P value†
Model intercept	1.53	< 0.001
Age group‡		
New seedlings§	0.00	—
Old seedlings	-0.54	< 0.001
Year		
1998§	0.00	—
1999	-0.45	0.03
2000	-0.88	< 0.001
2001	-0.73	< 0.001
2002	-0.28	0.18
2003	-0.46	0.03
2004	-0.45	< 0.001

\* Percentage annual death was arcsine-transformed prior to statistical modeling (Neter et al. 1990).

† Significance determined as a conditional t-test.

‡ New = 1997; old = established prior to 1997.

§ Referent group of seedlings for the regression model.



UC Natural Reserve System



**Acorns are a critical food for numerous wildlife species; the acorn woodpecker stores acorns in a dead “granary” tree. In this study, blue oak seedlings became established after about a decade and were more likely to survive than younger seedlings.**

ling stage (see page 16), we recommend that these efforts focus on established, older seedlings rather than on newly germinated ones.

R.L. Phillips was Range/Natural Resources and Livestock Advisor, UC Cooperative Extension (UCCE), Kern and Tulare counties (now retired); N.K. McDougald is Area Watershed Management Advisor, UCCE Madera and Fresno counties; D. McCreary is Area Natural Resource Specialist, Sierra Foothill Research and Extension Center, Browns Valley; and E.R. Atwill is Professor of Environmental Animal Health and Medical Ecology, School of Veterinary Medicine, UC Davis.

**References**

Allen-Diaz BH, Bartolome JW. 1992. Survival of *Quercus douglasii* (Fagaceae) seedlings under the influence of fire and grazing. *Madroño* 39:47–53.  
 Bartolome JW, Muick PC, McClaran MP. 1987. Natural regeneration of California hardwoods. In:

Plumb TR, Pillsbury TH (tech. coords.). Proc Symp on Multiple-Use Management of California Hardwood Resource. Nov 12–14, 1986, San Luis Obispo, CA. USDA Forest Service Gen Tech Rep PSW-100:26–31.  
 Giusti GA, McCreary DD, Standiford RB. 2005. *A Planner's Guide for Oak Woodlands* (9th ed.). UC ANR Pub 3491. 116 p.  
 Gordon DR, Rice KL, Welker JM. 1991. Soil water effects on blue oak seedling establishment. In: Standiford RB (tech. coord.). Proc Symp on Oak Woodlands and Hardwood Rangeland Management. Oct 31–Nov 2, 1990, Davis, CA. USDA Forest Service Gen Tech Rep PSW-126:54–8.  
 Jepson WL. 1910. *The Silva of California (Memoirs of the University of California)*. Berkeley, CA: The University Press. 480 p.  
 Neter J, Wasserman W, Kutner MH. 1990. *Applied Linear Statistical Models*. Richard D Irwin Pub. p 607–32.  
 Phillips RL, McDougald NK, Standiford RB, Frost WE. 1996. Blue oak seedlings may be older than they look. *Cal Ag* 51(3):17–9.  
 Phillips RL, McDougald NK, Standiford RB, et al. 1997. Blue oak regeneration in southern Sierra Nevada foothills. In: Pillsbury NH, Verner J, Tietje WD (tech. coords.). Proc Symp on Oak Woodlands: Ecology, Management and Urban Interface Issues. March 19–22, 1996, San Luis Obispo, CA. USDA Forest Ser-

vice Gen Tech Rep PSW-160:177–81.  
 Pinheiro JC, Bates DM. 2000. *Mixed-Effects Models in S and S-Plus*. New York: Springer. p 201–66.  
 Standiford RB, McDougald NK, Phillips RL, Nelson A. 1991. South Sierra oak regeneration weak in sapling stage. *Cal Ag* 45:12–4.  
 Swiecki TJ, Bernhardt EA, Arnold RA. 1991. Insect and disease impacts on blue oak acorns and seedlings. In: Standiford RB (tech. coord.). Proc Symp on Oak Woodlands and Hardwood Rangeland Management. Oct 31–Nov 2, 1990, Davis, CA. USDA Forest Service Gen Tech Rep PSW-126:149–55.  
 Swiecki TJ, Bernhardt EA, Drake C. 1997a. Factors affecting blue oak sapling recruitment. In: Pillsbury NH, Verner J, Tietje WD (tech. coords.). Proc Symp on Oak Woodlands: Ecology, Management and Urban Interface Issues. March 19–22, 1996, San Luis Obispo, CA. USDA Forest Service Gen Tech Rep PSW-GTR-160:157–67.  
 Swiecki TJ, Bernhardt EA, Drake C. 1997b. Stand-level status of blue oak sapling recruitment and regeneration. In: Pillsbury NH, Verner J, Tietje WD (tech. coords.). Proc Symp on Oak Woodlands: Ecology, Management and Urban Interface Issues. March 19–22, 1996, San Luis Obispo, CA. USDA Forest Service Gen Tech Rep PSW-GTR-160:147–56.  
 Wieslander AE. 1935. A vegetation type map of California. *Madroño* 3:140–4.