

UC San Diego

UC San Diego Previously Published Works

Title

Addendum: Impact of downward longwave radiative deficits on Antarctic sea-ice extent predictability during the sea ice growth period (2022 Environ. Res. Lett. 17 084008)

Permalink

<https://escholarship.org/uc/item/2sm0m5st>

Journal

Environmental Research Letters, 18(2)

ISSN

1748-9318

Authors

Cerovečki, Ivana
Sun, Rui
Bromwich, David H
[et al.](#)

Publication Date

2023-02-01

DOI

10.1088/1748-9326/acb162

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

ENVIRONMENTAL RESEARCH
LETTERS

ADDENDUM

OPEN ACCESS

RECEIVED
15 December 2022ACCEPTED FOR PUBLICATION
9 January 2023PUBLISHED
20 January 2023

Original content from
this work may be used
under the terms of the
[Creative Commons
Attribution 4.0 licence](#).

Any further distribution
of this work must
maintain attribution to
the author(s) and the title
of the work, journal
citation and DOI.



Addendum: Impact of downward longwave radiative deficits on Antarctic sea-ice extent predictability during the sea ice growth period (2022 *Environ. Res. Lett.* **17** 084008)

Ivana Cerovečki¹ , Rui Sun¹, David H Bromwich² , Xun Zou¹ , Matthew R Mazloff¹
and Sheng-Hung Wang² 

¹ Scripps Institution of Oceanography, University of California, San Diego, CA, United States of America

² Byrd Polar and Climate Research Center, The Ohio State University, Columbus, OH, United States of America



In our article and its supplemental material, we examined the causes for a systematic model overestimation of the Antarctic sea-ice edge. We concluded that this overestimation was not due to uncertainties in the sea ice or ocean models, but due to an underestimate of downwelling longwave radiation (DLW). We performed an extensive sensitivity analysis, testing various aspects of the simulation, such as the sea ice dynamics solver, the sea ice lead closing parameter h_0 , the ocean vertical diffusion coefficients, and different ocean initial conditions. However, all simulations overestimated the predicted sea-ice growth during the sea-ice growth period.

Since publication, however, we have found an important source of uncertainty in the sea ice model related to parameterized heat exchanges between ice and ocean (Maykut and McPhee 1995, Holland and Jenkins 1999, Kim *et al* 2006, Hunke *et al* 2017). In our Massachusetts Institute of Technology (MIT) general ocean circulation model (MITgcm) simulations presented in the manuscript, we used the parameters to simulate sea ice from Version 4 Release 4 of ECCO (Estimating the Circulation and Climate of the Ocean) global ocean and sea ice model (Forget *et al* 2015, ECCO Consortium 2020). In this implementation, sea ice can form before the ocean temperature in the top ocean model grid cell reaches the freezing point due to a parameterized limitation in the rate of available heat transfer from the ocean (Fenty 2010). Limiting the heat transfer rate from the ocean to ice is a common assumption. The heat exchange rate is determined by a constant times a friction velocity times the temperature above the freezing point (equation (5.29) in McPhee 2017). This limitation is often not considered until the ice has formed; however, the MITgcm always considers this limited heat availability. This is justified as the temperature of the upper grid cell represents a bulk heat content, and not the true 'Sea Surface Temperature'. We can increase

the ocean heat availability preventing sea ice formation until the upper ocean cell is at the freezing temperature by increasing a constant in the model referred to as 'SEAICE_mcPheePiston'. In doing so the sea ice growth rate can be reduced compared to the results shown in our article (figure 1).

Because available heat flux from the upper ocean model grid cell is limited in the case presented in the paper, sea ice will grow where this cell temperature is warmer than the freezing point. Sea ice growth can be slowed down by changing this parameterization and requiring that the upper grid cell fully reaches the freezing temperature in order for sea ice to form (figure 1). Figure 2 shows the sea ice extent and upper grid cell temperatures for three cases of different heat transfer rate on 31 May 2018. Figure 2 also shows these quantities in the European Centre for Medium-Range Weather Forecasts (ECMWF) forecast, providing evidence that upper ocean model temperature can be above freezing when ice forms in that system as well. Since the forecasts are modeling the bulk temperature in the upper grid cell, further work is needed to determine which solutions are a better representation of the actual oceanic conditions and to determine the uncertainty in handling this parameterization. This conclusion will depend on the observed temperature scales of variability.

The evidence in the article remains valid, showing that the ECMWF reanalysis 5th generation and other reanalyses underestimate DLW during the sea ice growth period in the Antarctic region. Comparison to observations implied large uncertainties in the reanalysis DLW radiation. We found a deficit of approximately 20 W m^{-2} when compared to the Ocean Observatories Initiative mooring data in the South Pacific at 54.08° S , 89.67° W . We also were not able to simulate sea ice growth consistent with observations without enhancing DLW. However, in this addendum, we now add the significant caveat that



the sea-ice model is also very sensitive to the parameterization of available heat content from the ocean. Although the uncertainty of the DLW radiation is still a major concern, we also now highlight the need for a better understanding of the sea ice-ocean energy budget in the marginal ice zone throughout the sea ice growing season. A dedicated effort could yield the necessary data to identify the cause of these model biases and improve Antarctic sea ice predictability.

ORCID iDs

Ivana Cerovečki <https://orcid.org/0000-0001-8979-9952>

David H Bromwich <https://orcid.org/0000-0003-4608-8071>

Xun Zou <https://orcid.org/0000-0003-1620-3198>
Sheng-Hung Wang <https://orcid.org/0000-0002-8057-835X>

References

- Fukumori I, Wang O, Fenty I, Forget G, Heimbach P and Ponte R M (ECCO Consortium) 2020 Synopsis of the ECCO central production global ocean and sea-ice state estimate (Version 4 Release 4) *Zenodo* (<https://doi.org/10.5281/ZENODO.3765929>)
- Fenty I G 2010 State estimation of the Labrador Sea with a coupled sea ice-ocean adjoint model *Doctoral Dissertation* Massachusetts Institute of Technology
- Forget G, Campin J M, Heimbach P, Hill C N, Ponte R M and Wunsch C 2015 ECCO version 4: an integrated framework for non-linear inverse modeling and global ocean state estimation *Geosci. Model Dev.* **8** 3071–104

- Holland D M and Jenkins A 1999 Modeling thermodynamic ice–ocean interactions at the base of an ice shelf *J. Phys. Oceanogr.* **29** 1787–800
- Hunke E, Lipscomb W, Jones P, Turner A, Jeffery N and Elliott S 2017 CICE, the Los Alamos sea ice model *CICE; 005315WKSTN00* (Los Alamos, NM: Los Alamos National Lab. (LANL))
- Kim J G, Hunke E C and Lipscomb W H 2006 Sensitivity analysis and parameter tuning scheme for global sea-ice modeling *Ocean Modelling* **14** 61–80
- Maykut G A and McPhee M G 1995 Solar heating of the Arctic mixed layer *J. Geophys. Res.* **100** 24691–703
- McPhee M G 2017 The sea ice–ocean boundary layer *Sea Ice* (Hoboken, NJ: Wiley) pp 138–59