

UC Irvine

UC Irvine Previously Published Works

Title

Adipocytokines in renal transplant recipients

Permalink

<https://escholarship.org/uc/item/2sr1h0x6>

Journal

Clinical Kidney Journal, 9(3)

ISSN

2048-8505

Authors

Nagy, Kristof

Nagaraju, Shankar Prasad

Rhee, Connie M

et al.

Publication Date

2016-06-01

DOI

10.1093/ckj/sfv156

Copyright Information

This work is made available under the terms of a Creative Commons Attribution-NonCommercial License, available at <https://creativecommons.org/licenses/by-nc/4.0/>

Peer reviewed



CKJ REVIEW

Adipocytokines in renal transplant recipients

Kristof Nagy¹, Shankar Prasad Nagaraju², Connie M. Rhee³, Zoltan Mathe¹ and Miklos Z. Molnar⁴

¹Department of Transplantation and Surgery, Semmelweis University, Budapest, Hungary, ²Department of Nephrology, Kasturba Medical College, Manipal University, Manipal, India, ³Harold Simmons Center for Chronic Disease Research and Epidemiology, Division of Nephrology and Hypertension, University of California Irvine, Orange, CA, USA and ⁴Division of Nephrology, Department of Medicine, University of Tennessee Health Science Center, Memphis, TN, USA

Correspondence to: Miklos Z. Molnar; E-mail: mzmolnar@uthsc.edu

Abstract

In the last two decades, perceptions about the role of body fat have changed. Adipocytes modulate endocrine and immune homeostasis by synthesizing hundreds of hormones, known as adipocytokines. Many studies have been investigating the influences and effects of these adipocytokines and suggest that they are modulated by the nutritional and immunologic milieu. Kidney transplant recipients (KTRs) are a unique and relevant population in which the function of adipocytokines can be examined, given their altered nutritional and immune status and subsequent dysregulation of adipocytokine metabolism. In this review, we summarize the recent findings about four specific adipocytokines and their respective roles in KTRs. We decided to evaluate the most widely described adipocytokines, including leptin, adiponectin, visfatin and resistin. Increasing evidence suggests that these adipocytokines may lead to cardiovascular events and metabolic changes in the general population and may also increase mortality and graft loss rate in KTRs. In addition, we present findings on the interrelationship between serum adipocytokine levels and nutritional and immunologic status, and mechanisms by which adipocytokines modulate morbidity and outcomes in KTRs.

Key words: adiponectin, kidney transplantation, leptin, resistin, visfatin

Introduction of adipocytokines

In healthy individuals, the role of adipocytokines on various outcomes has been widely studied. However, little is known about how impaired kidney function and renal replacement therapy modify these associations [1]. For example, few studies have examined the impact of adipocytokines on long-term outcomes, such as graft loss and death, in kidney transplant recipients (KTRs). In Table 1, we summarize all studies investigating the role of adipocytokines in KTR and chronic kidney disease (CKD) patients.

Leptin was first described as a biomarker of obesity two decades ago; it is a 16-kDa peptide product of the obese gene predominantly secreted by white adipose tissue with the primary function of modulating hunger and satiety [72, 73]. Leptin crosses the blood–brain barrier and inhibits neuropeptide Y neurons. It also activates the sympathetic neuron system in the hypothalamus and increases circulating sympathetic hormone levels [74–76]. Leptin acts upon the Ob leptin receptors and activates tyrosine kinase and intracellular pathways [72, 75].

Leptin's counterpart is adiponectin, which negatively correlates with nutritional parameters, suggesting that higher

Received: November 2, 2015. Accepted: December 18, 2015

© The Author 2016. Published by Oxford University Press on behalf of ERA-EDTA.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Table 1. Studies evaluating adipocytokines in renal transplant recipients and patients with CKD

Author (year)	Number of patients	RRT	Findings	Positive correlations	Negative correlations	Lack of correlations
(A) Leptin						
Agras et al. (2005) [2]	41	Tx.	Leptin seems to increase bone mass.	BMI, BMD, z score of BMD		
Agras et al. (2006) [3]	63	Tx.	Leptin has an effect on lymphoid stem cells.	CD34/7		CD34/7/8/4
Baczowski et al. (2000) [4]	28	Tx.	Imbalance between leptin and body weight also persists after renal transplantation			Cortisol, BMI
El Haggan et al. (2004) [5]	41	Tx.	Pretransplant leptin levels reduced after transplantation	Fat mass, CRP		Dietary intake
Fonseca et al. (2015) [6]	40	Tx.	Leptin levels are independently determined by graft function	Male, DGF		Acute rejection
Kagan et al. (2002) [7]	24	Tx.	Leptin shows correlation with gender, BMI, insulin and cortisol levels	Gender, BMI, cortisol, insulin		
Kayacan et al. (2003) [8]	34	Tx.	Pretransplant leptin levels reduced after transplantation and was not effected by alimentary intake	HOMA, fat mass		
Kokot et al. (1998) [9]	40	Tx	Pretransplant leptin levels reduced after transplantation	BMI	Age	
Kokot et al. (1999) [10]	nd.	Tx	Elevated leptin levels not only modulated by BMI	BMI		
Kovesdy et al. (2010) [11]	978	Tx	Leptin lowers the bone turnover independently from PTH	PTH	vitD	
Landt et al. (1998) [12]	29	Tx	Pretransplant leptin levels reduced after transplantation	BMI		Gender
Lee et al. (2010) [13]	55	Tx.	Leptin correlates with metabolic syndrome	MetSy, waistCX, BMI, fat mass, CRP		
Lee et al. (2014) [14]	74	Tx.	Leptin was positively associated with peripheral arterial stiffness among renal transplant recipients			
Malyszko et al. (2005) [15]	27	Tx.	Leptin is associated with graft function, but not related to BMD and bone metabolism	Body mass/fat, creatinine		Nutrition, BMD
Nicoletto et al. (2012) [16]	32	Tx.	Pretransplant leptin levels reduced after transplantation	Gender, BF, HOMA		
Rafeian-Kopaei et al. (2013) [17]	72	Tx	Leptin levels and duration of kidney transplant shows strong negative correlation	Gender	Duration of kidney Tx	Age, BMI, creatinine
Souza et al. (2007) [18]	32	Tx.	Pretransplant leptin levels reduced after transplantation	HOMA, fat mass		GFR
(B) Adiponectin						
Adamczak et al. (2007) [19]	228	Tx	ADPN levels are higher in RTR than in healthy controls, but lower than in HD patients		BMI, GFR, HOMA-IR	
Adamczak et al. (2011) [20]	88	Tx	Role of ADPN in LVH and atherosclerosis cannot be confirmed			
Alam et al. (2012) [21]	987	Tx	Elevated levels of ADPN increase mortality		GFR, BMI, abd.circ., CRP	CCI
Bayes et al. (2007) [22]	68	Tx	Atorvastatin therapy did not modulate ADPN levels	HDL	HOMA-IR, creatinin	
Bayes et al. (2005) [23]	199	Tx	ADPN were lower, BMI higher in patients who developed NODAT	TNF, BMI, PAPP-A	Insulin	Age, sex
Canas et al. (2012) [24]	157	Tx	ADPN has an inverse association with insulin resistance		HOMA-R, c-IMT	
Chitalia et al. (2010) [25]	43	Tx	ADPN levels do not predict the CV risk in RTR	hsCRP	GFR, BMI, Hgb, waist circumference	BP, smoking, lipids, DM
Chudek et al. (2003) [26]	44	HD/Tx	Kidney plays an important role in biodegradation of ADPN		HOMA-IR	
Chudek et al. (2013) [27]	372	Tx	ACE I/D polymorphism modulates ADPN levels	Female, ACE II genotype	BMI	
Fonseca et al. (2015) [6]	40	Tx	ADPN level is not only modified by early graft function	Male, DGF		Acute rejection
Ho et al. (2015) [28]	69	Tx	ADPN has negative correlation with arterial stiffness		DM, smoking, BMI, waist CX, BP, arterial stiffness	

Table 1. Continued

Author (year)	Number of patients	RRT	Findings	Positive correlations	Negative correlations	Lack of correlations
Idorn et al. (2012) [29]	57	Tx	ADPN level decreases after transplantation and does not predict NODAT		GFR, BMI, insulin	
Kaisar et al. (2009) [30]	137	Tx	Hypoadiponectinemia associated with CVD	HDL, female	BMI, MetSy, IGT, TG, CRP, GFR	
Kang et al. (2012) [31]	575	Tx	ADIPOQ rs1501299 is associated with PTDM in a sex-specific manner			
Kulshrestha et al. (2013) [32]	74	Tx	Patients with metabolic syndrome have lower ADPN levels after transplantation		Clinical events	
Lee et al. (2011) [33]	55	Tx	Body fat mass is an independent predictor of ADPN levels		Fat mass, waist CX, MetSy	
Leibowitz et al. (2013) [34]	35	Tx	ADPN in hypertensive patients is not a predictive factor for CVD			BMI, TG
Malyszko et al. (2005) [35]	82	Tx	ADPN seems to have defense mechanism against endothelial damage	CD146, thrombomodulin, creatinine	BMI, protein Z	
Nicoletto et al. (2013) [36]	270	Tx	TT genotype of ADPN increases the prevalence of NODAT			
Nishimura et al. (2009) [37]	98	Tx	TAC and ARB modulate ADPN levels and posttransplant ADPN levels correlate with NODAT		HOMA-IR	hsCRP
Prasad et al. (2012) [38]	129	Tx	ADPN level lower in South African population		GFR	
Roos et al. (2012) [39]	206	Tx	Pretransplant ADPN level predicts higher risk for graft loss		Graft loss	
Sethna et al. (2009) [40]	33	Tx	Lower ADPN levels associate with higher ambulatory BP		HT	
Shen et al. (2007) [41]	54	Tx	ADPN levels are higher in RTR than in healthy controls, but lower than in HD patients (AdipoR1/2)		HOMA-IR	
Shu et al. (2012) [42]	271	Tx	ADPN level is lower in patients with metabolic syndrome, even with lower GFR	HDL	GFR, MetSy, BMI	
Taherimahmoudi et al. (2010) [43]	67	Tx	ADPN levels are higher in RTR than in healthy controls, but lower than in HD patients. ADPN did not decrease immediately after transplantation			BMI, HOMA-R, GFR
Teplan et al. (2007) [44]	68	Tx	Immunosuppressive therapy could decrease BMI	Leptin	BMI	
Teplan et al. (2008) [45]	140	Tx	In obese RTR, ADMA is increased and ADPN levels are decreased		BMI	
Yilmaz et al. (2005) [46]	27	Tx	ADMA, hsCRP decreasing instantly after transplantation, not like FMD and ADPN (they change later on)			
Yu et al. (2011) [47]	398	Tx	SNP-45/276 of the ADPN gene were significantly associated with an increased risk for NODAT			
(C) Visfatin						
Axelsson et al. (2007) [48]	189	CKD	Elevated with higher CKD stages and may predict mortality	IL-6, hsCRP, VCAM	GFR	
Bessa et al. (2010) [49]	40	CKD	Visfatin is strongly associated with ED and flow-mediated dilatation	ICAM, VCAM, CRP, IL-6	FMD, GFR	
Carrero et al. (2009) [50]	246	CKD	Elevated visfatin is associated with anorexia	PEW	TG, Chol, albumin	BMI, leptin
Eleftheriadis et al. (2013) [51]	33	HD	Visfatin is elevated in HD patients and is connected with decreased demands for rHuEpo	TSAT, Hgb		DM, BMI, IL-6

Table continues

Table 1. Continued

Author (year)	Number of patients	RRT	Findings	Positive correlations	Negative correlations	Lack of correlations
Erten et al. (2008) [52]	31/30	HD/CAPD	In CAPD patients, visfatin is higher than in HD/healthy individuals	IL-6, TNF	Left ventricular diastolic function	Left ventricular mass index
Kato et al. (2009) [53]	68	HD	Visfatin shows a strong association with time spent on HD	Time on HD, hsCRP	Albumin	BMI, adiponectin, body fat
Lu et al. (2013) [54]	173	CKD	Visfatin level is significantly higher in CAD patients and correlates with E-selectin	CAD, hsCRP, BNP, WBC, LDL	GFR, albumin	
Mahmood et al. (2010) [55]	50	CKD	Higher than in healthy controls. No modulation by DM	Proteinuria	GFR	DM
Malyszko et al. (2009) [56]	100	Tx	Higher than in healthy controls	VCAM, CRP, PTH	GFR, albumin	Gender, comorbidities, medication
Malyszko et al. (2010) [57]	75/40	HD/CAPD	Clearance modulated by RRT type, visfatin could be the link between inflammation and adipocytokines	TG, hsCRP, IL-6, TNF, ICAM, VCAM, CD146, HD vin	GFR	
Mu et al. (2011) [58]	117	CKD	Visfatin may play an important role in uremia-related atherosclerosis	ED, hsCRP, TG, LDL	GFR, FMD, HDL	
Yilmaz et al. (2008) [59]	58	Tx	Visfatin levels decline after Tx	ED, hsCRP	GFR, FMD	Medication
Yilmaz et al. (2008) [60]	406	CKD	Visfatin is associated with ED independently from inflammation	ED, hsCRP	GFR, FMD	
(D) Resistin						
Akagun et al. (2014) [61]	69	HD	Increased in HD patients with failed renal allografts	TNF, IL-6, hsCRP	Albumin	
Chung et al. (2012) [62]	100	HD	Low resistin levels independently predict poor hospitalization-free survival			IL-6
Dan et al. (2014) [63]	96	CKD	Serum resistin is higher in the CKD population			
Filippidis et al. (2005) [64]	33	HD	HD does not effect the resistin levels, kidney plays a role in elimination, did not reduce insulin sensitivity			BMI, body fat, HOMA-R, insulin
Kawamura et al. (2010) [65]	3192	CKD	Serum resistin is higher in CKD population.	hsCRP, TG, HOMA	GFR, HDL	BMI
Kaynar et al. (2014) [66]	150	Tx	Resistin is not elevated in Tx patients	PEW		
Kielstein et al. (2003) [67]	30	HD	Resistin levels depend mainly on GFR and its levels do not modulate insulin sensitivity	Homocysteine, age	GFR	Insulin, leptin, BMI, waistCX
Malyszko et al. (2006) [68]	96	Tx	Kidney function is a major determinant of elevating resistin and inflammation	hsCRP, IL-6, RBCc, WBC, VCAM	GFR	
Marouga et al. (2013) [69]	80	CKD	Resistin may be a part of the reverse epidemiology phenomenon of CKD patients	TNF, hsCRP	Alb, GFR, Htc, BMI, leptin, HOMA	HOMA, BMI, cholesterine, leptin
Oltean et al. (2013) [70]	63	DBD	High resistin level in DABD causes delayed graft function			
Spoto et al. (2013) [71]	231	HD	Resistin predicts death depending on ADPN level	hsCRP	ADPN	Leptin, HOMA

abd.circ., abdominal circumference; ACE, angiotensin-converting enzyme; ADMA, asymmetric dimethylarginine; ADPN, adiponectin; Alb, albumin; ARB, angiotensin receptor blocker; BF, body fat; BMD, bone mineral density; BMI, body mass index; BP, blood pressure; CAD, coronary artery disease; CCI, chronic coronary insufficiency; CKD, chronic kidney disease; CRP, C-reactive protein; CVD, cardiovascular disease; DABD, donation after brain death; DGF, delayed graft function; DM, diabetes mellitus; ED, endothelial dysfunction; FMD, fibromuscular dysplasia; GFR, glomerular filtration rate; HD, hemodialysis; HDL, high density lipoprotein; Hgb, hemoglobin; HOMA, homeostasis model assessment; hsCRP, high-sensitivity C-reactive protein; ICAM, intracellular adhesion molecule; IGT, impaired glucose tolerance; IL, interleukin; LDL, low-density lipoprotein; LVH, left ventricular hypertrophy; MetSy, metabolic syndrome; NODAT, new-onset diabetes after transplantation; PAPP-A, pregnancy-associated plasma protein A; PEW, protein energy wasting; PTH, parathyroid hormone; RBC, red blood cell; RRT, renal replacement therapy; RTR, renal transplant recipients; SNP, single nucleotide polymorphism; TSAT, transferrin saturation; TG, triglyceride; TNF, tumor necrosis factor; Tx, transplantation; vitD, vitamin D; VCAM, vascular cell adhesion molecule; waistCX, waist circumference; WBC, white blood cell.

adiponectin levels are observed in healthy individuals. It is a 30-kDa plasma protein synthesized within adipose tissue, and it circulates in the blood as low and high molecular weight isoforms [78]. It activates the downstream signaling pathways through the AdipoR1 receptor, found in kidney cells and skeletal muscle, as well as the AdipoR2 receptor, located in the liver. In the general population, it is considered to be an anti-inflammatory adipocytokine that improves insulin sensitivity and offers cardioprotective benefits [79].

One of the largest adipocytokines is visfatin (52 kDa), which is largely synthesized and secreted by granulocytes and has recently been found to be an insulin sensitizer [56, 80]. Interestingly, endothelial cells within the uremic milieu have also been found to secrete visfatin [81]. While the exact pathways and functions of visfatin are still under investigation, it most likely activates insulin-like receptors and the tyrosine kinase pathway and has insulin-like, pro-inflammatory and anti-apoptotic actions [56, 80].

Resistin is another adipocytokine with metabolic functions and is known to cause insulin resistance. It is a 12.5-kDa cysteine-rich protein that is synthesized in different isoforms by macrophages [82–85]. It acts as a pro-inflammatory factor that increases the production of inflammatory cytokines and elevates the expression of cell adhesion molecules. While its exact biologic functions are still being elucidated, higher resistin levels have been associated with chronic inflammatory diseases such as rheumatoid arthritis and inflammatory bowel disease, and resistin may play a role in the pathophysiology of atherosclerosis and endothelial cell injury [86, 87].

There are other adipocytokines, such as apelin, chemerin and omentin, that will not be discussed in detail in this article. These hormones have been recently discovered, and they are primarily expressed in visceral adipose tissue. For example, apelin synthesis is upregulated by nutritional and inflammatory markers, and it is negatively correlated with endothelial dysfunction [88]. In KTRs, lower apelin levels were associated with underlying cardiovascular disease (CVD) [89]. Moreover, lower apelin levels were observed and were found to be associated with endothelial damage and inflammation [89]. In addition, chemerin affects adipocyte differentiation and insulin signaling, and, despite positive correlations with inflammatory parameters and dyslipidemia, elevated levels have been associated with a survival advantage in dialysis patients [90]. Lastly, omentin's gene locus has been linked to diabetes and it is negatively correlated with inflammatory and nutritional parameters and positively correlated with plasma adiponectin levels, suggesting a protective role [91, 92].

Less is known about adipocytokines' functions and associations with outcomes within KTRs. To better inform the field, we analyzed these associations using one of the largest prevalent KTR cohorts (MINIT-HU) with measured adipocytokines [93–98]. In this study, we collected sociodemographic information, clinical parameters, medical and transplant history and laboratory data from 993 prevalent KTRs who were followed within a single ambulatory transplant clinic at the Department of Transplantation and Surgery, Faculty of Medicine, Semmelweis University in Budapest, Hungary, during the period of 31 December 2006–31 December 2007 [93–98]. Table 2 shows the correlations of the measured adipocytokines in this population, which will be discussed later in this article.

Table 2. Correlations in 988 renal transplant recipients

	Leptin		Adiponectin		Resistin	
	Correlation coefficient	P-value	Correlation coefficient	P-value	Correlation coefficient	P-value
Demographic data						
Age	0.08	0.02	0.08	<0.01	−0.05	0.12
Charlson comorbidity index	0.03	0.43	0.13	<0.01	<0.01	0.89
ESRD time	−0.02	0.58	0.09	<0.01	0.1	<0.01
Kidney-related parameters						
eGFR	−0.22	<0.01	−0.25	<0.01	−0.45	<0.01
Inflammatory markers						
TNF- α	0.06	0.08	0.07	0.04	0.21	<0.01
CRP	0.09	<0.01	0.02	0.55	0.18	<0.01
IL-6	0.09	<0.01	0.08	0.01	0.11	<0.01
Nutritional parameters						
BMI	0.48	<0.01	−0.19	<0.01	−0.11	<0.01
Abdominal circumference	0.31	<0.01	−0.22	<0.01	−0.08	0.02
HDL cholesterol	−0.02	0.54	0.34	<0.01	−0.13	<0.01
LDL cholesterol	0.02	0.44	−0.01	0.70	−0.12	<0.01
Cholesterol	0.10	0.00	0.12	<0.01	−0.1	<0.01
Adipocytokines						
Leptin			−0.04	0.18	0.0472	0.1402
Resistin	0.05	0.14	0.13	<0.01		
Adiponectin	−0.04	0.18			0.13	<0.01
Transplantation-related data						
Cold ischemic time	0.06	0.07	0.07	0.03	0.07	0.04
PRA mean	0.04	0.24	0.05	0.14	0.06	0.06
HLA mismatch	−0.07	0.03	0.02	0.45	−0.04	0.17

ESRD, end-stage renal disease; eGFR, estimated glomerular filtration rate; CRP, C-reactive protein; TNF, tumor necrosis factor; IL, interleukin; BMI, body mass index; HDL, high-density lipoprotein; LDL, low-density lipoprotein; PRA, panel reactive antibodies; HLA, human leukocyte antigen.

Levels of adipocytokines in the serum

Adipocytokine concentrations are typically measured in serum, although normal ranges have not yet been defined in KTRs. Inflammation may potentially induce production, and given that the kidneys play a primary role in many of these hormones' clearance, impaired renal function may lead to higher circulating adipocytokine levels.

Prior data have shown that the kidney function has a modulating effect on leptin levels. Cumin *et al.* [99] demonstrated that leptin clearance is eliminated following bilateral nephrectomy in rats. Additionally, studies also suggest that leptin is filtered through the glomerulus and degraded in the proximal tubule by megalin [73, 100, 101]. In addition, small proportions of leptin have been observed in urine in animals as well as humans [102–104]. These findings indicate that in KTRs, leptin levels may be elevated in the context of impaired renal function and possibly by the uremic milieu [105, 106]. In the non-obese population, levels typically range from 5 to 10 ng/mL, while in obese subjects levels may be 10-fold higher in healthy persons with normal kidney function [107, 108]. In patients with impaired kidney function, hyperleptinemia with levels up to 490 ng/mL have been observed [109]. Some studies suggest that uremia may be associated with higher leptin levels, while others suggest that only nutritional parameters modulate these hormones [110–112].

In contrast, leptin's counterpart, adiponectin, is downregulated in patients with obesity. In diabetic populations, plasma concentrations typically range between 5 and 30 µg/mL, with higher levels observed in those who are female and of older age [113, 114]. Adiponectin is metabolized by the liver, and metabolites are eliminated by the kidneys [115]. Adiponectin molecules may be filtered via the glomerulus, and they are detected in renal arteriole muscle and endothelial cells, as well as within proximal and distal tubules. Proximal tubule cells can synthesize and secrete adiponectin and can therefore be measured in urine [116, 117]. In patients with impaired kidney function, serum adiponectin levels are elevated. The uremic milieu and pro-inflammatory cytokines impair adiponectin synthesis as biofeedback for impaired renal elimination [118].

While the pathways by which visfatin is metabolized and eliminated are still under investigation, it is thought that the kidney plays an important role in its biodegradation. In a study of metabolic syndrome patients, the mean \pm standard deviation of visfatin concentrations was 3.8 ± 8.8 ng/mL [119]. Visfatin levels are typically elevated in patients with CKD, including those undergoing hemodialysis, although a recent large cohort study suggested that levels only correlated with inflammation and insulin resistance and not with kidney function [50, 120]. Moreover, in a study of >3000 patients by Kocelak *et al.* [121], it was found that visfatin levels were not associated with kidney function, although the cohort was largely comprised of elderly patients. In a study of KTRs, higher visfatin concentrations were observed (37 ± 14 ng/mL), although higher levels were also observed in the healthy control group (28 ± 11 ng/mL) [56].

The normal range of resistin has not been defined, although in one study of healthy volunteers by Spoto *et al.* [122] it was found that levels ranged between 1.2 and 29.9 ng/mL. Some studies suggest that resistin functions as a pro-inflammatory cytokine and its serum level may be elevated by inflammatory parameters [123]. In CKD patients, strong correlations between resistin and kidney function have been observed [124]. Furthermore, Malyszko *et al.* [125] demonstrated that hemodialysis patients with residual kidney function had significantly lower resistin levels than those without residual kidney function, supporting the hypothesis that resistin is cleared via the kidney.

Based on these data (Table 1), there does not appear to be a strong, if any, correlation between leptin and glomerular filtration rate, although there does appear to be a correlation between leptin and nutritional parameters. In contrast, adiponectin levels are decreased in obese patients, but higher levels are observed with impaired kidney function. Kidney function, as well as inflammation, also appears to have a substantial impact on visfatin and resistin levels. Table 2 also indicates a correlation between adipocytokines and graft function.

Adipocytokines and nutritional status

Adipocytokines modulate appetite and nutritional status in KTR and CKD patients. Leptin is the primary hormone that regulates satiety and maintains body weight. In healthy subjects, an increase in body weight results in higher leptin levels and subsequent reduction in hunger [126]. Higher leptin levels may also lead to anorexia by binding to melanocortin-4 receptors (MC4-R), and inhibition of peripheral MC4-R can reduce muscle atrophy and improve anorexia via insulin-like growth factor-1 (IGF-1) [127]. As such, higher leptin concentrations due to impaired kidney function may contribute to protein energy wasting (PEW).

Adiponectin is considered to be an anti-inflammatory cytokine and also plays a key role in metabolic syndrome. In experimental studies, higher adiponectin levels have been shown to decrease high-density lipoprotein (HDL) levels, vis-à-vis apolipoprotein A1 and dysregulation of hepatocytes [128]. Elevated adiponectin levels may also lead to malnutrition via weight loss due to increased energy expenditure [129]. It has been observed that malnourished patients on renal replacement therapy have higher adiponectin levels [130].

Visfatin is a pro-inflammatory cytokine synthesized by macrophages embedded in visceral adipose tissue. In non-obese individuals, there is a similar degree of visfatin production in subcutaneous and visceral fat tissue, while in obese patients visceral synthesis is more prominent [131, 132]. At this time, the association between visfatin and nutritional status remains unclear [50, 132–136]. In a study by Carrero *et al.* [50], visfatin was not linked with nutritional parameters in Stage 5 CKD patients, although it was observed that higher serum visfatin levels were associated with increased satiety. Considering that CKD patients have higher visfatin levels, and that visfatin is associated with reduced appetite, it has been suggested that this hormone is part of the PEW syndrome.

Resistin was recently discovered as a hormone that causes insulin resistance in rodents, although these findings are still under investigation in humans. In obese rodents, higher resistin concentrations were observed, which positively correlated with dyslipidemia. It has also been suggested that a reduction of serum resistin levels with antibody neutralization might improve insulin sensitivity [137–139]. Obese individuals have higher resistin levels that persist even with weight gain or loss [140]. It has been suggested that resistin may link PEW to inflammation, since patients with a lower body mass index (BMI) have higher levels of inflammatory factors as well as resistin levels [66, 69].

Adipocytokines in inflammation and their effects on kidney cells

Figure 1 shows the potential mechanisms of adipocytokines on kidney cells. Leptin acts as a pro-inflammatory factor, stimulating the production of pro-inflammatory cytokines [141]. It activates neutrophil and monocyte cell phagocytosis and reactive




Fig. 1. (A) Effects of leptin and adiponectin on different types of cells. (B) Effects of visfatin and resistin on different types of cells. ATII, angiotensin II; CRP, C-reactive protein; ECM, extracellular matrix; eNOS, extracellular nitric oxide synthetase; ET1, endothelin-1; ICAM, intracellular adhesion molecule; IL, interleukin; iNOS, inducible nitric oxide synthetase; MCP, methyl-accepting chemotaxis protein; MMP, matrix metalloproteinase; Na, sodium; NF-kB, nuclear factor κB; NO, nitric oxide; PAI1, plasminogen activator inhibitor-1; RAAS, renin-angiotensin-aldosterone system; RAAS, renin-angiotensin-aldosterone system; TGF, transforming growth factor; TNF, tumor necrosis factor; VCAM, vascular cell adhesion molecule; VEGF, vascular endothelial growth factor; VSMC, vascular smooth muscle cell.

oxygen species through natural killer cells [142, 143]. Leptin also alters the adaptive immune system by activating CD4⁺ T lymphocytes and signaling negatively for CD25⁺ T regulatory cells [144]. It has been hypothesized that the inactivity of regulatory T cells might lead to graft loss since stimulated CD4⁺CD25⁺Foxp3 regulatory T cells prevent rejection, although this has not yet been confirmed in studies [145]. In addition, some *in vitro* studies suggest that stem cell incubation in a leptin-rich environment induces granulocyte formation [146].

The location of adipocytokine receptors bears particular relevance, given their central role in the activation of intracellular pathways. Leptin receptors are expressed in different areas of the kidney, including the inner medulla and vascular structures in the corticomedullary region [147–149]. Within endothelial cells, leptin increases sodium excretion and diuresis by decreasing sodium–potassium channel activity and reducing nitrogen monoxide (NO) production, inhibiting its protective role [150–154]. Within glomerular endothelial, mesangial and vascular smooth muscle

cells, leptin induces proliferation and size increases by activating the mitogen-activated protein kinase (MAPK) pathways [155–157]. Moreover, it induces hypertrophy through different factors causing thickening of the basement membrane. Lastly, leptin also activates the renin–angiotensin–aldosterone system (RAAS) [156, 158, 159]. Observational studies have not found an association between leptin and inflammation in KTRs (Tables 1 and 2).

In contrast to leptin, adiponectin functions as an anti-inflammatory cytokine. Adiponectin enhances anti-inflammatory cytokine synthesis, inhibits angiotensin II-induced inflammation and decreases albuminuria. Recent findings suggest that adiponectin activates nuclear factor κ B (NF- κ B) transcription. NF- κ B is a protein kinase that regulates the immune system through T-cell activity and plays an important role in acute rejection [160]. Furthermore, some but not all data suggest that adiponectin has regulatory effects on T cells. The anti- and pro-inflammatory effects may change according to the molecular isoform of adiponectin [161, 162].

Receptors of adiponectin are also present in kidney cells. Through AdipoR1 it activates the adenosine monophosphate (AMP)-activated protein kinase (AMPK) pathway, inducing protective processes. Within podocytes, adiponectin restores foot processes and inhibits apoptosis, fibrotic effects of TGF-1, nephrin and Nox4 synthesis. Within endothelial cells, it downregulates pro-inflammatory and vasoconstrictor factors and oxidative stress. Furthermore, it elevates endothelial nitric oxide synthase (NOS) concentration and controls anabolic processes [163]. In summary, adiponectin may have protective effects on kidney, although higher levels have not been associated with better graft survival in KTRs [21]. Indeed, adiponectin has demonstrated weak associations with inflammatory parameters in KTRs (Tables 1 and 2).

Resistin's and visfatin's exact receptors are still unknown. Both facilitate the synthesis of inflammatory cytokines within mononuclear and endothelial cells and inhibit the apoptosis of macrophages. They also promote the expression of cell adhesion molecules and chemotaxis [85, 164–168]. Within endothelial cells, they activate NF- κ B transcription factors that augment inducible NOS concentrations and increase the release of endothelin-1 and NO. Visfatin also promotes proliferation and capillary tube formation [81, 168–173]. Moreover, visfatin activates superoxide production, inducing glomerular permeability and RAAS [174–176]. Visfatin and resistin are important factors of inflammation in KTRs (Tables 1 and 2).

It should be noted that strong associations have been observed between visfatin and CD68⁺ macrophages [135]. CD68⁺ macrophage accumulation has been observed in acute rejection, although further investigation is needed to determine this relationship. Specifically for resistin, Akagun et al. [61] demonstrated that patients receiving hemodialysis with failed kidney allografts had significantly higher resistin levels than those without allografts, suggesting that resistin may play a role in chronic inflammation.

In conclusion, higher adipocytokines observed in the context of impaired kidney function may be due to not only reduced clearance, but also to kidney degradation through inflammatory pathways leading to death and graft loss in KTRs.

Adipocytokines role in transplant comorbidities

Pro-inflammatory adipocytokines may lead to hypertension, insulin resistance, CVD and chronic or acute inflammation [177]. In KTRs with impaired allograft function receiving immunosuppression therapy, adipocytokines may contribute to the development of posttransplant bone disease, anemia, cardiovascular

morbidity, new-onset diabetes after transplantation (NODAT) and cancer.

Given that the kidney is one of the major end-organs regulating mineral homeostasis, impaired kidney function may lead to metabolic bone disease. Leptin may in fact accelerate bone metabolism by activating and converting vascular muscle cells into osteoblasts [178]. In epidemiological studies, parathyroid hormone and leptin have shown a negative correlation, and higher leptin levels have been associated with reduced bone turnover [11, 179]. Additionally, leptin causes oversynthesis of fibroblast growth factor 23, subsequently altering the regulation of phosphate homeostasis [180].

Impaired kidney function is the main cause of secondary anemia because of reduced erythropoietin production. Interestingly, leptin receptors may be found in hematopoietic stem cells and higher leptin production has been observed in bone marrow [181]. In epidemiological studies, serum leptin levels are a stronger predictor of erythropoietin sensitivity than body fat and are strongly associated with hemoglobin [182–184]. Erythropoietin stimulating agents (ESAs) administered to CKD patients with higher leptin levels seem to be less effective [182]. A correlation was also observed between serum leptin and erythropoietin resistance in KTRs [96].

Cardiovascular disease is the leading cause of morbidity and mortality among KTRs. Adiponectin presumably has a protective role in the pathogenesis of CVD. Adiponectin receptors are present in endothelial and vascular muscle cells and may reduce oxidative stress, atherosclerosis, fibrosis and proliferation. In the general population, observational studies suggest that lower adiponectin levels are associated with higher cardiovascular mortality [185, 186]. For example, in the LANDMARK 2 study, there was an inverse correlation between adiponectin, inflammation and nutrition in KTRs; furthermore, male patients with lower adiponectin levels had higher CVD risk [30, 187]. However, other data in KTRs have shown positive correlations between higher adiponectin levels and lower survival rates [21]. In a study by Sahin et al. [188], adiponectin levels were significantly higher in patients receiving cyclosporine compared with those on tacrolimus. Malgorzewicz et al. [189] suggested that higher leptin levels in obese patients are a risk factor for CVD and chronic rejection. They also demonstrated that patients with a BMI >25 kg/m² had significantly higher leptin:adiponectin ratios, and those obese patients with higher leptin:adiponectin ratios had a greater risk for cardiovascular events. According to Nicoletto et al. [16], 5 years after transplantation, patients' leptin, insulin resistance and lipid profiles were similar to the pretransplantation period. This metabolic profile is possibly associated with the higher incidence of CVD observed in the late posttransplantation period. Visfatin and resistin play an important role in the development of endothelial dysfunction that may lead to CVD. For example, higher visfatin levels have been observed in patients with coronary artery disease and atherosclerotic plaques [190–192]. Additionally, Weikert et al. [193] demonstrated that patients in the highest resistin quartile had a 2-fold higher risk of myocardial infarction than patients in the lowest quartile. In a prospective study of 2739 participants from the Framingham Offspring Study followed for 6 years, Frankel et al. [194] demonstrated that higher resistin levels were associated with a higher risk of incident heart failure.

Epidemiological data suggest that the prevalence of NODAT ranges from 4 to 25% in KTRs [195]. In one observational study, lower pretransplantation adiponectin levels were an independent predictor of NODAT [196]. Recently, Romanowski et al. [197] found a significant association between the leptin gene allele

and posttransplant diabetes, and it is thought that the 276G/T adiponectin gene polymorphism is also responsible for NODAT. Shu *et al.* [198] also examined various adipocytokines in KTRs with and without metabolic syndrome and found that metabolic syndrome was associated with significantly higher leptin, adiponectin, resistin and visfatin levels. Observational studies have not yet demonstrated an association between visfatin and glucose metabolism, but in animal studies visfatin has been found to have a protective effect on pancreatic beta cells [119, 199–201]. While resistin's role in diabetes is still under investigation, Shu *et al.* [198] have demonstrated strong associations between the presence of the metabolic syndrome and higher serum resistin levels.

Microalbuminuria is an important hallmark of worsening kidney function. Yano *et al.* [202] found that adiponectin is associated with low-grade albuminuria in obese and lean non-diabetic patients. Following multivariable adjustment, urine albumin excretion was significantly higher among patients with lower versus higher adiponectin levels. Several other studies suggest that lower adiponectin levels correlate with albuminuria in obese and diabetic patients [203]. In an observational study including patients with type 1 diabetes mellitus, Prior *et al.* [204] demonstrated strong positive associations between serum adiponectin and plasma antioxidant status. This correlation was most relevant in patients with albuminuria, suggesting an association between higher adiponectin levels, antioxidant status and reduced oxidative cellular burden.

It is well established that immunosuppression therapy heightens the risk of malignant disorders [205]. In some forms of cancer, adiponectin receptors have been found within malignant cells [206]. Recent data have also shown that lower adiponectin levels are associated with a higher incidence of colorectal and prostate cancer [207]. In spite of these findings, locally elevated adiponectin levels have been observed in chondrosarcoma, where adiponectin has been shown to increase the expression of vascular endothelial growth factor [206].

Adipocytokines and outcomes in KTRs

There are limited data regarding the association between adipocytokines and outcomes in patients with end-stage renal disease. Epidemiological studies suggest that leptin declines after kidney transplantation and that it correlates with nutritional parameters. In CKD patients, lower leptin concentrations predict mortality, although there has been only one study examining the association between leptin, mortality and graft loss in KTRs [208, 209]. Similar to other nutritional factors, higher leptin levels are associated with higher mortality risk in patients undergoing hemodialysis [210]. Another notable finding is that leptin deficiency has been associated with prolonged graft survival, suggesting that lean individuals with lower leptin levels may benefit from kidney transplantation [209]. This hypothesis is supported by data from Fonseca *et al.* [6], who found that decreases in leptin levels pre- and posttransplantation were associated with less delayed graft function since higher levels may increase the risk of graft loss; however, these relations require more investigation.

There have been comparatively more observational data assessing the role of adiponectin in KTRs. Only a few articles have examined the association between adiponectin and mortality. Similar to studies of adiponectin and mortality in hemodialysis patients [211], we found that higher levels of adiponectin were associated with increased mortality, but not graft loss, despite the protective effects of adiponectin [21]. Roos *et al.* [39] found that

low pretransplant adiponectin levels predicted risk of graft loss, which may be due to atherogenic pathways. It should also be highlighted that, according to Fonseca *et al.* [6], early graft function is not a primary modifier of adiponectin levels since there were no significant differences in serum adiponectin levels among those with preserved versus impaired graft function.

A number of studies have examined visfatin in patients with impaired renal function (Table 1). However, only two articles have investigated its association with outcomes in KTRs, suggesting that levels decline after transplantation and correlate with the parameters of endothelial dysfunction [56, 59]. Axelsson *et al.* [48] demonstrated that higher visfatin levels were associated with higher mortality in CKD.

Numerous studies have examined the role of resistin in mortality among patients with impaired kidney function, although few in the KTR population (Table 1). Spoto *et al.* [122] analyzed the association between resistin and all-cause and cardiovascular mortality among KTRs and found significant associations between serum resistin and death, although serum adiponectin appeared to be a potent effect modifier of this relationship. In contrast, Chung *et al.* [212] found that lower serum resistin concentrations were associated with lower survival rates in patients on hemodialysis. Our results also suggest that higher resistin levels have a dose-dependent relationship with risk of graft loss and mortality even after multivariable adjustment in KTRs [213]. We found that for every 10 ng/mL elevation of serum resistin level there is a 30% increased mortality risk and 73% increased risk of graft loss [213].

One of the major limitations of these collective studies is that they have typically examined adipocytokine concentrations at a single point in time; therefore, they cannot examine changes in levels over time. In addition, most were of small sample size and were unable to examine cause-specific mortality. At this time, further studies of adipocytokines are needed in the field of transplantation.

Conclusion

Adipocytokine levels are often elevated in patients with kidney disease, including KTRs with impaired graft function. These cytokines have been associated with endothelial dysfunction, inflammation and malnutrition, which may contribute to the degradation of kidney tissue. Higher leptin levels might play a role in rejection through the regulation of macrophages. Adiponectin also has important regulatory effects on the adaptive immune system and may have a role in rejection. In addition, few studies have shown that these adipocytokines are independent predictors of survival in KTRs. Further studies are needed to understand the role of adipocytokines in the nutritional status, graft function and survival of KTRs.

Conflict of interest statement

None declared.

References

1. Tilg H, Moschen AR. Adipocytokines: mediators linking adipose tissue, inflammation and immunity. *Nat Rev Immunol* 2006; 6: 772–783
2. Agras PI, Baskin E, Saatci U *et al.* Relationship between leptin and bone mineral density in renal transplant recipients. *Transplant Proc* 2005; 37: 3106–3108

3. Agras PI, Saatci U, Baskin E et al. Hyperleptinemia and its relation with peripheral C34(+)CD7(+) stem cells in renal transplant recipients. *Transpl Immunol* 2006; 15: 241–245
4. Baczkowska T, Soin J, Soluch L et al. The role of leptin in body mass index increase in renal allograft recipients. *Transplant Proc* 2000; 32: 1331–1332
5. El Haggan W, Chauveau P, Barthe N et al. Serum leptin, body fat, and nutritional markers during the six months post-kidney transplantation. *Metabolism* 2004; 53: 614–619
6. Fonseca I, Oliveira JC, Santos J et al. Leptin and adiponectin during the first week after kidney transplantation: biomarkers of graft dysfunction? *Metabolism* 2015; 64: 202–207
7. Kagan A, Haran N, Leschinsky L et al. Serum concentrations of leptin in heart, liver and kidney transplant recipients. *Isr Med Assoc J* 2002; 4: 213–217
8. Kayacan SM, Yildiz A, Kazancioglu R et al. The changes in serum leptin, body fat mass and insulin resistance after renal transplantation. *Clin Transplant* 2003; 17: 63–68
9. Kokot F, Adamczak M, Wiecek A. Plasma leptin concentration in kidney transplant patients during the early post-transplant period. *Nephrol Dial Transplant* 1998; 13: 2276–2280
10. Kokot F, Adamczak M, Wiecek A et al. Plasma immunoreactive leptin and neuropeptide Y levels in kidney transplant patients. *Am J Nephrol* 1999; 19: 28–33
11. Kovesdy CP, Molnar MZ, Czira ME et al. Associations between serum leptin level and bone turnover in kidney transplant recipients. *Clin J Am Soc Nephrol* 2010; 5: 2297–2304
12. Landt M, Brennan DC, Parvin CA et al. Hyperleptinaemia of end-stage renal disease is corrected by renal transplantation. *Nephrol Dial Transplant* 1998; 13: 2271–2275
13. Lee MC, Lee CJ, Ho GJ et al. Hyperleptinemia positively correlated with metabolic syndrome in renal transplant recipients. *Clin Transplant* 2010; 24: E124–E1249
14. Lee MC, Chen YC, Ho GJ et al. Serum leptin levels positively correlate with peripheral arterial stiffness in kidney transplantation patients. *Transplant Proc* 2014; 46: 353–358
15. Malyszko J, Malyszko JS, Pawlak K et al. Correlations between leptin, body composition, bone mineral density, and bone metabolism in kidney transplant recipients. *Transplant Proc* 2005; 37: 2151–2153
16. Nicoletto BB, Souza GC, Goncalves LF et al. Leptin, insulin resistance, and metabolic changes 5 years after renal transplantation. *J Ren Nutr* 2012; 22: 440–449
17. Rafeian-Kopaei M, Nasri H. Serum leptin in renal transplant patients. *JRIP* 2013; 2: 55–57
18. Souza GC, Costa CA, Goncalves LF et al. Leptin serum levels in the first year post-renal transplantation. *Transplant Proc* 2007; 39: 439–440
19. Adamczak M, Szotowska M, Chudek J et al. Plasma adiponectin concentration in patients after successful kidney transplantation—a single-center, observational study. *Clin Nephrol* 2007; 67: 381–390
20. Adamczak M, Blach A, Kolonko A et al. Plasma adiponectin concentration and left ventricular hypertrophy in kidney transplant patients. *Clin Transplant* 2011; 25: 561–568
21. Alam A, Molnar MZ, Czira ME et al. Serum adiponectin levels and mortality after kidney transplantation. *Clin J Am Soc Nephrol* 2013; 8: 460–467
22. Bayes B, Granada ML, Pastor MC et al. Obesity, adiponectin and inflammation as predictors of new-onset diabetes mellitus after kidney transplantation. *Am J Transplant* 2007; 7: 416–422
23. Bayes B, Granada ML, Lauzurica R et al. Effect of low doses of atorvastatin on adiponectin, glucose homeostasis, and clinical inflammatory markers in kidney transplant recipients. *Transplant Proc* 2005; 37: 3808–3812
24. Canas L, Bayes B, Granada ML et al. Is adiponectin a marker of preclinical atherosclerosis in kidney transplantation? *Clin Transplant* 2012; 26: 259–266
25. Chitalia N, Raja RB, Bhandara T et al. Serum adiponectin and cardiovascular risk in chronic kidney disease and kidney transplantation. *J Nephrol* 2010; 23: 77–84
26. Chudek J, Adamczak M, Karkoszka H et al. Plasma adiponectin concentration before and after successful kidney transplantation. *Transplant Proc* 2003; 35: 2186–2189
27. Chudek J, Szotowska M, Karkoszka H et al. Genotypes of renin-angiotensin system and plasma adiponectin concentration in kidney transplant patients. *Ann Transplant* 2013; 18: 593–603
28. Ho GJ, Lee MC, Lee CJ et al. Hypoadiponectinemia correlates with arterial stiffness in kidney transplantation patients. *Clin Exp Nephrol* 2015; 19: 534–541
29. Idorn T, Hornum M, Bjerre M et al. Plasma adiponectin before and after kidney transplantation. *Transpl Int* 2012; 25: 1194–1203
30. Kaiser MO, Armstrong K, Hawley C et al. Adiponectin is associated with cardiovascular disease in male renal transplant recipients: baseline results from the LANDMARK 2 study. *BMC Nephrol* 2009; 10: 29
31. Kang ES, Magkos F, Kim BS et al. Variants of the adiponectin and adiponectin receptor-1 genes and posttransplantation diabetes mellitus in renal allograft recipients. *J Clin Endocrinol Metab* 2012; 97: E129–E135
32. Kulshrestha S, Ojo AO, Luan FL. Metabolic syndrome, vitamin D deficiency and hypoadiponectinemia among nondiabetic patients early after kidney transplantation. *Am J Nephrol* 2013; 37: 399–404
33. Lee MC, Lee CJ, Chou KC et al. Hypoadiponectinemia correlates with metabolic syndrome in kidney transplantation patients. *Transplant Proc* 2011; 43: 2601–2605
34. Leibowitz A, Peleg E, Ben-David A et al. Normal adiponectin levels in kidney transplant patients with hypertension. *Clin Transplant* 2013; 27: 562–566
35. Malyszko J, Malyszko JS, Brzosko S et al. Markers of endothelial cell activation/injury: CD146 and thrombomodulin are related to adiponectin in kidney allograft recipients. *Am J Nephrol* 2005; 25: 203–210
36. Nicoletto BB, Souza GC, Fonseca NK et al. Association between 276G/T adiponectin gene polymorphism and new-onset diabetes after kidney transplantation. *Transplantation* 2013; 96: 1059–1064
37. Nishimura K, Kishikawa H, Kato T et al. Tacrolimus and angiotensin receptor blockers associated with changes in serum adiponectin level in new-onset diabetes after renal transplantation: single-center cross-sectional analysis. *Transpl Int* 2009; 22: 694–701
38. Prasad GV, Vorobeichik L, Nash MM et al. Lower total and percent of high-molecular-weight adiponectin concentration in South Asian kidney transplant recipients. *Clin Kidney J* 2012; 5: 124–129
39. Roos M, Baumann M, Liu D et al. Low pre-transplant adiponectin multimers are associated with adverse allograft outcomes in kidney transplant recipients a 3-year prospective study. *Regul Pept* 2012; 178: 11–15
40. Sethna CB, Leonard MB, Gallagher PR et al. Serum adiponectin levels and ambulatory blood pressure monitoring in pediatric renal transplant recipients. *Transplantation* 2009; 88: 1030–1037

41. Shen YY, Charlesworth JA, Kelly JJ et al. The effect of renal transplantation on adiponectin and its isoforms and receptors. *Metabolism* 2007; 56: 1201–1208
42. Shu KH, Tsai IC, Ho HC et al. Serum adiponectin levels in renal transplant recipients with and without metabolic syndrome. *Transplant Proc* 2012; 44: 676–679
43. Taherimahmoudi M, Ahmadi H, Mehrsai A et al. Plasma adiponectin concentration and insulin resistance: role of successful kidney transplantation. *Transplant Proc* 2010; 42: 797–800
44. Teplan V, Schuck O, Stollova M et al. Obesity and adiponectin after kidney transplantation. *Acta Physiol Hung* 2007; 94: 149–157
45. Teplan V, Schuck O, Racek J et al. Asymmetric dimethylarginine and adiponectin after renal transplantation: role of obesity. *J Ren Nutr* 2008; 18: 154–157
46. Yilmaz MI, Saglam M, Caglar K et al. Endothelial functions improve with decrease in asymmetric dimethylarginine (ADMA) levels after renal transplantation. *Transplantation* 2005; 80: 1660–1666
47. Yu AR, Xin HW, Wu XC et al. Adiponectin gene polymorphisms are associated with posttransplantation diabetes mellitus in Chinese renal allograft recipients. *Transplant Proc* 2011; 43: 1607–1611
48. Axelsson J, Witasp A, Carrero JJ et al. Circulating levels of visfatin/pre-B-cell colony-enhancing factor 1 in relation to genotype, GFR, body composition, and survival in patients with CKD. *Am J Kidney Dis* 2007; 49: 237–244
49. Bessa SS, Hamdy SM, El-Sheikh RG. Serum visfatin as a non-traditional biomarker of endothelial dysfunction in chronic kidney disease: an Egyptian study. *Eur J Intern Med* 2010; 21: 530–535
50. Carrero JJ, Witasp A, Stenvinkel P et al. Visfatin is increased in chronic kidney disease patients with poor appetite and correlates negatively with fasting serum amino acids and triglyceride levels. *Nephrol Dial Transplant* 2010; 25: 901–906
51. Eleftheriadis T, Pissas G, Remoundou M et al. Increased visfatin in hemodialysis patients is associated with decreased demands for recombinant human erythropoietin. *Ren Fail* 2013; 35: 1399–1403
52. Erten Y, Ebinc FA, Ebinc H et al. The relationship of visfatin levels to inflammatory cytokines and left ventricular hypertrophy in hemodialysis and continuous ambulatory peritoneal dialysis patients. *Ren Fail* 2008; 30: 617–623
53. Kato A, Odamaki M, Ishida J et al. Relationship between serum pre-B cell colony-enhancing factor/visfatin and atherosclerotic parameters in chronic hemodialysis patients. *Am J Nephrol* 2009; 29: 31–35
54. Lu YC, Hsu CC, Yu TH et al. Association between visfatin levels and coronary artery disease in patients with chronic kidney disease. *Iran J Kidney Dis* 2013; 7: 446–452
55. Mahmood N, Junejo AM, Jamal Q et al. Association of visfatin with chronic kidney disease in a cohort of patients with and without diabetes. *J Pak Med Assoc* 2010; 60: 922–926
56. Malyszko J, Malyszko JS, Mysliwiec M. Visfatin, a new adipocytokine, is predominantly related to inflammation/endothelial damage in kidney allograft recipients. *Transplant Proc* 2009; 41: 150–153
57. Malyszko J, Malyszko JS, Mysliwiec M. Visfatin and endothelial function in dialyzed patients. *Nephrology* 2010; 15: 190–196
58. Mu J, Feng B, Ye Z et al. Visfatin is related to lipid dysregulation, endothelial dysfunction and atherosclerosis in patients with chronic kidney disease. *J Nephrol* 2011; 24: 177–184
59. Yilmaz MI, Saglam M, Carrero JJ et al. Normalization of endothelial dysfunction following renal transplantation is accompanied by a reduction of circulating visfatin/NAMPT. A novel marker of endothelial damage? *Clin Transplant* 2009; 23: 241–248
60. Yilmaz MI, Saglam M, Carrero JJ et al. Serum visfatin concentration and endothelial dysfunction in chronic kidney disease. *Nephrol Dial Transplant* 2008; 23: 959–965
61. Akagun T, Caliskan Y, Yazici H et al. Elevated resistin levels are associated with inflammation in hemodialysis patients with failed renal allografts. *Int J Artif Organs* 2014; 37: 358–363
62. Chung W, Jung ES, Shin D et al. Low resistin level is associated with poor hospitalization-free survival in hemodialysis patients. *J Korean Med Sci* 2012; 27: 377–381
63. Dan S, Aditya P, Banerjee P et al. Effect of chronic kidney disease on serum resistin level. *Niger J Clin Pract* 2014; 17: 735–738
64. Filippidis G, Liakopoulos V, Mertens PR et al. Resistin serum levels are increased but not correlated with insulin resistance in chronic hemodialysis patients. *Blood Purif* 2005; 23: 421–428
65. Kawamura R, Doi Y, Osawa H et al. Circulating resistin is increased with decreasing renal function in a general Japanese population: the Hisayama Study. *Nephrol Dial Transplant* 2010; 25: 3236–3240
66. Kaynar K, Kural BV, Ulusoy S et al. Is there any interaction of resistin and adiponectin levels with protein-energy wasting among patients with chronic kidney disease. *Hemodial Int* 2014; 18: 153–162
67. Kielstein JT, Becker B, Graf S et al. Increased resistin blood levels are not associated with insulin resistance in patients with renal disease. *Am J Kidney Dis* 2003; 42: 62–66
68. Malyszko J, Malyszko JS, Pawlak K et al. Resistin, a new adipokine, is related to inflammation and renal function in kidney allograft recipients. *Transplant Proc* 2006; 38: 3434–3436
69. Marouga A, Dalamaga M, Kastania AN et al. Correlates of serum resistin in elderly, non-diabetic patients with chronic kidney disease. *Clin Lab* 2012; 59: 1121–1128
70. Oltean S, Pullerits R, Floden A et al. Increased resistin in brain dead organ donors is associated with delayed graft function after kidney transplantation. *J Transl Med* 2013; 11: 233
71. Spoto B, Mattace-Raso F, Sijbrands E et al. Resistin and all-cause and cardiovascular mortality: effect modification by adiponectin in end-stage kidney disease patients. *Nephrol Dial Transplant* 2013; 28(Suppl 4): iv181–iv187
72. Attoub S, Noe V, Pirola L et al. Leptin promotes invasiveness of kidney and colonic epithelial cells via phosphoinositide 3-kinase-, Rho-, and Rac-dependent signaling pathways. *FASEB J* 2000; 14: 2329–2338
73. Hama H, Saito A, Takeda T et al. Evidence indicating that renal tubular metabolism of leptin is mediated by megalin but not by the leptin receptors. *Endocrinology* 2004; 145: 3935–3940
74. Yang R, Barouch LA. Leptin signaling and obesity cardiovascular consequences. *Circ Res* 2007; 101: 545–559
75. Bjørnbæk C, Buchholz RM, Davis SM et al. Divergent roles of SHP-2 in ERK activation by leptin receptors. *J Biol Chem* 2001; 276: 4747–4755
76. Simonds SE, Cowley MA, Enriori PJ. Leptin increasing sympathetic nerve outflow in obesity: a cure for obesity or a potential contributor to metabolic syndrome? *Adipocyte* 2012; 1: 177–181
77. Dhillon SS, Belsham DD. Leptin differentially regulates NPY secretion in hypothalamic cell lines through distinct intracellular signal transduction pathways. *Regul Pept* 2011; 167: 192–200

78. Shen YY, Hughes JT, Charlesworth JA et al. Adiponectin is present in the urine in its native conformation, and specifically reduces the secretion of MCP-1 by proximal tubular cells. *Nephrology (Carlton)* 2008; 13: 405–410
79. Liu M, Liu F. Regulation of adiponectin multimerization, signaling and function. *Best Pract Res Clin Endocrinol Metab* 2014; 28: 25–31
80. Stastny J, Bienertova-Vasku J, Vasku A. Visfatin and its role in obesity development. *Diabetes Metab Syndr* 2012; 6: 120–124
81. Romacho T, Villalobos LA, Cercas E et al. Visfatin as a novel mediator released by inflamed human endothelial cells. *PLoS One* 2013; 8: e78283
82. Aruna B, Ghosh S, Singh AK et al. Human recombinant resistin protein displays a tendency to aggregate by forming intermolecular disulfide linkages. *Biochemistry* 2003; 42: 10554–10559
83. Patel SD, Rajala MW, Rossetti L et al. Disulfide-dependent multimeric assembly of resistin family hormones. *Science* 2004; 304: 1154–1158
84. Kusminski CM, McTernan PG, Kumar S. Role of resistin in obesity, insulin resistance and type II diabetes. *Clin Sci (Lond)* 2005; 109: 243–256
85. Reilly MP, Lehrke M, Wolfe ML et al. Resistin is an inflammatory marker of atherosclerosis in humans. *Circulation* 2005; 111: 932–939
86. Fadda SM, Gamal SM, Elsaid NY et al. Resistin in inflammatory and degenerative rheumatologic diseases. Relationship between resistin and rheumatoid arthritis disease progression. *Z Rheumatol* 2013; 72: 594–600
87. Karmiris K, Koutroubakis IE, Xidakis C et al. Circulating levels of leptin, adiponectin, resistin, and ghrelin in inflammatory bowel disease. *Inflamm Bowel Dis* 2006; 12: 100–105
88. Fujie S, Sato K, Miyamoto-Mikami E et al. Reduction of arterial stiffness by exercise training is associated with increasing plasma apelin level in middle-aged and older adults. *PLoS One* 2014; 9: e93545
89. Malyszko J, Malyszko JS, Pawlak K et al. Apelin, a novel adipocytokine, in relation to endothelial function and inflammation in kidney allograft recipients. *Transplant Proc* 2008; 40: 3466–3469
90. Yamamoto T, Qureshi AR, Anderstam B et al. Clinical importance of an elevated circulating chemerin level in incident dialysis patients. *Nephrol Dial Transplant* 2010; 25: 4017–4023
91. Ruster C, Wolf G. Adipokines promote chronic kidney disease. *Nephrol Dial Transplant* 2013; 28(Suppl 4): iv8–iv14
92. Miyamoto S, Sharma K. Adipokines protecting CKD. *Nephrol Dial Transplant* 2013; 28(Suppl 4): iv15–iv22
93. Molnar MZ, Czira ME, Rudas A et al. Association of the malnutrition-inflammation score with clinical outcomes in kidney transplant recipients. *Am J Kidney Dis* 2011; 58: 101–108
94. Molnar MZ, Tabak AG, Alam A et al. Serum erythropoietin level and mortality in kidney transplant recipients. *Clin J Am Soc Nephrol* 2011; 6: 2879–2886
95. Kovesdy CP, Mucsi I, Czira ME et al. Association of serum phosphorus level with anemia in kidney transplant recipients. *Transplantation* 2011; 91: 875–882
96. Molnar MZ, Czira ME, Rudas A et al. Association between the malnutrition-inflammation score and post-transplant anaemia. *Nephrol Dial Transplant* 2011; 26: 2000–2006
97. Kovesdy CP, Czira ME, Rudas A et al. Body mass index, waist circumference and mortality in kidney transplant recipients. *Am J Transplant* 2010; 10: 2644–2651
98. Molnar MZ, Keszei A, Czira ME et al. Evaluation of the malnutrition-inflammation score in kidney transplant recipients. *Am J Kidney Dis* 2010; 56: 102–111
99. Cumin F, Baum HP, Levens N. Leptin is cleared from the circulation primarily by the kidney. *Int J Obes Relat Metab Disord* 1996; 20: 1120–1126
100. Zou Z, Chung B, Nguyen T et al. Linking receptor-mediated endocytosis and cell signaling evidence for regulated intramembrane proteolysis of megalin in proximal tubule. *J Biol Chem* 2004; 279: 34302–34310
101. Shah M, Bateria OY, Taupin V. ARH directs megalin to the endocytic recycling compartment to regulate its proteolysis and gene expression. *J Cell Biol* 2013; 202: 113–127
102. Meyer C, Robson D, Rackovsky N. Role of the kidney in human leptin metabolism. *Am J Physiol* 1997; 273: E903–E907
103. Cumin F, Baum HP, Levens N. Mechanism of leptin removal from the circulation by the kidney. *J Endocrinol* 1997; 155: 577–585
104. Lönnqvist F, Nordfors L, Jansson M et al. Leptin secretion from adipose tissue in women. Relationship to plasma levels and gene expression. *J Clin Invest* 1997; 99: 2398–2404
105. Aminzadeh MA, Pahl MV, Barton CH et al. Human uraemic plasma stimulates release of leptin and uptake of tumour necrosis factor- α in visceral adipocytes. *Nephrol Dial Transplant* 2009; 24: 3626–3631
106. Kalbacher E, Koppe L, Zarrouki B et al. Human uremic plasma and not urea induces exuberant secretion of leptin in 3T3-L1 adipocytes. *J Ren Nutr* 2011; 21: 72–75
107. Boden G, Chen X, Mozzoli M et al. Effect of fasting on serum leptin in normal human subjects. *J Clin Endocrinol Metab* 1996; 81: 3419–3423
108. Garibotto G, Russo R, Franceschini R et al. Inter-organ leptin exchange in humans. *Biochem Biophys Res Commun* 1998; 247: 504–509
109. Dagogo-Jack S, Ovalle F, Landt M. Hyperleptinemia in patients with end-stage renal disease undergoing continuous ambulatory peritoneal dialysis. *Perit Dial Int* 1998; 18: 34–40
110. Silva MI, Vale BS, Lemos CC et al. Body adiposity index assess body fat with high accuracy in nondialyzed chronic kidney disease patients. *Obesity (Silver Spring)* 2013; 21: 546–552
111. Ho KJ, Xue H, Mauro CR et al. Impact of uremia on human adipose tissue phenotype. *J Surg Res* 2013; 179: 175–182
112. Ruhl CE, Harris TB, Ding J et al. Body mass index and serum leptin concentration independently estimate percentage body fat in older adults. *Am J Clin Nutr* 2007; 85: 1121–1126
113. Sharma K, Ramachandrarao S, Qiu G et al. Adiponectin regulates albuminuria and podocyte function in mice. *J Clin Invest* 2008; 118: 1645–1656
114. Obata Y, Yamada Y, Takahi Y et al. Relationship between serum adiponectin levels and age in healthy subjects and patients with type 2 diabetes. *Clin Endocrinol (Oxf)* 2013; 79: 204–210
115. Tacke F, Wustefeld T, Horn R et al. High adiponectin in chronic liver disease and cholestasis suggests biliary route of adiponectin excretion in vivo. *J Hepatol* 2005; 42: 666–673
116. Perri A, Vizza D, Lofaro D et al. Adiponectin is expressed and secreted by renal tubular epithelial cells. *J Nephrol* 2013; 26: 1049–1054
117. von Eynatten M, Liu D, Hock C et al. Urinary adiponectin excretion a novel marker for vascular damage in type 2 diabetes. *Diabetes* 2009; 58: 2093–2099
118. Bruun JM, Lihn AS, Verdich C et al. Regulation of adiponectin by adipose tissue-derived cytokines: in vivo and in vitro investigations in humans. *Am J Physiol Endocrinol Metab* 2003; 285: E527–E533
119. Chedraui P, Perez-Lopez FR, Escobar GS et al. Circulating leptin, resistin, adiponectin, visfatin, adipisin and ghrelin levels

- and insulin resistance in postmenopausal women with and without the metabolic syndrome. *Maturitas* 2014; 79: 86–90
120. Nusken KD, Petrasch M, Rauh M et al. Active visfatin is elevated in serum of maintenance haemodialysis patients and correlates inversely with circulating HDL cholesterol. *Nephrol Dial Transplant* 2009; 24: 2832–2838
 121. Kocelak P, Olszanecka-Glinianowicz M, Owczarek A et al. Plasma visfatin/nicotinamide phosphoribosyltransferase (visfatin/NAMPT) concentration is not related to kidney function in elderly subjects. *Clin Chem Lab Med* 2015; 53: 793–799
 122. Spoto B, Mattace-Raso F, Sijbrands E et al. Resistin and all-cause and cardiovascular mortality: effect modification by adiponectin in end-stage kidney disease patients. *Nephrol Dial Transplant* 2013; 28(Suppl 4): iv181–iv187
 123. Fargnoli JL, Sun Q, Olenczuk D et al. Resistin is associated with biomarkers of inflammation while total and high-molecular weight adiponectin are associated with biomarkers of inflammation, insulin resistance, and endothelial function. *Eur J Endocrinol* 2010; 162: 281–288
 124. Axelsson J, Bergsten A, Qureshi AR et al. Elevated resistin levels in chronic kidney disease are associated with decreased glomerular filtration rate and inflammation, but not with insulin resistance. *Kidney Int* 2006; 69: 596–604
 125. Malyszko J, Malyszko JS, Kozminski P et al. Elevated resistin is related to inflammation and residual renal function in haemodialysed patients. *Nephrology (Carlton)* 2007; 12: 246–253
 126. Friedman JM. Leptin and the regulation of body weight. *Keio J Med* 2011; 60: 1–9
 127. Scholze A, Tepel M. Role of leptin in reverse epidemiology in chronic kidney disease. *Semin Dial* 2007; 20: 534–538
 128. Liu Q, Yuan B, Lo KA et al. Adiponectin regulates expression of hepatic genes critical for glucose and lipid metabolism. *Proc Natl Acad Sci USA* 2012; 109: 14568–14573
 129. Qi Y, Takahashi N, Hileman SM et al. Adiponectin acts in the brain to decrease body weight. *Nat Med* 2004; 10: 524–529
 130. Ekramzadeh M, Sohrabi Z, Salehi M et al. Adiponectin as a novel indicator of malnutrition and inflammation in hemodialysis patients. *Iran J Kidney Dis* 2013; 7: 304–308
 131. Curat CA, Wegner V, Sengenès C et al. Macrophages in human visceral adipose tissue: increased accumulation in obesity and a source of resistin and visfatin. *Diabetologia* 2006; 49: 744–747
 132. Berndt J, Kloting N, Kralisch S et al. Plasma visfatin concentrations and fat depot-specific mRNA expression in humans. *Diabetes* 2005; 54: 2911–2916
 133. Pagano C, Pilon C, Olivieri M et al. Reduced plasma visfatin/pre-B cell colony-enhancing factor in obesity is not related to insulin resistance in humans. *J Clin Endocrinol Metab* 2006; 91: 3165–3170
 134. Varma V, Yao-Borengasser A, Rasouli N et al. Human visfatin expression: relationship to insulin sensitivity, intramyocellular lipids, and inflammation. *J Clin Endocrinol Metab* 2007; 92: 666–672
 135. Chang YC, Chang TJ, Lee WJ et al. The relationship of visfatin/pre-B-cell colony-enhancing factor/nicotinamide phosphoribosyltransferase in adipose tissue with inflammation, insulin resistance, and plasma lipids. *Metabolism* 2010; 59: 93–99
 136. Wang P, van Greevenbroek MM, Bouwman FG et al. The circulating PBEF/NAMPT/visfatin level is associated with a beneficial blood lipid profile. *Pflugers Arch* 2007; 454: 971–976
 137. Rajala MW, Qi Y, Patel HR et al. Regulation of resistin expression and circulating levels in obesity, diabetes, and fasting. *Diabetes* 2004; 53: 1671–1679
 138. Sato N, Kobayashi K, Inoguchi T et al. Adenovirus-mediated high expression of resistin causes dyslipidemia in mice. *Endocrinology* 2005; 146: 273–279
 139. Steppan CM, Bailey ST, Bhat S et al. The hormone resistin links obesity to diabetes. *Nature* 2001; 409: 307–312
 140. Park HK, Ahima RS. Resistin in rodents and humans. *Diabetes Metab J* 2013; 37: 404–414
 141. Loffreda S, Yang SQ, Lin HZ et al. Leptin regulates proinflammatory immune responses. *FASEB J* 1998; 12: 57–65
 142. Lord GM, Matarese G, Howard JK et al. Leptin modulates the T-cell immune response and reverses starvation-induced immunosuppression. *Nature* 1998; 394: 897–901
 143. Zhao Y, Sun R, You L et al. Expression of leptin receptors and response to leptin stimulation of human natural killer cell lines. *Biochem Biophys Res Commun* 2003; 300: 247–252
 144. De Rosa V, Procaccini C, Cali G et al. A key role of leptin in the control of regulatory T cell proliferation. *Immunity* 2007; 26: 241–255
 145. Joffre O, Santolaria T, Calise D et al. Prevention of acute and chronic allograft rejection with CD4⁺CD25⁺Foxp3⁺ regulatory T lymphocytes. *Nat Med* 2008; 14: 88–92
 146. Umemoto Y, Tsuji K, Yang FC et al. Leptin stimulates the proliferation of murine myelocytic and primitive hematopoietic progenitor cells. *Blood* 1997; 90: 3438–3443
 147. Martínez-Anso E, Lostao MP, Martínez JA. Immunohistochemical localization of leptin in rat kidney. *Kidney Int* 1999; 55: 1129–1130
 148. Elis Yildiz S, Deprem T, Karadag Sari E et al. Immunohistochemical distribution of leptin in kidney tissues of melatonin treated diabetic rats. *Biotech Histochem* 2015; 90: 270–277
 149. Serradeil-Le Gal C, Raufaste D, Brossard G et al. Characterization and localization of leptin receptors in the rat kidney. *FEBS Lett* 1997; 404: 185–191
 150. Bełtowski J, Wójcicka G, Górny D et al. Human leptin administered intraperitoneally stimulates natriuresis and decreases renal medullary Na⁺, K⁺-ATPase activity in the rat—impaired effect in dietary-induced obesity. *Med Sci Monit* 2002; 8: BR221–BR229
 151. Jackson EK, Li P. Human leptin has natriuretic activity in the rat. *Am J Physiol* 1997; 272: F333–F338
 152. Vecchione C, Maffei A, Colella S et al. Leptin effect on endothelial nitric oxide is mediated through Akt-endothelial nitric oxide synthase phosphorylation pathway. *Diabetes* 2002; 51: 168–173
 153. Ortiz PA, Garvin JL. Role of nitric oxide in the regulation of nephron transport. *Am J Physiol Ren* 2002; 282: F777–F784
 154. Bełtowski J, Jamroz-Wisniewska A, Borkowska E et al. Up-regulation of renal Na⁺, K⁺-ATPase: the possible novel mechanism of leptin-induced hypertension. *Pol J Pharmacol* 2004; 56: 213–222
 155. Lee MP, Orlov D, Sweeney G. Leptin induces rat glomerular mesangial cell hypertrophy, but does not regulate hyperplasia or apoptosis. *Int J Obes (Lond)* 2005; 29: 1395–1401
 156. Zeidan A, Purdham DM, Rajapurohitam V et al. Leptin induces vascular smooth muscle cell hypertrophy through angiotensin II- and endothelin-1-dependent mechanisms and mediates stretch-induced hypertrophy. *J Pharmacol Exp Ther* 2005; 315: 1075–1084
 157. Allison MA, Jenny NS, McClelland RL et al. The associations of adipokines with selected markers of the renin-angiotensinogen-aldosterone system: the multi-ethnic study of atherosclerosis. *J Hum Hypertens* 2015; 29: 127–133
 158. Lee MP, Madani S, Sekula D et al. Leptin increases expression and activity of matrix metalloproteinase-2 and does not alter

- collagen production in rat glomerular mesangial cells. *Endocr Res* 2005; 31: 27–37
159. Han DC, Isono M, Chen S et al. Leptin stimulates type I collagen production in db/db mesangial cells: glucose uptake and TGF-beta type II receptor expression. *Kidney Int* 2001; 59: 1315–1323
 160. Vu D, Tellez-Corrales E, Sakharkar P et al. Impact of NF-kappaB gene polymorphism on allograft outcome in Hispanic renal transplant recipients. *Transpl Immunol* 2013; 28: 18–23
 161. Fantuzzi G. Adiponectin in inflammatory and immune-mediated diseases. *Cytokine* 2013; 64: 1–10
 162. Scotece M, Conde J, Lopez V et al. Adiponectin and leptin: new targets in inflammation. *Basic Clin Pharmacol Toxicol* 2014; 114: 97–102
 163. Christou GA, Kiortsis DN. The role of adiponectin in renal physiology and development of albuminuria. *J Endocrinol* 2014; 221: R49–R61
 164. Burnett MS, Lee CW, Kinnaird TD et al. The potential role of resistin in atherogenesis. *Atherosclerosis* 2005; 182: 241–248
 165. Adya R, Tan BK, Punj A et al. Visfatin induces human endothelial VEGF and MMP-2/9 production via MAPK and PI3K/Akt signalling pathways: novel insights into visfatin-induced angiogenesis. *Cardiovasc Res* 2008; 78: 356–365
 166. Wang P, Xu TY, Guan YF et al. Perivascular adipose tissue-derived visfatin is a vascular smooth muscle cell growth factor: role of nicotinamide mononucleotide. *Cardiovasc Res* 2009; 81: 370–380
 167. Moschen AR, Kaser A, Enrich B et al. Visfatin, an adipocytokine with proinflammatory and immunomodulating properties. *J Immunol* 2007; 178: 1748–1758
 168. Pirvulescu MM, Gan AM, Stan D et al. Subendothelial resistin enhances monocyte transmigration in a co-culture of human endothelial and smooth muscle cells by mechanisms involving fractalkine, MCP-1 and activation of TLR4 and Gi/o proteins signaling. *Int J Biochem Cell Biol* 2014; 50: 29–37
 169. Li Y, Zhang Y, Dorweiler B et al. Extracellular Nampt promotes macrophage survival via a nonenzymatic interleukin-6/STAT3 signaling mechanism. *J Biol Chem* 2008; 283: 34833–34843
 170. Kim SR, Bae YH, Bae SK et al. Visfatin enhances ICAM-1 and VCAM-1 expression through ROS-dependent NF-kappaB activation in endothelial cells. *Biochim Biophys Acta* 2008; 1783: 886–895
 171. Maggio AB, Wacker J, Montecucco F et al. Serum resistin and inflammatory and endothelial activation markers in obese adolescents. *J Pediatr* 2012; 161: 1022–1027
 172. Romacho T, Azcutia V, Vazquez-Bella M et al. Extracellular PBEF/NAMPT/visfatin activates pro-inflammatory signalling in human vascular smooth muscle cells through nicotinamide phosphoribosyltransferase activity. *Diabetologia* 2009; 52: 2455–2463
 173. Calabro P, Samudio I, Willerson JT et al. Resistin promotes smooth muscle cell proliferation through activation of extracellular signal-regulated kinase 1/2 and phosphatidylinositol 3-kinase pathways. *Circulation* 2004; 110: 3335–3340
 174. Huang Q, Guo Y, Zeng H et al. Visfatin stimulates a cellular renin-angiotensin system in cultured rat mesangial cells. *Endocr Res* 2011; 36: 93–100
 175. Song HK, Lee MH, Kim BK et al. Visfatin: a new player in mesangial cell physiology and diabetic nephropathy. *Am J Physiol Renal Physiol* 2008; 295: F1485–F1494
 176. Boini KM, Zhang C, Xia M et al. Visfatin-induced lipid raft redox signaling platforms and dysfunction in glomerular endothelial cells. *Biochim Biophys Acta* 2010; 1801: 1294–1304
 177. Briffa JF, McAinch AJ, Poronnik P et al. Adipokines as a link between obesity and chronic kidney disease. *Am J Physiol Renal Physiol* 2013; 305: F1629–F1636
 178. Hamrick MW, Ferrari SL. Leptin and the sympathetic connection of fat to bone. *Osteoporos Int* 2008; 19: 905–912
 179. Zhang J, Wang N. Leptin in chronic kidney disease: a link between hematopoiesis, bone metabolism, and nutrition. *Int Urol Nephrol* 2014; 46: 1169–1174
 180. Tsuji K, Maeda T, Kawane T et al. Leptin stimulates fibroblast growth factor 23 expression in bone and suppresses renal 1alpha,25-dihydroxyvitamin D3 synthesis in leptin-deficient mice. *J Bone Miner Res* 2010; 25: 1711–1723
 181. Gainsford T, Alexander WS. A role for leptin in hemopoieses? *Mol Biotechnol* 1999; 11: 149–158
 182. Axelsson J, Qureshi AR, Heimbürger O et al. Body fat mass and serum leptin levels influence epoetin sensitivity in patients with ESRD. *Am J Kidney Dis* 2005; 46: 628–634
 183. Nasri H. Association of serum leptin with anemia in maintenance hemodialysis patients. *Saudi J Kidney Dis Transpl* 2006; 17: 521–525
 184. Stenvinkel P, Heimbürger O, Lonnqvist F et al. Does the ob gene product leptin stimulate erythropoiesis in patients with chronic renal failure? *Kidney Int* 1998; 53: 1430–1431
 185. Kumada M, Kihara S, Sumitani S et al. Association of hypoadiponectinemia with coronary artery disease in men. *Arterioscler Thromb Vasc Biol* 2003; 23: 85–89
 186. Tamura T, Furukawa Y, Taniguchi R et al. Serum adiponectin level as an independent predictor of mortality in patients with congestive heart failure. *Circ J* 2007; 71: 623–630
 187. Gnancinska M, Malgorzewicz S, Lysiak-Szydłowska W et al. The serum profile of adipokines in overweight patients with metabolic syndrome. *Endokrynol Pol* 2010; 61: 36–41
 188. Sahin G, Akay OM, Uslu S et al. Association between endothelial and platelet function markers and adiponectin in renal transplanted recipients on cyclosporine and tacrolimus immunosuppression based therapy. *Nephrology (Carlton)* 2015; 20: 392–398
 189. Malgorzewicz S, Debska-Slizien A, Czajka B et al. Adipokines and nutritional status in kidney transplant recipients. *Transplant Proc* 2014; 46: 2622–2626
 190. Wang XH, Dou LZ, Gu C et al. Plasma levels of omentin-1 and visfatin in senile patients with coronary heart disease and heart failure. *Asian Pac J Trop Med* 2014; 7: 55–62
 191. Hung WC, Yu TH, Hsu CC et al. Plasma visfatin levels are associated with major adverse cardiovascular events in patients with acute ST-elevation myocardial infarction. *Clin Invest Med* 2015; 38: E100–E109
 192. Dahl TB, Yndestad A, Skjelland M et al. Increased expression of visfatin in macrophages of human unstable carotid and coronary atherosclerosis: possible role in inflammation and plaque destabilization. *Circulation* 2007; 115: 972–980
 193. Weikert C, Westphal S, Berger K et al. Plasma resistin levels and risk of myocardial infarction and ischemic stroke. *J Clin Endocrinol Metab* 2008; 93: 2647–2653
 194. Frankel DS, Vasan RS, D'Agostino RB Sr et al. Resistin, adiponectin, and risk of heart failure the Framingham offspring study. *J Am Coll Cardiol* 2009; 53: 754–762
 195. Pham PT, Pham PM, Pham SV et al. New onset diabetes after transplantation (NODAT): an overview. *Diabetes Metab Syndr Obes* 2011; 4: 175–186
 196. Bayes B, Lauzurica R, Granada ML et al. Adiponectin and risk of new-onset diabetes mellitus after kidney transplantation. *Transplantation* 2004; 78: 26–30

197. Romanowski M, Dziedziejko V, Maciejewska-Karłowska A et al. Adiponectin and leptin gene polymorphisms in patients with post-transplant diabetes mellitus. *Pharmacogenomics* 2015; 16: 1–9
198. Shu KH, Wu MJ, Chen CH et al. Serum adipokine levels in renal transplant recipients. *Transplant Proc* 2014; 46: 381–384
199. Olszanecka-Glinianowicz M, Owczarek A, Bozentowicz-Wikarek M et al. Relationship between circulating visfatin/NAMPT, nutritional status and insulin resistance in an elderly population—results from the PolSenior substudy. *Metabolism* 2014; 63: 1409–1418
200. Xiang RL, Mei M, Su YC et al. Visfatin protects rat pancreatic beta-cells against IFN-gamma-induced apoptosis through AMPK and ERK1/2 signaling pathways. *Biomed Environ Sci* 2015; 28: 169–177
201. Kim da S, Kang S, Moon NR et al. Central visfatin potentiates glucose-stimulated insulin secretion and beta-cell mass without increasing serum visfatin levels in diabetic rats. *Cytokine* 2014; 65: 159–166
202. Yano Y, Hoshida S, Ishikawa J et al. Differential impacts of adiponectin on low-grade albuminuria between obese and nonobese persons without diabetes. *J Clin Hypertens (Greenwich)* 2007; 9: 775–782
203. Othman M, Kawar B, El Nahas AM. Influence of obesity on progression of non-diabetic chronic kidney disease: a retrospective cohort study. *Nephron Clin Pract* 2009; 113: c16–c23
204. Prior SL, Tang TS, Gill GV et al. Adiponectin, total antioxidant status, and urine albumin excretion in the low-risk “golden years” type 1 diabetes mellitus cohort. *Metabolism* 2011; 60: 173–179
205. Kasiske BL, Snyder JJ, Gilbertson DT et al. Cancer after kidney transplantation in the United States. *Am J Transplant* 2004; 4: 905–913
206. Scheid MP, Sweeney G. The role of adiponectin signaling in metabolic syndrome and cancer. *Rev Endocr Metab Disord* 2014; 15: 157–167
207. Hebbard L, Ranscht B. Multifaceted roles of adiponectin in cancer. *Best Pract Res Clin Endocrinol Metab* 2014; 28: 59–69
208. Scholze A, Rattensperger D, Zidek W et al. Low serum leptin predicts mortality in patients with chronic kidney disease stage 5. *Obesity (Silver Spring)* 2007; 15: 1617–1622
209. Moraes-Vieira PM, Bassi EJ, Larocca RA et al. Leptin deficiency modulates allograft survival by favoring a Th2 and a regulatory immune profile [corrected]. *Am J Transplant* 2013; 13: 36–44
210. Zoccali C, Postorino M, Marino C et al. Waist circumference modifies the relationship between the adipose tissue cytokines leptin and adiponectin and all-cause and cardiovascular mortality in haemodialysis patients. *J Intern Med* 2011; 269: 172–181
211. Rhee CM, Nguyen DV, Moradi H et al. Association of adiponectin with body composition and mortality in hemodialysis patients. *Am J Kidney Dis* 2015; 66: 313–321
212. Chung W, Jung ES, Shin D et al. Low resistin level is associated with poor hospitalization-free survival in hemodialysis patients. *J Korean Med Sci* 2012; 27: 377–381
213. Nagy K, Ujszaszi A, Czira ME et al. Association between serum resistin level and outcomes in kidney transplant recipients. *Transpl Int* 2016; 29: 352–361