

UCSF

UC San Francisco Previously Published Works

Title

Accuracy and precision of estimation equations to predict net endogenous acid excretion using the Australian food database

Permalink

<https://escholarship.org/uc/item/2tw0s96q>

Journal

Nutrition & Dietetics, 74(3)

ISSN

1446-6368

Authors

Parmenter, Benjamin H

Slater, Gary J

Frassetto, Lynda A

Publication Date

2017-07-01

DOI

10.1111/1747-0080.12324

Peer reviewed

ORIGINAL RESEARCH

Accuracy and precision of estimation equations to predict net endogenous acid excretion using the Australian food database

Benjamin H. PARMENTER,¹ Gary J. SLATER¹ and Lynda A. FRASSETTO²¹Faculty of Health, Science and Engineering, University of the Sunshine Coast, Queensland, Australia and ²School of Medicine, University of California San Francisco (UCSF), California, USA**Abstract**

Aim: The gold standard of measurement for net endogenous acid production (NEAP) is net acid excretion (NAE), a test that is not readily available, and consequently, estimative equations by Remer and Manz and Frassetto *et al.* are often used. These equations rely on nutrient databases and it is recommended that their validity be assessed using a country's database before their application in research in that country. We sought to delineate the accuracy and precision of these estimation equations using the Australian food database.

Methods: In a double blind, randomised, cross-over fashion, healthy participants (n = 13) residing in regional Australia were exposed to varying net acid loads while they collected weighted food diaries and 24-hour urine samples for measurement of NAE.

Results: In comparison to the Frassetto *et al.* equations (equation one bias = -57.1 mEq/day, equation two bias = -32.8 mEq/day), only the Remer and Manz equation was accurate (bias = -5.4 mEq/day); however, all equations were imprecise.

Conclusions: Using the Australian database, the performance of these equations to predict NEAP appears equal to other databases; however, caveats apply in their application. For future research, the equation by Remer and Manz is preferential for group estimates. None of the equations are recommended for individual estimates.

Key words: acid, base, diet, estimation.

Introduction

The homeostatic regulation of acid–base status is essential for physiological functioning, and the clinical consequences of acid–base disorders are well known. Another factor known to influence physiological acid–base systems is the dietary acid load.¹ The ratio of diet-derived fixed acid to base is estimated by the potential renal acid load (PRAL), and when endogenously produced organic acid is factored, cumulatively, this determines the net endogenous acid production (NEAP) for an individual.¹ When diet chronically releases fixed acid in excess of fixed base, the surplus acid has been hypothesised to be a contributing factor to chronic disease.^{2,3} While an elevated (acid) NEAP does not result in

clinical metabolic acidosis,⁴ research is being conducted to determine its relationship to sarcopenia,^{5,6} gout,^{7,8} renal stones,^{9,10} metabolic syndrome,¹¹ chronic renal insufficiency,¹² late metabolic acidosis in preterm infants,¹³ hypertension,^{14,15} insulin resistance,^{16,17} non-alcoholic fatty liver disease,¹⁸ impaired sports performance^{19,20} and osteoporosis,^{21,22} while the involvement of NEAP in obesity²³ and cancer⁴ has been theorised.

For the research-focused investigation of NEAP, its accurate quantification is essential. To directly measure NEAP, 24-hour urinary net acid excretion (NAE) is the criterion method.^{24,25} However, measurement of NAE is a labour- and laboratory-intensive technique and is not readily available.²⁶ As such, estimative equations from dietary intake were developed. Remer and Manz's equation (NEAP_R) factors for endogenously produced organic acid plus the PRAL, that is, the ratio of dietary conjugate bases of potassium, magnesium and calcium to conjugate acids of phosphorus- and sulphate-containing protein.²⁷ Conversely, the Frassetto *et al.* equations (NEAP_F) were developed to allow faster calculations and utilise the more easily obtained inputs of dietary protein and potassium intake.²⁸ However, world experts have recommended that before these equations are applied within a country, their accuracy is assessed with that country's food

B.H. Parmenter, BNut&Diet (Hons), Student

G.J. Slater, PhD, Senior Lecturer

L.A. Frassetto, MD, Non-Senate Acad Emeritus (WOS), Associate Clinical Professor of Medicine

Correspondence: B.H. Parmenter, Faculty of Health, Science and Engineering, University of the Sunshine Coast, 90 Sippy Downs Dr, Sippy Downs, QLD 4556, Australia. Tel: +61 4 2353 6798. Email: b_p026@student.usc.edu.au

Accepted August 2016

database.^{24,29} To date, this has only been investigated using German²⁹ and American²⁸ databases. Consequently, for future research investigating NEAP within Australia, we sought to delineate the accuracy and precision of estimate NEAP_R and NEAP_F using the Australian food database. We hypothesise that estimate NEAP_R and NEAP_F would predict NAE to a similar degree when using the Australian food database as they do when using other databases.

Methods

The design of the trial is described in detail in our companion paper.³⁰ In summary, in a double blind, randomised, cross-over fashion, healthy participants (n = 13) residing in regional Australia were exposed to varying NEAP loads while they collected weighted food diaries and 24-hour urine samples. To vary the NEAP loads, the participants consumed a fruit and vegetable concentrate or placebo for three days each, with diet standardised throughout. The urine samples were collected under thymol and paraffin oil in portable cooler boxes with ice packs for the subsequent analysis of NAE, pH and creatinine. Compliance to 24-hour urine collections was measured by creatinine index; those with a daily excretion <0.1 mmol/kg body weight were excluded.³¹ Diet compositions were determined using Foodworks Professional (Xyris, Brisbane, Australia) using the Australian food database (NUTTAB 2010 Australian Government Nutrient Database, Canberra, Australia). When food items were missing, nutrients were entered according to nutrient information on the food label before proceeding to estimate NEAP_R and two NEAP_F equations (Table 1). The methods were approved by the University of the Sunshine Coast's Human Ethics Committee (reference number: S/14/70), and all participants signed informed consent. The trial was designed to adhere to the Consolidated Standards of Reporting Trials (CONSORT) guidelines, and the trial protocol was registered with the Australian New Zealand Clinical Trials Registry (ACTRN12616000417482).

For the assessment of physique, in the week prior to commencing the study, volunteers presented to the laboratory in

a rested state after an overnight fast. After bladder evacuation, participants had their stretch stature measured to the nearest 0.1 cm using a technique previously described,³² with a Harpenden stadiometer (Holtain Limited, Crymch, UK) and undertook air-displacement plethysmography for assessment of body composition (BODPOD®; COSMED USA, Inc., Concord, CA, USA). The manufacturer's recommended procedure was followed,³³ and a predicted thoracic lung volume was used.³⁴ During the procedure, the participants wore skin-tight clothing, a silicone cap and removed all metal objects. Following the procedure, body mass index (BMI) was calculated conventionally, while the system's software (Version 5.3.2; COSMED USA, Inc., Concord, CA, USA) estimated fat mass and fat-free mass using the model defined by the Siri equation.³⁵

The 24-hour urine samples were analysed on the day of their return for urine pH, creatinine and NAE. For the determination of pH, samples were analysed by a calibrated labChem pH meter (TPS Pty Ltd, Melbourne, Australia) at ambient temperature and 36.5 ± 0.5 °C. Creatinine was measured spectrophotometrically by a colorimetric detection kit Cat no. 09151410 (Enzo Life Sciences, Farmingdale NY, USA) using a multi-scan GO plate reader (ThermoFisher Scientific, Waltham, MA, USA). Ammonium was measured using methods of Rice *et al.*³⁶ by flow injection analysis using a Lachat QuikChem 8000 (Lachat Instruments, Milwaukee, WI, USA). Titratable acids and bicarbonate were measured in duplicate by using the methods of Litkowski and Wilson³⁷ with the following modifications. Briefly, after addition of 0.05 M hydrogen chloride to 20 mL of urine samples, simmering occurred for 20 minutes at 90 °C. Concurrently, a blank was run in the same manner using MilliQ water (Millipore Corp., Bedford, MA, USA). After samples cooled, bicarbonate was determined as the amount of 0.05 M sodium hydroxide necessary to titrate back to the initial pH at ambient temperature less the corresponding value for the blank. Thereafter, titratable acid was determined as the amount of 0.05 M sodium hydroxide necessary to titrate to pH 7.4 from the bicarbonate endpoint. The mean of the duplicates was used for the calculation of NAE (Table 1), and the intra-assay CV was 8.0%.

Table 1 Equations to estimate the potential renal acid load, the net endogenous acid production and to calculate net acid excretion

Equation	Formula
NEAP _F (mEq/day)	Equation 1 = $(0.91 \times \text{protein (g/day)}) - (0.57 \times \text{potassium (mEq/day)}) + 21$ Equation 2 = $(54.5 \times \text{protein (g/day)}) / \text{potassium (mEq/day)} - 10.2$
NEAP _R (mEq/day)	= PRAL (mEq/day) + organic acids _{anthro}
PRAL (mEq)	= $0.488 \times \text{protein (g/day)} + 0.0366 \times \text{phosphorus (mg/day)} - 0.0205 \times \text{potassium (mg/day)} - 0.0263 \times \text{magnesium (mg/day)} - 0.0125 \times \text{calcium (mg/day)}$
Organic acids _{anthro} , (mEq/day)	= body surface area $\times 41 / 1.73$
Body surface area (m ²)	= $0.007184 - \text{height (cm)}^{0.725} - \text{weight (kg)}^{0.425}$
NAE (mEq)	= titratable acids (mEq) + ammonium (mEq) - bicarbonate (mEq)

Anthro, anthropometrical; NAE, net acid excretion; NEAP, estimate net endogenous acid production where subscript R pertains to the equation by Remer and Manz²⁷ and subscript F pertains to the equations by Frassetto *et al.*²⁸; PRAL, potential renal acid load.

Table 2 Characteristics of participants completing the study

	All participants	Males	Females
n	13	6	7
Age (years)	35 ± 13	30 ± 9	39 ± 16
Height (cm)	172 ± 7	177 ± 6	166 ± 3
Weight (kg)	73 ± 10	77 ± 7	69 ± 11
Fat mass (kg)	19 ± 10	12 ± 7	25 ± 9
Fat-free mass (kg)	54 ± 11	65 ± 4	44 ± 4
BMI (kg/m ²)	25 ± 3	24 ± 1	25 ± 5

Values are means ± SD.

Table 3 Net acid excretion and estimation of the net endogenous acid production using the Australian food database during the placebo period, supplement period and for all observations

	n	Placebo	Supplement	Combined placebo and supplement
NAE (mEq/day)	10	40 ± 19	7 ± 25	23 ± 27
NEAP _R (mEq/day)	10	48 ± 26	8 ± 26	28 ± 33
NAE (mEq/day)	11	38 ± 18	12 ± 30	25 ± 27
NEAP _{F1} (mEq/day)	11	85 ± 55	79 ± 55	82 ± 53
NEAP _{F2} (mEq/day)	11	63 ± 34	53 ± 28	57 ± 30

Values are means ± SD.

NAE, net acid excretion; NEAP, estimate net endogenous acid production where subscript R pertains to the equation by Remer and Manz²⁷ and subscripts F¹ and F² pertain to equations one and two by Frassetto *et al.*²⁸.

To statistically assess the agreement between estimated and analysed NEAP, the Bland–Altman method for repeat non-constant observations was used (i.e. one observation during the placebo period and one observation during the supplement period on the same participant).³⁸ Equations were computed using dietary data from the concurrent 24-hour urine collection periods. The distribution of the differences was checked and outliers ($\pm > 3.0$ SD of the difference) excluded.³⁹ Those with missing data were excluded.³⁸ A priori limits of agreement were set at ± 15 mEq/day as this would permit a reasonable estimation of NEAP in individuals and also account for imprecision in the criterion method. All statistical analyses were completed

using MedCalc (MedCalc Software; MedCalc, Mariakerke, Belgium).

Results

Of the 16 enrolled individuals, 13 completed the study; their characteristics are described in Table 2. Conversely, their dietary intake and a CONSORT diagram are presented in our companion paper.³⁰ During phase two, one participant returned an incomplete 24-hour urine sample, while another failed to replicate their food and fluid from phase one; both were excluded from analysis. One outlier was identified in the dataset of NEAP_R, while none were identified in either NEAP_F. The mean ± SD of NAE, NEAP_R and NEAP_F during the placebo period, supplement period and for all observations is presented in Table 3, while Table 4 presents the Bland–Altman bias and limits of agreement for all observations. NEAP_R showed the best agreement with an acceptable accuracy; however, all equations were beyond the a priori limits of agreement to predict NAE in individuals.

Discussion

To our knowledge, this is the first time the estimate NEAP equations have been assessed using the Australian food database. This is important because it impacts the choice of equation potentially used in future research conducted in Australia. Using the Australian database, the performance of these estimate NEAP equations appears similar to their performance reported using other databases. However, known imprecisions within the equations themselves are apparent, which precludes their application to individuals. Of the equations investigated, NEAP_R appears more accurate for group estimates, and its use is, therefore, preferentially encouraged.

Using the Australian food database, the performance of the estimate NEAP equations appears similar to their performance reported using other databases. Similar to the German database, our results show a reasonable group estimate using NEAP_R, evident by the small bias (−5.4 mEq).²⁹ Compared to the American database, investigations were completed under steady-state conditions, which are known to increase the magnitude of accuracy.^{28,40} Taken together, there is no reason to suggest that the equations performance is altered when using the Australian database.

Table 4 Bland–Altman agreement between measured 24-hour net acid excretion and net endogenous acid estimation equations in repeat non-constant observations in participants consuming non-steady-state diets using the Australian food database

	n	Bias	95% CI for the bias	Limits of agreement	95% CI for the upper limit of agreement	95% CI for the lower limit of agreement
NEAP _R (mEq/day)	10	−5.4	−19.8, 9.0	−54.2, 43.3	18.3, 68.3	−79.1, −29.2
NEAP _{F1} (mEq/day)	11	−57.1	−82.0, −32.2	−146.6, 32.4	10.7, 75.5	−189.7, −103.5
NEAP _{F2} (mEq/day)	11	−32.8	−48.6, −16.9	−90.7, 25.1	−2.3, 52.5	−118.1, −63.3

NEAP, estimate net endogenous acid production where subscript R pertains to the equation by Remer and Manz²⁷ and subscripts F¹ and F² pertain to equations one and two by Frassetto *et al.*²⁸

However, NEAP_R was imprecise, which precludes its application to individuals. This may be because of known errors within the equation surrounding dietary protein and exogenous organic acids.²⁹ To account for this, Sebastian *et al.*⁴¹ developed another equation that incorporates dietary cystine and methionine intake. Yet, as the Australian database has limited amino acid records, this equation was not computed. That said, no equations currently factor for exogenous organic acids. As such, researchers may wish to revise the equations by factoring for the variability in protein composition as well as exogenous organic acids, perhaps based on food groups.²⁹

The inaccuracy of the NEAF_F equations may have been caused by the experimental conditions. That is, the predominance of the supplements base was delivered by a large calcium dose (1.7 g/day); however, both NEAF_F equations do not factor for calcium. Consequently, it appears that these equations are less robust under such circumstances, and NEAF_F is not universally applicable. Moreover, negative-feedback control of endogenous acid production may have influenced NAE in some individuals, further contributing to the equations' imprecision.⁴² To this end, it may be most beneficial for future investigations to collect not only urinary PRAL (electrolytes measured in 24-hour urines) but also total urinary organic acids alongside dietary data so as to aid in the elucidation of the issues associated with endogenous and exogenous organic acids and NAE prediction. This may be pivotal in the interpretation of investigations delineating the relationship between NAE and various chronic degenerative diseases as NAE includes both organic acid components, wherein the exogenous component may provide an acid-forming yet health-supporting effect *in vivo*.⁴³

The limitations of the present study include measurement error within the NAE technique, the use of weighted food diaries and small sample size. However, our CV (8.0%) for NAE was below all other studies that reported a CV (10.1–10.9%).^{44,45} Consequently, this appears to be a common limitation. The use of free-living weighted food diaries may have introduced error (e.g. under-reporting); however, it may also provide re-assurance that the NEAP_R equation has a reasonable capacity to perform group estimates under free-living conditions. Finally, the small sample size impacts the analysis by the creation of large confidence intervals around the upper and lower limits of agreement. Yet, given that estimations of NEAP are not utilised in a critical clinical setting, the data adequately serves to illustrate which method is preferential for research purposes in Australia. In conclusion, the estimate NEAP equations studied should not be applied to individuals, and NEAP_R is preferential for group estimates.

Funding source

Supplements donated in kind by Morlife, Gold Coast, Australia (<http://www.morlife.com/>). Morlife had no role in any aspect of study design, data collection, interpretation or presentation.

Conflict of interest

The authors have no conflicts of interest to declare.

Authorship

We thank GS and BP for the project conception; BP for conducting the research and obtaining donations, Dr Peter Brooks for developing the analytical methods, Daryle Sullivan for the measurement of ammonium and LF for methodological contributions; Michael Nielsen for sourcing essential reagents and materials; BP for performing the statistics and writing the paper; and GS and LF for critical revision of the manuscript and study supervision. All authors have read and approved the final manuscript. The content has not been published or submitted for publication elsewhere.

References

- 1 Poupin N, Calvez J, Lassale C, Chesneau C, Tome D. Impact of the diet on net endogenous acid production and acid-base balance. *Clin Nutr* 2012; **31**: 313–21.
- 2 Pizzorno J, Frassetto LA, Katzinger J. Diet-induced acidosis: is it real and clinically relevant? *Br J Nutr* 2010; **103**: 1185–94.
- 3 Adeva MM, Souto G. Diet-induced metabolic acidosis. *Clin Nutr* 2011; **30**: 416–21.
- 4 Robey IF. Examining the relationship between diet-induced acidosis and cancer. *Nutr Metab* 2012; **9**: 72.
- 5 Frassetto L, Morris RC Jr, Sebastian A. Potassium bicarbonate reduces urinary nitrogen excretion in postmenopausal women. *J Clin Endocrinol Metab* 1997; **82**: 254–9.
- 6 Dawson HB, Harris SS, Ceglia L. Alkaline diets favor lean tissue mass in older adults. *Am J Clin Nutr* 2008; **87**: 662–5.
- 7 Kanbara A, Hakoda M, Seyama I. Urine alkalization facilitates uric acid excretion. *Nutr J* 2010; **9**: 45.
- 8 Kanbara A, Miura Y, Hyogo H, Chayama K, Seyama I. Effect of urine pH changed by dietary intervention on uric acid clearance mechanism of pH-dependent excretion of urinary uric acid. *Nutr J* 2012; **11**: 39.
- 9 Reddy ST, Wang CY, Sakhaee K, Brinkley L, Pak CY. Effect of low-carbohydrate high-protein diets on acid-base balance, stone-forming propensity, and calcium metabolism. *Am J Kidney Dis* 2002; **40**: 265–74.
- 10 Trinchieri A, Maletta A, Lizzano R, Marchesotti F. Potential renal acid load and the risk of renal stone formation in a case-control study. *Eur J Clin Nutr* 2013; **67**: 1077–80.
- 11 Bahadoran Z, Mirmiran P, Khosravi H, Azizi F. Associations between dietary acid-base load and cardiometabolic risk factors in adults: the Tehran Lipid and Glucose Study. *Endocrinol Metab* 2014; **30**: 201–7.
- 12 Banerjee T, Crews DC, Wesson DE *et al.* Dietary acid load and chronic kidney disease among adults in the United States. *BMC Nephrol* 2014; **15**: 137.
- 13 Kalhoff H, Manz F, Kiwull P, Kiwull-Schone H. Food mineral composition and acid-base balance in preterm infants. *Eur J Nutr* 2007; **46**: 188–95.
- 14 Murakami K, Sasaki S, Takahashi Y, Uenishi K. Association between dietary acid-base load and cardiometabolic risk factors in young Japanese women. *Br J Nutr* 2008; **100**: 642–51.

- 15 Engberink MF, Bakker SJ, Brink EJ *et al.* Dietary acid load and risk of hypertension: the Rotterdam Study. *Am J Clin Nutr* 2012; **95**: 1438–44.
- 16 Fagherazzi G, Vilier A, Bonnet F *et al.* Dietary acid load and risk of type 2 diabetes: the E3N-EPIC cohort study. *Diabetologia* 2014; **57**: 313–20.
- 17 Xu H, Jia T, Huang X *et al.* Dietary acid load, insulin sensitivity and risk of type 2 diabetes in community-dwelling older men. *Diabetologia* 2014; **57**: 1561–8.
- 18 Chan R, Wong VW-S, Chu WC-W *et al.* Higher estimated net endogenous acid production may be associated with increased prevalence of nonalcoholic fatty liver disease in Chinese adults in Hong Kong. *PLoS ONE* 2015; **10**: doi: 10.1371/journal.pone.0122406.
- 19 Hietavala EM, Puurinen R, Kainulainen H, Mero AA. Low-protein vegetarian diet does not have a short-term effect on blood acid-base status but raises oxygen consumption during submaximal cycling. *J Int Soc Sports Nutr* 2012; **9**: 50–9.
- 20 Caciano SL, Inman CL, Gockel-Blessing EE, Weiss EP. Effects of dietary acid load on exercise metabolism and aerobic exercise performance. *J Sports Sci Med* 2015; **14**: 364.
- 21 Lambert H, Frassetto L, Moore J *et al.* The effect of supplementation with alkaline potassium salts on bone metabolism: a meta-analysis. *Osteoporos Int* 2015; **26**: 1311–8.
- 22 Fenton TR, Tough SC, Lyon AW, Eliasziw M, Hanley DA. Causal assessment of dietary acid load and bone disease: a systematic review & meta-analysis applying Hill's epidemiologic criteria for causality. *Nutr J* 2011; **10**: 41.
- 23 Berkemeyer S. Acid-base balance and weight gain: are there crucial links via protein and organic acids in understanding obesity? *Med Hypotheses* 2009; **73**: 347–56.
- 24 Frassetto LA, Lanham New SA, Macdonald HM *et al.* Standardizing terminology for estimating the diet-dependent net acid load to the metabolic system. *J Nutr* 2007; **137**: 1491–2.
- 25 Scialla JJ, Appel LJ, Astor BC *et al.* Estimated net endogenous acid production and serum bicarbonate in African Americans with chronic kidney disease. *Clin J Am Soc Nephrol* 2011; **6**: 1526–32.
- 26 Welch AA, Mulligan A, Bingham SA, Khaw KT. Urine pH is an indicator of dietary acid-base load, fruit and vegetables and meat intakes: results from the European Prospective Investigation into Cancer and Nutrition (EPIC)-Norfolk population study. *Br J Nutr* 2008; **99**: 1335–43.
- 27 Remer T, Manz F. Estimation of the renal net acid excretion by adults consuming diets containing variable amounts of protein. *Am J Clin Nutr* 1994; **59**: 1356–61.
- 28 Frassetto LA, Todd KM, Morris RC Jr, Sebastian A. Estimation of net endogenous noncarbonic acid production in humans from diet potassium and protein contents. *Am J Clin Nutr* 1998; **68**: 576–83.
- 29 Remer T, Dimitriou T, Manz F. Dietary potential renal acid load and renal net acid excretion in healthy, free-living children and adolescents. *Am J Clin Nutr* 2003; **77**: 1255–60.
- 30 Parmenter BH, Slater GJ, Frassetto LA. Spot-testing urine pH, a novel dietary biomarker? A randomized cross-over trial. *Nutr Diet* 2016; doi: 10.1111/1747-0080.12325.
- 31 Remer T, Neubert A, Maser-Gluth C. Anthropometry-based reference values for 24-h urinary creatinine excretion during growth and their use in endocrine and nutritional research. *Am J Clin Nutr* 2002; **75**: 561–9.
- 32 Marfell-Jones MJ, Stewart AD, de Ridder JH. *International standards for anthropometric assessment*. Wellington, New Zealand: International Society for the Advancement of Kinanthropometry, 2012.
- 33 Dempster P, Aitkens S. A new air displacement method for the determination of human body composition. *Med Sci Sports Exerc* 1995; **27**: 1692–7.
- 34 Crapo R, Morris A, Clayton P, Nixon C. Lung volumes in healthy nonsmoking adults. *Bull Eur Physiopathol Respir* 1981; **18**: 419–25.
- 35 Siri WE. Body composition from fluid spaces and density: analysis of methods. In: Brozek J, Hanschel A, eds. *Techniques for Measuring Body Composition*. Washington, DC: National Academy of Sciences, National Research Council Washington, DC, 1961; 223–4.
- 36 Rice EW, Baird RB, Eaton AD, Clesceri LS. *Standard Methods for the Examination of Water and Wastewater*. Washington, DC: American Public Health Association, 2012.
- 37 Litkowski L, Wilson TL. Effect of protein on titrimetry of bicarbonate, titratable acid, and ammonium in urine. *Clin Chem* 1979; **25**: 362–5.
- 38 Bland JM, Altman DG. Agreement between methods of measurement with multiple observations per individual. *J Biopharm Stat* 2007; **17**: 571–82.
- 39 Bland JM, Altman D. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* 1986; **327**: 307–10.
- 40 Kurtz I, Maher T, Hulter HN, Schambelan M, Sebastian A. Effect of diet on plasma acid-base composition in normal humans. *Kidney Int* 1983; **24**: 670–80.
- 41 Sebastian A, Frassetto LA, Sellmeyer DE, Merriam RL, Morris RC Jr. Estimation of the net acid load of the diet of ancestral preagricultural *Homo sapiens* and their hominid ancestors. *Am J Clin Nutr* 2002; **76**: 1308–16.
- 42 Epstein FH, Hood VL, Tannen RL. Protection of acid-base balance by pH regulation of acid production. *N Engl J Med* 1998; **339**: 819–26.
- 43 Remer T, Berkemeyer S, Manz F. Acid base considerations in stone-age farming sweet potato eaters, modern-day sweet potato eaters, and high-protein consumers. *Open Nutr J* 2008; **2**: 23–8.
- 44 Jajoo R, Song L, Rasmussen H, Harris SS, Dawson-Hughes B. Dietary acid-base balance, bone resorption, and calcium excretion. *J Am Coll Nutr* 2006; **25**: 224–30.
- 45 Ceglia L, Harris SS, Abrams SA, Rasmussen HM, Dallal GE, Dawson-Hughes B. Potassium bicarbonate attenuates the urinary nitrogen excretion that accompanies an increase in dietary protein and may promote calcium absorption. *J Clin Endocrinol Metab* 2009; **94**: 645–53.