

# Lawrence Berkeley National Laboratory

## Recent Work

### Title

SYNTHESIS, STRUCTURE, AND CHARACTERIZATION OF A MIXED-VALENCE MN(III) , MN(IV) DIMU-OXO COMPLEX WITH A MACROCYCLIC TETRAAZA LIGAND

### Permalink

<https://escholarship.org/uc/item/2tx8v984>

### Authors

Brewer, K.J.

Calvin, M.

Otvos, J.W.

### Publication Date

1988-07-01

UC 401

LBL-25645

Preprint

c.1



# Lawrence Berkeley Laboratory

UNIVERSITY OF CALIFORNIA

## CHEMICAL BIODYNAMICS DIVISION

Submitted to Journal of the American Chemical Society

**Synthesis, Structure, and Characterization of a  
Mixed-Valence Mn(III), Mn(IV) di- $\mu$ -Oxo Complex  
with a Macrocyclic Tetraaza Ligand**

K.J. Brewer, M. Calvin, J.W. Otvos, and L.O. Spreer

July 1988


RECEIVED  
LAWRENCE  
BERKELEY LABORATORY

OCT 11 1988

LIBRARY AND  
DOCUMENTS SECTION

**For Reference**

Not to be taken from this room



Prepared for the U.S. Department of Energy under Contract Number DE-AC03-76SF00098.

LBL-25645  
c.1

## **DISCLAIMER**

This document was prepared as an account of work sponsored by the United States Government. While this document is believed to contain correct information, neither the United States Government nor any agency thereof, nor the Regents of the University of California, nor any of their employees, makes any warranty, express or implied, or assumes any legal responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by its trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States Government or any agency thereof, or the Regents of the University of California. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States Government or any agency thereof or the Regents of the University of California.

Synthesis, Structure, and Characterization of a Mixed-Valence  
Mn(III), Mn(IV) di- $\mu$ -Oxo Complex with a Macrocyclic Tetraaza Ligand

Karen J. Brewer, Melvin Calvin, John W. Otvos

Department of Chemistry and Lawrence Berkeley Laboratory

University of California, Berkeley, California 94720

and

Larry O. Spreer

Department of Chemistry, University of the Pacific

Stockton, California 95211

Abstract

The complex  $[(14\text{-aneN}_4)\text{MnO}]_2^{3+}$  (14-aneN<sub>4</sub> = 1,4,8,11-tetraazacyclo-tetradecane) has been prepared and characterized. The crystal structure of this mixed-valence (III,IV) compound shows discrete Mn(III) and Mn(IV) centers. Comparison of our crystallographic data with the data for the three previously reported Mn(III), Mn(IV) di- $\mu$ -oxo complexes indicates that while the external ligands are varied, the Mn-O distances in the Mn<sub>2</sub>O<sub>2</sub> core remain relatively unchanged ( $\pm 0.03$  Å). The X-band EPR experiments at 4K exhibit a 16-line spectrum centered at  $g = 2$ . This 16-line spectrum is attributed to the overlap of hyperfine splitting with two chemically different Mn nuclei ( $I = 5/2$ ) with one hyperfine coupling constant being roughly twice the magnitude of the other. This complex is a Robin and Day class II mixed-valence species. Detailed studies of the formation of this binuclear Mn species indicate that the bridging oxygens come from solvent water and that the bridge formation occurs with both manganese ions in the III oxidation state. Cyclic voltammetry of the complex exhibits a reversible one-electron oxidation ( $E_{1/2} = +1.0$  V vs SCE) to the IV,IV level as well as a reversible one

electron reduction ( $E_{1/2} = -0.12$  V vs SCE) to the III,III level. Spectroelectrochemical experiments have been performed, which help to establish the assignments of the electronic spectrum of the III,IV complex. The III,IV complex exhibits three main absorption maxima in the visible region ( $\lambda_{\text{max}}^{\text{abs}} = 538(\text{sh}), 550, 650$  nm) as well as a tail that extends into the near infrared. The bands at 538 and 550 nm are assigned to dd transitions involving the Mn(IV) and Mn(III) centers, respectively. The band at 650 nm is assigned to a ligand-to-metal charge transfer transition,  $\text{oxo} \rightarrow \text{Mn(IV)}$ . The tail into the near infrared is due to an intervalence transition, which underlies the intense 650 nm charge transfer band. Complexes such as 1 are of interest as models for multiple electron redox catalysts like the oxygen evolving complex (OEC) in photosystem II.

## Introduction

A dominant feature of the inorganic chemistry of manganese is the variety of complexes that can be made with different oxidation levels of the manganese. Complexes have been prepared which contain manganese in the 3-, 2-, 1-, 0, 1+, 2+, 3+, 4+, 5+, 6+ and 7+ oxidation states.<sup>1</sup> Living systems have utilized this property in that manganese-containing enzymes are involved in many diverse redox functions.<sup>2</sup> This has led to a great deal of recent work in attempting to develop synthetic model manganese compounds that mimic properties of natural systems.<sup>3</sup> Of particular interest has been the development of multinuclear manganese complexes. The biological function that has generated the most interest is the photosynthetic oxygen-evolving complex of photosystem II in which it is generally believed that four manganese atoms are involved.<sup>4</sup> These manganese atoms are in close proximity to one another. Much of the available data can be interpreted by a dimer of dimers structure<sup>5</sup> in which the two dimers are electronically isolated from each other. The first good model compounds for the short Mn--Mn distance and multiline signal in the EPR spectrum of photosystem II were the mixed valence di-μ-oxo species  $[L_2MnO_2MnL_2]^{3+}$  (L = bipyridine or phenanthroline).<sup>6</sup> Recently two new Mn(III),Mn(IV) binuclear di-μ-oxo complexes have been reported.<sup>7,8</sup> In addition, a number of trinuclear<sup>9</sup> and tetranuclear<sup>10</sup> manganese compounds involving at least one bridging oxo group have been described. Most of the synthetic procedures for these aggregated manganese species involve oxidation of manganese(II) precursors. No information about the mechanism of the formation of these multinuclear species has been presented, however. Also, the origin of the oxygen-bridging atom in this type of compound has not previously been estab-

blished because most synthetic routes have involved oxidants that can act as oxygen atom donors. Evidence is described herein that supports a scheme involving oxidation of Mn(II) to Mn(III), followed by dimer formation with water as the source for bridging oxygen.

We recently reported a new mixed-valence Mn(III), Mn(IV) di- $\mu$ -oxo compound with a macrocyclic tetradentate ligand, 1,4,8,11-tetraazacyclotetradecane,<sup>11</sup>  $[(14\text{-aneN}_4)\text{MnO}]_2^{3+}$ , 1. Our interest in this and other mixed-valence binuclear manganese species is their potential as multiple-electron redox catalysts that may promote the oxidation of water. The electrochemical oxidation of complex 1 in dry acetonitrile produces the IV,IV complex. If water is then added, the oxidized form of the complex reacts with water with partial regeneration of the III,IV complex. This chemistry is still being investigated.

Compound 1 is one of the few species that have been described with a cis-configuration for a fourteen-membered macrocycle. The vast majority of complexes of 14-aneN<sub>4</sub> previously prepared exist with the ligand in a planar or trans configuration. Even in the case of a large metal atom such as osmium(III), the ligand 14-aneN<sub>4</sub> adopts a planar, trans geometry to form trans-[Os(14-aneN<sub>4</sub>)Cl<sub>2</sub>]<sup>+</sup>.<sup>12</sup> Another interesting observation about the compound  $[(14\text{-aneN}_4)\text{MnO}]_2^{3+}$  is that it self-assembles from dilute solution. In contrast to most synthetic procedures of multinuclear metal complexes, which require very concentrated solutions (ca. 0.1 mol L<sup>-1</sup>), our complex can be prepared from millimolar solutions. These facts and other experiments described herein point out the remarkable stability of the Mn<sub>2</sub>O<sub>2</sub> core. This core is particularly stable with the Mn in the III and IV oxidation levels.

### Experimental Section

Visible spectra were recorded on a Hewlett-Packard 8450A Diode Array Spectrometer with 2 nm resolution or on a Varian 2300. Solutions used for these measurements were millimolar solutions in acetonitrile solvent. Typical spectra on the 8450A were the average of ten scans.

X-band EPR spectra were recorded on a Varian Model E109 Spectrometer equipped with a low-temperature dewar (Air Products, Ltd.) A dilute solution of the complex (ca.  $10^{-5}$  mol L<sup>-1</sup>) in 2:1 ethylene glycol: water (v:v) was frozen into a glass at 77K in liquid nitrogen and then cooled to 4K with helium. All EPR spectra were recorded at 4K.

Infrared spectra were recorded on a Nicolet 5DX FT Infrared Spectrometer System. Solid samples were used as KBr pellets (2-4 wt%) prepared in a Wilk's minipress. Typical spectra were an average of twenty scans.

Electrochemistry was performed using a PAR Model 175 Potentiostat equipped with a PAR Model 173 Controller. Dry acetonitrile was used as the solvent and tetrabutylammonium hexafluorophosphate (TBAH) (0.1 M) was used as the electrolyte. Solutions used for cyclic voltammograms were of millimolar concentration in the electroactive species. A nonaqueous Ag/Ag<sup>+</sup> electrode was used as the reference electrode (potential vs. Ag/Ag<sup>+</sup> + 0.34 V = potential vs. SCE). A glassy carbon disk working electrode and platinum wire counter electrode were used for cyclic voltammetry.

Spectroelectrochemistry was performed in a cell constructed from a quartz cuvette with a quartz fritted compartment separating the working and counter electrode units (see Fig. 1). The counter electrode was Pt foil and the working electrode was a Pt mesh cylinder mechanically



supported by two Pt strips. The light beam of the spectrometer passed between the two Pt strips. Stirring was accomplished by bubbling with argon. With this cell, one-electron controlled potential oxidation or reduction of a 1 millimolar solution of 1 in acetonitrile (with 0.1 M TBAH as supporting electrolyte) could be obtained in 6 minutes. The potential of the cell was controlled by a PAR Model 175 Potentiostat equipped with a PAR Model 173 Controller and a Model 179 Digital Coulometer.

### Materials and Methods

1,4,7,11-Tetraazacyclotetradecane was purchased from Strem Chemicals and used without further purification. Acetonitrile was UV grade from Burdick and Jackson dried over activated molecular sieves. Water was purified using a Millipore-MilliQ system. All other solvents were HPLC grade or the best commercially available grade, and were used without further purification.  $\text{Mn}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$  was purchased from Fluka and  $\text{Mn}(\text{CF}_3\text{SO}_3)_2 \cdot \text{CH}_3\text{CN}$  was synthesized by the reaction of manganese powder with trifluoromethane sulfonic acid which was diluted in  $\text{CH}_3\text{CN}$ . (CAUTION: This reaction is extremely exothermic and produces hydrogen gas. The triflic acid should be diluted at least 100-fold in acetonitrile and the manganese powder added slowly under nitrogen flow to flush away hydrogen as it is formed.)

Synthesis of  $[\text{14-aneN}_4]\text{MnO}]_2(\text{ClO}_4)_3 \cdot 2\text{H}_2\text{O}$  (1). A solution of 0.42 g (2.1 mmols) of 14-aneN<sub>4</sub> in 20 ml of water was added to a solution of 0.73 g (2.9 mmols) of  $\text{Mn}(\text{ClO}_4)_2 \cdot 6\text{H}_2\text{O}$  in 8 ml of water. A small quantity of reddish-brown hydrous manganese oxide formed and was removed by vacuum filtration. The resulting yellow-brown solution was allowed to remain undisturbed in contact with the atmosphere for several days. A

dark green microcrystalline product formed, which was collected by vacuum filtration and washed with diethyl ether to remove excess water. (CAUTION: The solid explodes upon heating. Perchlorate salts are often explosive under such conditions as mild shock and heat.) Calculated for  $[\text{Mn}(\text{C}_{10}\text{H}_{24}\text{N}_4)\text{O}]_2(\text{ClO}_4)_3 \cdot 2\text{H}_2\text{O}$  Mn, 12.53%; C, 27.39%; H, 5.98%; N, 12.78%; Cl, 12.13%. Found: Mn, 12.2%; C, 28.40%; H, 5.65%; N, 12.81%; Cl, 12.06%.

Crystal Growth and Analysis. Attempts to grow diffraction-quality crystals of the perchlorate salt of 1 produced blade-type crystals with insufficient thickness. Good crystals were obtained by adding a 10-fold excess of  $\text{Mn}(\text{CF}_3\text{SO}_3)_2$  to an aqueous solution of the macrocyclic ligand and allowing the green solution to stand undisturbed in a refrigerator for a week. The resultant shiny dark green crystals rapidly lose water of crystallization in a dry atmosphere, and it was necessary to seal them in thin-wall capillaries with a small amount of the mother liquor to prevent decomposition. A crystal with dimensions 0.20 x 0.22 x 0.40 mm was mounted in an Enraf-Nonius CAD-4 diffractometer and 5,432 unique intensity data were collected at room temperature.

The structure was solved by Patterson methods and refined via standard least-squares and Fourier techniques. One of the triflate anions was disordered through rotation around the carbon-sulfur bond and this was modeled with a combination of isotropic and anisotropic motions. In a difference Fourier map calculated following the refinement of all non-hydrogen atoms with anisotropic thermal parameters, peaks were found corresponding to the positions of all of the hydrogen atoms on the ligands. Hydrogen atoms were assigned idealized locations. They were included in structure factor

calculations, but not refined. The crystal was found to be monoclinic, belonging to space group  $P2_1/n$ , with  $a = 12.620(2)$  Å,  $b = 14.442(2)$  Å,  $c = 22.806(2)$  Å,  $\beta = 91.651^\circ$ ,  $V = 4.155$  Å<sup>3</sup>,  $d_{\text{calc}} = 1.64$  g/cm<sup>3</sup>. The R index for the data for which  $F^2 > 3\sigma(F^{2h})$  (4178 data) was 0.0481. The R value for all 5,432 data was .0916. Additional crystallographic data can be obtained as supplementary material.


## Results and Discussion

### Synthesis

Compound 1 is only the fourth species<sup>6a,7,8</sup> reported with a  $\text{Mn}_2\text{O}_2$  core and with crystallographically discrete Mn(III) and Mn(IV) atoms. Also, several mixed-valence complexes with manganese atoms bridged by two oxo groups and one acetate group have recently been reported.<sup>9,10,13</sup> Various synthetic routes have been used to make these compounds, but most involve oxidation of Mn(II) precursors. Oxidation has been accomplished by persulfate, permanganate, hydrogen peroxide or dioxygen. No suggestions have been put forward regarding the mechanism of formation of the  $\text{Mn}_2\text{O}_2$  core. This four-membered ring appears to be particularly stable and the steps in its assembly may have relevance to the nature of the oxygen-evolving center of photosystem II. Earlier work with <sup>18</sup>O-labeled water on the bipyridine analogue of 1 indicated that the oxo groups can be exchanged with solvent water in aqueous solution by recrystallization at elevated (80°C) temperature.<sup>14</sup> However, the original source of these bridging oxygens has previously not been established.

In the initial synthesis of the  $\text{Mn}_2\text{O}_2$  unit, oxidizing agents ( $\text{S}_4\text{O}_6^{2-}$ ,  $\text{MnO}_4^-$ ,  $\text{H}_2\text{O}_2$ ,  $\text{O}_2$ ) have been used that could donate oxygen atoms to form the bridge. However, our work supports the synthetic mechanism given in Scheme 1 in which water is the source of the bridging oxygens:

## Scheme 1



In this scheme, the initial step involves ligation of the  $\text{Mn}^{2+}$  ion by the macrocycle. This ligation has a large effect on the oxidation potential of the metal. An aqueous solution containing 1:1 equivalents of  $\text{Mn}^{2+}$  and macrocycle has a quasi-reversible oxidation wave in its CV at about  $-0.1$  V, while an aqueous solution of  $\text{Mn}(\text{ClO}_4)_2$  shows an irreversible oxidation at  $+0.8$  V. A solution containing only the macrocycle exhibits a broad, multielectron irreversible wave at  $+1.0$  V. A thoroughly deaerated solution of  $\text{Mn}^{2+}$  and the macrocyclic ligand is essentially colorless. Addition of dioxygen or other suitable oxidants produces a colored  $\text{MnL}^{3+}$  species.

Evidence establishing that the oxo bridges come from  $\text{H}_2\text{O}$  is provided by an electrochemical synthesis procedure. An acetonitrile solution was prepared that contained 2% by volume water, was 10 millimolar in  $\text{Mn}(\text{CF}_3\text{SO}_3)_2$ , 10 millimolar in ligand, and 100 millimolar in tetrabutylammonium hexafluorophosphate. This solution was deaerated and then oxidized at  $+0.34$  V (Pt gauze electrode, SCE reference). The colorless solution developed a green color over about 2 hours and gave the characteristic electronic spectrum of the III,IV di-oxo binuclear complex (see Fig. 3 and below). The only available source for the bridging oxo groups in this synthesis is water. This indicates that the role of dioxygen or other potentially oxygen atom donating oxidants in a conventional synthesis is simply that of an oxidizing agent.

This experiment also highlights the unusual tendency for formation of the  $\text{Mn}_2\text{O}_2$  core. It is remarkable that a binuclear species such as 1 will self-assemble from 10 millimolar solutions. The details of the actual bridge formation (step 3) remain unclear. Presumably the  $\text{MnL}^{3+}$  monomeric species contains two solvent molecules to raise the coordination number to six. In aqueous solution, these ligands would be water. At some point the water ligands must deprotonate to form the isolated complex 1.

The isolated complex  $[(14\text{-aneN}_4)\text{MnO}]_2^{3+}$  contains the 14-ane $\text{N}_4$  ligand coordinated in a cis fashion. In the literature, the vast majority of metal complexes isolated with the 14-ane $\text{N}_4$  macrocycle adopt the planar trans geometry.<sup>15</sup> In fact, even a metal atom as large as osmium(III) forms a trans- $[\text{Os}(14\text{-aneN}_4)\text{Cl}_2]^+$  complex.<sup>12</sup> Thus, it is quite surprising that we isolate a cis-coordinated 14-ane $\text{N}_4$  manganese complex. The di- $\mu$ -oxo  $\text{Mn}_2\text{O}_2$  core must trap the ligand in a cis-coordinated geometry. This points once again to the exceptional stability of this  $\text{Mn}_2\text{O}_2$  core.

An alternative route to the product of Scheme 1 might be to require oxidation of part of the monomeric  $\text{MnL}^{+3}$  to manganese +4 species, followed by dimerization involving Mn(III) and Mn(IV) monomers. Several observations argue against such a process. The first is that the cyclic voltammetry of 1 (see below) shows that it is much easier to oxidize the III,III dimer to the III,IV dimer than it is to oxidize the Mn(II)L monomer to the Mn(III) L monomer and to further oxidize the Mn(III)L monomer to the Mn(IV)L monomer. In fact, the electrochemistry of an aqueous solution containing  $\text{Mn}(\text{ClO}_4)_2$  and L exhibits no oxidation waves attributable to the conversion of Mn(III)L to Mn(IV)L prior to +1.5 V

vs. SCE. Another observation consistent with Scheme 1 is the relative kinetic inertness of the III,III dimer. In a spectroelectrochemistry experiment it was possible to reduce 1 to the III,III level and then to reoxidize to the original III,IV dimer (see below). The rate of the re-oxidation step was the same as the reduction which implies that the III,III dimer remains intact and does not immediately dissociate into monomer.

### Structure

Compound 1 is the fourth example<sup>16</sup> of a mixed-valence Mn(III),Mn(IV) di- $\mu$ -oxo species with crystallographically discrete Mn(III) and Mn(IV) atoms (Fig. 2). The ligands occupying the other coordination sites of Mn in the known compounds are bipyridine (bpy),<sup>6a</sup> N,N'-bis(2-methylpyridyl)ethane-1,2-diamine(bispicen),<sup>7</sup> 2,2'2''-tri-aminotriethylene (tren)<sup>8</sup> and the present 14-aneN<sub>4</sub> macrocycle. A comparison of the bond distances and bond angles of the compounds reveals the structural integrity of the Mn<sub>2</sub>O<sub>2</sub> core. The longest Mn(III)-oxo bond length is 1.861 Å in the present compound (Fig. 2) while the shortest is 1.833 Å in the bispicen; the range of Mn(IV)-oxo bond length is from 1.793 Å in the bispicen to 1.774 Å in the tren. Thus, the Mn<sub>2</sub>O<sub>2</sub> ring structure is not very sensitive to changes in the ligand environment of the Mn atoms. This fact is also reflected in the similarity of the optical and EPR spectra that have been reported for these complexes.

The structure of the [Mn(14-aneN<sub>4</sub>)O]<sub>2</sub><sup>3+</sup> ion is shown in Fig. 2. The structure is that of discrete Mn(III) and Mn(IV) ions with a planar di( $\mu$ -oxo)dimanganese core. The three-carbon segments of the ligands for the two manganese atoms are eclipsed rather than staggered as might have been predicted from steric considerations. The average Mn(III)-O bond

length is 1.862 Å versus 1.788 Å for Mn(IV)-O. There is a large difference between the axial Mn(III)-N bond distances, 2.349 Å average, and the equatorial Mn(III)-N average of 2.122 Å. This reflects the expected Jahn-Teller distortion for a high spin Mn(III)  $d^4$  ion.<sup>17</sup> The axial Mn(III)-N bond is exceptionally long, probably because of the degree of flexibility inherent in the 14-member macrocycle. Interestingly, the average bond lengths for the equatorial nitrogens are very similar (Mn(III)-N<sub>eq</sub> average 2.122 Å versus Mn(IV)-N<sub>eq</sub> average 2.113 Å). This similarity<sup>7</sup> has also been observed in the mixed-valence Mn(III),Mn(IV) di- $\mu$ -oxo compound with N,N'-bis(2-pyridylmethyl)ethane-1,2-diamine as a ligand. The authors claim that the similarity indicates partial delocalization in their compound. However, there are a number of factors that affect the equilibrium distances of the equatorial nitrogens of the two manganese atoms. The first is the oxidation state of the manganese; manganese(III)-nitrogen bonds are expected to be longer than manganese(IV)-nitrogen bonds. Further, the Jahn-Teller distortion of the Mn(III) will affect the equatorial as well as the axial bond lengths. With Jahn-Teller distortion the axial bonds lengthen and the equatorial bonds shorten. Another factor present is the oxo effect which acts to lengthen bonds trans to an oxygen group.<sup>18</sup> This trans-oxygen effect has been observed in the bpy<sup>6a</sup> and tren<sup>8</sup> compounds. The oxo effect is evident in the present case as manifested by the longer Mn(IV)-N bond trans to the  $\mu$ -oxo groups (2.113 Å) compared to 2.065 Å for the average axial distance. A final factor affecting Mn-N equatorial distances is the fact that the coordinating nitrogens are incorporated into a macrocycle, which puts constraints on movement. With all these factors influencing the observed bond lengths it does not

seem necessary to invoke delocalization in the  $\text{Mn}_2\text{O}_2$  core to explain similar Mn-N equatorial bond distances for the two Mn centers in 1.

#### Infrared Spectrum

The infrared spectrum of 1 was taken in a KBr pellet and exhibits a strong band at  $679\text{ cm}^{-1}$ . Bands in this region have been attributed to a vibration of the  $\text{Mn}_2\text{O}_2$  core.<sup>14</sup> When the synthesis reaction was run in 51%  $^{18}\text{O}$ -labeled water, the dark green crystalline product thus obtained showed a splitting of this band into two peaks, one at  $681\text{ cm}^{-1}$  and the other at  $670\text{ cm}^{-1}$ . This confirms assignment of the  $679\text{ cm}^{-1}$  band as due to  $\text{Mn}_2\text{O}_2$ . The bpy analogue of 1 shows the same behavior.<sup>14</sup>

#### EPR Spectrum

The EPR spectrum of 1 in 2:1 ethylene glycol:water at  $8^\circ\text{K}$  is very similar, feature for feature, with those reported for the bpy and tren analogues of 1 (Fig. 4). The 16-line spectrum is characteristic of strongly exchange-coupled ions of spin 2 [Mn(III)] and spin 3/2 [Mn(IV)], where the two isolated ions have nearly equal hyperfine coupling constants. The spectrum also shows a small asymmetry and a resolution of additional lines at the high field end of the spectrum because the hyperfine interaction is not small in comparison with the Zeeman interaction.<sup>6b</sup>

#### Electrochemistry

The cyclic voltammogram of 1 in dry  $\text{CH}_3\text{CN}$  is shown in Fig. 5. There is a reversible oxidation at +1.00 V and a reversible reduction at -0.12 V vs. SCE ( $E_p = 59\text{ mV}$ ,  $i_p^a/i_p^c = 1.0$ ). These oxidation and reduction steps are more reversible than those reported for the bpy, phen, bispicen and tren analogues of 1. This increased reversibility is



presumably due to the macrocyclic effect of the 14-aneN<sub>4</sub> ligand. The comproportionation constant  $K_c = [\text{III,IV}]^2 / [\text{III,III}][\text{IV,IV}]$  can be calculated<sup>19</sup> from the difference of the two potentials ( $\log K_c = 16.9 \Delta E_{1/2}$ ). The value for 1 is  $10^{19}$ , which is exceptionally large. The value reported for the phenanthroline analogue is  $10^{16}$  and values of  $10^{10}$  for the bispicen and  $10^{13}$  for the tren can be calculated from reported electrochemical data.

### Spectroelectrochemistry

The optical spectrum of 1 is given in Fig. 3. This spectrum is similar in some features to that observed for the  $[(\text{bpy})_2\text{MnO}]_2^{3+}$  compound. Electronic spectra of completely localized mixed-valence binuclear compounds (class I)<sup>20</sup> are expected to be the sum of the individual spectra of the components. In class II, or partially delocalized compounds, a new low-energy band appears, which is frequently attributed to a metal-to-metal charge transfer (intervalence) band.<sup>21</sup> A tentative assignment of the bands of 1 can be made from this classification scheme, spectra of monomeric Mn(III) and Mn(IV) complexes and prior assignment of the spectrum of the bpy complex.<sup>14</sup> The intense absorption tailing from the UV is assigned to ligand-to-metal charge transfer bands. There is a weak shoulder at roughly 560 nm and an asymmetric peak at 556 nm; somewhat similar features in the spectrum of the bpy complex have been attributed to d-d transitions.<sup>14</sup> A peak with an extinction coefficient of  $760 \text{ M}^{-1} \text{ cm}^{-1}$  is observed at 646 nm. This peak compares in position and intensity to charge transfer bands assigned to oxygen-to-Mn(IV) charge transfer transitions in monomeric compounds.<sup>22</sup> However, a band in this region for the bpy dimer has been

assigned to an oxo-to-Mn(III) CT transition.<sup>14</sup> The spectrum of the bpy dimer has a broad band at about 800 nm which has no parallel in either Mn(III) or Mn(IV) monomeric compounds and is attributed to an intervalence transition. Such a band is reported as being absent in the tren Mn(III)-Mn(IV) mixed valence complex although no absorption spectrum is shown.<sup>8</sup> In the present case no distinct near-IR peak appears but the 646 peak has a broad tail extending into this region. Compound 1 has an extinction coefficient of  $125 \text{ M}^{-1} \text{ cm}^{-1}$  at 800 nm. This broadened absorption may be due to an intervalence band that underlies other types of transitions.

Spectroelectrochemistry was employed to learn more about the electronic transitions and to investigate the stabilities of oxidized and reduced forms of 1. In dry acetonitrile it was possible both to reduce and to oxidize the III,IV mixed-valence compound by one electron. In both cases the process can be reversed by changing electrolysis potentials, and more than 95% regeneration of the original mixed-valence species can be obtained.

Reduction of a green 1 millimolar solution of 1 in  $\text{CH}_3\text{CN}$  at a platinum electrode ( $E = -0.26 \text{ V}$  vs SCE,  $0.1 \text{ M}$  TBAH) leads to an essentially colorless solution after passage of one equivalent of charge (Fig. 6a). When the potential is changed to  $0.34 \text{ V}$  vs. SCE, the current is reversed, and after one equivalent of oxidation, the original spectrum of the III,IV compound is obtained. In our cell the reduction and re-oxidation steps both take about six minutes. The ease of the re-oxidation step is taken as evidence that the reduced species retains the bridging oxo groups and is presumably the III,III compound. Oxidation of monomeric complexes to form the di- $\mu$ -oxo complex occurs on

a much longer time scale under similar conditions (ca. 2 hours). The much lower extinction coefficients for the III,III complex are consistent with d to d transitions and support the argument that the 648 nm band in the spectrum of 1 is an oxo-to-Mn(IV) CT transition.

This assignment of the 648 nm peak is supported by the spectrum that is obtained when a green 1.0 millimolar solution of the III,IV species is oxidized by one equivalent at +1.29 V versus SCE (Fig. 6b). A change from deep green to a more yellow green is discernible to the eye during the oxidation. This resulting solution can also be reduced rapidly to the original compound by lowering the potential to +0.34 V. The presumed IV,IV compound has lower intensity in the 650 to 800 nm region compared to the spectrum of 1, which is consistent with a broad IT band being present in this region in the mixed-valence compound and disappearing in the isoivalent IV,IV complex. The peak at 648 nm in 1 is shifted to 630 nm in the IV,IV compound and increases in intensity. This corroborates the assignment of the 648 nm band in 1 to an oxo-to-Mn(IV) charge transfer. In the process of the oxidation of the mixed-valence III,IV species, the peak at 555 nm decreases while a peak at 540 nm grows in. This feature at 540 nm is in the same region as transitions assigned as d-d in monomeric Mn(IV) Schiff base complexes.<sup>23</sup> A shoulder near 540 nm is also observed in the spectrum of the Mn(III),Mn(IV) mixed-valence compound, which is presumably due to a d-d transition associated with the Mn(IV) moiety.

The  $[(\text{phen})_2\text{MnO}]_2(\text{ClO}_4)_4$  compound (IV,IV) has been isolated<sup>24</sup> and characterized by X-ray crystallography.<sup>14</sup> Interestingly this species is reddish-brown in contrast to our observation that  $[(14\text{-aneN}_4)\text{MnO}]_2^{4+}$  is green. We have confirmed in a spectroelectrochemical oxidation in


acetonitrile that the visible spectrum of the red  $[(\text{phen})_2\text{MnO}]_2^{4+}$  consists of an intense absorption at 400 nm with weak shoulders at about 440, 510, and 675 nm. The reversibility of the oxidation of the phen binuclear species is not as complete as that of the 14-aneN<sub>4</sub> macrocycle. The difference in the visible spectra of the two Mn(IV),Mn(IV) species is presumably due to the difference in capping ligands. With the aliphatic secondary amine macrocycle (14-aneN<sub>4</sub>) a ligand-to-metal charge transfer (LMCT) band dominates the spectrum at wavelengths below 500 nm. With the more easily oxidized phenanthroline ligand this LMCT band has moved to longer wavelengths (beginning at ca. 650 nm) and thus the observed color is different.

Our spectroelectrochemical experiments are then consistent with the following assignments: Charge transfer bands, probably  $L \longrightarrow \text{Mn(IV)}$ , tail into the visible region from the UV. These appear in the spectrum of 1 and are shifted to lower energy in the oxidized IV,IV species. A peak at 540 nm for the IV,IV (shoulder in spectrum of 1) is attributable to a d-d transition for Mn(IV). A peak at 550 nm in the mixed-valence compound is probably a d-d transition centered at Mn(III). A prominent band at 648 nm in the spectrum of 1 (shifted to 630 nm in the IV,IV) is assigned to an oxo-to-Mn(IV) LMCT transition. A broad, underlying intervalence band causes a long tail into the near IR for the mixed-valence compound. This absorption feature is lost in the spectrum of the IV,IV species.


### Summary and Conclusions

The complex  $[(14\text{-aneN}_4)\text{MnO}]_2^{3+}$  is only the fourth mixed-valence complex prepared with a  $\text{Mn}_2\text{O}_2$  core that shows crystallographically discrete Mn(III) and Mn(IV) metal centers.<sup>6a,7,8</sup> The Mn-O distances are nearly equal ( $\pm 0.03$  Å) in all four complexes, while the Mn-N distances vary ( $\pm 0.2$  Å). This points to the integrity of the  $\text{Mn}_2\text{O}_2$  core. Other compounds with different capping ligands may be expected to be reported in the future. The tendency of manganese to form di- $\mu$ -oxo complexes of this type is quite interesting. The ligand 14-aneN<sub>4</sub> would normally coordinate in a planar trans geometry;<sup>12,15</sup> however, in our reaction the propensity of manganese to form the  $\text{Mn}_2\text{O}_2$  core leads to the isolation of a complex with a cis-coordinated 14-aneN<sub>4</sub> ligand.

The electrochemistry of our complex  $[(14\text{-aneN}_4)\text{MnO}]_2^{3+}$  is characterized by two reversible one-electron processes:



$$E_{1/2} = -0.12 \text{ V vs. SCE}$$



$$E_{1/2} = +1.00 \text{ V vs. SCE}$$

The 1.12 V range of stability of the mixed-valence complex is unusually large with respect to other examples in the literature.<sup>25</sup> Both the reduction of the III,IV complex to the III,III state and the oxidation to the IV,IV state are reversible in dry acetonitrile. This is in marked contrast to the electrochemical behavior of other binuclear Mn(III), Mn(IV) complexes in the literature.

The accessibility of the three oxidation states--III,III, III,IV and IV,IV--makes it possible to perform spectroelectrochemical experiments and make assignments of the electronic spectra. The results

of these experiments indicate that the intense band at 646 nm ( $\epsilon = 760 \text{ M}^{-1} \text{ cm}^{-1}$ ) in the III,IV complex is due to an oxygen-to-Mn(IV) ligand-to-metal charge transfer (LMCT) transition. This is in contrast to its previous assignment as an oxygen-to-Mn(III) LMCT band in the bipyridine complex.<sup>14</sup> Additionally, spectroelectrochemical results indicate that the tail into the near infrared is due to an intervalence transition that underlies the charge transfer band.

The complex  $[(14\text{-aneN}_4)\text{MnO}]_2^{3+}$  is remarkably stable. The spectrum of the complex in aqueous solution is unchanged at pH = 7.0 for 30 minutes. Acetonitrile solutions of the complex are stable for days in contact with the atmosphere. The use of this macrocyclic ligand for the formation of  $\text{Mn}_2\text{O}_2$  type complexes not only provides stability to the Mn(III),Mn(IV) binuclear complex but also to the one-electron reduced and one-electron oxidized forms of the complex.

With the use of electrochemical synthetic techniques, we have established that the bridging oxygens in the di- $\mu$ -oxo core come from water. This information is important since the incorporation of water oxygens into a manganese complex is desirable for water oxidation via such manganese complexes, and it may have implications for the oxygen evolving complex (OEC) in photosynthetic plants.

Acknowledgment: We owe special thanks to Dr. Melvin P. Klein for his cooperation during the use of his instrumentation for EPR. Also, we would like to thank Dr. F. J. Hollander for X-ray analysis and many helpful discussions about the results. This work was supported by the Office of Energy Research, Office of Basic Energy Sciences, Division of Chemical Sciences of the Department of Energy under contract DE-AC03-76SF00098.

References

1. Cotton, F. A., Wilkinson, G. Advanced Inorganic Chemistry, Fourth Ed., Wiley, New York 1980.
2. Lawrence, G. D., Sawyer, D. T. Coord. Chem. Rev., 27, 1978, 173-193.
3. For leading references, see: (a) Sheats, J. E., Czernuszewicz, R. S., Dismukes, G. C., Rheingold, A. L., Stubbe, J., Armstrong, W. A., R. H. Beer, Lippard, S. J., J. Amer. Chem. Soc., 1987, 109, 1435-1444. (b) Mabud, B., Tuchagnes, J.-P., Hwang, Y. T. Hendrickson, D. N., ibid, 1985, 107, 2801-2802. (c) Wiegardt, K., Bossek, U., Bonvoisi, J., Beauvillain, P., Girerd, J. J., Nuber, B., Weiss, J., Heinze, J., Angew. Chem., Int. Ed. Engl., 1986, 25, 1030-1033. (d) Gohdes, J. W., Armstrong, W. H., Inorg. Chem., 1988, 27, 1842-1846.
4. Dismukes, G. C., Photochem. Photobiol., 1986, 43, 99-115.
5. Kuwabara, T. Miyao, M. Murata, N., Biochim. Biophys. Acta, 1985, 806 283-289.
6. (a) Plaksin, P. M., Stoufer, R. C., Mathew, M., Palenik, G. J., J. Amer. Chem. Soc., 1972, 94, 2121-2122. (b) Cooper, S. R., Dismukes, G. C., Klein, M.P., Calvin, M., ibid, 1978, 100, 7248-7252.
7. Collins, M. A., Hodgson, D. J., Michelsen, K., Towle, D. K., J. Chem. Soc., Chem. Commun., 1987, 1659-2660.
8. Hagen, K. S., Armstrong, W. H., Inorg. Chem., 1988, 27, 967-969.
9. Li, X., Kessissoglou, D. P., Kirk, M. L., Bender, G. J., Pecoraro, V. L., ibid, 1988, 27, 1-3.
10. (a) Kulawiec, R. L., Crabtree, R. H., Brudvig, G. H., Schulte, G. K.,

- ibid, 1988, 27, 1309-1311. (b) Vincent, J. B., Christmas, C., Huffman, J. C., Christou, G., Chang, H.-R., Hendrickson, D. N. J. Chem. Soc. Chem. Commun 1987, 236-238.
11. Brewer, K. J., Liegeois, A., Otvos, J. W., Calvin, M., Spreer, L. O., J. Chem. Soc., Chem. Commun , in press.
  12. Che, C.-M., Cheng, W.-K., Lai, T.-F., Poon, C.-K., Mak, T.C.W., Inorg. Chem., 1987, 26, 1683-1689.
  13. (a) Wieghardt, K., Bossek, U., Zsolnai, L., Huttner, G., Blondin, G., Gireerd, J.-J., Babonneau, F., J. Chem. Soc., Chem. Commun., 1987, 651-653. (b) Bashkin, J.S., Schake, A. R., Vincent, J. B., Chang, H.-R., Li, Q., Huffman, J. C., Christou, G., Hendrickson, D. N. J. Chem. Soc., Chem. Commun. 1988, 700-702.
  14. Cooper, S. R., Calvin, M., J. Amer. Chem. Soc., 1977, 99, 6623-6630.
  15. See for example: Fabbrizzi, L., Comments. Inorg. Chem., 1985, 4(1), 33-54. Busch, D. H., Martin, L. Y., Sperati, C. R., J. Amer. Chem. Soc., 1977, 99, 2968-2981. Hancock, R. D., Thom, V. J., Shaikjee, M. S., Inorg. Chem., 1986, 25, 2992-3000 and references therein.
  16. Stebler, et al. have interpreted the orthophenanthroline analogue of 1 as a disordered system with discrete Mn(III) and Mn(IV) sites. Stebler, M., Lundi, A., Burgi, H.-B., ibid, 1986, 25, 4743-4750.
  17. Cotton, F. A., Wilkinson, G., Advanced Inorganic Chemistry, 4th ed., Wiley, New York, 1980, pp 678-682.
  18. Armstrong, W. H., Spool, A., Papaefthymiou, G. C., Frankel, R. B., Lippard, S. J., J. Amer. Chem. Soc., 1984, 106, 3653-3667.
  19. Tinemans, A. H. A., Timmer, K., Reinten, M., Kraaijkamp, J. G., Alberts, A. H., van der Linden, J. G. M., Schmitz, J. E. J.,



- Saaman, A. A., Inorg. Chem., 1981, 20, 3698-
20. Robin, M. B., Day, P., Adv. Inorg. Chem. Radiochem., 1967, 10,  
247-422.
21. Hush, N. S., Prog. Inorg. Chem., 1967, 8, 357-444.
22. Okawa, H., Nakamura, M., Kida, S., Bull. Chem. Soc. Jpn., 1982, 466-  
470.
23. Keissoglou, D. P., Li, X., Butler, W. M., Pecoraro, V. L. Inorg.  
Chem., 1987, 26, 2487-2492.
24. Goodwin, H. A., Sylva, R. N., Aust. J. Chem., 1967, 20, 619-637.
25. Richardson, D. E., Taube, H., Coordination Chem. Rev., 1984, 60,  
107-129.

Figure Captions

Figure 1. Spectroelectrochemical cell constructed of quartz utilizing a 1 cm quartz cuvette. The light beam of the spectrophotometer passes between two platinum strips and below the platinum mesh cylinder of the large surface area working electrode. The counter electrode compartment is separated by two quartz frits from the working electrode compartment. The working electrode compartment was deoxygenated by bubbling through a teflon needle and the other compartments by septa and needle techniques.

Figure 2. Structure of the cation  $[(14\text{-aneN}_4)\text{MnO}]_2^{3+}$  with 50% probability ellipsoids and showing the atom labeling scheme. Hydrogen atoms were located but are not shown. Pertinent bond distances (Å) and angles (deg): Mn(III)-Mn(IV) 2.741(1); Mn(III)-O1 1.855(2); Mn(III)-O2 1.868(2); Mn(IV)-O1 1.786(5); Mn(IV)-O2 1.790(2); Mn(III)-N1 2.336(3); Mn(III)-N2 2.129(2); Mn(III)-N3 2.362(3); Mn(III)-N4 2.116(3); Mn(IV)-N5 2.051(2); Mn(IV)-N6 2.111(3); Mn(IV)-N7 2.049(3); Mn(IV)-N8 2.115(3); O1-Mn(III)-O2 80.6; O1-Mn(IV)-O2 84.6; Mn(III)-O1-Mn(IV) 97.7; Mn(III)-O2-Mn(IV) 97.1; O1-Mn(III)-N1 95.8; O1-Mn(IV)-N5 95.6.


Figure 3. Electronic spectrum of  $[(14\text{-aneN}_4)\text{MnO}]_2^{3+}$  in acetonitrile. Assignments are discussed in the text: IT, Intervalence transition; CT, O  $\rightarrow$  Mn(IV) charge transfer; d-d, probably Mn(III).

Figure 4. X-band EPR spectrum of  $[(14\text{-aneN}_4)\text{MnO}]_2^{3+}$  in 2:1 ethylene glycol:water at 8K with the following instrument settings: Microwave power 2mW; modulation amplitude 10 G; modulation frequency 100 kHz.

Figure 5. Cyclic voltammogram of  $[(14\text{-aneN}_4)\text{MnO}]_2(\text{CF}_3\text{SO}_3)_3$  in  $\text{CH}_3\text{CN}$ ,  $0.1 \text{ M } [(\text{C}_4\text{H}_9)_4\text{N}]\text{PF}_6$ . Glassy carbon disk working electrode, Pt wire counter electrode,  $\text{Ag}/\text{Ag}^+$  reference electrode. Scan rate 100 mv/sec.



Figure 6a. Spectroelectrochemical results. — original spectrum of  $[(14\text{-aneN}_4)\text{MnO}]_2(\text{CF}_3\text{SO}_3)_3$  in  $\text{CH}_3\text{CN}$  ( $0.1 \text{ M } (\text{C}_4\text{H}_9\text{N})\text{PF}_6$ ); — — — after reduction with 1.05 equivalents at  $-0.5 \text{ V } (\text{Ag}/\text{Ag}^+)$  at a Pt mesh electrode, total elapsed time, 8 minutes; — • — after reoxidation with 1.05 equivalent at  $0.0 \text{ V}$ , total elapsed time 6 minutes.

Figure 6b. Spectroelectrochemical results. — original spectrum of  $[(14\text{-aneN}_4)\text{MnO}]_2(\text{CF}_3\text{SO}_3)_3$  in  $\text{CH}_3\text{CN}$  ( $0.1 \text{ M } (\text{C}_4\text{H}_9\text{N})\text{PF}_6$ ); — — — after oxidation with 1.05 equivalents at  $0.95 \text{ V } (\text{Ag}/\text{Ag}^+)$  Pt mesh electrode, total elapsed time 6 minutes; — • — after rereduction with 1.05 equivalents at  $0.0 \text{ V}$ , total elapsed time, 6 minutes.




XBL 887 2498

Fig. 1




XBL 887 2499

Fig. 2




XBL 887-2500

Fig. 3



| g = 2.0

Fig. 4



XBL 887-2502

Fig. 5







Fig. 6a

$[(14\text{-ane } N_4)\text{MnO}]_2(\text{ClO}_4)_3$  in  $\text{CH}_3\text{CN}$




Fig. 6b

*LAWRENCE BERKELEY LABORATORY  
TECHNICAL INFORMATION DEPARTMENT  
UNIVERSITY OF CALIFORNIA  
BERKELEY, CALIFORNIA 94720*