

UCLA

UCLA Previously Published Works

Title

Enzyme-free synthesis of natural phospholipids in water

Permalink

<https://escholarship.org/uc/item/2wv815zm>

Journal

Nature Chemistry, 12(11)

ISSN

1755-4330

Authors

Liu, Luping
Zou, Yike
Bhattacharya, Ahanjit
[et al.](#)

Publication Date

2020-11-01

DOI

10.1038/s41557-020-00559-0

Peer reviewed



Published in final edited form as:

Nat Chem. 2020 November ; 12(11): 1029–1034. doi:10.1038/s41557-020-00559-0.

Enzyme-free synthesis of natural phospholipids in water

Luping Liu¹, Yike Zou², Ahanjit Bhattacharya¹, Dongyang Zhang¹, Susan Q. Lang³, K. N. Houk², Neal K. Devaraj¹.✉

¹Department of Chemistry and Biochemistry, University of California, San Diego, CA, USA.

²Department of Chemistry and Biochemistry, University of California, Los Angeles, CA, USA.

³School of the Earth, Ocean, and Environment, University of South Carolina, Columbia, SC, USA.

Abstract

All living organisms synthesize phospholipids as the primary constituent of their cell membranes. Enzymatic synthesis of diacylphospholipids requires preexisting membrane-embedded enzymes. This limitation has led to models of early life in which the first cells used simpler types of membrane building blocks and has hampered integration of phospholipid synthesis into artificial cells. Here we demonstrate an enzyme-free synthesis of natural diacylphospholipids by transacylation in water, which is enabled by a combination of ion pairing and self-assembly between lysophospholipids and acyl donors. A variety of membrane-forming cellular phospholipids have been obtained in high yields. Membrane formation takes place in water from natural alkaline sources such as soda lakes and hydrothermal oceanic vents. When formed vesicles are transferred to more acidic solutions, electrochemical proton gradients are spontaneously established and maintained. This high-yielding non-enzymatic synthesis of natural phospholipids in water opens up new routes for lipid synthesis in artificial cells and sheds light on the origin and evolution of cellular membranes.

Cellular membranes composed of glycerophospholipids are found in all living organisms. Bacterial and eukaryotic membranes consist of diacylphospholipids, in which two ester linkages connect a polar head group to two hydrophobic tails¹. Cells synthesize diacylphospholipids through enzymatic acylation of lysophospholipids². As several enzymes involved in diacylphospholipid biosynthesis must themselves be membrane-bound for enzyme activity, it is unclear whether phospholipid membranes could have formed in the

Reprints and permissions information is available at www.nature.com/reprints.

✉ **Correspondence and requests for materials** should be addressed to N.K.D. ndevaraj@ucsd.edu.

Author contributions

L.L. and N.K.D. conceived the project. L.L. designed and performed the synthetic experiments. L.L., A.B. and D.Z. performed microscopy experiments. Y.Z. and K.N.H. performed the theoretical study. L.L., Y.Z., A.B., D.Z., S.Q.L., K.N.H. and N.K.D. analysed the data. L.L., Y.Z. and N.K.D. wrote the manuscript. S.Q.L. and K.N.H. assisted in writing and editing the manuscript.

Online content

Any methods, additional references, Nature Research reporting summaries, source data, extended data, supplementary information, acknowledgements, peer review information; details of author contributions and competing interests; and statements of data and code availability are available at <https://doi.org/10.1038/s41557-020-00559-0>.

Competing interests

The authors declare no competing interests.

Supplementary information is available for this paper at <https://doi.org/10.1038/s41557-020-00559-0>.

absence of advanced enzymatic machinery^{3–6}. Using wet–dry cycling, an enzyme-free reaction between ether mono-glyceride, a short-chain fatty acid, and phosphate in the presence of dicyanamide can yield glycerophospholipids that possess an ether linkage and a relatively short acyl chain⁷. Acylation of glycerophosphates can take place in the presence of a large excess of activated acyl imidazole derivatives, but the reactions require organic co-solvent due to the hydrophobicity of the acylating precursors and result in unnatural diacylphospholipids^{8,9}. Here we demonstrate that a combination of ion pairing and self-assembly allows transacylation reactions of lysophospholipids with acyl donors to afford natural diacylphospholipids in water. The high-yielding aqueous synthesis of natural phospholipids provides new possible routes to the origin of cell membranes and suggests that diacylphospholipids may have been incorporated before the evolution of complex biochemical machinery^{10,11}. Furthermore, we believe that the use of ion pairing and self-assembly in water to dramatically accelerate reactions and control reaction selectivity will have applications in green chemistry, for example, in the aqueous synthesis of peptides and glycans.

Results and discussion

In cells, lysophospholipid esterification requires acyl thioesters and the action of an acyltransferase. Inspired by acyltransferase reactions in lipid biosynthesis², and past hypotheses on the role of thioesters in the origins of life¹², we questioned whether synthetic acyl thioesters **2**, in lieu of acyl coenzyme A, could acylate 1-acyl-2-hydroxy-*sn*-glycero-3-phosphocholines **1** in water to give the desired natural phosphatidylcholines (PCs) **3** (Fig. 1a). An ionized polar head group on **2** would facilitate transacylation by increasing the long-chain acyl thioester's solubility in water and driving the self-assembly of micelles, possibly in combination with the lysophospholipid reactant. We initially investigated the synthesis of a naturally occurring phospholipid, 1,2-dioleoyl-*sn*-glycero-3-phosphocholine **3a** (DOPC)¹³, in aqueous solution via acylation of lysophospholipid 1-oleoyl-2-hydroxy-*sn*-glycero-3-phosphocholine **1a** (Fig. 1a). A water-soluble anionic thioester, sodium 2-(oleoylthio)ethane-1-sulfonate **2a**, was chosen as an oleoyl donor¹⁴. Only trace conversion was observed (<1%) and addition of acylation catalysts to accelerate the transacylation reaction did not have a substantial effect (Supplementary Table 1); however, to enable spontaneous membrane formation in water, acylation would have to be high-yielding and occur readily at near stoichiometric ratios of starting materials to prevent amphiphilic precursors from disrupting membrane formation.

To improve the yield of the desired diacylphospholipids, we sought a way to increase the yield of transacylation while minimizing the hydrolysis of the acyl donor. Transacylation reactions between acyl donors and alcohols proceed via a tetrahedral intermediate (Fig. 1b)^{15,16}. We reasoned that the observed poor reactivity in the transacylation reaction is probably due to the inability of acylation reagent **2a** to lower ΔG for the tetrahedral intermediate. We therefore sought to synthesize an acyl donor that would stabilize the reaction intermediate. In principle, a positively charged leaving group on the acyl donor **2** would lead to a favourable Coulombic interaction¹⁷ with the negatively charged phosphate group of lysophospholipid **1** (Fig. 1b), thus stabilizing intermediate **4** and accelerating the anticipated transacylation reaction through ion pairing. Based on this hypothesis, we

prepared 2-(oleoylthio)-*N,N,N*-trimethylethan-1-aminium chloride **2b**, which contains a positively charged quaternary amine head group (Fig. 1c). Precursor **2b** is accessible in one synthetic step from oleic acid and thiocholine. Compound **2b** could also be obtained using prebiotically relevant condensing agents such as dicyandiamide (Supplementary Fig. 1)¹⁸. Lysophospholipids such as lysophosphatidic acid can be obtained by the hydrolysis of cyclophosphate amphiphiles, which have been shown to be prebiotically synthesized from glycerol, fatty acid and diamidophosphate in the presence of imidazole¹⁹. Furthermore, modification of phosphatidic acid head group by choline under prebiotically relevant conditions has also been demonstrated²⁰. We evaluated the esterification reaction of lysophospholipid **1a** (0.5 mM) with **2b** (0.75 mM) in alkaline bicarbonate buffer (pH = 10.6) in the absence of additional catalyst; 94% conversion of lysophospholipid **1a** to DOPC **3a** was obtained after 5 h at 37 °C (Fig. 1c). By contrast, under the same reaction conditions, we observed only trace acylation of **1a** with either anionic **2a** or the charge-neutral acylation reagent **2c** (<1% conversion).

Our initial results suggested that ion pairing is involved in the aqueous transacylation of 2-lysophospholipid. Density functional theory calculations were performed to better understand the effect of ion pairing on lysophospholipid acylation²¹. The *n*-butyryl group was employed in place of the oleoyl group as a model system (Fig. 2). Calculations were performed at D3-B3LYP/6-311+ +G(2*d,p*)/B3LYP/6-31G (*d,p*) level with the Truhlar–Cramer empirical model (Solvation Model Density, SMD) for water²². Tetrahedral **INT1** and **INT2** resemble the corresponding 1,2-addition transition state and predict kinetic barriers; G for the formation of the tetrahedral intermediate **INT1** from model acylation reagent 2-(butyrylthio) ethane-1-sulfonate was calculated as 24.1 kcal mol⁻¹, whereas for **INT2** from 2-(butyrylthio)-*N,N,N*-trimethylethan-1-aminium it was 16.5 kcal mol⁻¹ (Fig. 2a). These computational results predict a higher reactivity of the quaternary ammonium-containing reagent analogous to **2b** in the transacylation reaction, in line with our experimental observations. Further, the optimized structures of the reaction intermediates indicate that intramolecular Coulombic interactions are important for the stability of the tetrahedral intermediate (Fig. 2b). In **INT1**, the sulfonate side chain is distal to the phosphate group, which results from the electrostatic repulsion of these two negatively charged groups. By contrast, the optimized structure of **INT2** shows electrostatic attraction between the negatively charged phosphate group and the two positively charged quaternary ammonium groups. Our experimental and theoretical findings are consistent with the hypothesis that a positively charged head group on **2** stabilizes the reaction intermediate, resulting in efficient transacylation.

We next explored the range of reaction conditions under which diacylphospholipids could be synthesized. The transacylation reaction between lysophospholipid **1a** and acyl donor **2b** proceeded rapidly and in high yield in alkaline bicarbonate buffers (pH 9.5–10.6) (Table 1)²³. Good reaction yields were also obtained in Na₂CO₃/NaHCO₃ buffer over a longer time period at pH = 8.8 (63% yield, 24 h, entry 3). By contrast, trace product was obtained using phosphate buffered saline at pH = 7.4 (<1% yield, 24 h, entry 1) or HEPES buffer at pH = 7.5 (<1% yield, 24 h, entry 2). Based on these results, we tested if the reaction could proceed in more complex naturally derived alkaline water samples. Numerous models have proposed

that life may have started near alkaline hydrothermal vents or soda lakes^{24,25}. We obtained water samples from the Lost City hydrothermal field (LCHF, pH = 9.1) and Mono Lake (pH = 10)—a Californian soda lake—to test whether synthesis of phospholipids could take place in such environments. Acylation of **1a** with **2b** to form phospholipid **3a** took place in water from the LCHF (42% yield, 48 h, entry 4) and Mono Lake water (76% yield, 5 h, entry 6). These results suggest that natural alkaline water sources would have been privileged environments for thioester mediated acylation of lysophospholipids.

Having optimized reaction conditions (entry 7, Table 1), we investigated the generality of the technique by attempting to synthesize a variety of naturally occurring phospholipids. We explored the acylation of multiple lysophosphatidylcholines with acylation agents of different chain lengths and compositions. Good acylation yields were achieved when **2b** was reacted with lysophospholipids that possess either unsaturated (oleoyl) or saturated (palmitoyl and stearoyl) acyl chains (**3b** and **3c**) (Table 2). Changing the acyl donor to a saturated palmitoyl group through the use of 2-(palmitoyl thio)-*N,N,N*-trimethylethan-1-aminium chloride **2d** led to similar reactivities (**3d–3f**) and enabled the synthesis of dipalmitoylphosphatidylcholine, a key component of pulmonary surfactant²⁶. We also tested whether we could synthesize phosphatidic acids, as they are universal intermediates in glycerophospholipid biosynthesis²⁷. Compared with lysophosphatidylcholine, which has a neutral zwitterionic headgroup, lysophosphatidic acid has a negatively charged phosphate headgroup, and acylation of the sodium salt of 1-oleoyl-2-hydroxy-*sn*-glycero-3-phosphate with **2b** led to a 55% yield of phosphatidic acid **3g**. We hypothesized that the lower efficiency might be due to the less facile deprotonation of the 2-hydroxy group of lysophosphatidate, or possibly a higher G for the tetrahedral intermediate. We further attempted to obtain various diacylphospholipids such as phosphatidylethanolamine, phosphatidylserine and phosphatidylglycerol by oleoylating the corresponding lysophospholipids; however, the reactivity of acylation reagent **2b** is insufficient for these lysolipids and no considerable yields of the desired diacylphospholipids were achieved (Supplementary Figs. 3–5).

This transacylation reaction turns out to be highly chemoselective, being unperturbed by the presence of non-amphiphilic biomolecules that possess nucleophilic functional groups. Under the optimized reaction conditions, no oleoylation products were observed when combining reagent **2b** with either serine, threonine or glycerophosphocholine as substrates (Supplementary Figs. 6–8), and less than <1% of acetylation product **3o** was observed when a non-amphiphilic acyl donor acetylthiocholine chloride was used for the acylation of **1a** (Supplementary Fig. 9). Excellent yields of product **3a** were obtained in the oleoylation reaction of lysolipid **1a** in the presence of serine (88% yield, 5 h, Supplementary Fig. 10) or disodium phosphate (91% yield, 5 h, Supplementary Fig. 11), but less than 10% yield was achieved with an additive sodium dodecyl sulfate, which is a well-known detergent (Supplementary Fig. 12). On the basis of these findings, we believe that the mixed micelles formed by the preorganization of the amphiphilic reagents accelerates the reaction rate and increases the selectivity of lipid synthesis. With this in mind, we next investigated shorter single-chain lipid precursors, which are more prebiotically plausible reactants but require higher concentrations to form micelles than longer chain

precursors. When we reacted the ten-carbon chain lysophospholipid 1-decanoyl-2-hydroxy-*sn*-glycero-3-phosphocholine **1e** (0.5 mM) with decanoyl donor **2e** 2-(decanoylthio)-*N,N,N*-trimethylethan-1-aminium chloride (0.75 mM) at 37 °C for 5 h, no 1,2-didecanoyl-*sn*-glycero-3-phosphocholine was observed (Supplementary Fig. 13); however, when we carried out a reaction between lysophospholipid **1e** (5 mM) near its critical micelle concentration (Supplementary Fig. 2C) and decanoyl donor **2e** (7.5 mM) above its critical micelle concentration (Supplementary Fig. 2F), 83% yield of 1,2-didecanoyl-*sn*-glycero-3-phosphocholine was obtained (Supplementary Fig. 13), providing further evidence that mixed micelle formation is necessary for the transacylation reaction to occur.

Enabling the synthesis of diacylphospholipids through esterification reactions in water should result in the spontaneous de novo formation of cell-like lipid membranes. This is due to the expected micelle to lamellar transition as single-chain lysolipids are converted to diacylphospholipids. We used time-lapse fluorescence microscopy to observe the de novo formation of vesicles during the optimized reaction (Fig. 3a). Neither lysophospholipid **1a** nor the oleoyl thioester **2b** formed vesicles when incubated alone in Na₂CO₃/NaHCO₃ buffer of pH = 8.8 at 37 °C (Supplementary Fig. 14). No visible structures were observed after initial mixing of **1a** (0.5 mM) and **2b** (0.75 mM) in Na₂CO₃/NaHCO₃ buffer of pH = 8.8 at 37 °C (Fig. 3a and Supplementary Fig. 15); however, after 10 min tubular structures appeared, and after 30 min spherical vesicles were observed (Supplementary Figs. 15 and 16). Likewise, large vesicles were formed when **1a** (0.5 mM) and **2b** (0.75 mM) were mixed in water collected from the LCHF (Fig. 3b and Supplementary Fig. 17A). Membrane-bound vesicles were also detected on mixing the reactants in Mono Lake water, and transmission electron microscopy was able to confirm the generation of vesicles (Fig. 3c and Supplementary Fig. 18C). We reasoned that both fusion of submicron-scale vesicles and growth of small vesicles through addition of nearby newly formed phospholipid led to the observed cell-sized vesicles (Supplementary Fig. 19).

All living organisms harness proton gradients across phospholipid membranes to generate energy. Vesicles formed from simpler amphiphiles such as fatty acids are unable to maintain proton gradients, suggesting that diacylphospholipid membranes appeared early in the origin of life²⁸. De novo phospholipid vesicle formation in alkaline solution results in an alkaline interior of formed vesicles. If the exterior solution is then exchanged for a more acidic media, a proton gradient might spontaneously form. To test whether this was possible, a water-soluble pH indicator dye, 8-hydroxypyrene-1, 3,6-trisulfonic acid (HPTS), was added before the reaction of **1a** with **2b** in the alkaline Mono Lake water (pH = 10); HPTS is highly fluorescent at alkaline pH, but >99% fluorescence is quenched at pH = 4.6 (Supplementary Fig. 20). After 5 h of reaction, fluorescence microscopy indicated that HPTS was successfully encapsulated in the in situ formed membrane vesicles (Supplementary Fig. 21A). To test whether a proton gradient could be maintained, the vesicle media was exchanged to citrate buffer (pH = 4.6) using spin-filtration. Fluorescence intensity was initially maintained, indicating the spontaneous formation of a proton gradient. Fluorescence intensity slowly diminished by 80% over 2 h (Fig. 3d). This finding demonstrates that diacylphospholipid membranes generated in alkaline water sources are capable of forming and maintaining a proton gradient over hours. It is tempting to speculate

that similar phenomena may have occurred in the early origin of membranes, perhaps as alkaline hydrothermal vent water was diluted in the more acidic water of the Hadean ocean^{29,30}. Such a scenario may have driven selection pressures for developing primitive catalysts for maintaining such proton gradients, which could have eventually been utilized to form chemical energy.

Methods

Synthesis of acyl thioesters.

4-Dimethylaminopyridine (15 mg, 0.12 mmol) and *N*-(3-dimethylaminopropyl)-*N'*-ethylcarbodiimide hydrochloride (EDC•HCl) (248 mg, 1.3 mmol) were added to a solution of fatty acid (1.3 mmol) in CH₂Cl₂ (10 ml) at 0 °C under argon; the reaction mixture was stirred at 0 °C for 30 min. The resulting solution was added dropwise to thiocholine chloride (188 mg, 1.2 mmol) at -78 °C under argon and the reaction was warmed to room temperature and stirred for 4 h. The reaction solvent was removed under reduced pressure. The residue was purified by column chromatography on silica gel. Full experimental details and characterization of new compounds can be found in the Supplementary Information.

Enzyme-free transacylation reactions between lysophospholipids and acyl donors to afford diacylphospholipids in water.

A freshly prepared Na₂CO₃/NaHCO₃ buffer of pH = 10.6 (8 ml) was added to a mixture of lysophospholipid (4 μmol) and acylation reagent (6 μmol) at 37 °C unless specified otherwise. The reaction mixture was tumbled at 37 °C for 5 h. The resulting mixture was adjusted to pH 7.0 and concentrated in vacuo. Purification was performed by column chromatography or preparative thin layer chromatography on silica gel. Full experimental details and characterization of new compounds can be found in the Supplementary Information.

pH gradient decay in the de novo formed vesicles.

The reaction was carried out by adding HPTS (0.25 mM) before the reaction of lysophospholipid **1a** (0.5 mM) with 2-(oleoylthio)-*N,N,N*-trimethylethan-1-aminium chloride **2b** (0.75 mM) in the presence of Mono Lake water (pH = 10). The reaction mixture was tumbled at 37 °C for 5 h. The media of vesicles was then exchanged with citrate buffer (pH = 4.6, in the same osmolarity as Mono Lake water) using spin-filtration, spontaneously generating a transmembrane pH gradient. Samples of the reaction mixture were taken at different time points, and fluorescence micrographs of vesicles were collected to monitor the maintenance of the pH gradient in the de novo-formed vesicles. Full experimental details and characterization of vesicles can be found in the Supplementary Information.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

N.K.D. acknowledges financial support for this work provided by the National Science Foundation (grant no. CHE-1254611) and the National Institutes of Health (grant no. DP2DK111801). K.N.H. acknowledges National Science Foundation (grant no. CHE-1764328), the National Institutes of Health, National Institute of General Medical Sciences (grant no. R01 GM109078), for financial support of this research. S.Q.L. acknowledges financial support for this work provided by National Science Foundation (grant no. OCE-1536702). Computation time was provided by the UCLA Institute for Digital Research and Education (IDRE).

Data availability

The data that support the findings of this study are available within the paper and its Supplementary Information.

References

1. Jackowski S, Cronan JE Jr & Rock CO Biochemistry of Lipids, Lipoproteins and Membranes Vol. 20, 80–81 (Elsevier, 1991).
2. Lands WEM Metabolism of glycerolipides: a comparison of lecithin and triglyceride synthesis. *J. Biol. Chem* 231, 883–888 (1958). [PubMed: 13539023]
3. Schmidli PK, Schurtenberger P & Luisi PL Liposome-mediated enzymatic synthesis of phosphatidylcholine as an approach to self-replicating liposomes. *J. Am. Chem. Soc* 113, 8127–8130 (1991).
4. Deamer DW & Boatman DE An enzymatically driven membrane reconstitution from solubilized components. *J. Cell Biol* 84, 461–467 (1980). [PubMed: 6769929]
5. Morris-Natschke SL et al. Synthesis of phosphocholine and quaternary amine ether lipids and evaluation of in vitro antineoplastic activity. *J. Med. Chem* 36, 2018–2025 (1993). [PubMed: 8336340]
6. Harayama T et al. Lysophospholipid acyltransferases mediate phosphatidylcholine diversification to achieve the physical properties required in vivo. *Cell Metabol.* 20, 295–305 (2014).
7. Hargreaves WR, Mulvihill SJ & Deamer DW Synthesis of phospholipids and membranes in prebiotic conditions. *Nature* 266, 78–80 (1977). [PubMed: 840303]
8. Fernandez-Garcia C & Powner MW Selective acylation of nucleosides, nucleotides, and glycerol-3-phosphocholine in water. *Synlett* 28, 78–83 (2017).
9. Bonfio C et al. Length-selective synthesis of acylglycerol-phosphates through energy-dissipative cycling. *J. Am. Chem. Soc* 141, 3934–3939 (2019). [PubMed: 30767518]
10. Szostak JW, Bartel DP & Luisi PL Synthesizing life. *Nature* 409, 387–390 (2001). [PubMed: 11201752]
11. Budin I & Szostak JW Physical effects underlying the transition from primitive to modern cell membranes. *Proc. Natl Acad. Sci. USA* 108, 5249–5254 (2011). [PubMed: 21402937]
12. de Ouve C The beginnings of life on earth. *American Scientist* 83, 428–437 (1995).
13. Ejsinga CS et al. Global analysis of the yeast lipidome by quantitative shotgun mass spectrometry. *Proc. Natl Acad. Sci. USA* 106, 2136–2141 (2009). [PubMed: 19174513]
14. Brea RJ, Cole CM & Devaraj NK In situ vesicle formation by native chemical ligation. *Angew. Chem. Int. Ed* 53, 14102–14105 (2014).
15. Bender ML Mechanisms of catalysis of nucleophilic reactions of carboxylic acid derivatives. *Chem. Rev* 60, 53–113 (1960).
16. McClelland RA & Santry LJ Reactivity of tetrahedral intermediates. *Acc. Chem. Res* 16, 394–399 (1983).
17. Raynal M, Ballester P, Vidal-Ferran A & van Leeuwen PWNM Supramolecular catalysis. Part 1: non-covalent interactions as a tool for building and modifying homogeneous catalysts. *Chem. Soc. Rev* 43, 1660–1733 (2014). [PubMed: 24356298]
18. Steinman G & Cole MN Synthesis of biologically pertinent peptides under possible primordial conditions. *Proc. Natl Acad. Sci. USA* 58, 735–742 (1967). [PubMed: 5233470]

19. Toparlak D, Karki M, Egas Ortuno V, Krishnamurthy R & Mansy SS Cyclophospholipids increase protocellular stability to metal ions. *Small* 16, 1903381 (2019).
20. Rao M, Eichberg J & Oró J Synthesis of phosphatidylcholine under possible primitive Earth conditions. *J. Mol. Evol* 18, 196–202 (1982). [PubMed: 7097779]
21. Frisch MJ et al. Gaussian 09, Revision A.02 (Gaussian Inc, 2016).
22. Marenich AV, Cramer CJ & Truhlar DG Universal solvation model based on solute electron density and on a continuum model of the solvent defined by the bulk dielectric constant and atomic surface tensions. *J. Phys. Chem. B* 113, 6378–6396 (2009). [PubMed: 19366259]
23. Delory DE & King EJ A sodium carbonate-bicarbonate buffer for alkaline phosphatases. *Biochem. J* 39, 245 (1945).
24. Kempe S & Degens ET An early soda ocean? *Chem. Geol* 53, 95–108 (1985).
25. Martin W & Russell MJ On the origin of biochemistry at an alkaline hydrothermal vent. *Phil. Trans. R. Soc. B* 362, 1887–1925 (2007). [PubMed: 17255002]
26. Perkins WR et al. Role of lipid polymorphism in pulmonary surfactant. *Science* 273, 330–332 (1996). [PubMed: 8662513]
27. Athenstaedt K & Daum G Phosphatidic acid, a key intermediate in lipid metabolism. *Eur. J. Biochem* 266, 1–16 (1999). [PubMed: 10542045]
28. Chen IA & Szostak JW Membrane growth can generate a transmembrane pH gradient in fatty acid vesicles. *Proc. Natl Acad. Sci. USA* 101, 7965–7970 (2004). [PubMed: 15148394]
29. Morese JW & Mackenzie FT Hadean ocean carbonate geochemistry. *Aquat. Geochem* 4, 301–319 (1998).
30. Macleod G, McKeown C, Hall AJ & Russell MJ Hydrothermal and oceanic pH conditions of possible relevance to the origin of life. *Orig. Life Evol. Biospheres* 24, 19–41 (1994).




Fig. 2 |. Predicted effects of thioester charge on phospholipid synthesis.

a, Energetics from B3LYP-D3 density functional theory calculations on transacylation reactions. Tetrahedral intermediates **iNt1** and **iNt2** resemble the transition states of the corresponding addition steps. **iNt1** is destabilized by over 7 kcal mol⁻¹ compared with **iNt2**. **b**, Optimized structures of reaction intermediates. The negatively charged sulfonate group of **iNt1** is distal from the phosphate group to avoid a disfavoured charge repulsion interaction, whereas the positively charged side chain of **iNt2** moves into a geometry that will increase the favorable interaction with the phosphate group to stabilize the tetrahedral intermediate. Distances are shown in ångströms.




Fig. 3 | Enzyme-free formation of phospholipid membranes.

The reaction was carried out by mixing 1-oleoyl-2-hydroxy-*sn*-glycero-3-phosphocholine **1a** (0.5 mM) and oleoylation reagent **2b** (0.75 mM) in different solvents at 37 °C. **a**, Fluorescence micrographs are shown: Na₂CO₃/NaHCO₃ buffer (pH = 8.8) was used as a solvent; samples were taken from the reaction mixture at different time points and were stained by using 0.1 mol% Nile red dye. Scale bars, 10 μm. **b**, A fluorescence micrograph is shown: Lost City vent fluid was used as a solvent; the sample was taken from the reaction mixture after 48 h and was stained by using 0.1 mol% Nile red dye. Scale bar, 5 μm. **c**, A negative staining transmission electron micrograph is shown: mono Lake water was used as a solvent; the sample was taken from the reaction mixture after 5 h. Scale bar, 200 nm. **d**, A fluorescence micrograph of vesicles formed in mono Lake water (pH = 10) containing the pH indicator dye HPTS, two hours after the external media was exchanged to citrate buffer (pH = 4.6). Scale bar, 10 μm.

Table 1 |Aqueous synthesis of phospholipid **3a** using different water sources

The reaction scheme shows the synthesis of phospholipid **3a** from lysophospholipid **1a** and acylation reagent **2b**. **1a** is a lysophospholipid with a phosphate group and a hydroxyl group. **2b** is an acylation reagent with a thiol group. The reaction is carried out in a buffer at 37 °C, resulting in the formation of **3a**, a phospholipid with two fatty acid chains, and a thiolate byproduct.

Entry	Solvent	pH	Time (h)	Yield (%)
1	PBS	7.4	24	<1 ^a
2	HEPES	7.5	24	<1 ^a
3	Na ₂ CO ₃ /NaHCO ₃	8.8	24	63 ^a
4	Lost City vent fluid	9.1	48	42 ^a
5	Na ₂ CO ₃ /NaHCO ₃	9.5	5	73 ^a
6	Mono Lake water	10	5	76 ^a
7	Na ₂ CO ₃ /NaHCO ₃	10.6	5	88 ^b


Reactions were carried out with lysophospholipid **1a** (0.5 mM) and acylation reagent **2b** (0.75 mM) to afford phospholipid **3a** in different aqueous sources at 37 °C. PBS, phosphate buffered saline.

^aHPLC yield.

^bIsolated yield.

Table 2 |

Scope of ion pairing-enabled synthesis of natural phospholipids



Reactions were carried out with lysophospholipids **1** (0.5 mM) and acylation reagents **2** (0.75 mM) to afford phospholipid **3** in $\text{Na}_2\text{CO}_3/\text{NaHCO}_3$ buffer (pH = 10.6) at 37 °C for 5 h.

^a Isolated yield.

^b 1.5 mM of reagent **2b** was used.