

UC San Diego

UC San Diego Previously Published Works

Title

Is Increased Sleep Responsible for Reductions in Myocardial Infarction During the COVID-19 Pandemic?

Permalink

<https://escholarship.org/uc/item/2xw1n6b4>

Authors

Advani, Ira
Gunge, Deepti
Banks, Sarah
et al.

Publication Date

2020-09-01

DOI

10.1016/j.amjcard.2020.06.027

Peer reviewed



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Is Increased Sleep Responsible for Reductions in Myocardial Infarction During the COVID-19 Pandemic?

The COVID-19 pandemic caused by the highly contagious SARS-CoV-2 virus has had devastating consequences across the globe. However, multiple clinics and hospitals have experienced a decrease in rates of acute myocardial infarction and corresponding cardiac catheterization lab activations, raising the question: Has the risk of myocardial infarction decreased during COVID? Sleep deprivation is known to be an independent risk factor for myocardial infarction, and sleep has been importantly impacted during the pandemic, possibly due to the changes in work-home life leading to a lack of structure. We conducted a social media-based survey to assess potential mechanisms underlying the observed improvement in risk of myocardial infarction. We used validated questionnaires to assess sleep patterns, tobacco consumption and other important health outcomes to test the hypothesis that increases in sleep duration may be occurring which have a beneficial impact on health. We found that the COVID-19 pandemic led to shifts in day/night rhythm, with subjects waking up 105 minutes later during the pandemic ($p < 0.0001$). Subjects also reported going to sleep 41 minutes later during the pandemic ($p < 0.0001$). These shifts led to longer duration of sleep during the COVID-19 pandemic. Before the pandemic, subjects reported sleeping 6.8 hours per night, which rose to 7.5 hours during the pandemic, a 44 minute or 11% increase ($p < 0.0001$). We acknowledge the major negative health impact of the global pandemic but would advocate for using this crisis to improve the work and sleep habits of the general population, which may lead to overall health benefits for our society. Published by Elsevier Inc. (Am J Cardiol 2020;00:1–2)

The COVID pandemic has had devastating consequences globally including impact on health and economics. However, some hospitals have seen a 38–49% decrease in presentations of acute myocardial infarction (MI).^{1–3} Whether this finding represents patients' avoiding medical care due to COVID, which would lead to greater numbers of cardiovascular deaths and higher rates of post-MI complications, versus a true reduction in incidence of MI via unknown mechanisms is unclear.^{1–4} Sleep deprivation is common in today's 24/7 society and has been independently associated with risk of incident MI. Furthermore, sleep has been importantly impacted during the pandemic, perhaps due to a lack of regular daily structure, and sleep deprivation is an independent risk factor for MI.⁵ We conducted a social media survey to assess potential mechanisms underlying observed reductions in MI. We used validated questionnaires to assess sleep patterns, inhalant use, anxiety and depression. We sought to test the hypothesis that sleep duration has increased in the time of COVID and may beneficially impact overall health.

Methods

Participants were recruited to participate in online surveys through widespread Twitter, Facebook, Craigslist and Reddit advertisements. Participants

were incentivized to complete both pre- and during-pandemic online surveys based on random lottery. We received responses to both surveys from 135 individuals of which 131 included complete data. Of these, 56% identified as female, 43% as male, and 1.5% as non-binary, with an age-range of 14 to 64 years.

Results

We found several effects of the pandemic on sleep. First, mean sleep onset was delayed from 11:30 PM to 12:11 AM (41 minutes, $p < 0.0001$). Second, mean awakening was delayed from 7:00 AM to 8:45 AM (105 minutes, $p < 0.0001$). Third, sleep duration increased by 11% (44 minutes, $p < 0.0001$). This finding was only true for non-inhalant users (non-smokers, nonvapers; $p < 0.0001$). Fourth, we observed no change in subjective sleep based on overall Pittsburgh Sleep Quality Index score. With regards to other potential cardiometabolic risk factors, we observed no decrease in tobacco and vaping consumption, with, if anything, consumption somewhat increased during the pandemic. These findings demonstrate possible increased sleep duration and delayed circadian phase without a major change in sleep quality. The impact of these changes on cardiometabolic health is unclear but would be predicted to be beneficial.

With regards to gender, females had longer sleep latency during the pandemic (28 vs 21 minutes, 95%CI 0.4046 to 13.69), suggesting that they may suffer from greater anxiety due to the pandemic and shelter-in-place orders. Both females and males had increased sleep duration (34 and 50 minutes, respectively). The greater increase for males is due to their average sleep duration being 6 hours 40 minutes prior to the pandemic, while females slept 6 hours 54 minutes prior. During the pandemic, both females and males reported sleeping 7 hours 30 minutes.

Discussion

In conclusion, we believe our findings are important as the increased sleep duration observed during the COVID pandemic may have important health benefits. Short sleep duration is common in today's 24/7 society. Sleep deprivation is also an epidemic and has been associated with incident MI, impaired glucose tolerance, increased obesity risk and pneumonia.^{5–8} The reported alleviation of sleep deprivation is likely to have important health benefits for some patients. In particular, males had a noteworthy increase in the duration of sleep (6 hours 40 minutes prior to the pandemic, increased to 7 hours 30 minutes in the setting of

stay-in-place orders) which may reduce their cardiovascular disease risk.

Regarding the reported reductions in MI, several mechanisms seem plausible. In theory patients may be reluctant to leave their homes and thus some patients with chest pain may stay home rather than being assessed in the emergency room.^{9,10} That is, the rate of MI may not have changed as much as might be suggested by the reduced numbers being seen in the emergency room. Others have suggested that work-related stress may be reduced during the stay-at-home orders. The global pandemic is a stressful time and indeed we found elevated Hospital Anxiety Depression Scale anxiety scores of 7.42 in this cohort. The third potential explanation which seems likely based on our findings is that the increased sleep duration that has been facilitated by the stay-at-home orders has contributed to reduction in sleep-deprivation associated cardiac risk. The fact that men are at increased risk of MI compared to women and reported greater improvements in sleep deprivation, based on our data, is consistent with our purported mechanism.

Our study has limitations. First, we used a prospective sample based on social media and thus have no real way to assess whether there was participation bias in those who responded. Thus, our conclusions are limited to the population studied. Second, we relied on self-report and thus some inaccuracy is possible if not likely. However, we expect such misclassification to be random and thus should bias towards the null hypothesis. Third, we did not assess other cardiometabolic risk factors such as lipids or body weight, but do not believe these improved during the pandemic. In fact,

many have referred to the “COVID 10” or “Quarantine 15” to reflect the increase in body weight that many experienced during the stay-at-home orders. Despite these limitations we view our findings as potentially important and worthy of further mechanistic research.

We would advocate for using this global crisis to improve work and sleep habits of the general population, which may lead to overall health benefits for our society.

Disclosures

The authors have no conflicts of interest to disclose. ResMed provided a philanthropic donation to UC San Diego.

Ira Advani, BS^{a,b}

Deepti Gunge, BS^{a,b}

Sarah Banks, PhD^c

Sagar Mehta^{a,b}

Kenneth Park, BS^{a,b}

Mitul Patel, MD^d

Atul Malhotra, MD^b

Laura E. Crotty Alexander, MD^{a,b,*}

^a Pulmonary Critical Care Section, Veterans Affairs (VA) San Diego Healthcare System, La Jolla, California

^b Division of Pulmonary, Critical Care and Sleep Medicine, Department of Medicine, University of California San Diego (UCSD), La Jolla, California

^c Department of Neurosciences, UCSD, La Jolla, California

^d Division of Cardiovascular Medicine, Department of Medicine, UCSD, La Jolla, California

2 June 2020

15 June 2020

1. Solomon MD, McNulty EJ, Rana JS, Leong TK, Lee C, Sung S-H, Ambrosy AP, Sidney S, Go AS. The Covid-19 pandemic and the incidence of acute myocardial infarction. *N Engl J Med* 2020.

2. Garcia S, Albaghdadi MS, Meraj PM, Schmidt C, Garberich R, Jaffer FA, Dixon S,

Rade JJ, Tannenbaum M, Chambers J, Huang PP, Henry TD. Reduction in ST-segment elevation cardiac catheterization laboratory activations in the United States during COVID-19 pandemic. *J Am Coll Cardiol* 2020;75:2871–2872.

3. Oriol Rodríguez-Leor BCA, Soledad O, Javier MM, José Ramón R, Ramón LP, Ana S, Ángel C, Rafael R, Ignacio C, Armando Pérez DP, Raúl M. Impact of the COVID-19 pandemic on care activity in interventional cardiology in Spain. *REC Interv Cardiol* 2020;2:82–89.

4. Huet F, Prieur C, Schurtz G, Gerbaud E, Manzo-Silberman S, Vanzetto G, Elbaz M, Tea V, Mercier G, Lattuca B, Duflos C, Roubille F. One train may hide another: acute cardiovascular diseases could be neglected because of the COVID-19 pandemic. *Arch Cardiovasc Dis* 2020;113:303–307.

5. Ayas NT, White DP, Manson JE, Stampfer MJ, Speizer FE, Malhotra A, Hu FB. A prospective study of sleep duration and coronary heart disease in women. *Arch Intern Med* 2003;163:205–209.

6. Mesarwi O, Polak J, Jun J, Polotsky VY. Sleep disorders and the development of insulin resistance and obesity. *Endocrinol Metab Clin North Am* 2013;42:617–634.

7. Mullington JM, Haack M, Toth M, Serrador JM, Meier-Ewert HK. Cardiovascular, inflammatory, and metabolic consequences of sleep deprivation. *Prog Cardiovasc Dis* 2009;51:294–302.

8. Patel SR, Malhotra A, Gao X, Hu FB, Neuman MI, Fawzi WW. A prospective study of sleep duration and pneumonia risk in women. *Sleep* 2012;35:97–101.

9. De Rosa S, Spaccarotella C, Basso C, Calabro MP, Curcio A, Filardi PP, Mancone M, Mercurio G, Muscoli S, Nodari S, Pedrinelli R, Sinagra G, Indolfi C, Societa Italiana di C, the CCUAig. Reduction of hospitalizations for myocardial infarction in Italy in the COVID-19 era. *Eur Heart J* 2020;41:2083–2088.

10. Tam CF, Cheung KS, Lam S, Wong A, Yung A, Sze M, Lam YM, Chan C, Tsang TC, Tsui M, Tse HF, Siu CW. Impact of coronavirus disease 2019 (COVID-19) outbreak on ST-segment-elevation myocardial infarction care in Hong Kong, China. *Circ Cardiovasc Qual Outcomes* 2020;13:e006631.

<https://doi.org/10.1016/j.amjcard.2020.06.027>