

UC Irvine

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health

Title

Tuberculous Pleural Effusion

Permalink

<https://escholarship.org/uc/item/2zk6n5q9>

Journal

Western Journal of Emergency Medicine: Integrating Emergency Care with Population Health, 13(4)

ISSN

1936-900X

Authors

Schlesinger, Shira A
Perera, Philips

Publication Date

2012

DOI

10.5811/westjem.2011.9.6846

Supplemental Material

<https://escholarship.org/uc/item/2zk6n5q9#supplemental>

Copyright Information

Copyright 2012 by the author(s). This work is made available under the terms of a Creative Commons Attribution-NonCommercial License, available at
<https://creativecommons.org/licenses/by-nc/4.0/>

Peer reviewed

Tuberculous Pleural Effusion

Shira A. Schlesinger, MD, MPH Los Angeles County + USC Medical Center, Department of Emergency Medicine,
Philips Perera, MD Los Angeles, California

Supervising Section Editor: Sean Henderson, MD

Submission history: Submitted June 29, 2011; Revision received August 15, 2011; Accepted September 19, 2011

Reprints available through open access at http://escholarship.org/uc/uciem_westjem

DOI: 10.5811/westjem.2011.9.6846

Pleural effusions are a common finding in emergency departments, with cytologic analysis traditionally required for definitive diagnosis. This article describes a classic sonographic appearance of tuberculous pleural effusion. [West J Emerg Med. 2012;13(4):313–314.]

CASE

A 33-year-old male with no previous medical history presented for evaluation of a left pleural effusion detected on chest radiograph at a local clinic. The patient had visited the clinic the previous week for flu-like symptoms that had since resolved. He denied productive cough, recent weight loss, or night sweats. His only risk factor for tuberculosis was recent immigration. On exam, the patient was a thin, comfortable-appearing man with an intermittent nonproductive cough, normal vitals, and normal pulse oximetry. Chest auscultation demonstrated decreased breath sounds at the left lung base. Bedside emergency department (ED) ultrasound revealed pleural thickening adjoining a complex pleural effusion with multiple thin septations (see video; online only). The patient was placed in respiratory isolation and admitted for tuberculosis treatment following acid-fast bacilli positive sputums.

DISCUSSION

Approximately 13,000 cases of tuberculosis are reported in the United States each year. Foreign-born and racial/ethnic minorities continue to bear a disproportionate burden of the disease.¹ ED physicians are likely to have primary contact with these and other individuals unlikely to receive timely care from other settings.

Pleural effusions are associated with fluid overload, tuberculosis, and malignancy, among other conditions. Previous authors have subdivided effusions into 4 types by sonographic appearance: anechoic, homogeneously echogenic, complex septated, and complex nonseptated.^{2,3} Studies and guidelines applying this scheme have demonstrated ultrasound to be a useful diagnostic aid, particularly in differentiating

tuberculous from other etiologies.^{4,5} Pleural thickening and a complex septated pattern, with fibrinous strands in the pleural space producing a weblike or branching appearance, has been strongly associated with tuberculosis.^{6–8} Chen et al⁹ found a 96% specificity for tuberculous pleural effusions when differentiating between tuberculosis and malignancy.

Bedside ultrasound examination of effusions is best performed using a combination of high- and low-frequency probes. A higher-frequency (10 MHz) probe gives a more detailed view of the effusion, while the 3-MHz lower-frequency probe provides a wider view. The probe should be positioned along the lateral chest wall over the effusion.

In evaluating a patient with a pleural effusion, increasing pretest probability of a specific etiology may eliminate unnecessary invasive procedures. Ultrasound appearance of a tuberculous pleural effusion in patients with low-to-moderate suspicion for the disease will assist in appropriate allocation of ED resources and rapid isolation from the general public. With the growing availability of bedside ultrasound, knowledge of this common appearance of tuberculous effusions can assist providers in rapidly stratifying and advancing care of otherwise challenging patients.

Address for Correspondence: Shira A. Schlesinger, MD, MPH, Los Angeles County + USC Medical Center, Department of Emergency Medicine, 1200 N State St, Rm 1011, Los Angeles, CA 90033.
E-mail: shira.schlesinger@gmail.com

Conflicts of Interest: By the WestJEM article submission agreement, all authors are required to disclose all affiliations,

funding, sources, and financial or management relationships that could be perceived as potential sources of bias. The authors disclosed none.

REFERENCES

1. Centers for Disease Control and Prevention. Trends in tuberculosis—United States, 2008. *MMRW*. 2009;58:249–253.
2. Yang PC, Luh KT, Chang DB, et al. Value of sonography in determining the nature of pleural effusion: analysis of 320 cases. *Am J Roentgenol*. 1992;159:29–33.
3. Lai YF, Su MC, Weng HH, et al. Sonographic septation: a predictor of sequelae of tuberculous pleurisy after treatment. *Thorax*. 2009;64:806–809.
4. Beckh S, Bolcskei PL, Lessnau KD. Real-time chest ultrasonography: a comprehensive review for the pulmonologist. *Chest*. 2002;122:1759–1773.
5. Hooper C, Lee G, Maskell N. Investigation of a unilateral pleural effusion in adults: British Thoracic Society pleural disease guideline 2010. *Thorax*. 2010;65:Sii4–17.
6. Tshibwabwa ET, Richenberg JL, Aziz ZA. Lung radiology in the tropics. *Clin Chest Med*. 2002;23:309–328.
7. Martinez OC, Serrano BV, Romero RR. Real-time ultrasound evaluation of tuberculous pleural effusions. *J Clin Ultrasound*. 1989;17:407–410.
8. Akhan O, Demirkazik FB, Ozmen MN, et al. Tuberculous pleural effusions: ultrasonic diagnosis. *J Clin Ultrasound* 1992;20:461–465.
9. Chen HJ, Hsu WH, Tu CY, et al. Sonographic septation in lymphocyte-rich exudative pleural effusions: a useful diagnostic predictor for tuberculosis. *Am J Ultrasound Med*. 2006;25:857–863.