

UCSF

UC San Francisco Previously Published Works

Title

Trans-activation, post-transcriptional maturation, and induction of antibodies to HERV-K (HML-2) envelope transmembrane protein in HIV-1 infection

Permalink

<https://escholarship.org/uc/item/2zx5d71z>

Journal

Retrovirology, 11(1)

ISSN

1742-4690

Authors

Michaud, Henri-Alexandre

de Mulder, Miguel

SenGupta, Devi

et al.

Publication Date

2014-12-01

DOI

10.1186/1742-4690-11-10

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed



RESEARCH

Open Access

Trans-activation, post-transcriptional maturation, and induction of antibodies to HERV-K (HML-2) envelope transmembrane protein in HIV-1 infection

Henri-Alexandre Michaud¹, Miguel de Mulder¹, Devi SenGupta¹, Steven G Deeks², Jeffrey N Martin³, Christopher D Pilcher², Frederick M Hecht², Jonah B Sacha^{4,5} and Douglas F Nixon^{1,6*}

Abstract

Background: Human Endogenous Retroviruses (HERVs) comprise about 8% of the human genome and have lost their ability to replicate or to produce infectious particles after having accumulated mutations over time. We assessed the kinetics of expression of HERV-K (HML-2) Envelope mRNA transcript and surface unit (SU) and transmembrane (TM) subunit proteins during HIV-1 infection. We also mapped the specificity of the humoral response to HERV-K (HML-2) Envelope protein in HIV-1 infected subjects at different stages of disease, and correlated the response with plasma viral load.

Results: We found that HIV-1 modified HERV-K (HML-2) Env mRNA expression, resulting in the expression of a fully N-glycosylated HERV-K (HML-2) envelope protein on the cell surface. Serological mapping of HERV-K (HML-2) envelope protein linear epitopes revealed two major immunogenic domains, one on SU and another on the ectodomain of TM. The titers of HERV-K (HML-2) TM antibodies were dramatically increased in HIV-1 infected subjects ($p < 0.0001$). HIV-1 infected adults who control HIV-1 in the absence of therapy ("elite" controllers) had a higher titer response against TM compared to antiretroviral-treated adults ($p < 0.0001$) and uninfected adults ($p < 0.0001$).

Conclusions: These data collectively suggest that HIV-1 infection induces fully glycosylated HERV-K (HML-2) envelope TM protein to which antibodies are induced. These anti-HERV-K (HML-2) TM antibodies are a potential marker of HIV-1 infection, and are at higher titer in elite controllers. HERV-K (HML-2) envelope TM protein may be a new therapeutic target in HIV-1 infection.

Keywords: HIV, Antibody, HERV, Endogenous retroviruses, Transmembrane, Envelope, Elite controllers, Alternative transcripts

Background

Human endogenous retroviruses (HERVs) comprise about 8% of the human genome [1]. Their ability to replicate or produce infectious particles is inhibited by host restriction [2,3] and they are now considered to be stably integrated, largely silent, and transmitted in a Mendelian fashion [4]. However, HERV-K (HML-2), which is the most recently integrated of the HERV families, exhibits more ability to

express proteins than other older HERV families [5,6]. The genome of HERV-K (HML-2), the gag, pol, pro, and env genes, are flanked by two Long Terminal Repeats (LTR,) and it is possible to express all the viral proteins under specific conditions [2,7]. HERV expression has been associated with autoimmune diseases [8-13] and cancers [14-19], and in these settings mRNA transcripts and proteins are found in blood or tissues. Despite their status as self-antigens, translated HERV products can induce an immune response that correlates with disease progression or regression in some cancers [20-25].

During HIV-1 infection we have previously shown re-activation of HERV-K (HML-2) [26,27]. The mechanisms leading to HERV-K (HML-2) expression are still being

* Correspondence: dnixon@gwu.edu

¹Division of Experimental Medicine, Department of Medicine, University of California, San Francisco, USA

⁶Microbiology, Immunology & Tropical Medicine, School of Medicine & Health Sciences, The George Washington University, Ross Hall 502, 2300 Eye Street, NW, Washington, DC 20037, USA

Full list of author information is available at the end of the article

elucidated but HIV-1 Vif and Tat protein have been implicated [27,28]. These studies strengthen the concept that HIV-1 specifically induces the transcription of HERV-K (HML-2) mRNA which results in the expression of HERV-K (HML-2) proteins in HIV-1 infected cells. We have shown that an anti-HERV-K (HML-2) cellular immune response is generated in HIV-1 infected patients (HIV^{POS}), significantly increased in elite controllers, and HERV-K specific T-cell clones can eliminate HIV-1 infected cells *in vitro* [26,27,29]. Whether HERV-K reactivation in HIV-1 infection leads to an anti-HERV-K (HML-2) antibody response is controversial [30].

The goal of this study was to identify B-cell epitopes present on the HERV-K (HML-2) Env protein, and ascertain how HIV-1 infection impacts anti-HERV humoral immunity. HERV-K (HML-2) Env is composed of three proteins, the signal peptide (SP), the surface unit (SU) and the transmembrane (TM) protein [31]. We identified two major immunogenic domains of HERV-K (HML-2) Env, one on SU and another on the ectodomain of TM. We found that HIV-1 preferentially modified the anti-HERV-K HML-2 TM antibody response. While the anti-SU antibody titer was largely unchanged between HIV-1 infected (HIV^{POS}) or uninfected subjects (HIV^{NEG}), the titers of anti-TM antibodies were dramatically increased in HIV^{POS}. Although the anti-HERV-K (HML-2) TM response correlated with HIV-1 plasma viral load in viremic non-controllers and was reduced during efficient HAART treatment, elite controllers who naturally suppress HIV-1 viremia have a higher titer antibody response against HERV-K (HML-2) TM compared to HIV^{NEG} or HAART-suppressed patients. We determined that HIV-1 infection modifies HERV-K (HML-2) Env mRNA expression, which leads to a fully N-glycosylated HERV-K (HML-2) transmembrane envelope protein on the cell surface. These data are consistent with our overall conceptual model that HIV-1 infection changes HERV-K (HML-2) expression and protein production within an infected cell.

Results

Identification of two linear antibody epitopes in HERV-K (HML-2) Envelope

To identify the immunogenic domains on HERV-K (HML-2) Env, we first mapped humoral linear epitopes on the HERV-K (HML-2) Env protein using a set of 172 "15mer" HERV-K (HML-2) Env peptides, in a peptide-based ELISA assay. We used sera from four HIV-1 infected subjects (HIV^{POS}) and two HIV-1 seronegative low risk healthy donors (HIV^{NEG}). We found two strong immunogenic domains, one with homology to a domain previously described on the HERV-K (HML-2) Env SU protein [9] (see Discussion), and one novel epitope on TM (Figure 1A). We then compared the responses using recombinant HERV SU or TM proteins to individual

peptides and observed concordance. Using HERV-K (HML-2) SU or TM recombinant proteins, and sera from HIV-1 positive or negative subjects, we observed a concordance of responsiveness between recombinant protein and peptide (Figure 1B). We confirmed that the humoral response directed against HERV-K (HML-2) Env was restricted to two linear epitopes, one on the SU protein and the other to a novel epitope on the TM protein, in both HIV^{NEG} and HIV^{POS} subjects.

Comparison of the HERV-K (HML-2) envelope SU and TM humoral responses in healthy donors and HIV-1-infected subjects

Using the two amino-acid sequences identified, we performed a serological screen of HERV-K (HML-2) Env responses in 80 chronically infected, untreated HIV-1-infected subjects (HIV^{POS}) and 40 HIV^{NEG} subjects, in a cross sectional study (subject characteristics are detailed in Table 1). No differences in titer for anti-HERV-K (HML-2) SU responses between HIV^{POS} and HIV^{NEG} were observed, with a mean of 1:180 and 1:190 respectively (Figure 2A). However, compared to uninfected HIV^{NEG}, the antibody response to HERV-K (HML-2) TM was significantly increased in HIV^{POS} subjects ($p < 0.0001$), with a mean of titer of 1:450 and 1:1370 respectively (Figure 2B). A comparison of within each HIV^{POS} subject anti-HERV-K (HML-2) TM and -SU responses indicated an exclusivity of response specificity, as those making anti-HERV-K (HML-2) TM responses in general did not make anti-HERV-K (HML-2) SU responses and vice versa (Figure 2C).

To determine whether any relationship existed between anti-HERV-K (HML-2) Env antibody responses and clinical progression, we assessed responses among 40 elite controllers and 40 viremic non-controllers. Although no significant difference was detected for the anti-HERV-K (HML-2) SU response (data not shown), non-controllers had a significantly greater titer of anti-HERV-K (HML-2) TM antibodies (mean 1:1600) compared to controllers (mean 1:1100). Both controllers and non-controllers had a greater anti-HERV-K (HML-2) TM response compared to the HIV^{NEG} subjects (mean 1:450) (Figure 2D). We then examined responses during antiretroviral treatment, and found that treatment did not significantly modify the anti-HERV-K (HML-2) SU titer (mean 1:190 off-HAART versus 1:230 on-HAART). However, successful treatment was associated with a decrease in the anti-HERV-K (HML-2) TM response (mean 1:1370 off-HAART versus 1:440 on-HAART), to a titer similar to that found in seronegative healthy donors (Figure 2E,F). We examined the potential relationships between responses, time from infection, plasma viremia and CD4+ T cell count. While there was no correlation between the anti-HERV-K (HML-2) SU response and plasma viremia or CD4+ T cell count, there was a direct correlation between the anti-HERV-K (HML-2)




Figure 1 Identification of two linear epitopes on HERV-K (HML-2) Env. (A) 4 sera from chronically HIV-1 infected patients (HIV^{POS}; black) and 2 sera from seronegative low risk healthy donors (HIV^{neg}; grey) were used for antibody epitope identification by ELISA. The 3 sub-units signal peptide (SP), surface-unit protein (SU) and trans-membrane proteins (TM) are represented by 172 redundant 15mers. The lines represent the average of duplicate signals (OD) for each individual. (B) Sera from patients have antibodies reacting only with the SU-peptide (HIV#1 and #2), only with the TM-peptide (HIV#3 and #4), or negative for both epitopes (HIV#5, #6 and HIV^{neg} #1), were used to confirm the signal obtained with the peptide-based ELISA. Plain columns represent the peptides and the hatched columns represent the recombinant protein; SU in white and TM in gray. The ELISA was performed in duplicate, and the error bars represent the SEM.

TM humoral response and the plasma viral load ($p = 0.0083$) (Figure 3A). Furthermore, there was an inverse correlation between the anti-HERV-K (HML-2) TM humoral response and CD4⁺ T cell count ($p = 0.003$) (Figure 3B). These cross sectional data, from 120 HIV^{POS} subjects, showed that only the response against HERV-K HML-2

TM is modified during HIV-1 infection. Although the anti-HERV-K (HML-2) TM response correlated with the presence or absence of HIV-1 viremia in non-controllers, HAART suppressed and HIV^{neg} subjects, a strong anti-HERV-K (HML-2) TM response was still detected in elite controllers, despite the absence of detectable viremia.

Table 1 Characteristics of study subjects


Participant category (n)	Median age (yr [IQR])	Gender ^a		Median CD4 ⁺ T cell count (cells/mm ³ [IQR])	Median HIV-1 viral load (copy/ml [IQR])
		M	F		
Elite controllers (40)	50 [44.25-55.75]	24	15	771 [500-1108]	<50
HAART suppressor (40)	50.50 [43.75-54.75]	31	8*	609 [452-807]	<50
Viremic non controllers (40)	39.50 [32.25-49]	35	3*+	444 [362-625]	40,474 [21,322-83,318]
HIV-1 negative (80) ^b	18-30 (18)	40	40	n/a	
	31-50 (24)				
	51-70 (34)				
	>70 (4)				

^aF, female; M, male.

^bSome information about the CD4⁺ T-cell count was not available (n/a). Only a range of age was available.

*One patient in this cohort was transgender male to female.


+One patient in this cohort was intersex.



Detection of HERV-K (HML-2) envelope specific B-cells


To assess the presence and frequency of HERV-K (HML-2) Env specific B-cells in one healthy donor and one HIV-1 infected subject (OP-115), we used SU and TM epitope-specific peptides conjugated to biotin in a flow cytometry

assay. For OP-115 we used three time points: an early time point (day 116 post-infection), a later time point (day 352 post-infection), and one time point when the patient was on anti-retroviral treatment (day 383 post-infection). The frequency of both anti-HERV-K (HML-2) SU and anti-



HERV-K (HML-2) TM-specific B cells among the CD19+ population in OP-115 were greater than those observed in the uninfected subject (4.11% vs. 0.73% for SU and 3.46% vs. 0.44% for TM for the early time point) (Figure 4A). However, in OP-115, the frequency of anti-HERV-K (HML-2) SU B cells remained stable over time, while the

frequency of anti-HERV-K (HML-2) TM B cells increased with sustained high viral load, before decreasing during antiretroviral therapy, with 3.46% of specific B-cells at day 116, 5.31% at day 352 and 3.62% at day 383 (Figure 4A). Using membrane IgD and CD27 expression to distinguish the maturity of the anti-HERV-K (HML-2) SU and anti-



HERV-K (HML-2) TM-specific B cells, we observed that the anti-HERV-K (HML-2) TM- and -SU-specific B cells had similar status at day 116 (Figure 4B). However, as the frequency of HERV-K (HML-2) TM-specific B cells increased over time, the subset of unswitched and switched memory B cells also increased two fold (Figure 4B). During antiretroviral therapy, the frequency of the specific anti-HERV-K (HML-2) TM B cells and the percentage of memory cells decreased to baseline post-infection level, on par with the anti-HERV-K (HML-2) TM antibody response observed in HAART treated patients. These results corroborate the serological studies, and demonstrate the induction of both anti-HERV-K (HML-2) TM B cells and anti-HERV-K (HML-2) TM antibody responses after HIV-1 infection, which decrease in frequency and titer respectively with successful antiretroviral therapy.

A longitudinal study of anti-HERV-K (HML-2) humoral response before and after HIV-1 infection

To better understand the effect of HIV-1 infection on HERV-K (HML-2) Env antibody responses, we monitored the anti-SU and anti-TM antibody responses in four patients in whom samples were available before and after HIV-1 infection. We found the anti-SU response decreased or remained undetectable after infection (Figure 5A) whilst the anti-TM response increased or remained stable (Figure 5B). The mean difference of antibody titer (after infection – before infection) showed a significant difference between TM and SU (Figure 5C) and confirmed the results of the cross-sectional study. These two anti-HERV-K (HML-2) Env antibody responses are differentially modulated during HIV-1 infection, and this occurs from the earliest stage of infection. We then monitored one subject




Figure 5 Kinetics of anti-HERV-K Env antibodies after HIV-1 infection. **A,B** Comparison of the anti-SU (**A**) and anti-TM (**B**) response before and after HIV-1 infection in 4 patients. The time between the two time points did not exceed 1 month. A 1:200 sera dilution was used for this assay. **C** The evaluation of the kinetics of the response was determined by the subtraction of the OD of the time point after infection by the OD of the time point before infection. A negative result meant a decrease of the response while a positive result meant an increased response. **D,E** Total IgG (**D**) or IgM (**E**) anti-HERV-K TM and SU were assayed by ELISA on the same plate for 5 time points (1:200 dilution) for the patient OP-1830. The thin black line represents the CD4+ T cell count (cells/mm³) and the hatched line the log₁₀ of the HIV-1 copy number/mm³. Statistical analyses were performed using the Mann & Whitney *t*-test; **p* < 0.05. Results are typical of three independent experiments.

(OP-1830) before and during acute infection, and after antiviral treatment. This subject had a rapid rise in HIV-1 plasma viremia which peaked at day 42, followed by a decrease when the subject started antiretroviral drug therapy at day 76 (Figure 5D). Before infection (day 0 on the graph), the subject had low anti-HERV-K (HML-2) SU and -TM antibody titers. These levels remained static at the time of the first detection of HIV-1 plasma viremia, estimated to be day 10 after infection (Figure 5D). At day 42, the anti-HERV-K (HML-2) TM response increased in parallel with HIV-1 plasma viremia, while the level of anti-SU antibodies decreased below the level of detection. After initiation of antiretroviral treatment, the anti-HERV-K (HML-2) TM response decreased in parallel with the decay of plasma viremia, but the anti-HERV-K (HML-2) SU response reappeared to reach the pre-infection titer, corroborating our previous observation (Figure 2E,F and Figure 4A,B). We observed that the anti-HERV-K (HML-2) TM IgM peaked at the zenith of HIV-1 viremia at day 42 while the anti-HERV-K (HML-2) SU IgM peak was detected after the treatment initiation, at day 76. These observations suggested that the TM response is tightly associated to active HIV-1 replication. Furthermore, the induction of anti-HERV-K (HML-2) TM IgM response during the peak of HIV-1 viremia suggested that HERV-K (HML-2) Env TM protein is preferentially expressed during infection.

Evidence of HERV-K (HML-2) envelope TM trans-activation and post-transcriptional maturation

To investigate whether HIV-1 modified HERV-K (HML-2) Env TM or SU protein expression, we designed primers and probes to detect HERV-K (HML-2) Env transcripts, with one pair of primers amplifying a domain coding for the HERV-K (HML-2) SU epitope (SU_{primers}), and another amplifying a domain coding for the HERV-K (HML-2) TM epitope (TM_{primers}). We measured HERV-K (HML-2) Env mRNA expression *in vitro* in a time course experiment at d0, d1 and d2 after HIV-1 infection of PBMCs. The expression of HERV-K (HML-2) Env was compared to β -actin mRNA expression. At d0, no HERV-K (HML-2) mRNA expression was detected. At d1, HERV-K (HML-2) Env transcripts were detected using the two pairs of primers (SU_{primers} and TM_{primers}) at a similar level. At d2, we observed an increased of transcription of HERV-K (HML-2) Env, but the amplicons detected by the TM_{primers} were overrepresented compared to the transcripts detected by SU_{primers} (Figure 6A). Cumulative data obtained from independent experiments showed that the quantity of transcripts detected using TM_{primers} was significantly superior (mean fold of 77 [4–130]) (Figure 6B). These *in vitro* experiments suggest that HERV-K (HML-2) Env mRNA is expressed under a truncated or mutated form in HIV-1 infected cells.


We then measured HERV-K (HML-2) Env expression using different primers designed for nested PCR amplifying domains coding for either SU or TM in plasma before and after HIV-1 infection in subject OP-1830. Nested PCR was performed on plasma-isolated viral RNA. There was a strong signal at day 42 (peak of HIV-1 viremia and anti-TM IgM) when TM_{primers} were used, while no HERV-K (HML-2) Env expression was detected using SU primers (Figure 6C). At d104, after treatment initiation and decrease in HIV-1 plasma viral load, HERV-K (HML-2) Env mRNA expression was detected with either TM or SU primers (Figure 6A). These results corroborated the *in vitro* results and suggested that the mechanisms involved in the modification of HERV-K (HML-2) mRNA expression *in vitro* occurred *in vivo* as well. These results suggest that HIV-1 induces HERV-K (HML-2) TM protein expression in infected cells.

To investigate HERV-K (HML-2) Env proteins expression in HIV-1-infected cells, we used Hela-T4 cells that are permissive to HIV-1. A commercially available antibody, previously described as detecting the precursor protein (70-100 KDa) and the mature form (about 40 KDa) of TM [32,33] was used to follow HERV-K (HML-2) Env protein expression (Figure 6D). Uninfected cells, used as control for basal protein expression, showed not TM expression (column 1, Figure 6D). Transfected cells with a plasmid coding for the whole HERV-K (HML-2) envelope were used as positive controls (column 2, Figure 6D). HIV-1 infection induced HERV-K (HML-2) Env precursor and, at a lower level, TM protein expression (column 3, Figure 6B). Using the N-glycosylation inhibitor Tunicamycin, we observed that the HERV-K (HML-2) Env protein expressed after HIV-1 infection had a lower MW and inhibited the expression of the mature form of TM (column 4, Figure 6B). These results indicate that HIV-1 infection induces the expression of a fully N-glycosylated HERV-K (HML-2) Env precursor protein and the transmembrane glycoprotein.

A study which reconstituted a functional TM from a consensus HERV-K sequence showed that the N-glycosylation state affects the membrane location and function of TM [32]. Using immunofluorescence on non-permeabilized cells we confirmed the expression of HERV-K (HML-2) Env TM on the cell surface of infected PBMCs (Figure 6E). Taken together, these data support the hypothesis that HIV-1 induces HERV-K (HML-2) Env TM expression on the cell surface.

Discussion

In this study we investigated how infection with HIV-1 influences HERV-K (HML-2) Env mRNA and protein expression, and assessed the antibody response to the two HERV-K (HML-2) Env subunits, the surface unit (SU) and the transmembrane (TM) protein. We found discordant



(See figure on previous page.)

Figure 6 Evidence of trans-activation and post-transcriptional modification of HERV-K (HML-2) Env TM. **A)** HERV-K (HML-2) Env mRNA expression was detected using primers designed to bind the SU domain or TM domain (SU_{primers} or TM_{primers} respectively) at 0, 1 and 2 days post infection. Copies of HERV-K (HML-2) Env detected by TM_{primers} (plain line) increased by the time of infection compared to the copies detected by SU_{primers} (dashed line). The graph is a representative experiment of 3 individual independent experiments. Copy number was determined as described previously, and β -Actin was used as reference gene [52]. **B)** The graph represents cumulative data from 3 independent experiments and shows the ratio of copies of HERV-K (HML-2) Env detected by TM_{primers}/copies detected by SU_{primers} from HIV-1_{LAI} infected PBMCs 2 days post-infection. Similar data were obtained using primary isolates (91US_4/R5 tropic and BK132/X4 tropic). **C)** Nested PCR. The figure shows the amplicons obtained after the second round of PCR (around 500pb). TM mRNA is over-expressed during the peak of viremia at d42. Antiviral treatment induced the transcription of both SU and TM mRNA at a similar level. HIV^{neg}: HIV-1 seronegative low risk donor; OP-1830: HIV-1 seroconverter patient (d0: before infection; d12, d42, d76: after infection without treatment, d104: after treatment); water: the non-template well. **(D)** Assumed precursor proteins at 75 to 90 kDa and TM subunits at 32 to 38 kDa are visible. HeLa-T4 cells are infected by HIV-1LAI in presence (4) or not (3) of 10 μ l/ml of tunicamycin. HERK-Env transfected cells (2) were used as positive controls for HERV-K TM expression. Uninfected-untransfected cells were used as control for endogenous HERV-K basal expression (1). **(E)** Representative images of HERV-K TM extracellular expression on PBMCs. HERM-1811-5 (anti-TM) mouse monoclonal antibody and goat anti-mouse Alexa555 (red) were used to detect extracellular expression of HERV-K (HML-2) TM.

antibody responses against the two HERV-K (HML-2) envelope protein domains, SU and TM, and determined that HIV-1 influence was at the level of mRNA expression. We showed that HIV-1 infected elite controllers had higher titers of anti-HERV-K (HML-2) TM antibodies, compared to HIV-1 infected patients on effective HAART or healthy uninfected donors. These findings suggest that the anti-HERV-K (HML-2) TM humoral response may play a role in HIV-1 pathogenesis.

The presence and quantity of the antibody response against HERV-K (HML-2) during HIV-1 infection is controversial [30]. Some studies have shown an increase of antibodies against HERV-K (HML-2) [34,35], while others saw no differences between HIV-1-infected and non-infected patients [20,36,37]. We had different results from others [34,35], which might be due to technical differences in methods. The first study describing an anti-HERV-K (HML-2) antibody response showed no difference between HIV-1 infected and uninfected patients [37]. They assessed the antibody response by western blot, detecting sera positive for recombinant SU. They found fifteen percent of HIV-1 infected patients were seropositive for SU, similar to our findings. Another study mapped the anti-SU response in healthy subjects and patients with autoimmune disorders and found a similar epitope on SU [9] (Additional file 1: Figure S1A). In one study, an anti-HERV-K (HML-2) TM antibody had been generated after immunization with a recombinant TM protein, and was directed to a similar epitope present on the ectodomain [33]. This observation, and the mapping obtained in a preceding study [9], agree with our findings that the antibody responses against HERV-K (HML-2) SU and TM are restricted to one peptide for each subunit.

We determined that the origin of the anti-HERV-K (HML-2) TM B-cell and humoral response during HIV-1 infection was the induction of the extracellular expression of the gp36, the transmembrane HERV-K (HML-2) envelope

protein subunit. We showed that *in vitro* HIV-1 infection led to the expression of the glycosylated HERV-K (HML-2) Env precursor. The state of glycosylation is crucial for the production of a functional glycoprotein HERV-K (HML-2) Env TM gp36, and a previous study had shown that fully glycosylated gp36 could reach the cellular membrane [32].

HERV-K (HML-2) SU and TM proteins are translated from a unique mRNA coding for the protein precursor. The detection of this unique mRNA using either SU_{primers} or TM_{primers} by qPCR gave non-concordant results. This could be explained by a difference of the primers binding efficiency. Similar number of HERV-K (HML-2) Env mRNA copies were detected using TM_{primers} or SU_{primers} at d1 post-infection *in vitro* and, *in vivo* with different set of TM_{primers} and SU_{primers}, when the HIV-1 viral activity was suppressed by the treatment. This suggests that TM_{primers} and SU_{primers} have equivalent efficiency. The increase of HERV-K (HML-2) Env mRNA copies detected by TM_{primers} only at d2 post-infection *in vitro* and d42 at the peak of viremia *in vivo* strongly suggests that an aberrant or truncated mRNA is generated during high HIV-1 transcriptional activity. It is well described that HIV-1 or lentiviral vector integration induces alternative splicing and aberrant transcripts [38-40]. Comparative analysis of transcriptomic profiles between HIV-1 infected and uninfected primary T cells revealed that a large number of host genes is virus-induced [38]. These virus-induced sequences are not restricted to coding mRNA but also affect non-coding mRNA [41]. Using next-generation sequencing, it has been determined that the number of altered host gene copies is low at 12 hours post-infection, but dramatically increases at the peak of HIV-1 transcriptional activity after 24 hours post-infection *in vitro* [41]. The chronology of HIV-1-induced host gene transcription supports our finding that the induction of an alternative HERV-K (HML-2) Env mRNA, undetectable with SU_{primers}, occurs at the peak of HIV-1 transcriptional activity. The anti-SU antibody response observed in

different pathological contexts, such as cancers, is often associated and correlated with an increase in SU protein expression (reviewed in [42]). In our study, our results suggest that HERV-K (HML-2) SU protein expression is poorly induced, because of the low mRNA expression induced by HIV-1. That would explain why we, and others, did not detect an increase in anti-HERV-K (HML-2) SU IgG titer. An alternative HERV-K (HML-2) Env mRNA may code for a dysfunctional SU protein that could be quickly degraded. This hypothesis would support the induction of a specific anti-SU IgM and B-cell responses, but the low level of protein induction would not be enough to induce and maintain a long lasting IgG response.

Although the presence of viral mRNA in the blood of patients suggests the release of viral particles by infected cells [28,43], the absence of a functional HERV-K (HML-2) SU protein makes the release of infectious HERV-K (HML-2) particles unlikely. However, we cannot exclude HERV-K (HML-2) mRNA inclusion in HIV-1 viral particles.

The amino acid sequence we identified as DWNTS (in which the N residue is known to be glycosylated [32]) is nestled between the two cysteines flanking the ectodomain (Additional file 1: Figure S1). This sequence is conserved among HERV-K families (Additional file 1: Figure S1), and is recognized by sera independent of disease stage. There is some sequence homology between the HERV-K (HML-2) TM epitope and an HCV polyprotein (Additional file 1: Figure S1). However, we found no significant difference between HIV-1 positive HCV positive or HIV-1 positive HCV negative subjects in their anti-HERV-K (HML-2) TM response (Additional file 2: Figure S2). The SU epitope also shares some limited homology with the HIV-1 Tat protein (Additional file 1: Figure S1), but the absence of significant increase of anti-HERV-K (HML-2) SU response in HIV-1 infected patients or controllers suggests a limited impact of any potential cross-reactive anti-Tat activity.

To understand the potential role of HERV-K (HML-2) during HIV-1 disease progression, we compared anti-HERV-K (HML-2) TM responses in elite controllers and individuals on effective antiretroviral therapy. Although the level of viremia in these two groups are low, the level of residual HIV-1 replication (as compared to simply production) is known to be higher in controllers than antiretroviral drug treated subjects. We found higher anti-HERV-K (HML-2) TM antibody responses in controllers, suggesting ongoing rounds of HIV-1 replication drive HERV-K (HML-2) expression. An alternative explanation is that a higher level of immune activation in elite controllers compared to antiretroviral-treated subjects or healthy donors, might drive antibody production independent of the level of HERV-K protein production, although as we observed no differences in SU titer this explanation is unlikely. The ratio of anti-HERV-K (HML-2)

TM/total IgG and anti-SU confirmed the cross-sectional study, and showed that the TM humoral response was not increased due to polyclonal expansion. Thus, elite controllers have anti-HERV-K (HML-2) TM antibody production even in the absence of detectable viremia (Additional file 3: Figure S3).

Although the only prior study that demonstrated TM protein expression was during pregnancy in the extravillous cytotrophoblast, TM seems to be absent from the cellular membrane at a normal physiological state [33,42]. This raises the possibility that HERV-K (HML-2) may prove to be a marker of an HIV-1 infected cell, and thus could be considered as an HIV-1-Associated Neo Antigen (HANA).

In assessing a potential role for anti-HERV-K (HML-2) TM antibodies in pathogenesis, we observed a correlation between the anti-HERV-K (HML-2) TM response and loss of CD4⁺ T cells for viremic non-controllers suggesting that during viremic rebound anti-TM antibodies could target HERV-K (HML-2) Env TM-expressing HIV-1 infected CD4⁺ cells and accelerate CD4⁺ T cell depletion through mechanisms such as antibody-dependent cell-mediated cytotoxicity or complement-dependent cytotoxicity. However, the presence of an intermediate high anti-HERV-K (HML-2) TM level in elite controllers in the absence of detectable viremia suggests that the antibody response could also play a role in the control of virus through similar mechanisms.

The expression of the HERV-K (HML-2) Env TM protein during HIV-1 infection could have a direct role in immunopathogenesis. Some studies have shown that retroviral transmembrane glycoproteins such as HERV-K (HML-2) TM contain an immunosuppressive domain that inhibits lymphocyte proliferation and plays a role in the immune escape [33,42,44-47]. The increase of expression in viral non-controllers might be linked to progression of the disease. Antibodies present in elite controllers might help to inhibit a TM immunosuppressive effect.

The use of HERV-K (HML-2) proteins as tumor- or viral-associated antigens has already been investigated in different models. HERV-K (HML-2) Env elicits antibodies in patients with breast cancer [19,24] and melanoma [21]. A mouse monoclonal antibody directed against SU showed strong anti-tumor activity *in vitro* and *in vivo* in a mouse model [25,45,48]. In melanoma, pancreatic or prostate cancer, the tumor-associated antigen HERV-K-MEL has been proposed as a specific target for tumor cells [16,23,49]. One concern with a HERV-based approach to vaccination or therapeutic treatment is autoimmunity and immunopathogenesis. However, we have demonstrated the immunogenicity and safety of an endogenous retrovirus vaccine in non-human primates. The vaccine, a combination of DNA and adenovirus coding for Simian-ERV-K, induced specific T-cell and B-cell responses, including an anti-TM

antibody response [50]. Thus, ERV-K Env-specific antibodies may not be inherently detrimental to the host due to the restricted expression of ERV-K Env.

Conclusions

In summary, we have found a novel HERV-K (HML-2) Envelope TM neo-antigen over-expressed in the context of an HIV-1 infected cell, and this stimulates a specific antibody response against HERV-K (HML-2) Envelope TM. Our determination that naturally aviremic HIV-1 infected “elite” controllers have higher titers of this antibody compared to HAART-treated aviremic patients suggests that this antibody response may be of importance in viral control. We speculate that anti-HERV (HML-2) Env TM antibodies could also target infected HIV-1 infected latent “reservoir” cells. As this epitope is highly conserved and not subject to mutation, these findings could lead to a new approach to HIV-1 vaccines or immunotherapy.

Methods

Study populations

Samples of peripheral blood mononuclear cells (PBMCs) were selected from participants in two different San Francisco-based HIV-1-infected cohorts: OPTIONS (n = 5) and SCOPE (n = 120). Samples from HIV-1-negative controls were obtained from the Blood Center of the Pacific of San Francisco (n = 80). The study was approved by the local institutional review board (University of California San Francisco Committee on Human Research), research conducted according to the Declaration of Helsinki, and individuals gave written informed consent. Studies were performed on cryopreserved PBMCs and sera.

PBMC and sera samples were obtained from the following categories of chronically HIV-1-infected individuals: 40 elite controllers (EC: naive for treatment, undetectable viral load for two years, CD4 > 350); 40 highly active antiretroviral therapy (HAART: Viremic suppressed with undetectable viral load for at least two years, CD4 > 350), and 40 untreated virologic non-controllers (naive for treatment, viral load >2000 copies/mL).

HIV-1_{LAI} stock virus

Stocks of HIV-1_{LAI}, a CXCR4-tropic laboratory strain, were obtained from the AIDS Research and Reference Reagent Program and amplified on stimulated PBMCs for 7 days [51,52]. HIV-1-infected cells were pelleted at 3,000 rpm for 20 min, and supernatant fluid was passed through a 0.2- μ m filter and frozen in aliquots at -80°C. The titers of stocks were determined using TZM cells [53-56].

Ex vivo mRNA isolation nested RT-PCR

Viral mRNA was isolated from 140 μ L of plasma using QIAamp® Viral RNA Mini Kit from Qiagen. mRNA obtained were directly used in One-Step RT-PCR Kit (Qiagen)

according to the manufacturer. Briefly, 5 μ l of viral RNA equivalent to 15 μ l of plasma was reverse transcribed at 50°C for 30 min. The 1st round PCR was performed in 20 cycles consisting of 94°C for 30 sec; an annealing step at 55°C for both pairs of outer primers (SU Forward: GTATCAATGGTGGTAAGTCTCC; SU Reverse: CACTGCAATTAAAGTAAAAAT; TM Forward: GCCATTTTATACTRTCGTCCTAA; TM reverse: GACAAAACCRCCATC GTACTCAT) for 30 sec; and an extension step of 90 sec at 72°C. PCR product were next diluted at 1:50 and used as template for the 2nd round PCR. The second round was performed using Phusion® High-Fidelity PCR Master Mix in 35 cycles consisting of 98°C for 10 sec; an annealing step at 60°C for both pairs of inner primers (SU Forward: TGGATAATCCTATAGAARTAT; SU Reverse: TATGTTTGTCTAAACTTTCTGT; TM Forward: GCTGTAGCAGGAGTTGCATTG; TM reverse: TAATTGTAGTACTTCCAATGGTC) for 30 sec; and an extension step of 60 sec at 72°C. PCR products were separated on 1% agarose gels.

In vitro mRNA isolation and Q-PCR

mRNA was isolated from PBMCs using RNeasy Mini Kit (Qiagen) with on-column DNase treatment (Qiagen RNase-Free DNase Set) and eluted in 30 μ l of RNase-free water according to the manufacturer. A second step of DNase treatment using TURBO™ DNase (Life Technologies) was performed to eliminate efficiently DNA contaminants. qPCR was performed using TaqMan® One-Step RT-PCR or TaqMan® Universal PCR Mastermix for no-RT control. PCR was performed with one step at 48°C for 30 minutes, 95°C for 10 minutes and 40 cycles consisting of 95°C for 15 sec; an annealing/extension step at 60°C for both pairs of primers (SU Forward CCTGCAGTCCAAAATTGGTT; SU Reverse GCCACACATTCTTCCCAAAC; SU Probe CTCAGGCCACGGGTAAATTA; TM Forward GTTGCCTAAAGCCCCCTTAT; TM Reverse CCCTCTCTTGCTCTCACCAG; TM Probe AATTGGCAACACCGTATTCTG; β -Actin Forward GAGCGGGCTACAGCTT; β -Actin Reverse TCCTTAATGTCACGCACGATTT; β -Actin PROBE ACCACCACGGCCGAGCGG) for 60 sec. Thermal cycling was performed using a StepOne™ Real-Time PCR System (Applied Biosystems). Data was analyzed using StepOne™ Software (Applied Biosystems). Gene expression and fold induction was determined using the comparative Ct method [57].

Plasmid and recombinant proteins

HERV-K (HML-2) Env nucleotide sequence of HERV-K (HML-2) 102 was cloned in a pCDNA3.1 plasmid by CellFree Sciences Co. (Japan). The sequence presents high homology (>98%) with the major HERV-K (HML-2) family members as presented in Additional file 1: Figure S1.

SU and TM sequences were cloned in a pGAEx vector (GENEART vector). Both proteins were produced in HEK293

cells and purified 6 days post transfection by Ni-HiTrap columns by GENEART, Burlingame, CA.

Transfection and HIV-1_{LAI} infection/N-glycosylation inhibition

Hela-T4 [58] were plated in 12 well plates at 0.8×10^6 cells/well and transfected with HERV-K (HML-2) Env coding plasmid using Lipofectamine™ 2000 (Invitrogen) according to the manufacturer's protocol. Briefly, plasmid and Lipofectamine™ were mixed at a 1:2 ratio for 20 min at RT, and incubated with cells for 16 h. Untransfected and transfected cells were washed with PBS and infected with HIV-_{LAI} with 10 µg/ml of tunicamycin (Sigma) or grow medium for 16 h.

Western blot

Hela T4 cells were lysed in presence of anti-protease cocktail (Sigma) in n-Dodecyl β-D maltoside (Sigma) diluted at 0.1 mg/ml according the manufacturer protocol in dH₂O/0.05 M-TRIS HCl/0.15 M-NaCl lysis buffer. Prior to loading, the samples were mixed with lamely buffer 2× and boiled at 95°C for 5 minutes. Approximately, 25 µg of total proteins assayed with BCA Proteins assay kit (Thermo Scientific) were loaded on 4-16% gradient precast gels (Pierce). PVDF (Biorad) membranes were blocked 1 h at RT in PBS/0.05%-Tween 20/10%-non fat dry milk and incubated with mouse monoclonal anti-HERV-K TM HERM-1811-5 (Austral Biologicals) in PBS/0.05%-Tween 20/5%-nonfat dry milk at 1/1000 over-night at 4°C. Membranes were then washed 3 times in PBS/0.1%-Tween 20 and incubated with an HRP-conjugated anti-mouse (Abcam) 2 hours at room temperature. After 6 washes, the membranes were incubated with ECL-plus (GE Healthcare) and exposed at different time points on Kodak® Biomax™ MR film.

PBMC infection

Fresh PBMCs were isolated by standard Ficoll-Hypaque density gradient centrifugation on fresh blood samples and immediately cryopreserved in fetal calf serum (HyClone, Logan, UT) containing 10% DMSO (Sigma Aldrich, St. Louis, MO) in liquid nitrogen. PBMCs were stimulated with 2 µg/ml of phytohemagglutinin (PHA-L; Sigma, St. Louis, MO) for 48 h in RPMI-10% FBS complemented with IL-2 70U/ml before the addition of HIV-1_{LAI} at a MOI of 0.005 in RPMI-10% FBS.

Immunofluorescence

PBMCs were fixed in PBS/PFA 2% (Electron Microscopy Sciences) and coated on slides using Cytospin (300 rpm 2 minutes), washed twice with PBS and blocked 15 minutes in PBS/0.5%-BSA/10%-Goat serum. Cells were incubated with mouse anti-HERV-K TM (HERM-1811-5, Austral Biologicals) at 1:50, dilution in a humidified dark

room at room temperature for 1 hour and then washed 5 times with PBS. Then, the cells were incubated with an Alexa-555 anti-mouse IgG (Invitrogen) at 1:500 in a humidified dark room at room temperature for 1 hour and then washed 5 times with PBS. DNA was then stained with DAPI (0.5 µg/ml) for 5 minute at RT and the cells washed 5 times in PBS and 1 time with dH₂O. Coverslips are next mounted with PermaFluor (Thermo Electron Corporation) on microscope slides (Fischerbrand) and dried in the dark. Slides were analyzed on a LEICA DM6000B microscope and photos acquired on Image-pro 6.2 (Scientific Computing).

ELISA

A set of 172 overlapping “15-mer” HERV-K (HML-2) Env peptides (JPT Peptide Technologies, Berlin, Germany) were used to comprehensively map the antibody response. Positive signals were confirmed by peptides produced by two other companies (New England Peptide and Gene Script). Two peptides corresponding to the immunodominant epitopes defined on SU (RPKKGKPCPKKEIPKES) and TM (HRFQLQCDWNTSDFC) were used for the whole study (Gene Script). 96 microtiter wells plate (Nunc-Immuno Plate MaxiSorp Surface) were coated for 1 hour at 37°C with peptides at 10 µg/ml in PBS or over-night at 4°C with recombinant protein (GeneArt) at 5 µg/ml in PBS. Plates were then washed 3 times with 200 µL of PBS/0.05%-Tween 20 and blocked with 100 µL of blocking buffer (PBS/2.5%-BSA) at room temperature (RT). The samples were diluted in blocking buffer and incubated 2 h at RT in duplicates. Plates were then washed 3 times with 200 µL of PBS/0.05%-Tween 20. An anti-human IgG or anti-human IgM HRP-conjugated secondary antibody was diluted at 1:1000 in blocking buffer and incubated at RT for 1 hour. Plates were then washed 6 times with 200 µL of PBS/0.05%-Tween 20 and incubated for 10 minutes with 100 µL of TMB (Invitrogen). Addition of 50 µL H₂SO₄ 2 M stopped the reaction. The plates were read at 450 nm and 690 nm for the background on a plate reader. Background from 690 nm uncoated wells and PBS-BSA as negative controls was subtracted from the mean absorbance of the coated wells. Titers were determined as the reverse of the highest dilution of each serum sample that gives an optical density superior than its respective negative control average. Detection of anti-SU antibodies. ODs were normalized with serum from a high responder in a standard curve. The STDEV intra experiment was less than 7%. Detection of anti-TM antibodies. Sera were used at 1:400. OD were normalized with serum from a high responder in a standard curve. The STDEV intra experiment was less than 4%.

Tetramer preparation and B-cell staining

Adapted from Franz et al. [59].

All tetramers were prepared freshly for each experiment. Biotinylated-SU or -TM peptides (Gene Script) were incubated with premium-grade phycoerythrin-labeled streptavidin (Molecular Probes) for at least 20 minutes on ice at a molar ratio of 4:1. Before cell staining, tetramer preparations were centrifuged for 10 minutes at maximum speed to remove aggregates.

PBMCs were isolated by standard Ficoll-Hypaque density gradient centrifugation on fresh blood samples and immediately cryopreserved in fetal calf serum (HyClone, Logan, UT) containing 10% DMSO (Sigma Aldrich, St. Louis, MO) in liquid nitrogen. The cryopreserved cells were stored in liquid nitrogen until they were used.

Cells were thawed, washed, counted, and resuspended in PBS/5% FCS. For memory B-cell labeling, cells were enriched with the use of Human B Cell Enrichment Kit (Miltenyi). After enrichment, cells were adjusted to a density of 5×10^6 cells/mL and stained with SU-Tet or TM-Tet and incubated on ice for 30 minutes with intermittent gentle vortexing. Cells were co-stained with, IgD-APC (BD Biosciences), CD27-PacificBlue (BD Biosciences), and CD19-APC-H7 (BD Biosciences) for an additional 20 minutes on ice. LIVEDEAD[®] Fixable Dead Cell Stain Kit was used to discriminate live and dead cells. Cells were washed and stored in 2% paraformaldehyde at 4°C until acquisition on the LSR-II flow cytometer.

For all flow cytometry experiments, data were acquired with an LSR-II system (Becton Dickinson). At least 100,000 events were collected and analyzed with FlowJo software, version 9.0 (Tree Star, Ashland, OR).

Statistical analyses

Humoral responses assayed by ELISA were compared between groups using the Kruskal-Wallis, Dunn's multiple comparison, or two-tailed Mann-Whitney *t* tests. Linear regression and Spearman correlation analyses were used to measure associations between humoral response and HIV-1 viral load or CD4+ T cells count. All tests were conducted using GraphPad Prism, version 5.00 (GraphPad Software, San Diego, CA), with the statistical significance of the findings set at a *p* value of less than 0.05.

Additional files

Additional file 1: Figure S1. Epitope mapping and comparison homology studies. A, and B), Identification of the residues recognized by sera on SU and TM peptides. A set of peptides with the single mutation X to A were used in a peptide-based ELISA to determine which amino acids (aa) were recognized. The graphics represent the % of binding of the sera on the mutated peptide compared to the original peptide $[(\text{OD mutated peptide}/\text{OD original peptide}) \times 100]$. The original peptide corresponds to the peptide that gave the better signal for the mapping (see method section). The precise aa sequence was determined by reduction of binding. For both epitopes, sera from infected and healthy donor were used (6 for SU and 9 for TM). The graphs show the average of the 6 and 9 sera used respectively for SU and TM mapping. The results

show that the sequences recognized were PKEIPKE for SU epitope and DWNTS for the TM epitope. C) Comparison of the sequence of SU and TM peptides among the different main HERV-K (HML-2) families. Dots represent the same aa as the original sequence; (-) represents an aa deletion; a letter represents a mutation. On the left are the names of the HERV-K (HML-2) families. Corresponding PubMed access numbers: K101 (AF164609.1); K102 (AF164610.1); K103 (AF164611.1); K104 (AF164612.1); K107 (AF164613.1); K108 (AF164614.1); K109 (AF164615.1); K113 (AY037928.1); K115 (AY037929.1) D) Major homology of sequence between SU and TM sequence with other viruses. Underscored letters are the epitopes determined in Additional file 1: Figure S1A; homologies are represented in bold. The homologies were determined by using Blast tool from the PubMed website.

Additional file 2: Figure S2. Anti-TM response in HCV positive patients. Results from the cross sectional studies were analyzed with respect to the HCV serostatus of the 120 HIV-1 infected patients. 47 were identified as HCV positive and 69 as HCV negative. This clinical information was not available for 4 patients. The statistical significance of data between the two groups was established using the Mann & Whitney *T* test.

Additional file 3: Figure S3. Ratio anti-HERV-K (HML-2) Env antibodies/total IgG. The total IgG level was assayed using a human IgG ELISA kit (Mabtech) according to the provider's protocol. The sera were diluted at 1/4000000. A ratio was calculated as follows: anti-SU or anti-TM titer (OD)/total IgG (OD). *n* = 40 for each group. The ratio difference shows the humoral responses against SU and TM do not result from a nonspecific polyclonal expansion. The statistical significance of data between the different groups was established using ANOVA Kruskal-Wallis and Dunn's Multiple Comparison. The figure shows the representative results of three independent experiments. A *p* value <0.05 was considered as significant. **p* < 0.05, ***p* < 0.01, ****p* < 0.001.

Competing interests

H-AM, SGD, JBS and DFN are listed on a patent application filed by UCSF in respect of this work.

Authors' contributions

H-AM designed and performed the experiments and wrote the paper, MM helped design and perform the experiments, DS helped design and perform the experiments, SD helped design the experiments and provided clinical samples, JNM provided clinical samples, JJK helped design the experiments, JBS helped design the experiments and co-wrote the paper, DFN helped design the experiments, and co-wrote the paper. All authors helped edit the paper. All authors read and approved the final manuscript.

Authors' information

H-AM and MM are postdoctoral fellows in DFN's laboratory at UCSF. DS is an Adjunct Assistant professor in DFN's laboratory. SGD is Professor of Medicine at PHP/AIDS Division, UCSF. JNM is Professor of Epidemiology and Biostatistics, UCSF. CDP is Associate Professor of Medicine at PHP/AIDS Division. FMH is Professor of Medicine at PHP/AIDS Division, UCSF. JBS is Assistant Professor at OSHU. DFN is Adjunct Professor of Medicine, Division of Experimental Medicine, UCSF, and Chair, Department of Microbiology, Immunology & Tropical Medicine at GW University, Washington, DC.

Acknowledgements

We thank Dennis Burton, Nancy L Haigwood, Pheroze Joshi, Neil Sheppard, James Kobie, Jeffrey Milush, Andre Raposo, R. Brad Jones and Mario Ostrowski for helpful discussions, and Pandey Suchitra, Gerald Spotts and Lisa Loeb for help in providing clinical specimens and data. We gratefully acknowledge reagents supplied from the NIH AIDS Research and Reference Reagent Program. This work was supported in part by grant number 108248-51-RGRL from amfAR, The Foundation for AIDS Research, The Bill & Melinda Gates Foundation (BMGF:01434000363), the Peter and Shelagh Godsoe Family Foundation through the AIDS Research Institute at UCSF, Pfizer Inc. through a sponsored research agreement, and funds from the National Institutes of Health (AI076059 and AI084113). This research was supported in part by a grant from the National Institutes of Health, University of California, San Francisco-Gladstone Institute of Virology & Immunology Center for AIDS Research, P30-AI027763 and the District of Columbia Developmental Center for AIDS Research (P30AI087714). The Options study is funded in part by

NIAID (P01 AI071713). The SCOPE cohort is SCOPE supported in part by the NIAID (RO1 AI087145, K24AI069994, U19AI0961090), the UCSF/Gladstone Institute of Virology & Immunology CFAR (grant number P30 AI027763), the UCSF Clinical and Translational Research Institute Clinical Research Center (UL1 RR024131), the Center for AIDS Prevention Studies (P30 MH62246), and the CFAR Network of Integrated Systems (R24 AI067039). This work was supported in part by NIH OD 011092. The following reagents were obtained through the AIDS Research and Reference Reagent Program, Division of AIDS, NIAID, NIH: 1) HeLa T4⁺ from Dr. Richard Axel [54]. 2) T2M-bl from Dr. John C. Kappes, Dr. Xiaoyun Wu and Tranzyme Inc [51,55-57]. 3) HIV-1_{LAI} from Dr. Jean-Marie Bechet and Dr. Luc Montagnier [58,59].

Author details

¹Division of Experimental Medicine, Department of Medicine, University of California, San Francisco, USA. ²HIV/AIDS Program, Department of Medicine, University of California, San Francisco, USA. ³Department of Epidemiology and Biostatistics, University of California, San Francisco, USA. ⁴Vaccine & Gene Therapy Institute, Oregon Health & Science University, Beaverton, OR 97006, USA. ⁵Oregon National Primate Research Center, Oregon Health & Science University, Beaverton, OR 97006, USA. ⁶Microbiology, Immunology & Tropical Medicine, School of Medicine & Health Sciences, The George Washington University, Ross Hall 502, 2300 Eye Street, NW, Washington, DC 20037, USA.

Received: 8 August 2013 Accepted: 7 January 2014

Published: 28 January 2014

References

- Lander ES, Linton LM, Birren B, Nusbaum C, Zody MC, Baldwin J, Devon K, Dewar K, Doyle M, FitzHugh W, et al: **Initial sequencing and analysis of the human genome.** *Nature* 2001, **409**:860–921.
- Lee YN, Bieniasz PD: **Reconstitution of an infectious human endogenous retrovirus.** *PLoS Pathog* 2007, **3**:e10.
- Goff SP: **Retrovirus restriction factors.** *Mol Cell* 2004, **16**:849–859.
- Subramanian RP, Wildschutte JH, Russo C, Coffin JM: **Identification, characterization, and comparative genomic distribution of the HERV-K (HML-2) group of human endogenous retroviruses.** *Retrovirology* 2011, **8**:90.
- Kurth R, Bannert N: **Beneficial and detrimental effects of human endogenous retroviruses.** *Int J Cancer* 2010, **126**:306–314.
- Jha AR, Nixon DF, Rosenberg MG, Martin JN, Deeks SG, Hudson RR, Garrison KE, Pillai SK: **Human endogenous retrovirus K106 (HERV-K106) was infectious after the emergence of anatomically modern humans.** *PLoS One* 2011, **6**:e20234.
- Lower R, Boller K, Hasenmaier B, Korbmacher C, Muller-Lantzsch N, Lower J, Kurth R: **Identification of human endogenous retroviruses with complex mRNA expression and particle formation.** *Proc Natl Acad Sci USA* 1993, **90**:4480–4484.
- Ehlhardt S, Seifert M, Schneider J, Ojak A, Zang KD, Mehraein Y: **Human endogenous retrovirus HERV-K(HML-2) Rec expression and transcriptional activities in normal and rheumatoid arthritis synovia.** *J Rheumatol* 2006, **33**:16–23.
- Herve CA, Lugli EB, Brand A, Griffiths DJ, Venables PJ: **Autoantibodies to human endogenous retrovirus-K are frequently detected in health and disease and react with multiple epitopes.** *Clin Exp Immunol* 2002, **128**:75–82.
- Lai OY, Chen H, Michaud HA, Hayashi G, Kuebler PJ, Hultman GK, Ariza ME, Williams MV, Batista MD, Nixon DF, et al: **Protective effect of human endogenous retrovirus K dUTPase variants on psoriasis susceptibility.** *J Invest Dermatol* 2012, **132**:1833–1840.
- Pereyra F, Jia X, McLaren PJ, Telenti A, de Bakker PI, Walker BD, Ripke S, Brumme CJ, Pulit SL, Carrington M, et al: **The major genetic determinants of HIV-1 control affect HLA class I peptide presentation.** *Science* 2010, **330**:1551–1557.
- Sicat J, Sutkowski N, Huber BT: **Expression of human endogenous retrovirus HERV-K18 superantigen is elevated in juvenile rheumatoid arthritis.** *J Rheumatol* 2005, **32**:1821–1831.
- Armitage AE, Katzourakis A, de Oliveira T, Welch JJ, Belshaw R, Bishop KN, Kramer B, McMichael AJ, Rambaut A, Iversen AK: **Conserved footprints of APOBEC3G on Hypermutated human immunodeficiency virus type 1 and human endogenous retrovirus HERV-K(HML2) sequences.** *J Virol* 2008, **82**:8743–8761.
- Buscher K, Trefzer U, Hofmann M, Sterry W, Kurth R, Denner J: **Expression of human endogenous retrovirus K in melanomas and melanoma cell lines.** *Cancer Res* 2005, **65**:4172–4180.
- Goering W, Ribarska T, Schulz WA: **Selective changes of retroelement expression in human prostate cancer.** *Carcinogenesis* 2011, **32**:1484–1492.
- Ishida T, Obata Y, Ohara N, Matsushita H, Sato S, Uenaka A, Saika T, Miyamura T, Chayama K, Nakamura Y, et al: **Identification of the HERV-K gag antigen in prostate cancer by SEREX using autologous patient serum and its immunogenicity.** *Cancer Immunol* 2008, **8**:15.
- Katoh I, Mirova A, Kurata S, Murakami Y, Horikawa K, Nakakuki N, Sakai T, Hashimoto K, Maruyama A, Yonaga T, et al: **Activation of the long terminal repeat of human endogenous retrovirus K by melanoma-specific transcription factor MITF-M.** *Neoplasia* 2011, **13**:1081–1092.
- Schanab O, Humer J, Gleiss A, Mikula M, Sturlan S, Grunt S, Okamoto I, Muster T, Pehamberger H, Waltenerberger A: **Expression of human endogenous retrovirus K is stimulated by ultraviolet radiation in melanoma.** *Pigment Cell Melanoma Res* 2011, **24**:656–665.
- Wang-Johanning F, Frost AR, Jian B, Azerou R, Lu DW, Chen DT, Johanning GL: **Detecting the expression of human endogenous retrovirus E envelope transcripts in human prostate adenocarcinoma.** *Cancer* 2003, **98**:187–197.
- Boller K, Janssen O, Schuldes H, Tonjes RR, Kurth R: **Characterization of the antibody response specific for the human endogenous retrovirus HTDV/HERV-K.** *J Virol* 1997, **71**:4581–4588.
- Hahn S, Ugurel S, Hanschmann KM, Strobel H, Tondera C, Schadendorf D, Lower J, Lower R: **Serological response to human endogenous retrovirus K in melanoma patients correlates with survival probability.** *AIDS Res Hum Retroviruses* 2008, **24**:717–723.
- Rakoff-Nahoum S, Kuebler PJ, Heymann JJ, MES, Ortiz GM, GSO, Barbour JD, Lenz J, Steinfeld AD, Nixon DF: **Detection of T lymphocytes specific for human endogenous retrovirus K (HERV-K) in patients with seminoma.** *AIDS Res Hum Retroviruses* 2006, **22**:52–56.
- Schiavetti F, Thonnard J, Colau D, Boon T, Coulie PG: **A human endogenous retroviral sequence encoding an antigen recognized on melanoma by cytolytic T lymphocytes.** *Cancer Res* 2002, **62**:5510–5516.
- Wang-Johanning F, Radvanyi L, Rycaj K, Plummer JB, Yan P, Sastry KJ, Piyathilake CJ, Hunt KK, Johanning GL: **Human endogenous retrovirus K triggers an antigen-specific immune response in breast cancer patients.** *Cancer Res* 2008, **68**:5869–5877.
- Zhao J, Rycaj K, Geng S, Li M, Plummer JB, Yin B, Liu H, Xu X, Zhang Y, Yan Y, et al: **Expression of human endogenous retrovirus type K envelope protein Is a novel candidate prognostic marker for human breast cancer.** *Genes Cancer* 2011, **2**:914–922.
- Garrison KE, Jones RB, Meiklejohn DA, Anwar N, Ndhlovu LC, Chapman JM, Erickson AL, Agrawal A, Spotts G, Hecht FM, et al: **T cell responses to human endogenous retroviruses in HIV-1 infection.** *PLoS Pathog* 2007, **3**:e165.
- Jones RB, Garrison KE, Mujib S, Mihajlovic V, Aidarus N, Hunter DV, Martin E, John VM, Zhan W, Faruk NF, et al: **HERV-K-specific T cells eliminate diverse HIV-1/2 and SIV primary isolates.** *J Clin Invest* 2012, **122**(2):4473–49.
- Contreras-Galindo R, Kaplan MH, Contreras-Galindo AC, Gonzalez-Hernandez MJ, Ferlenghi I, Giusti F, Lorenzo E, Gitlin SD, Dosik MH, Yamamura Y, Markovitz DM: **Characterization of human endogenous retroviral elements in the blood of HIV-1-infected individuals.** *J Virol* 2012, **86**:262–276.
- SenGupta D, Tandon R, Vieira RG, Ndhlovu LC, Lown-Hecht R, Ormsby CE, Loh L, Jones RB, Garrison KE, Martin JN, et al: **Strong human endogenous retrovirus-specific T cell responses are associated with control of HIV-1 in chronic infection.** *J Virol* 2011, **85**:6977–6985.
- van der Kuyl AC: **HIV infection and HERV expression: a review.** *Retrovirology* 2012, **9**:6.
- Dewannieux M, Blaise S, Heidmann T: **Identification of a functional envelope protein from the HERV-K family of human endogenous retroviruses.** *J Virol* 2005, **79**:15573–15577.
- Hanke K, Kramer P, Seeher S, Beimforde N, Kurth R, Bannert N: **Reconstitution of the ancestral glycoprotein of human endogenous retrovirus k and modulation of its functional activity by truncation of the cytoplasmic domain.** *J Virol* 2009, **83**:12790–12800.
- Kammerer U, Germeyer A, Stengel S, Kapp M, Denner J: **Human endogenous retrovirus K (HERV-K) is expressed in villous and extravillous cytotrophoblast cells of the human placenta.** *J Reprod Immunol* 2011, **91**:1–8.
- Laderoute MP, Giulivi A, Larocque L, Bellfroy D, Hou Y, Wu HX, Fowke K, Wu J, Diaz-Mitoma F: **The replicative activity of human endogenous retrovirus K102 (HERV-K102) with HIV viremia.** *Aids* 2007, **21**:2417–2424.

35. Lower R, Lower J, Kurth R: **The viruses in all of us: characteristics and biological significance of human endogenous retrovirus sequences.** *Proc Natl Acad Sci USA* 1996, **93**:5177–5184.
36. Goedert JJ, Sauter ME, Jacobson LP, Vessella RL, Hilgartner MW, Leitman SF, Fraser MC, Mueller-Lantsch NG: **High prevalence of antibodies against HERV-K10 in patients with testicular cancer but not with AIDS.** *Cancer Epidemiol Biomarkers Prev* 1999, **8**:293–296.
37. Vogetseder W, Dumfahrt A, Mayersbach P, Schonitzer D, Dierich MP: **Antibodies in human sera recognizing a recombinant outer membrane protein encoded by the envelope gene of the human endogenous retrovirus K.** *AIDS Res Hum Retroviruses* 1993, **9**:687–694.
38. Imbeault M, Giguere K, Ouellet M, Tremblay MJ: **Exon level transcriptomic profiling of HIV-1-infected CD4(+) T cells reveals virus-induced genes and host environment favorable for viral replication.** *PLoS Pathog* 2012, **8**:e1002861.
39. Moiani A, Mavilio F: **Alternative splicing caused by lentiviral integration in the human genome.** *Methods in enzymology* 2012, **507**:155–169.
40. Moiani A, Paleari Y, Sartori D, Mezzadra R, Miccio A, Cattoglio C, Cocchiarella F, Lidonnici MR, Ferrari G, Mavilio F: **Lentiviral vector integration in the human genome induces alternative splicing and generates aberrant transcripts.** *J Clin Invest* 2012, **122**:1653–1666.
41. Chang ST, Sova P, Peng X, Weiss J, Law GL, Palermo RE, Katze MG: **Next-generation sequencing reveals HIV-1-mediated suppression of T cell activation and RNA processing and regulation of noncoding RNA expression in a CD4+ T cell line.** *mBio* 2011, **2**:e00134–11.
42. Hohn O, Hanke K, Bannert N: **HERV-K(HML-2), the best preserved family of HERVs: endogenization, expression, and implications in health and disease.** *Frontiers in oncology* 2013, **3**:246.
43. Boller K, Schonfeld K, Lischer S, Fischer N, Hoffmann A, Kurth R, Tonjes RR: **Human endogenous retrovirus HERV-K113 is capable of producing intact viral particles.** *J Gen Virol* 2008, **89**:567–572.
44. Haraguchi S, Good RA, Cianciolo GJ, Engelman RW, Day NK: **Immunosuppressive retroviral peptides: immunopathological implications for immunosuppressive influences of retroviral infections.** *J Leukoc Biol* 1997, **61**:654–666.
45. Kraus B, Fischer K, Buchner SM, Wels WS, Lower R, Sliva K, Schnierle BS: **Vaccination directed against the human endogenous retrovirus-K envelope protein inhibits tumor growth in a murine model system.** *PLoS One* 2013, **8**:e72756.
46. Mangeney M, Heidmann T: **Tumor cells expressing a retroviral envelope escape immune rejection in vivo.** *Proc Natl Acad Sci USA* 1998, **95**:14920–14925.
47. Oostendorp RA, Schaaper WM, Post J, Meloen RH, Scheper RJ: **Synthetic hexapeptides derived from the transmembrane envelope proteins of retroviruses suppress N-formylpeptide-induced monocyte polarization.** *J Leukoc Biol* 1992, **51**:282–288.
48. Wang-Johanning F, Rycaj K, Plummer JB, Li M, Yin B, Frerich K, Garza JG, Shen J, Lin K, Yan P, *et al*: **Immunotherapeutic potential of anti-human endogenous retrovirus-K envelope protein antibodies in targeting breast tumors.** *J Natl Cancer Inst* 2012, **104**:189–210.
49. Schmitz-Winnenthal FH, Galindo-Escobedo LV, Rimoldi D, Geng W, Romero P, Koch M, Weitz J, Krempien R, Niethammer AG, Beckhove P, *et al*: **Potential target antigens for immunotherapy in human pancreatic cancer.** *Cancer Lett* 2007, **252**:290–298.
50. Sacha JB, Kim IJ, Chen L, Ullah JH, Goodwin DA, Simmons HA, Schenkman DI, von Pelchrzim F, Gifford RJ, Nimityongskul FA, *et al*: **Vaccination with cancer- and HIV infection-associated endogenous retrotransposable elements is safe and immunogenic.** *J Immunol* 2012, **189**:1467–1479.
51. Barre-Sinoussi F, Chermann JC, Rey F, Nugeyre MT, Chamaret S, Gruest J, Dauguet C, Axler-Blin C, Vezinet-Brun F, Rouzioux C, *et al*: **Isolation of a T-lymphotropic retrovirus from a patient at risk for acquired immune deficiency syndrome (AIDS).** *Science* 1983, **220**:868–871.
52. Wain-Hobson S, Vartanian JP, Henry M, Chenciner N, Cheynier R, Delassus S, Martins LP, Sala M, Nugeyre MT, Guetard D, *et al*: **LAV revisited: origins of the early HIV-1 isolates from Institut Pasteur.** *Science* 1991, **252**:961–965.
53. Platt EJ, Bilkska M, Kozak SL, Kabat D, Montefiori DC: **Evidence that ecotropic murine leukemia virus contamination in TZM-bl cells does not affect the outcome of neutralizing antibody assays with human immunodeficiency virus type 1.** *J Virol* 2009, **83**:8289–8292.
54. Takeuchi Y, McClure MO, Pizzato M: **Identification of gammaretroviruses constitutively released from cell lines used for human immunodeficiency virus research.** *J Virol* 2008, **82**:12585–12588.
55. Wei X, Decker JM, Liu H, Zhang Z, Arani RB, Kilby JM, Saag MS, Wu X, Shaw GM, Kappes JC: **Emergence of resistant human immunodeficiency virus type 1 in patients receiving fusion inhibitor (T-20) monotherapy.** *Antimicrob Agents Chemother* 2002, **46**:1896–1905.
56. Platt EJ, Wehrly K, Kuhmann SE, Chesebro B, Kabat D: **Effects of CCR5 and CD4 cell surface concentrations on infections by macrophagotropic isolates of human immunodeficiency virus type 1.** *J Virol* 1998, **72**:2855–2864.
57. Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F: **Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes.** *Genome biology* 2002, **3**:RESEARCH0034.
58. Maddon PJ, Dalgleish AG, McDougal JS, Clapham PR, Weiss RA, Axel R: **The T4 gene encodes the AIDS virus receptor and is expressed in the immune system and the brain.** *Cell* 1986, **47**:333–348.
59. Franz B, May KF Jr, Dranoff G, Wucherpfennig K: **Ex vivo characterization and isolation of rare memory B cells with antigen tetramers.** *Blood* 2011, **118**:348–357.

doi:10.1186/1742-4690-11-10

Cite this article as: Michaud *et al*: Trans-activation, post-transcriptional maturation, and induction of antibodies to HERV-K (HML-2) envelope transmembrane protein in HIV-1 infection. *Retrovirology* 2014 **11**:10.

Submit your next manuscript to BioMed Central and take full advantage of:

- Convenient online submission
- Thorough peer review
- No space constraints or color figure charges
- Immediate publication on acceptance
- Inclusion in PubMed, CAS, Scopus and Google Scholar
- Research which is freely available for redistribution

Submit your manuscript at
www.biomedcentral.com/submit

