

**UCLA**

**UCLA Previously Published Works**

**Title**

Whole-mount skeletal staining

**Permalink**

<https://escholarship.org/uc/item/3007h7vh>

**Authors**

Rigueur, D

Lyons, KM

**Publication Date**

2014

**DOI**

10.1007/978-1-62703-989-5-9

Peer reviewed

- [Journal List](#)
- [HHS Author Manuscripts](#)
- PMC5384832



[Methods Mol Biol.](#) Author manuscript; available in PMC 2017 Apr 7.

Published in final edited form as:

[Methods Mol Biol.](#) 2014; 1130: 113–121.

doi: [10.1007/978-1-62703-989-5\\_9](https://doi.org/10.1007/978-1-62703-989-5_9)

PMCID: PMC5384832

NIHMSID: NIHMS853686

PMID: [24482169](#)

## Whole-Mount Skeletal Staining

[Diana Rigueur](#) and [Karen M. Lyons](#)

### Abstract

#### 1 Introduction

---

The formation of bone occurs through two processes: endochondral and intramembranous. Endochondral bone formation occurs after cells in mesenchymal condensations differentiate into chondrocytes. These cells secrete a cartilaginous extracellular matrix rich in proteoglycans, glycosaminoglycans (GAGs), and collagen (types II and X). Chondrocytes undergo a process of stratified differentiation and apoptosis, enabling the replacement of cartilage by bone. In contrast, during intramembranous ossification, mesenchymal cells differentiate directly into bone-forming osteoblasts. Alcian blue and Alizarin red stain cartilage and bone, respectively. As a cationic dye, Alcian blue binds strongly to sulfated GAGs and glycoproteins, while Alizarin red, an anionic dye, binds to cationic metals such as calcium [1]. Because cartilage contains higher concentrations of GAGs than any other tissue, it binds more Alcian blue. However, other tissues, such as the skin, contain GAGs and other glycoproteins that bind Alcian blue. In contrast, because 99 % of the calcium in the body is localized in bone, Alizarin red is highly specific for bone. The selective staining properties of Alcian blue and Alizarin red and their use in whole-mount skeletal preparation have been widespread for over a century [2]. The protocol has been refined over the years, and there are variations employing enzymatic steps and different fixatives [2–5]. The protocol described here is used in our laboratory. Most laboratories use very similar protocols.

As a brief overview of the process, specimens are prepared by removing skin, organs, and brown fat. They are then dehydrated and fixed in 95 % ethanol. For further removal of fatty tissue and tissue permeabilization, specimens are exposed to acetone and then consecutively transferred to Alcian blue and Alizarin red staining solutions. Concurrent with Alizarin red staining, exposure to potassium hydroxide (KOH) hydrolyzes soft tissue, leading to transparency and allowing visualization of stained skeletal elements. The procedure can be adjusted depending on the size/age of the specimens.

## 2 Materials

---

### 2.1 Solutions

Use analytical grade reagents.

1. Phosphate-buffered saline (PBS): 137 mM NaCl, 10 mM phosphate, 2.7 mM KCl, pH 7.4. To prepare 1 L of 1× PBS pour 800 mL of distilled water into a beaker. Add in order 8 g of NaCl, 0.2 g of KCl, 1.44 g of Na<sub>2</sub>HPO<sub>4</sub>, 0.24 g of KH<sub>2</sub>PO<sub>4</sub>. Adjust the pH to 7.4 with HCl. Add distilled water to a total volume of 1 L.
2. 100 % Ethanol (EtOH).
3. 100 % Acetone ((CH<sub>3</sub>)<sub>2</sub>CO).
4. Glycerol (C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).
5. 1 % Potassium hydroxide (KOH): 1 % (w/v) KOH, 99 % dH<sub>2</sub>O. To make a 200 mL solution, weigh 2 g of KOH pellets. Gradually add to a beaker containing 200 mL of deionized water (*see* [Note 1](#)).
6. Alcian blue stain: 0.03 % (w/v), 80 % EtOH, 20 % (glacial) acetic acid. To make a 200 mL solution, weigh 0.06 g of Alcian blue 8GX and place in a beaker, add 160 mL of 100 % EtOH and 40 mL of 100 % glacial acetic acid (*see* [Notes 2](#) and [3](#)).
7. Alizarin red stain: 0.005 % (w/v) in 1 % (w/v) KOH. To make a 200 mL solution, weigh 10 mg of Alizarin red and add to 200 mL of 1 % KOH (*see* [Notes 4](#) and [5](#)).

### 2.2 Tools/Supplies

1. Dissecting microscope and photographic equipment.
2. Diapers.
3. Hot water bath.
4. Forceps.
5. Scalpel.
6. Glass scintillation vials, 15 mL, or 50 mL conical tubes.


## 3 Methods

---

### 3.1 Staining of Skeletal Elements: Mid-Gestation Stages (E12.5–E16.5)

1. Collect embryos following euthanization of timed pregnant females and place in 1× PBS.
2. Remove the extraembryonic membranes encircling the embryo. Remove the eyes while in PBS [[3–7](#)]. For early-stage embryos (E12.5–E14.5), it is not necessary to remove the skin or eviscerate the embryo. For older embryos, eviscerate as follows: initiate skin removal by creating a horizontal slit in the skin at the abdomen using dissecting scissors. Then pull apart using your fingers or forceps. Continue to tug the skin gently apart. Repeat by making a vertical slit in the dorsal skin of the embryo and pulling the edges apart. To remove skin from the arms, make a vertical slit on the ventral and dorsal surfaces of the fore and hind limbs. Peel the skin using forceps.

3. Place the embryos in glass scintillation vials containing 70 % ethanol to fix overnight at 4 °C (see [Note 6](#)).
4. Remove the 70 % EtOH and replace with 95 % EtOH for 1 h.
5. Remove the 95 % EtOH and replace with acetone overnight at room temperature (see [Note 7](#)).
6. Remove the acetone and replace with Alcian blue stain for 1–4 h (see [Note 8](#)).
7. Remove the Alcian blue stain and replace with Alizarin red stain for 3–4 h (see [Note 9](#)).
8. Remove the Alizarin red stain and replace with 1 % KOH for 12 h to overnight for clearing of the embryo [[5](#), [7–9](#)] (see [Note 10](#)).
9. Transfer the embryos to a 50 % glycerol:50 % (1 %) KOH solution at room temperature until tissue appears transparent.
10. Once cleared, transfer the specimen to 100 % glycerol for long-term storage [[5](#), [7–9](#)]. See [Fig. 1](#).




[Fig. 1](#)

Alcian blue and Alizarin red staining of (a) E14.5 and (b) E16.5 embryos. Images were taken using bright field optics. Following adjustment of color levels using Photoshop, the embryos were pasted onto a *white* background

### 3.2 Late-Gestation and Early-Postnatal Stages: E16.5–P21

1. Wash and then scald the specimen in hot tap water for 20–30 s at 65 °C to facilitate maceration (permeabilization) of the tissue and removal of skin [9] (see [Note 11](#)).
2. Remove eyes, all skin, internal organs, adipose tissue, and bubbles from the body cavity [7, 10] (see [Notes 12](#) and [13](#)).
3. Fix the embryos in 95 % EtOH overnight at room temperature [5, 7–9].
4. Place the samples in acetone overnight at room temperature (see [Note 7](#)).
5. Stain for cartilage by submerging the embryo in a glass scintillation vial containing at least enough Alcian blue solution to cover the embryo. Incubate the sample overnight at room temperature [5, 7–9] (see [Note 14](#)).
6. Destain embryos by initially washing them in two changes of 70 % EtOH and then incubating them in 95 % EtOH overnight [5] (see [Note 15](#)).
7. To pre-clear the tissue, remove the 95 % EtOH and add 1 % KOH solution for 1 h at room temperature [7].
8. Then remove the KOH solution, and replace it with Alizarin red solution for 3–4 h at room temperature. To slow down the staining, place the samples at 4 °C overnight [5, 7, 9] (see [Note 16](#)).
9. Replace the Alizarin red solution with a 50 % glycerol:50 % (1 %) KOH solution. You can incubate the specimen in this solution at room temperature or 4 °C until the excess red color is removed and the specimen is cleared [7].
10. Transfer sample in 100 % glycerol for long-term storage [7–11]. See [Fig. 2](#).



[Fig. 2](#)

Alcian blue and Alizarin red staining of a P0 pup. Image was processed as in [Fig. 1](#)

### 3.3 Later Postnatal and Adult Stages (3 Weeks Old and Over)

1. Euthanize the mouse and spray with 70 % EtOH.
2. Remove the skin (*see* Subheading 3.1, **step 2**, and [Notes 12](#) and [13](#)).
3. Remove eyes and visceral organs.
4. Remove as much adipose and other excess tissue as possible (*see* [Note 17](#)).
5. Place in two changes of 95 % EtOH overnight at room temperature to dehydrate and fix the specimen.
6. Replace the solution with 100 % acetone for 2 days to further fix the specimen and to remove adipose tissue (*see* [Note 7](#)).
7. Incubate the specimen in Alcian blue solution for 1–3 days (*see* [Notes 8](#) and [14](#)).
8. Destain embryos by initially washing them in two changes of 70 % EtOH and then incubating them in 95 % EtOH overnight [[5](#), [7](#)] (*see* [Note 15](#)).
9. Replace with 1 % KOH for 4 h at room temperature to overnight at 4 °C to pre-clear the specimens.
10. Replace the 1 % KOH with Alizarin red solution, and incubate the specimens for 2–5 days (*see* [Notes 16](#) and [17](#)).
11. Transfer the specimen into 1 % KOH as a clearing solution (*see* [Note 18](#)).
12. For long-term storage, keep the specimen in 100 % glycerol. *See* [Fig. 3](#).







### [Fig. 3](#)

Alcian blue and Alizarin red staining of a 4-week-old postnatal mouse. The image was processed as in [Fig. 1](#)

## 3.4 Preparing for Imaging

1. Carefully place the stained skeleton in a clear 60 mm or 10 cm plastic or glass dish using a scooping utensil (*see* [Note 19](#)).
2. Fill the plate with 100 % glycerol, and allow Schlieren patterns to settle (*see* [Note 20](#)).
3. Place the specimen under a dissecting microscope utilizing bright field optics and a white background.
4. If necessary, trim away excess tissue (*see* [Note 21](#)).

## Footnotes

---

<sup>1</sup>KOH is highly caustic to skin. Lab coat, gloves, and goggles should be worn when working with this chemical. KOH dissolves glass, so this solution should be stored in a plastic container or made fresh.

<sup>2</sup>Mix the Alcian blue solution very well; this may take 30 min to an hour. Then filter the solution. Otherwise, unequal and patchy distribution of the dye will lead to uneven staining of skeletal preparations.

<sup>3</sup>This solution can be stored for several weeks but works best when freshly prepared. It should be filtered before each use.

<sup>4</sup>To stain bone a purple–red appearance, use the formula stated above. If a more orange–red appearance is preferred, you can use Alizarin red diluted in EtOH: Alizarin red: 0.05 % (w/v) in 95 % (w/v) EtOH. Then digest in KOH until specimen clears.

<sup>5</sup>Alizarin red solutions work best when prepared freshly.

<sup>6</sup>The embryos can be transferred to 15 mL conical tubes or glass scintillation vials. We prefer the latter because it is easier to visualize the extent of staining.

<sup>7</sup>Acetone acts as a fixative, permeabilizes cell membranes to permit penetration of the stain, and dissolves fat.

<sup>8</sup>The skin of young embryos is porous enough to permit penetration of the Alcian blue stain following treatment with acetone. However, long-term exposure to Alcian blue stain will result in a nearly irreversible over-staining of skin and other soft tissues, making it difficult to see the underlying skeletal elements. If over-staining occurs, placing the embryo in 20 % glacial acetic acid can allow some leaching of the stain from soft tissues.

<sup>9</sup>Incubate the specimens in Alizarin red until the bone turns red. Over-staining will lead to irreversible uptake of the stain by muscle.

<sup>10</sup>If the sample is kept in KOH too long, the embryo will disintegrate and/or skeletal elements will disarticulate. The length of time the specimen should be maintained in KOH depends on its size, but at the end of the incubation, the specimen should be nearly transparent.

<sup>11</sup>For older pups with hair, omit this step and just spray specimen generously with 70 % EtOH to dampen the hair to facilitate dissection.

<sup>12</sup>One of the most commonly encountered problems with cleared skeletal preparations is insufficient staining of the digits due to failure to remove the skin around them. One way to remove the skin from hands and feet is to make a cut with a scalpel along the dorsal surface of the hand/foot and then along each finger. Then use the forceps to pinch the skin between the fingers to pull it off.

<sup>13</sup>Ensure removal of fat, organs, and bubbles from the body cavity; they impair staining by Alcian blue. It is important to remove the brown fat behind the neck and between the scapulas because it is pigmented and does not clear well. Moreover, Alcian blue stains fat.

<sup>14</sup>The entire specimen should turn blue, along with the cartilage. Weak staining can be attributed to improper evisceration, old solution, or inadequate exposure to Alcian blue. If weak staining is observed, keep the specimen in solution for a longer period of time, but this introduces a risk of over-staining.

<sup>15</sup>Keep changing the 70 % EtOH until it shows no blue coloration. This dehydration will fix the Alcian blue solution in cartilage and help destain surrounding soft tissues.

<sup>16</sup>Samples can be incubated in Alizarin red solution overnight at room temperature; however, avoid longer incubations as over-staining can occur, and the KOH can lead to disarticulation.

<sup>17</sup>If the solution becomes murky, after its 2–3-day incubation, transfer the specimen to 1 % KOH solution. For an older mouse, 2 % KOH exposure is acceptable; however, beware that hydrolysis is faster. Change the solution daily until no longer murky and the tissue clears. Then place in a 50 % glycerol:50 % (1 %) KOH solution for complete clearance.

<sup>18</sup>Decreasing gradients of 1 % KOH to glycerol can be used to further clear the specimen; however, for long-term storage, ensure that the specimen is stored in glycerol; otherwise, the KOH will cause the skeleton to disarticulate.

<sup>19</sup>Take a plastic disposable 1 mL pipette and cut off the top half of the bulb lengthwise—this creates a scoop, with the dropper forming a handle.

<sup>20</sup>The viscosity of glycerol facilitates positioning of the skeleton for imaging; however, Schlieren patterns result from differences in solution concentrations of glycerol or salt content. It can take several hours to overnight for these to disappear. Storing the specimens in a large enough volume of 100 % glycerol or graded concentrations of glycerol to EtOH, to completely submerge the specimen when the contents of the vial are transferred to the dish will eliminate this problem.

<sup>21</sup>Removing excess tissue at this stage makes a big difference to the final quality of the image. For larger embryos and postnatal stages, forceps may be sufficient. For mid-gestation embryos, a set of 4-in. straight micro scissors is an excellent investment.